

FIRST AID FOR THE[®]

USMLE[®]
STEP 1

2020

SMTEBOOKS
30th ANNIVERSARY EDITION

Ketabton.com

Time-proven blueprint for Step 1 success

1,300+ must-know concepts with many new high-yield facts

1,200+ color photos and illustrations, expanded and revised

Student-proven exam strategies designed to boost your score

**Mc
Graw
Hill**

TAO LE ■ VIKAS BHUSHAN ■ MATTHEW SOCHAT ■ VAISHNAVI VAIDYANATHAN

FIRST AID FOR THE®

USMLE STEP 1 2020

TAO LE, MD, MHS

Founder, ScholarRx
Associate Clinical Professor, Department of Medicine
University of Louisville School of Medicine

VIKAS BHUSHAN, MD

Boracay

MATTHEW SOCHAT, MD
Fellow, Department of Hematology/Oncology
St. Louis University School of Medicine



VAISHNAVI VAIDYANATHAN, MD

Resident, Department of Pediatric Neurology
Barrow Neurological Institute at Phoenix Children's Hospital

SARAH SCHIMANSKY, MB BCH BAO

Resident, Department of Ophthalmology
Royal United Hospitals Bath

JORDAN ABRAMS

St. George's University School of Medicine
Class of 2020

KIMBERLY KALLIANOS, MD

Assistant Professor, Department of Radiology and Biomedical Imaging
University of California, San Francisco School of Medicine



New York / Chicago / San Francisco / Athens / London / Madrid / Mexico City
Milan / New Delhi / Singapore / Sydney / Toronto

Copyright © 2020 by Tao Le and Vikas Bhushan. All rights reserved. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written permission of the publisher.

ISBN: 978-1-26-046205-0

MHID: 1-26-046205-6

The material in this eBook also appears in the print version of this title: ISBN: 978-1-26-046204-3,
MHID: 1-26-046204-8.

eBook conversion by codeMantra

Version 1.0

All trademarks are trademarks of their respective owners. Rather than put a trademark symbol after every occurrence of a trademarked name, we use names in an editorial fashion only, and to the benefit of the trademark owner, with no intention of infringement of the trademark. Where such designations appear in this book, they have been printed with initial caps.

McGraw-Hill Education eBooks are available at special quantity discounts to use as premiums and sales promotions or for use in corporate training programs. To contact a representative, please visit the Contact Us page at www.mhprofessional.com.

Notice

Medicine is an ever-changing science. As new research and clinical experience broaden our knowledge, changes in treatment and drug therapy are required. The authors and the publisher of this work have checked with sources believed to be reliable in their efforts to provide information that is complete and generally in accord with the standards accepted at the time of publication. However, in view of the possibility of human error or changes in medical sciences, neither the authors nor the publisher nor any other party who has been involved in the preparation or publication of this work warrants that the information contained herein is in every respect accurate or complete, and they disclaim all responsibility for any errors or omissions or for the results obtained from use of the information contained in this work. Readers are encouraged to confirm the information contained herein with other sources. For example and in particular, readers are advised to check the product information sheet included in the package of each drug they plan to administer to be certain that the information contained in this work is accurate and that changes have not been made in the recommended dose or in the contraindications for administration. This recommendation is of particular importance in connection with new or infrequently used drugs.

TERMS OF USE

This is a copyrighted work and McGraw-Hill Education and its licensors reserve all rights in and to the work. Use of this work is subject to these terms. Except as permitted under the Copyright Act of 1976 and the right to store and retrieve one copy of the work, you may not decompile, disassemble, reverse engineer, reproduce, modify, create derivative works based upon, transmit, distribute, disseminate, sell, publish or sublicense the work or any part of it without McGraw-Hill Education's prior consent. You may use the work for your own noncommercial and personal use; any other use of the work is strictly prohibited. Your right to use the work may be terminated if you fail to comply with these terms.

THE WORK IS PROVIDED "AS IS." MCGRAW-HILL EDUCATION AND ITS LICENSORS MAKE NO GUARANTEES OR WARRANTIES AS TO THE ACCURACY, ADEQUACY OR COMPLETENESS OF OR RESULTS TO BE OBTAINED FROM USING THE WORK, INCLUDING ANY INFORMATION THAT CAN BE ACCESSED THROUGH THE WORK VIA HYPERLINK OR OTHERWISE, AND EXPRESSLY DISCLAIM ANY WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. McGraw-Hill Education and its licensors do not warrant or guarantee that the functions contained in the work will meet your requirements or that its operation will be uninterrupted or error free. Neither McGraw-Hill Education nor its licensors shall be liable to you or anyone else for any inaccuracy, error or omission, regardless of cause, in the work or for any damages resulting therefrom. McGraw-Hill Education has no responsibility for the content of any information accessed through the work. Under no circumstances shall McGraw-Hill Education and/or its licensors be liable for any indirect, incidental, special, punitive, consequential or similar damages that result from the use of or inability to use the work, even if any of them has been advised of the possibility of such damages. This limitation of liability shall apply to any claim or cause whatsoever whether such claim or cause arises in contract, tort or otherwise.

Dedication

To the contributors to this and past editions, who took time to share their knowledge, insight, and humor for the benefit of students and physicians everywhere.



This page intentionally left blank

Contents

Contributing Authors	vii	General Acknowledgments	xv
Associate Authors	viii	How to Contribute	xvii
Faculty Advisors	ix	How to Use This Book	xix
Thirtieth Anniversary Foreword	xi	Selected USMLE Laboratory Values	xx
Preface	xiii	First Aid Checklist for the USMLE Step 1	xxii
Special Acknowledgments	xiv		

▶ SECTION I GUIDE TO EFFICIENT EXAM PREPARATION 1

Introduction	2	Test-Taking Strategies	22
USMLE Step 1—The Basics	2	Clinical Vignette Strategies	23
Defining Your Goal	12	If You Think You Failed	24
Learning Strategies	13	Testing Agencies	24
Timeline for Study	16	References	25
Study Materials	20		

▶ SECTION I SUPPLEMENT SPECIAL SITUATIONS 27

▶ SECTION II HIGH-YIELD GENERAL PRINCIPLES 29

How to Use the Database	30	Pathology	205
Biochemistry	33	Pharmacology	229
Immunology	95	Public Health Sciences	255
Microbiology	123		

▶ SECTION III HIGH-YIELD ORGAN SYSTEMS 275

Approaching the Organ Systems	276	Neurology and Special Senses	489
Cardiovascular	279	Psychiatry	553
Endocrine	325	Renal	577
Gastrointestinal	357	Reproductive	611
Hematology and Oncology	403	Respiratory	659
Musculoskeletal, Skin, and Connective Tissue	445	Rapid Review	689

▶ SECTION IV TOP-RATED REVIEW RESOURCES 711

How to Use the Database	712	Biochemistry	716
Question Banks and Books	714	Cell Biology and Histology	716
Web and Mobile Apps	714	Microbiology and Immunology	717
Comprehensive	715	Pathology	717
Anatomy, Embryology, and Neuroscience	715	Pharmacology	718
Behavioral Science	716	Physiology	718
Abbreviations and Symbols	719	Index	749
Image Acknowledgments	727	About the Editors	808

Contributing Authors

MAJED H. ALGHAMDI, MBBS

Resident, Joint Program of Preventive Medicine
Jeddah, Saudi Arabia

LILIT ASLANYAN

New York Institute of Technology College of Osteopathic Medicine
Class of 2020

HUMOOD BOQAMBAR, MB BCh BAO

Assistant Registrar, Department of Orthopaedic Surgery
Farwaniya Hospital

WEELIC CHONG

Sidney Kimmel Medical College at Thomas Jefferson University
MD/PhD Candidate

KRISTINA DAMISCH

University of Iowa Carver College of Medicine
Class of 2020

YUMI KOVIC, MD

Resident, Department of Psychiatry
University of Massachusetts Medical School

KAITLYN MELNICK, MD

Resident, Department of Neurological Surgery
University of Florida College of Medicine, Gainesville

MARY KATHERINE MONTES de OCA, MD

Resident, Department of Obstetrics and Gynecology
Duke University Hospital

SCOTT MOORE, DO

Assistant Professor of Medical Laboratory Sciences
Weber State University

VASILY OVECHKO, MD

Resident, Department of Surgery
Russian Medical Academy of Continuous Professional Education

VIVEK PODDER

MBBS Student
Tairunnessa Memorial Medical College and Hospital, Bangladesh

CONNIE QIU

Lewis Katz School of Medicine at Temple University
MD/PhD Candidate

IMAGE AND ILLUSTRATION TEAM

CAROLINE COLEMAN

Emory University School of Medicine
Class of 2020

MATTHEW HO ZHI GUANG

University College Dublin (MD), DFCI (PhD)
MD/PhD Candidate

VICTOR JOSE MARTINEZ LEON, MD

Central University of Venezuela

ALIREZA ZANDIFAR, MD

Research Fellow
Isfahan University of Medical Sciences, Iran

Associate Authors

HUZAIFA AHMAD, MD

Resident, Department of Medicine
Georgetown University Hospital/MedStar Washington Hospital Center

ALEXANDER R. ASLESEN

Kirkville College of Osteopathic Medicine
Class of 2020

ANUP K. BHATTACHARYA, MD

Resident, Mallinckrodt Institute of Radiology
Washington University School of Medicine

ANUP CHALISE, MBBS

Resident, Department of General Surgery
Nepal Medical College and Teaching Hospital

ASHTEN R. DUNCAN, MPH

University of Oklahoma-Tulsa School of Community Medicine
Class of 2021

SARINA KOILPILLAI

St. George's University School of Medicine
Class of 2020

LAUREN N. LESSOR, MPH, MD

Resident, Department of Pediatrics
Mercy Health – St. Vincent Medical Center

ROHAN BIR SINGH, MD

Fellow, Department of Ophthalmology
Massachusetts Eye and Ear
Harvard Medical School

IMAGE AND ILLUSTRATION TEAM

YAMNA JADOON, MD

Research Associate
Aga Khan University

DANA M. JORGENSON

Chicago College of Osteopathic Medicine
Class of 2020

MITCHELL A. KATONA

University of Texas Health Science Center, Long School of Medicine
Class of 2020

TAYLOR MANEY, MD

Resident, Department of Anesthesiology
Brigham and Women's Hospital

Faculty Advisors

DIANA ALBA, MD

Clinical Instructor
University of California, San Francisco School of Medicine

MARK A.W. ANDREWS, PhD

Professor of Physiology
Lake Erie College of Osteopathic Medicine at Seton Hill

MARIA ANTONELLI, MD

Assistant Professor, Division of Rheumatology
MetroHealth Medical Center, Case Western Reserve University

HERMAN SINGH BAGGA, MD

Urologist, Allegheny Health Network
University of Pittsburgh Medical Center Passavant

SHIN C. BEH, MD

Assistant Professor, Department of Neurology & Neurotherapeutics
UT Southwestern Medical Center at Dallas

JOHN R. BUTTERLY, MD

Professor of Medicine
Dartmouth Geisel School of Medicine

SHELDON CAMPBELL, MD, PhD

Professor of Laboratory Medicine
Yale School of Medicine

BROOKS D. CASH, MD

Professor of Medicine, Division of Gastroenterology
University of South Alabama School of Medicine

SHIVANI VERMA CHMURA, MD

Adjunct Clinical Faculty, Department of Psychiatry
Stanford University School of Medicine

BRADLEY COLE, MD

Assistant Professor of Basic Sciences
Loma Linda University School of Medicine

LINDA S. COSTANZO, PhD

Professor, Physiology & Biophysics
Virginia Commonwealth University School of Medicine

MANAS DAS, MD, MS

Director, Clinical Anatomy, Embryology, and Histology
University of Massachusetts Medical School

ANTHONY L. DeFRANCO, PhD

Professor, Department of Microbiology and Immunology
University of California, San Francisco School of Medicine

CHARLES S. DELA CRUZ, MD, PhD

Associate Professor, Department of Pulmonary and Critical Care Medicine
Yale School of Medicine

SAKINA FARHAT, MD

Consulting Gastroenterologist
State University of New York Downstate Medical Center

CONRAD FISCHER, MD

Associate Professor, Medicine, Physiology, and Pharmacology
Touro College of Medicine

RAYUDU GOPALAKRISHNA, PhD

Associate Professor, Department of Integrative Anatomical Sciences
Keck School of Medicine of University of Southern California

RYAN C.W. HALL, MD

Assistant Professor, Department of Psychiatry
University of South Florida School of Medicine

LOUISE HAWLEY, PhD

Immediate Past Professor and Chair, Department of Microbiology
Ross University School of Medicine

JEFFREY W. HOFMANN, MD, PhD

Resident, Department of Pathology
University of California, San Francisco School of Medicine

CLARK KEBODEAUX, PharmD

Clinical Assistant Professor, Pharmacy Practice and Science
University of Kentucky College of Pharmacy

KRISTINE KRAFTS, MD

Assistant Professor, Department of Basic Sciences
University of Minnesota School of Medicine

MATTHEW KRAYBILL, PhD

Clinical Neuropsychologist
Cottage Health, Santa Barbara, California

GERALD LEE, MD

Assistant Professor, Departments of Pediatrics and Medicine
Emory University School of Medicine

KACHIU C. LEE, MD, MPH

Assistant Clinical Professor, Department of Dermatology
The Warren Alpert Medical School of Brown University

WARREN LEVINSON, MD, PhD

Professor, Department of Microbiology and Immunology
University of California, San Francisco School of Medicine

JAMES LYONS, MD

Professor of Pathology and Family Medicine
Alabama College of Osteopathic Medicine

PETER MARKS, MD, PhD

Center for Biologics Evaluation and Research
US Food and Drug Administration

DOUGLAS A. MATA, MD, MPH

Brigham Education Institute and Brigham and Women's Hospital
Harvard Medical School

VICKI M. PARK, PhD, MS

Assistant Dean
University of Tennessee College of Medicine

SOROUGH RAIS-BAHRAMI, MD

Assistant Professor, Departments of Urology and Radiology
University of Alabama at Birmingham School of Medicine

SASAN SAKIANI, MD

Fellow, Transplant Hepatology
Cleveland Clinic

MELANIE SCHORR, MD

Assistant in Medicine
Massachusetts General Hospital

SHIREEN MADANI SIMS, MD

Chief, Division of Gynecology, Gynecologic Surgery, and Obstetrics
University of Florida School of Medicine

NATHAN W. SKELLEY, MD

Assistant Professor, Department of Orthopaedic Surgery
University of Missouri, The Missouri Orthopaedic Institute

HOWARD M. STEINMAN, PhD

Assistant Dean, Biomedical Science Education
Albert Einstein College of Medicine

SUPORN SUKPRAPRUT-BRAATEN, PhD

Director of Research, Graduate Medical Education
Unity Health, Searcy, Arkansas

RICHARD P. USATINE, MD

Professor, Dermatology and Cutaneous Surgery
University of Texas Health Science Center San Antonio

J. MATTHEW VELKEY, PhD

Assistant Dean, Basic Science Education
Duke University School of Medicine

TISHA WANG, MD

Associate Clinical Professor, Department of Medicine
David Geffen School of Medicine at UCLA

SYLVIA WASSERTHEIL-SMOLLER, PhD

Professor Emerita, Department of Epidemiology and Population Health
Albert Einstein College of Medicine

ADAM WEINSTEIN, MD

Assistant Professor, Pediatric Nephrology and Medical Education
Geisel School of Medicine at Dartmouth

ABHISHEK YADAV, MBBS, MSc

Associate Professor of Anatomy
Geisinger Commonwealth School of Medicine

KRISTAL YOUNG, MD

Clinical Instructor, Department of Cardiology
Huntington Hospital, Pasadena, California

Thirtieth Anniversary Foreword

Our exam experiences remain vivid in our minds to this day as we reflect on 30 years of *First Aid*. In 1989, our big idea was to cobble together a “quick and dirty” study guide so that we would never again have to deal with the USMLE Step 1. We passed, but in a Faustian twist, we now relive the exam yearly while preparing each new edition.

Like all students before us, we noticed that certain topics tended to appear frequently on examinations. So we compulsively bought and rated review books and pored through a mind-numbing number of “recall” questions, distilling each into short facts. We had a love-hate relationship with mnemonics. They went against our purist desires for conceptual knowledge, but remained the best way to absorb the vocabulary and near-random associations that unlocked questions and eponyms.

To pull it all together, we used a then “state-of-the-art” computer database (Paradox/MS DOS 4) that fortuitously limited our entries to 256 characters. That length constraint (which predated Twitter by nearly two decades) imposed extreme brevity. The three-column layout created structure—and this was the blueprint upon which *First Aid* was founded.

The printed, three-column database was first distributed in 1989 at the University of California, San Francisco. The next year, the official first edition was self-published under the title *High-Yield Basic Science Boards Review: A Student-to-Student Guide*. The following year, our new publisher dismissed the *High-Yield* title as too confusing and came up with *First Aid for the Boards*. We thought the name was a bit cheesy, but it proved memorable. Interestingly, our “High-Yield” name resurfaced years later as the title of a competing board review series.

We lived in San Francisco and Los Angeles during medical school and residency. It was before the Web, and before med students could afford cell phones and laptops, so we relied on AOL e-mail and bulky desktops. One of us would drive down to the other person’s place for multiple weekends of frenetic revisions fueled by triple-Swiss white chocolate lattes from the Coffee Bean & Tea Leaf, with R.E.M. and the Nusrat Fateh Ali Khan playing in the background. Everything was marked up on 11- by 17-inch “tearsheets,” and at the end of the marathon weekend we would converge at the local 24-hour Kinko’s followed by the FedEx box near LAX (10 years before these two great institutions merged). These days we work with our online collaborative platform A.nnotate, GoToMeeting, and ubiquitous broadband Internet, and sadly, we rarely get to see each other.









What hasn’t changed, however, is the collaborative nature of the book. Thousands of authors, editors, and contributors have enriched our lives and made this book possible. Most helped for a year or two and moved on, but a few, like Ted Hon, Chirag Amin, and Andi Fellows, made lasting contributions. Like the very first edition, the team is always led by student authors who live and breathe (and fear) the exam, not professors years away from that reality.

We’re proud of the precedent that *First Aid* set for the many excellent student-to-student publications that followed. More importantly, *First Aid* itself owes its success to the global community of medical students and international medical graduates (IMGs) who each year contribute ideas, suggestions, and new content. In the early days, we used book coupons and tear-out business reply mail forms. These days, we get many thousands of comments and suggestions each year via our blog FirstAidTeam.com and A.nnotate.

At the end of the day, we don't take any of this for granted. Students are expected to synthesize an ever increasing amount of information, and we have a bigger challenge ahead of us to try to keep *First Aid* indispensable to students and IMGs. We want and need your participation in the *First Aid* community. (See How to Contribute, p. xvii.) With your help, we hope editing *First Aid* will continue to be just as fun and rewarding as the past 30 years have been.

Louisville Tao Le
Boracay Vikas Bhushan

First Aid for the USMLE Step 1 Through the Years



Preface

With the 30th edition of *First Aid for the USMLE Step 1*, we continue our commitment to providing students with the most useful and up-to-date preparation guide for the USMLE Step 1. This edition represents an outstanding revision in many ways, including:

- 50 entirely new or heavily revised high-yield topics reflecting evolving trends in the USMLE Step 1.
- Reorganization of high-yield topics in Pharmacology, Endocrine, and Reproductive chapters for improved study.
- Extensive text revisions, new mnemonics, clarifications, and corrections curated by a team of more than 30 medical student and resident physician authors who excelled on their Step 1 examinations and verified by a team of expert faculty advisors and nationally recognized USMLE instructors.
- Updated with 178 new and revised diagrams and illustrations as part of our ongoing collaboration with USMLE-Rx and ScholarRx (MedIQ Learning, LLC).
- Updated with 75 new and revised photos to help visualize various disorders, descriptive findings, and basic science concepts. Additionally, revised imaging photos have been labeled and optimized to show both normal anatomy and pathologic findings.
- Updated study tips on the opening page of each chapter.
- Improved integration of clinical images and illustrations to better reinforce and learn key anatomic concepts.
- Improved organization and integration of text, illustrations, clinical images, and tables throughout for focused review of high-yield topics.
- Revised and expanded ratings of current, high-yield review resources, with clear explanations of their relevance to USMLE review.
- Real-time Step 1 updates and corrections can be found exclusively on our blog, www.firstaidteam.com.

We invite students and faculty to share their thoughts and ideas to help us continually improve *First Aid for the USMLE Step 1* through our blog and collaborative editorial platform. (See *How to Contribute*, p. xvii.)

<i>Louisville</i>	Tao Le
<i>Boracay</i>	Vikas Bhushan
<i>St. Louis</i>	Matthew Sochat
<i>Phoenix</i>	Vaishnavi Vaidyanathan
<i>Bristol</i>	Sarah Schimansky
<i>New York City</i>	Jordan Abrams
<i>San Francisco</i>	Kimberly Kallianos

Special Acknowledgments

This has been a collaborative project from the start. We gratefully acknowledge the thousands of thoughtful comments, corrections, and advice of the many medical students, international medical graduates, and faculty who have supported the authors in our continuing development of *First Aid for the USMLE Step 1*.

For support and encouragement throughout the process, we are grateful to Thao Pham, Jinky Flang, and Jonathan Kirsch, Esq. Thanks to Louise Petersen for organizing and supporting the project. Thanks to our publisher, McGraw-Hill, for the valuable assistance of its staff, including Bob Boehringer, Jeffrey Herzich, and Christina Thomas.

We are also very grateful to Dr. Fred Howell and Dr. Robert Cannon of Textensor Ltd for providing us extensive customization and support for their powerful Annotate.co collaborative editing platform (www.annotate.co), which allows us to efficiently manage thousands of contributions. Thanks to Dr. Richard Usatine and Dr. Kristine Krafts for their outstanding image contributions. Thanks also to Jean-Christophe Fournet (www.humpath.com), Dr. Ed Uthman, and Dr. Frank Gaillard (www.radiopaedia.org) for generously allowing us to access some of their striking photographs.

For exceptional editorial leadership, enormous thanks to Kathleen Naylor, Christine Diedrich and Emma Underdown. Thank you to our USMLE-Rx/ScholarRx team of editors, Jessie Schanzle, Ruth Kaufman, Janene Matragrano, Susan Mazik, Isabel Nogueira, Sharon Prevost, Jen Shimony, and Hannah Warnshuis. Special thanks to our indexer Dr. Anne Fifer. We are also grateful to our medical illustrator, Hans Neuhart, for his creative work on the new and updated illustrations. Lastly, tremendous thanks to Graphic World, especially Anne Banning, Sandy Brown, Gary Clark, and Cindy Geiss.

<i>Louisville</i>	Tao Le
<i>Boracay</i>	Vikas Bhushan
<i>St. Louis</i>	Matthew Sochat
<i>Phoenix</i>	Vaishnavi Vaidyanathan
<i>Bristol</i>	Sarah Schimansky
<i>New York City</i>	Jordan Abrams
<i>San Francisco</i>	Kimberly Kallianos

General Acknowledgments

Each year we are fortunate to receive the input of thousands of medical students and graduates who provide new material, clarifications, and potential corrections through our website and our collaborative editing platform. This has been a tremendous help in clarifying difficult concepts, correcting errata from the previous edition, and minimizing new errata during the revision of the current edition. This reflects our long-standing vision of a true, student-to-student publication. We have done our best to thank each person individually below, but we recognize that errors and omissions are likely. Therefore, we will post an updated list of acknowledgments at our website, www.firstaidteam.com/bonus/. We will gladly make corrections if they are brought to our attention.

For submitting contributions and corrections, many thanks to Raed Ababneh, Antara Afrin, Rasim Agaev, Vanya Aggarwal, Ataa Ahmed, Hasan Alarouri, Basim Ali, Muhammad Faizan Ali, Moatasem Al-Janabi, Mohamed Almahmodi, Chima Amadi, Arman Amin, Jacqueline Aredo, Ranya Baddourah, Daniel Badin, Nida Bajwa, Dileni Bandarage, Jerrin Bawa, Esra Bayram, Craig Beavers, Jacqueline Bekhit, Matthias Bergmann, Stephanie Biecker, Aaron Birnbaum, Prateek Bommu, Nathaniel Borochoy, Susan Brands, Olivia W. Brooks, Meghan Brown, Stanley Budzinski, Kevin Budziszewski, Pavel Burski, Elisa M. Cairns, Sergio Camba, Katie Carsky, Esteban Casasola, Marielys Castro, Jesse Chait, Bliss Chang, Santosh Cherian, Heewon Choi, Charilaos Chourpiliadis, Maruf Chowdhury, Matthew J. Christensen, Matthew Yat Hon Chung, Alexander Ciaramella, Dillon Clancy, Sofija Conic, M. Marwan Dabbagh, Parag Das, Ketan Dayma, Elmer De Camps, Charles de Leeuw, Xavier De Pena, Christopher DeAngelo, Elliott Delgado, Anthony DeMarinis, Stacy Diaz, Evan Dishion, Nicola Helen Duzak, Emily Edwards, Alec Egan, Mohamed Elashwal, Osama El-Gabalawy, Matthew Eli, Awab Elnaeem, Sally El Sammak, Dylan Erwin, Stephanie Estevez-Marin, Gray Evans, Najat Fadlallah, Aria Fariborzi, Richard Ferro, Adam Fletcher, Kimberly A. Foley, Kyle Fratta, Samantha Friday, Nikhila Gandrakota, Siva Garapati, Nicolas Curi Gawlinski, Joanna Georgakas, Beth Anne George, Ashley Ghaemi, E. Sophia Gonzalez, Justin Graff, Gabriel Graham, Donovan Griggs, David Gruen, Gursewak Hadday, Jacqueline Hairston, Hunter Harrison, Gull Shahmir Hasnat, Maximillan Hawkins, Grecia Haymee, Briana Hernandez, Robin Hilder, Tammy Hua, Derrek Humphries, Audrey Hunt, Nanki Hura, Danny Ibrahim, Jyothik Varun Inampudi, Hnin Ingyin, Maham Irfan, Mina Iskandar, Kritika Iyer, Christina Jacobs, Arpit Jain, Neil K. Jain, Ala Jamal, Natalie Jansen, Jordan Jay, Mohammad Jmasi, Colton Junod, Talia Kamdjou, Filip Kaniski, Lydia Kaoutzani, Panagiotis Kaparalios, Srikrishna Karnatapu, Patrick Keller, Olivia Keller-Baruch, Cameron Kerl, Ahmed Ali Khan, Sara Khan, Shaima Khandaker, Samir Khouzam, Sonya Klein, Elana Kleinman, Andrew Ko, Soheil Kooraki, Anna Kukharchuk, Dennis Vu Kulp, Anil A. Kumar, Julie Kurek, Chloe Lahoud, Mike Lawandy, Ramy Lawandy, Jessica Lazar, Andrea Leal-Lopez, Lynda Lee, Chime Lhatso, Christine Lin, Benjamin Lodge, Soon Khai Low, Estefanía Henríquez Luthje, Lisa-Qiao MacDonald, Divya Madhavarapu, Mahir Mameledzija, Keerer Mann, Rajver Mann, Nadeen Mansour, Yusra Mansour, Bridget Martinez, Ahmad Mashlah, Rick Mathews, Amy McGregor, Alexandra & Joshua Medeiros & Fowler, Viviana Medina, Areeka Memon, Pedro G. R. Menicucci, Ben Meyers, Stephan A. Miller, Fatima Mirza, Murli Mishra, Elana Molcho, Guarina Molina, John Moon, Nayla Mroueh, Neha Mylarapu, Behnam Nabavizadeh, Moeko Nagatsuka, Ghazal Naghibzadeh, Alice Nassar, Nadya Nee, Lucas Nelson, Zach Nelson, Monica Nemat, Kenneth Nguyen, Michael Nguyen, Christian

Nieves, Nyia Njamfa, Ahmed Noor, Kyle Nyugen, Ahamd Obeidat, Gerald Olayan, Anndres Olson, Hasaan Omar, Daniel Ortiz, Michael O'Shea, Zonghao Pan, Vasilis Sebastian Paraschos, Christopher Parrino, Janak Patel, Vanisha Patel, Cyril Patra, Rita Paulis, Dmytro Pavlenko, Nancy A. Pina, Alexander Polyak, Jackeline Porto, Shannon D. Powell, Jacob Pruett, Laith Rahabneh, Kamleshun Ramphul, Janhvi Rana, Nidaa Rasheed, Abdul Sattar Raslan, Tomas Ream, Rashelle Ripa, Amanda Michelle Ritchie, Helio Manuel Grullón Rodríguez, Sarah Rohrig, Gessel Romero, Alexander Rose, Rachel Rose, Erica Rubin, Areesha Saati, Jeffrey Sackey, Raza H. Sagarwala, Chhavi Saini, Sergii Sakhno, Allie Sakowicz, Shadia Saleh, Roshun Sangani, Dhruv Sarwal, Abeer Sarwar, M. Sathyanarayanan, Neetu Scariya, Tonio Felix Schaffert, Melissa Schechter, Kathryn Scheinberg, Emma Schnuckle, Emma Schulte, Taylor Schweigert, Lee Seifert, Sheila Serin, Deeksha Seth, Omid Shafaat, Nirav Shah, Samir K. Shah, Wasif Nauman Shah, Muhanad Shaib, Ahmed Shakir, Purnima Sharma, Tina Sharma, Kayla Sheehan, Dr. Priya Shenwai, Sami Shoura, Kris Sifeldeen, Akhand Singh, Manik Inder Singh, Ramzi Y. Skaik, Samantha A. Smith, Timothy Smith, Emilie Song, Hang Song, Shichen Song, Luke Sorensen, Charles Starling, Jonathan Andrew Stone, Nathan Stumpf, Johnny Su, Bahaa Eddine Succar, Saranya Sundaram, Steven Svoboda, Clara Sze, Olive Tang, Brian Tanksley, Omar Tayh, Joshua Taylor, Valerie Teano, Warren Teltser, Steffanie Camilo Tertulien, Roger Torres, Michael Trainer, Andrew Trinh, Aalap K. Trivedi, Georgeanna Tsoumas, Elizabeth Tsui, Cem Turam, Methma Udawatta, Daramfon Udofia, Adaku Ume, Rio Varghese, Judith Vásquez, Earl Vialpando, Sagar Vinayak, Phuong Vo, Habiba Wada, Jason Wang, Tiffany Wang, Zoe Warczak, Mitchell Waters, Rachel Watson, Elizabeth Douglas Weigel, Rabbi Michael Weingarten, Kaystin Weisenberger, Aidan Woodthorpe, Mattia Wruble, Angela Wu, Catherine Xie, Rebecca Xu, Nicholas Yeisley, Sammy Yeroushalmi, Melissas Yuan, Sahil Zaveri, and Yolanda Zhang.

How to Contribute

This version of *First Aid for the USMLE Step 1* incorporates thousands of contributions and improvements suggested by student and faculty advisors. We invite you to participate in this process. Please send us your suggestions for:

- Study and test-taking strategies for the USMLE Step 1
- New facts, mnemonics, diagrams, and clinical images
- High-yield topics that may appear on future Step 1 exams
- Personal ratings and comments on review books, question banks, apps, videos, and courses

For each new entry incorporated into the next edition, you will receive **up to a \$20 Amazon.com gift card** as well as personal acknowledgment in the next edition. Significant contributions will be compensated at the discretion of the authors. Also, let us know about material in this edition that you feel is low yield and should be deleted.

All submissions including potential errata should ideally be supported with hyperlinks to a dynamically updated Web resource such as UpToDate, AccessMedicine, and ClinicalKey.

We welcome potential errata on grammar and style if the change improves readability. Please note that *First Aid* style is somewhat unique; for example, we have fully adopted the *AMA Manual of Style* recommendations on eponyms (“We recommend that the possessive form be omitted in eponymous terms”) and on abbreviations (no periods with eg, ie, etc). We also avoid periods in tables unless required for full sentences. Kindly refrain from submitting “style errata” unless you find specific inconsistencies with the *AMA Manual of Style*.

The preferred way to submit new entries, clarifications, mnemonics, or potential corrections with a valid, authoritative reference is via our website: www.firstaidteam.com.

This website will be continuously updated with validated errata, new high-yield content, and a new online platform to contribute suggestions, mnemonics, diagrams, clinical images, and potential errata.

Alternatively, you can email us at: firstaid@scholarrx.com.

Contributions submitted by **May 15, 2020**, receive priority consideration for the 2021 edition of *First Aid for the USMLE Step 1*. We thank you for taking the time to share your experience and apologize in advance that we cannot individually respond to all contributors as we receive thousands of contributions each year.

▶ NOTE TO CONTRIBUTORS

All contributions become property of the authors and are subject to editing and reviewing. Please verify all data and spellings carefully. Contributions should be supported by at least two high-quality references.

Check our website first to avoid duplicate submissions. In the event that similar or duplicate entries are received, only the first complete entry received with valid, authoritative references will be credited. Please follow the style, punctuation, and format of this edition as much as possible.

▶ JOIN THE FIRST AID TEAM

The *First Aid* author team is pleased to offer part-time and full-time paid internships in medical education and publishing to motivated medical students and physicians. Internships range from a few months (eg, a summer) up to a full year. Participants will have an opportunity to author, edit, and earn academic credit on a wide variety of projects, including the popular *First Aid* series.

For 2020, we are actively seeking passionate medical students and graduates with a specific interest in improving our medical illustrations, expanding our database of medical photographs, and developing the software that supports our crowdsourcing platform. We welcome people with prior experience and talent in these areas. Relevant skills include clinical imaging, digital photography, digital asset management, information design, medical illustration, graphic design, tutoring, and software development.

Please email us at firstaid@scholarrx.com with a CV and summary of your interest or sample work.

How to Use This Book

CONGRATULATIONS: You now possess the book that has guided nearly two million students to USMLE success for 30 years. With appropriate care, the binding should last the useful life of the book. Keep in mind that putting excessive flattening pressure on any binding will accelerate its failure. If you purchased a book that you believe is defective, please **immediately** return it to the place of purchase. If you encounter ongoing issues, you can also contact Customer Service at our publisher, McGraw-Hill Education, at <https://www.mheducation.com/contact.html>.

START EARLY: Use this book as early as possible while learning the basic medical sciences. The first semester of your first year is not too early! Devise a study plan by reading Section I: Guide to Efficient Exam Preparation, and make an early decision on resources to use by checking Section IV: Top-Rated Review Resources. Note that *First Aid* is neither a textbook nor a comprehensive review book, and it is not a panacea for inadequate preparation.

CONSIDER FIRST AID YOUR ANNOTATION HUB: Annotate material from other resources, such as class notes or comprehensive textbooks, into your book. This will keep all the high-yield information you need in one place. Other tips on keeping yourself organized:

- For best results, use fine-tipped ballpoint pens (eg, BIC Pro+, Uni-Ball Jetstream Sports, Pilot Drawing Pen, Zebra F-301). If you like gel pens, try Pentel Slicci, and for markers that dry almost immediately, consider Staedtler Triplus Fineliner, Pilot Drawing Pen, and Sharpies.
- Consider using pens with different colors of ink to indicate different sources of information (eg, blue for USMLE-Rx Step 1 Qmax, green for UWorld Step 1 Qbank).
- Choose highlighters that are bright and dry quickly to minimize smudging and bleeding through the page (eg, Tombow Kei Coat, Sharpie Gel).
- Many students de-spine their book and get it 3-hole-punched. This will allow you to insert materials from other sources, including curricular materials.

INTEGRATE STUDY WITH CASES, FLASH CARDS, AND QUESTIONS: To broaden your learning strategy, consider integrating your *First Aid* study with case-based reviews (eg, *First Aid Cases for the USMLE Step 1*), flash cards (eg, First Aid Flash Facts), and practice questions (eg, the USMLE-Rx Step 1 Qmax). Read the chapter in the book, then test your comprehension by using cases, flash cards, and questions that cover the same topics. Maintain access to more comprehensive resources (eg, *First Aid for the Basic Sciences: General Principles and Organ Systems* and First Aid Express videos) for deeper review as needed.

PRIME YOUR MEMORY: Return to your annotated Sections II and III several days before taking the USMLE Step 1. The book can serve as a useful way of retaining key associations and keeping high-yield facts fresh in your memory just prior to the exam. The Rapid Review section includes high-yield topics to help guide your studying.

CONTRIBUTE TO FIRST AID: Reviewing the book immediately after your exam can help us improve the next edition. Decide what was truly high and low yield and send us your comments. Feel free to send us scanned images from your annotated *First Aid* book as additional support. Of course, always remember that **all examinees are under agreement with the NBME to not disclose the specific details of copyrighted test material.**

Selected USMLE Laboratory Values

* = Included in the Biochemical Profile (SMA-12)

Blood, Plasma, Serum	Reference Range	SI Reference Intervals
* Alanine aminotransferase (ALT, GPT at 30°C)	8–20 U/L	8–20 U/L
Amylase, serum	25–125 U/L	25–125 U/L
* Aspartate aminotransferase (AST, GOT at 30°C)	8–20 U/L	8–20 U/L
Bilirubin, serum (adult)		
Total // Direct	0.1–1.0 mg/dL // 0.0–0.3 mg/dL	2–17 μmol/L // 0–5 μmol/L
* Calcium, serum (Total)	8.4–10.2 mg/dL	2.1–2.8 mmol/L
* Cholesterol, serum (Total)	Rec: < 200 mg/dL	< 5.2 mmol/L
* Creatinine, serum (Total)	0.6–1.2 mg/dL	53–106 μmol/L
Electrolytes, serum		
Sodium (Na ⁺)	136–145 mEq/L	136–145 mmol/L
Chloride (Cl ⁻)	95–105 mEq/L	95–105 mmol/L
* Potassium (K ⁺)	3.5–5.0 mEq/L	3.5–5.0 mmol/L
Bicarbonate (HCO ₃ ⁻)	22–28 mEq/L	22–28 mmol/L
Magnesium (Mg ²⁺)	1.5–2 mEq/L	0.75–1.0 mmol/L
Gases, arterial blood (room air)		
P _{O₂}	75–105 mm Hg	10.0–14.0 kPa
P _{CO₂}	33–45 mm Hg	4.4–5.9 kPa
pH	7.35–7.45	[H ⁺] 36–44 nmol/L
* Glucose, serum	Fasting: 70–110 mg/dL 2-h postprandial: < 120 mg/dL	3.8–6.1 mmol/L < 6.6 mmol/L
Growth hormone – arginine stimulation	Fasting: < 5 ng/mL provocative stimuli: > 7 ng/mL	< 5 μg/L > 7 μg/L
Osmolality, serum	275–295 mOsm/kg	275–295 mOsm/kg
* Phosphatase (alkaline), serum (p-NPP at 30°C)	20–70 U/L	20–70 U/L
* Phosphorus (inorganic), serum	3.0–4.5 mg/dL	1.0–1.5 mmol/L
Prolactin, serum (hPRL)	< 20 ng/mL	< 20 μg/L
* Proteins, serum		
Total (recumbent)	6.0–7.8 g/dL	60–78 g/L
Albumin	3.5–5.5 g/dL	35–55 g/L
Globulins	2.3–3.5 g/dL	23–35 g/L
Thyroid-stimulating hormone, serum or plasma	0.5–5.0 μU/mL	0.5–5.0 mU/L
* Urea nitrogen, serum (BUN)	7–18 mg/dL	1.2–3.0 mmol/L
* Uric acid, serum	3.0–8.2 mg/dL	0.18–0.48 mmol/L

(continues)

XX

Cerebrospinal Fluid	Reference Range	SI Reference Intervals
Glucose	40–70 mg/dL	2.2–3.9 mmol/L
Hematologic		
Erythrocyte count	Male: 4.3–5.9 million/mm ³ Female: 3.5–5.5 million/mm ³	4.3–5.9 × 10 ¹² /L 3.5–5.5 × 10 ¹² /L
Erythrocyte sedimentation rate (Westergen)	Male: 0–15 mm/h Female: 0–20 mm/h	0–15 mm/h 0–20 mm/h
Hematocrit	Male: 41–53% Female: 36–46%	0.41–0.53 0.36–0.46
Hemoglobin, blood	Male: 13.5–17.5 g/dL Female: 12.0–16.0 g/dL	2.09–2.71 mmol/L 1.86–2.48 mmol/L
Hemoglobin, plasma	1–4 mg/dL	0.16–0.62 μmol/L
Leukocyte count and differential		
Leukocyte count	4,500–11,000/mm ³	4.5–11.0 × 10 ⁹ /L
Segmented neutrophils	54–62%	0.54–0.62
Band forms	3–5%	0.03–0.05
Eosinophils	1–3%	0.01–0.03
Basophils	0–0.75%	0–0.0075
Lymphocytes	25–33%	0.25–0.33
Monocytes	3–7%	0.03–0.07
Mean corpuscular hemoglobin	25.4–34.6 pg/cell	0.39–0.54 fmol/cell
Mean corpuscular volume	80–100 μm ³	80–100 fL
Partial thromboplastin time (activated)	25–40 seconds	25–40 seconds
Platelet count	150,000–400,000/mm ³	150–400 × 10 ⁹ /L
Prothrombin time	11–15 seconds	11–15 seconds
Reticulocyte count	0.5–1.5% of red cells	0.005–0.015
Sweat		
Chloride	0–35 mmol/L	0–35 mmol/L
Urine		
Creatinine clearance	Male: 97–137 mL/min Female: 88–128 mL/min	
Osmolality	50–1,400 mOsmol/kg H ₂ O	
Proteins, total	< 150 mg/24 h	< 0.15 g/24 h

First Aid Checklist for the USMLE Step 1

This is an example of how you might use the information in Section I to prepare for the USMLE Step 1. Refer to corresponding topics in Section I for more details.

- Years Prior** — Use top-rated review resources for first-year medical school courses.
 Ask for advice from those who have recently taken the USMLE Step 1.
- Months Prior** — Review computer test format and registration information.
 Register six months in advance.
 Carefully verify name and address printed on scheduling permit. Make sure the name on scheduling permit matches the name printed on your photo ID.
 Go online for test date ASAP.
 Define your exam goals (pass comfortably, beat the mean, ace the test)
 Set up a realistic timeline for study. Cover less crammable subjects first.
 Evaluate and choose study materials (review books, question banks).
 Use a question bank to simulate the USMLE Step 1 to pinpoint strengths and weaknesses in knowledge and test-taking skills.
- Weeks Prior** — Do another test simulation in a question bank.
 Assess how close you are to your goal.
 Pinpoint remaining weaknesses. Stay healthy (exercise, sleep).
 Verify information on admission ticket (eg, location, date).
- One Week Prior** — Remember comfort measures (loose clothing, earplugs, etc).
 Work out test site logistics (eg, location, transportation, parking, lunch).
 Print or download your Scheduling Permit and Scheduling Confirmation to your phone.
- One Day Prior** — Relax.
 Lightly review short-term material if necessary. Skim high-yield facts.
 Get a good night's sleep.
- Day of Exam** — Relax.
 Eat breakfast.
 Minimize bathroom breaks during exam by avoiding excessive morning caffeine.
- After Exam** — Celebrate, regardless of how well you feel you did.
 Send feedback to us on our website at www.firstaidteam.com.

SECTION I

Guide to Efficient Exam Preparation

“I don’t love studying. I hate studying. I like learning. Learning is beautiful.”

—Natalie Portman

“Finally, from so little sleeping and so much reading, his brain dried up and he went completely out of his mind.”

—Miguel de Cervantes Saavedra, *Don Quixote*

“Sometimes the questions are complicated and the answers are simple.”

—Dr. Seuss

“He who knows all the answers has not been asked all the questions.”

—Confucius

“The expert in anything was once a beginner.”

—Helen Hayes

“It always seems impossible until it’s done.”

—Nelson Mandela

▶ Introduction	2
▶ USMLE Step 1—The Basics	2
▶ Defining Your Goal	12
▶ Learning Strategies	13
▶ Timeline for Study	16
▶ Study Materials	20
▶ Test-Taking Strategies	22
▶ Clinical Vignette Strategies	23
▶ If You Think You Failed	24
▶ Testing Agencies	24
▶ References	25

▶ INTRODUCTION

Relax.

This section is intended to make your exam preparation easier, not harder. Our goal is to reduce your level of anxiety and help you make the most of your efforts by helping you understand more about the United States Medical Licensing Examination, Step 1 (USMLE Step 1). As a medical student, you are no doubt familiar with taking standardized examinations and quickly absorbing large amounts of material. When you first confront the USMLE Step 1, however, you may find it all too easy to become sidetracked from your goal of studying with maximal effectiveness. Common mistakes that students make when studying for Step 1 include the following:

- Starting to study (including *First Aid*) too late
- Starting to study intensely too early and burning out
- Starting to prepare for boards before creating a knowledge foundation
- Using inefficient or inappropriate study methods
- Buying the wrong resources or buying too many resources
- Buying only one publisher's review series for all subjects
- Not using practice examinations to maximum benefit
- Not understanding how scoring is performed or what the score means
- Not using review books along with your classes
- Not analyzing and improving your test-taking strategies
- Getting bogged down by reviewing difficult topics excessively
- Studying material that is rarely tested on the USMLE Step 1
- Failing to master certain high-yield subjects owing to overconfidence
- Using *First Aid* as your sole study resource
- Trying to prepare for it all alone

▶ The test at a glance:

- 8-hour exam
- Up to a total of 280 multiple choice items
- 7 test blocks (60 min/block)
- Up to 40 test items per block
- 45 minutes of break time, plus another 15 if you skip the tutorial

In this section, we offer advice to help you avoid these pitfalls and be more productive in your studies.

▶ USMLE STEP 1—THE BASICS

The USMLE Step 1 is the first of three examinations that you must pass in order to become a licensed physician in the United States. The USMLE is a joint endeavor of the National Board of Medical Examiners (NBME) and the Federation of State Medical Boards (FSMB). The USMLE serves as the single examination system for US medical students and international medical graduates (IMGs) seeking medical licensure in the United States.

The Step 1 exam includes test items that can be grouped by the organizational constructs outlined in Table 1 (in order of tested frequency).

TABLE 1. Frequency of Various Constructs Tested on the USMLE Step 1.*

Competency	Range, %	System	Range, %
Medical knowledge: applying foundational science concepts	52–62	General principles	13–17
Patient care: diagnosis	20–30	Behavioral health & nervous systems/special senses	9–13
Patient care: management	7–12	Respiratory & renal/urinary systems	9–13
Practice-based learning & improvement	5–7	Reproductive & endocrine systems	9–13
Communication/professionalism	2–5	Blood & lymphoreticular/immune systems	7–11
Discipline	Range, %	Multisystem processes & disorders	7–11
Pathology	45–52	Musculoskeletal, skin & subcutaneous tissue	6–10
Physiology	26–34	Cardiovascular system	6–10
Pharmacology	16–23	Gastrointestinal system	5–9
Biochemistry & nutrition	14–24	Biostatistics & epidemiology/population health	5–7
Microbiology & immunology	15–22	Social sciences: communication skills/ethics	3–5
Gross anatomy & embryology	11–15		
Histology & cell biology	9–13		
Behavioral sciences	8–12		
Genetics	5–9		

*Percentages are subject to change at any time. www.usmle.org

How Is the Computer-Based Test (CBT) Structured?

The CBT Step 1 exam consists of one “optional” tutorial/simulation block and seven “real” question blocks of up to 40 questions per block with no more than 280 questions in total, timed at 60 minutes per block. A short 11-question survey follows the last question block. The computer begins the survey with a prompt to proceed to the next block of questions.

Once an examinee finishes a particular question block on the CBT, he or she must click on a screen icon to continue to the next block. Examinees **cannot** go back and change their answers to questions from any previously completed block. However, changing answers is allowed **within** a block of questions as long as the block has not been ended and if time permits.

What Is the CBT Like?

Given the unique environment of the CBT, it's important that you become familiar ahead of time with what your test-day conditions will be like. You can access a 15-minute tutorial and practice blocks at <http://orientation.nbme.org/Launch/USMLE/STPF1>. This tutorial interface is very similar to the one you will use in the exam; learn it now and you can skip taking it during the exam, giving you up to 15 extra minutes of break time. You can gain experience with the CBT format by taking the 120 practice questions (3 blocks with 40 questions each) available online or by signing up for a practice session at a test center for a fee.

► **Keyboard shortcuts:**

- *A, B, etc—letter choices*
- *Enter or spacebar—move to next question*
- *Esc—exit pop-up Calculator and Notes windows*

► *Heart sounds are tested via media questions. Make sure you know how different heart diseases sound on auscultation.*

► *Be sure to test your headphones during the tutorial.*

► *Familiarize yourself with the commonly tested lab values (eg, Hgb, WBC, platelets, Na⁺, K⁺).*

► **Illustrations on the test include:**

- *Gross specimen photos*
- *Histology slides*
- *Medical imaging (eg, x-ray, CT, MRI)*
- *Electron micrographs*
- *Line drawings*

► *Ctrl-Alt-Delete are the keys of death during the exam. Don't touch them at the same time!*

For security reasons, examinees are not allowed to bring any personal electronic equipment into the testing area. This includes both digital and analog watches, iPods, tablets, calculators, cell phones, and electronic paging devices. Examinees are also prohibited from carrying in their books, notes, pens/pencils, and scratch paper. Food and beverages are also prohibited in the testing area. The testing centers are monitored by audio and video surveillance equipment. However, most testing centers allot each examinee a small locker outside the testing area in which he or she can store snacks, beverages, and personal items.

Questions are typically presented in multiple choice format, with 4–5 possible answer options. There is a countdown timer on the lower left corner of the screen as well. There is also a button that allows the examinee to mark a question for review. If a given question happens to be longer than the screen (which occurs very rarely), a scroll bar will appear on the right, allowing the examinee to see the rest of the question. Regardless of whether the examinee clicks on an answer choice or leaves it blank, he or she must click the “Next” button to advance to the next question.

The USMLE features a small number of media clips in the form of audio and/or video. There may even be a question with a multimedia heart sound simulation. In these questions, a digital image of a torso appears on the screen, and the examinee directs a digital stethoscope to various auscultation points to listen for heart and breath sounds. The USMLE orientation materials include several practice questions in these formats. During the exam tutorial, examinees are given an opportunity to ensure that both the audio headphones and the volume are functioning properly. If you are already familiar with the tutorial and planning on skipping it, first skip ahead to the section where you can test your headphones. After you are sure the headphones are working properly, proceed to the exam.

The examinee can call up a window displaying normal laboratory values. In order to do so, he or she must click the “Lab” icon on the top part of the screen. Afterward, the examinee will have the option to choose between “Blood,” “Cerebrospinal,” “Hematologic,” or “Sweat and Urine.” The normal values screen may obscure the question if it is expanded. The examinee may have to scroll down to search for the needed lab values. You might want to memorize some common lab values so you spend less time on questions that require you to analyze these.

The CBT interface provides a running list of questions on the left part of the screen at all times. The software also permits examinees to highlight or cross out information by using their mouse. There is a “Notes” icon on the top part of the screen that allows students to write notes to themselves for review at a later time. Finally, the USMLE has recently added new functionality including text magnification and reverse color (white text on black background). Being familiar with these features can save time and may help you better view and organize the information you need to answer a question.

For those who feel they might benefit, the USMLE offers an opportunity to take a simulated test, or “CBT Practice Session” at a Prometric center. Students are eligible to register for this three-and-one-half-hour practice session after they have received their scheduling permit.

The same USMLE Step 1 sample test items (120 questions) available on the USMLE website, www.usmle.org, are used at these sessions. **No new items will be presented.** The practice session is available at a cost of \$75 (or more if taken outside of the US and Canada) and is divided into a short tutorial and three 1-hour blocks of ~40 test items each. Students receive a printed percent-correct score after completing the session. **No explanations of questions are provided.**

You may register for a practice session online at www.usmle.org. A separate scheduling permit is issued for the practice session. Students should allow two weeks for receipt of this permit.

How Do I Register to Take the Exam?

Prometric test centers offer Step 1 on a year-round basis, except for the first two weeks in January and major holidays. The exam is given every day except Sunday at most centers. Some schools administer the exam on their own campuses. Check with the test center you want to use before making your exam plans.

US students can apply to take Step 1 at the NBME website. This application allows you to select one of 12 overlapping three-month blocks in which to be tested (eg, April–May–June, June–July–August). Choose your three-month eligibility period wisely. If you need to reschedule outside your initial three-month period, you can request a one-time extension of eligibility for the next contiguous three-month period, and pay a rescheduling fee. The application also includes a photo ID form that must be certified by an official at your medical school to verify your enrollment. After the NBME processes your application, it will send you a scheduling permit.

The scheduling permit you receive from the NBME will contain your USMLE identification number, the eligibility period in which you may take the exam, and two additional numbers. The first of these is known as your “scheduling number.” You must have this number in order to make your exam appointment with Prometric. The second number is known as the “candidate identification number,” or CIN. Examinees must enter their CINs at the Prometric workstation in order to access their exams. However, you will not be allowed to bring your permit into the exam and will be asked to copy your CIN onto your scratch paper. Prometric has no access to the codes. **Make sure to bring a paper or electronic copy of your permit with you to the exam!** Also bring an unexpired, government-issued photo ID that includes your signature (such as a driver’s license or passport). Make sure the name on your photo ID exactly matches the name that appears on your scheduling permit.

▶ You can take a shortened CBT practice test at a Prometric center.

▶ The Prometric website will display a calendar with open test dates.

► *The confirmation emails that Prometric and NBME send are not the same as the scheduling permit.*

► *Test scheduling is done on a “first-come, first-served” basis. It’s important to schedule an exam date as soon as you receive your scheduling permit.*

► *Register six months in advance for seating and scheduling preference.*

Once you receive your scheduling permit, you may access the Prometric website or call Prometric’s toll-free number to arrange a time to take the exam. You may contact Prometric two weeks before the test date if you want to confirm identification requirements. Although requests for taking the exam may be completed more than six months before the test date, examinees will not receive their scheduling permits earlier than six months before the eligibility period. The eligibility period is the three-month period you have chosen to take the exam. Most medical students choose the April–June or June–August period. Because exams are scheduled on a “first-come, first-served” basis, it is recommended that you book an exam date on the Prometric website as soon as you receive your permit. Prometric will provide appointment confirmation on a print-out and by email. Be sure to read the latest *USMLE Bulletin of Information* for further details.

What If I Need to Reschedule the Exam?

You can change your test date and/or center by contacting Prometric at 1-800-MED-EXAM (1-800-633-3926) or www.prometric.com. Make sure to have your CIN when rescheduling. If you are rescheduling by phone, you must speak with a Prometric representative; leaving a voicemail message will not suffice. To avoid a rescheduling fee, you will need to request a change at least 31 calendar days before your appointment. Please note that your rescheduled test date must fall within your assigned three-month eligibility period.

When Should I Register for the Exam?

You should plan to register as far in advance as possible ahead of your desired test date (eg, six months), but, depending on your particular test center, new dates and times may open closer to the date. Scheduling early will guarantee that you will get either your test center of choice or one within a 50-mile radius of your first choice. For most US medical students, the desired testing window is in June, since most medical school curricula for the second year end in May or June. Thus, US medical students should plan to register before January in anticipation of a June test date. The timing of the exam is more flexible for IMGs, as it is related only to when they finish exam preparation. Talk with upperclassmen who have already taken the test so you have real-life experience from students who went through a similar curriculum, then formulate your own strategy.

Where Can I Take the Exam?

Your testing location is arranged with Prometric when you book your test date (after you receive your scheduling permit). For a list of Prometric locations nearest you, visit www.prometric.com.

How Long Will I Have to Wait Before I Get My Scores?

The USMLE reports scores in three to four weeks, unless there are delays in score processing. Examinees will be notified via email when their scores are available. By following the online instructions, examinees will be able to view, download, and print their score report online for ~120 days after score notification, after which scores can only be obtained through requesting an official USMLE transcript. Additional information about score timetables and accessibility is available on the official USMLE website.

What About Time?

Time is of special interest on the CBT exam. Here's a breakdown of the exam schedule:

15 minutes	Tutorial (skip if familiar with test format and features)
7 hours	Seven 60-minute question blocks
45 minutes	Break time (includes time for lunch)

The computer will keep track of how much time has elapsed on the exam. However, the computer will show you only how much time you have remaining in a given block. Therefore, it is up to you to determine if you are pacing yourself properly (at a rate of approximately one question per 90 seconds).

The computer does not warn you if you are spending more than your allotted time for a break. You should therefore budget your time so that you can take a short break when you need one and have time to eat. You must be especially careful not to spend too much time in between blocks (you should keep track of how much time elapses from the time you finish a block of questions to the time you start the next block). After you finish one question block, you'll need to click to proceed to the next block of questions. If you do not click within 30 seconds, you will automatically be entered into a break period.

Break time for the day is 45 minutes, but you are not required to use all of it, nor are you required to use any of it. You can gain extra break time (but not extra time for the question blocks) by skipping the tutorial or by finishing a block ahead of the allotted time. Any time remaining on the clock when you finish a block gets added to your remaining break time. Once a new question block has been started, you may not take a break until you have reached the end of that block. If you do so, this will be recorded as an "unauthorized break" and will be reported on your final score report.

Finally, be aware that it may take a few minutes of your break time to "check out" of the secure resting room and then "check in" again to resume testing, so plan accordingly. The "check-in" process may include fingerprints, pocket checks, and metal detector scanning. Some students recommend pocketless clothing on exam day to streamline the process.

► Gain extra break time by skipping the tutorial or finishing a block early.

► Be careful to watch the clock on your break time.

If I Freak Out and Leave, What Happens to My Score?

Your scheduling permit shows a CIN that you will need to enter to start your exam. Entering the CIN is the same as breaking the seal on a test book, and you are considered to have started the exam when you do so. However, no score will be reported if you do not complete the exam. In fact, if you leave at any time from the start of the test to the last block, no score will be reported. The fact that you started but did not complete the exam, however, will appear on your USMLE score transcript. Even though a score is not posted for incomplete tests, examinees may still get an option to request that their scores be calculated and reported if they desire; unanswered questions will be scored as incorrect.

The exam ends when all question blocks have been completed or when their time has expired. As you leave the testing center, you will receive a printed test-completion notice to document your completion of the exam. To receive an official score, you must finish the entire exam.

What Types of Questions Are Asked?

► Nearly three fourths of Step 1 questions begin with a description of a patient.


All questions on the exam are **one-best-answer multiple choice items**. Most questions consist of a clinical scenario or a direct question followed by a list of five or more options. You are required to select the single best answer among the options given. There are no “except,” “not,” or matching questions on the exam. A number of options may be partially correct, in which case you must select the option that best answers the question or completes the statement. Additionally, keep in mind that experimental questions may appear on the exam, which do not affect your score.

How Is the Test Scored?

Each Step 1 examinee receives an electronic score report that includes the examinee’s pass/fail status, a three-digit test score, a bar chart comparing the examinee’s performance to that of other examinees’, and a graphic depiction of the examinee’s performance by physician task, discipline and organ system.

The USMLE score report highlights the examinee’s strength and weaknesses by providing an overview of their performance by physician task, discipline and organ system compared to their overall performance on the exam. Each of the questions (minus experimental questions) is tagged according to any or all relevant content areas. Yellow-colored boxes (lower, same, higher) on your score report indicate your performance in each specific content area relative to your overall performance on the exam. This is often a direct consequence of the total number of questions for each physician task, discipline or system, which is indicated by percentage range after each specified content area on the score report (see Figure 1).

FIGURE 1. Samples from the USMLE Step 1 Performance Profile.



The NBME provides a three-digit test score based on the total number of items answered correctly on the examination, which corresponds to a particular percentile (see Figure 2). Your three-digit score will be qualified by the mean and standard deviation of US and Canadian medical school first-time examinees.


Since some questions may be experimental and are not counted, it is possible to get different scores for the same number of correct answers. In 2018, the mean score was 230 with a standard deviation of 19.

The passing score for Step 1 is 194. The NBME does not report the minimum number of correct responses needed to pass, but estimates that it is roughly 60–70%. The NBME may adjust the minimum passing score in the future, so please check the USMLE website or www.firstaidteam.com for updates.

According to the USMLE, medical schools receive a listing of total scores and pass/fail results plus group summaries by discipline and organ system. Students can withhold their scores from their medical school if they wish. Official USMLE transcripts, which can be sent on request to residency programs, include only total scores, not performance profiles.

► The mean Step 1 score for US medical students continues to rise, from 200 in 1991 to 230 in 2018.

FIGURE 2. Score and Percentile for First-time Step 1 Takers.



Consult the USMLE website or your medical school for the most current and accurate information regarding the examination.

What Does My Score Mean?

The most important point with the Step 1 score is passing versus failing. Passing essentially means, “Hey, you’re on your way to becoming a fully licensed doc.” As Table 2 shows, the majority of students pass the exam, so remember, we told you to relax.

TABLE 2. Passing Rates for the 2017–2018 USMLE Step 1.²

	2017		2018	
	No. Tested	% Passing	No. Tested	% Passing
Allopathic 1st takers	20,353	96%	20,670	96%
Repeaters	1,029	67%	941	67%
Allopathic total	21,382	94%	21,611	95%
Osteopathic 1st takers	3,786	95%	4,092	96%
Repeaters	49	76%	44	73%
Osteopathic total	3,835	95%	4,136	96%
Total US/Canadian	25,217	94%	25,747	94%
IMG 1st takers	14,900	78%	14,332	80%
Repeaters	2,303	41%	2,111	44%
IMG total	17,203	73%	16,443	75%
Total Step 1 examinees	42,420	85%	42,190	86%

Beyond that, the main point of having a quantitative score is to give you a sense of how well you've done on the exam and to help schools and residencies rank their students and applicants, respectively.

Official NBME/USMLE Resources

The NBME offers a Comprehensive Basic Science Examination (CBSE) for practice that is a shorter version of the Step 1. The CBSE contains four blocks of 50 questions each and covers material that is typically learned during the basic science years. Scores range from 45 to 95 and correlate with a Step 1 equivalent (see Table 3). The standard error of measurement is approximately 3 points, meaning a score of 80 would estimate the student's proficiency is somewhere between 77 and 83. In other words, the actual Step 1 score could be predicted to be between 218 and 232. Of course, these values do not correlate exactly, and they do not reflect different test preparation methods. Many schools use this test to gauge whether a student is expected to pass Step 1. If this test is offered by your school, it is usually conducted at the end of regular didactic time before any dedicated Step 1 preparation. If you do not encounter the CBSE before your dedicated study time, you need not worry about taking it. Use the information to help set realistic goals and timetables for your success.

The NBME also offers six forms of Comprehensive Basic Science Self-Assessment (CBSSA). Students who prepared for the exam using this web-based tool reported that they found the format and content highly indicative of questions tested on the actual exam. In addition, the CBSSA is a fair predictor of USMLE performance (see Table 4). The test interface, however, does not match the actual USMLE test interface, so practicing with these forms alone is not advised.

The CBSSA exists in two formats: standard-paced and self-paced, both of which consist of four sections of 50 questions each (for a total of 200 multiple choice items). The standard-paced format allows the user up to 75 minutes to complete each section, reflecting time limits similar to the actual exam. By contrast, the self-paced format places a 5-hour time limit on answering all multiple choice questions. Every few years, a new form is released and an older one is retired, reflecting changes in exam content. Therefore, the newer exams tend to be more similar to the actual Step 1, and scores from these exams tend to provide a better estimation of exam day performance.

Keep in mind that this bank of questions is available only on the web. The NBME requires that users start and complete the exam within 90 days of purchase. Once the assessment has begun, users are required to complete the sections within 20 days. Following completion of the questions, the CBSSA provides a performance profile indicating the user's relative strengths and weaknesses, much like the report profile for the USMLE Step 1 exam. The profile is scaled with an average score of 500 and a standard deviation of 100. In addition to the performance profile, examinees will be informed of the number of questions answered incorrectly. You will have the ability to review the text of the incorrect question with the correct answer.

TABLE 3. CBSE to USMLE Score Prediction.

CBSE Score	Step 1 Equivalent
≥ 94	≥ 260
92	255
90	250
88	245
86	240
84	235
82	230
80	225
78	220
76	215
74	210
72	205
70	200
68	195
66	190
64	185
62	180
60	175
58	170
56	165
54	160
52	155
50	150
48	145
46	140
≤ 44	≤ 135

► Practice questions may be easier than the actual exam.

TABLE 4. CBSSA to USMLE Score Prediction.

CBSSA Score	Approximate USMLE Step 1 Score
150	155
200	165
250	175
300	186
350	196
400	207
450	217
500	228
550	238
600	248
650	259
700	269
750	280
800	290

Explanations for the correct answer, however, will not be provided. The NBME charges \$60 for assessments with expanded feedback. The fees are payable by credit card or money order. For more information regarding the CBSE and the CBSSA, visit the NBME’s website at www.nbme.org.

The NBME scoring system is weighted for each assessment exam. While some exams seem more difficult than others, the score reported takes into account these inter-test differences when predicting Step 1 performance. Also, while many students report seeing Step 1 questions “word-for-word” out of the assessments, the NBME makes special note that no live USMLE questions are shown on any NBME assessment.

Lastly, the International Foundations of Medicine (IFOM) offers a Basic Science Examination (BSE) practice exam at participating Prometric test centers for \$200. Students may also take the self-assessment test online for \$35 through the NBME’s website. The IFOM BSE is intended to determine an examinee’s relative areas of strength and weakness in general areas of basic science—not to predict performance on the USMLE Step 1 exam—and the content covered by the two examinations is somewhat different. However, because there is substantial overlap in content coverage and many IFOM items were previously used on the USMLE Step 1, it is possible to roughly project IFOM performance onto the USMLE Step 1 score scale. More information is available at <http://www.nbme.org/ifom/>.


► DEFINING YOUR GOAL

It is useful to define your own personal performance goal when approaching the USMLE Step 1. Your style and intensity of preparation can then be matched to your goal. Furthermore, your goal may depend on your school’s requirements, your specialty choice, your grades to date, and your personal assessment of the test’s importance. Do your best to define your goals early so that you can prepare accordingly.

The value of the USMLE Step 1 score in selecting residency applicants remains controversial, and some have called for less emphasis to be placed on the score when selecting or screening applicants.³ For the time being, however, it continues to be an important part of the residency application, and it is not uncommon for some specialties to implement filters that screen out applicants who score below a certain cutoff. This is more likely to be seen in competitive specialties (eg, orthopedic surgery, ophthalmology, dermatology, otolaryngology). Independent of your career goals, you can maximize your future options by doing your best to obtain the highest score possible (see Figure 3). At the same time, your Step 1 score is only one of a number of factors that are assessed when you apply for residency. In fact, many residency programs value other criteria such as letters of recommendation, third-year clerkship grades, honors, and research experience more than a high score on Step 1. Fourth-year medical students who have recently completed the residency application process can be a valuable resource in this regard.

► *Some competitive residency programs place more weight on Step 1 scores when choosing candidates to interview.*

► *Fourth-year medical students have the best feel for how Step 1 scores factor into the residency application process.*

FIGURE 3. Median USMLE Step 1 Score by Specialty for Matched US Seniors.^{a,b}

▶ LEARNING STRATEGIES

Many students feel overwhelmed during the preclinical years and struggle to find an effective learning strategy. Table 5 lists several learning strategies you can try and their estimated effectiveness for Step 1 preparation based on the literature (see References). These are merely suggestions, and it's important to take your learning preferences into account. Your comprehensive learning approach will contain a combination of strategies (eg, elaborative interrogation followed by practice testing, mnemonics review using spaced repetition, etc). Regardless of your choice, the foundation of knowledge you build during your basic science years is the most important resource for success on the USMLE Step 1.

▶ *The foundation of knowledge you build during your basic science years is the most important resource for success on the USMLE Step 1.*

HIGH EFFICACY

Practice Testing

Also called “retrieval practice,” practice testing has both direct and indirect benefits to the learner.⁴ Effortful retrieval of answers does not only identify weak spots—it directly strengthens long-term retention of material.⁵ The more effortful the recall, the better the long-term retention. This advantage has been shown to result in higher test scores and GPAs.⁶ In fact, research has shown a positive correlation between the number of boards-style practice questions completed and Step 1 scores among medical students.⁷

▶ *Research has shown a positive correlation between the number of boards-style practice questions completed and Step 1 scores among medical students.*

Practice testing should be done with “interleaving” (mixing of questions from different topics in a single session). Question banks often allow you to intermingle topics. Interleaved practice helps learners develop their ability to focus on the relevant concept when faced with many possibilities. Practicing topics in massed fashion (eg, all cardiology, then all dermatology) may seem intuitive, but there is strong evidence that interleaving correlates with longer-

TABLE 5. Effective Learning Strategies.

EFFICACY	STRATEGY	EXAMPLE RESOURCES
<i>High efficacy</i>	Practice testing (retrieval practice)	UWorld Qbank NBME Self-Assessments USMLE-Rx QMax Kaplan Qbank
	Distributed practice	USMLE-Rx Flash Facts Anki Firecracker Memorang Osmosis
<i>Moderate efficacy</i>	Mnemonics	<i>Pre-made:</i> SketchyMedical Picmonic <i>Self-made:</i> Mullen Memory
	Elaborative interrogation/ self-explanation	
	Concept mapping	Coggle FreeMind XMind MindNode
<i>Low efficacy</i>	Rereading	
	Highlighting/underlining	
	Summarization	

term retention and increased student achievement, especially on tasks that involve problem solving.⁵

In addition to using question banks, you can test yourself by arranging your notes in a question-answer format (eg, via flash cards). Testing these Q&As in random order allows you to reap the benefit of interleaved practice. Bear in mind that the utility of practice testing comes from the practice of information retrieval, so simply reading through Q&As will attenuate this benefit.

Distributed Practice

Also called “spaced repetition,” distributed practice is the opposite of massed practice or “cramming.” Learners review material at increasingly spaced out intervals (days to weeks to months). Massed learning may produce more short-term gains and satisfaction, but learners who use distributed practice have better mastery and retention over the long term.^{5,9}

Flash cards are a simple way to incorporate both distributed practice and practice testing. Studies have linked spaced repetition learning with flash

cards to improved long-term knowledge retention and higher exam scores.^{6,8,10} Apps with automated spaced-repetition software (SRS) for flash cards exist for smartphones and tablets, so the cards are accessible anywhere. Proceed with caution: there is an art to making and reviewing cards. The ease of quickly downloading or creating digital cards can lead to flash card overload (it is unsustainable to make 50 flash cards per lecture!). Even at a modest pace, the thousands upon thousands of cards are too overwhelming for Step 1 preparation. Unless you have specific high-yield cards (and have checked the content with high-yield resources), stick to pre-made cards by reputable sources that curate the vast amount of knowledge for you.

If you prefer pen and paper, consider using a planner or spreadsheet to organize your study material over time. Distributed practice allows for some forgetting of information, and the added effort of recall over time strengthens the learning.

MODERATE EFFICACY

Mnemonics

A “mnemonic” refers to any device that assists memory, such as acronyms, mental imagery (eg, keywords with or without memory palaces), etc. Keyword mnemonics have been shown to produce superior knowledge retention when compared with rote memorization in many scenarios. However, they are generally more effective when applied to memorization-heavy, keyword-friendly topics and may not be broadly suitable.⁵ Keyword mnemonics may not produce long-term retention, so consider combining mnemonics with distributed, retrieval-based practice (eg, via flash cards with SRS).

Self-made mnemonics may have an advantage when material is simple and keyword friendly. If you can create your own mnemonic that accurately represents the material, this will be more memorable. When topics are complex and accurate mnemonics are challenging to create, pre-made mnemonics may be more effective, especially if you are inexperienced at creating mnemonics.¹¹

Elaborative Interrogation/Self-Explanation

Elaborative interrogation (“why” questions) and self-explanation (general questioning) prompt learners to generate explanations for facts. When reading passages of discrete facts, consider using these techniques, which have been shown to be more effective than rereading (eg, improved recall and better problem-solving/diagnostic performance).^{5,12,13}

Concept Mapping

Concept mapping is a method for graphically organizing knowledge, with concepts enclosed in boxes and lines drawn between related concepts.

► *Studies have linked spaced repetition learning with flash cards to improved long-term knowledge retention and higher exam scores.*

► *Elaborative interrogation and self-explanation prompt learners to generate explanations for facts, which improves recall and problem solving.*

Creating or studying concept maps may be more effective than other activities (eg, writing or reading summaries/outlines). However, studies have reached mixed conclusions about its utility, and the small size of this effect raises doubts about its authenticity and pedagogic significance.¹⁴

LOW EFFICACY

Rereading

While the most commonly used method among surveyed students, rereading has not been shown to correlate with grade point average.⁹ Due to its popularity, rereading is often a comparator in studies on learning. Other strategies that we have discussed (eg, practice testing) have been shown to be significantly more effective than rereading.

Highlighting/Underlining

Because this method is passive, it tends to be of minimal value for learning and recall. In fact, lower-performing students are more likely to use these techniques.⁹ Students who highlight and underline do not learn how to actively recall learned information and thus find it difficult to apply knowledge to exam questions.

Summarization

While more useful for improving performance on generative measures (eg, free recall or essays), summarization is less useful for exams that depend on recognition (eg, multiple choice). Findings on the overall efficacy of this method have been mixed.⁵

▶ TIMELINE FOR STUDY


Before Starting

Your preparation for the USMLE Step 1 should begin when you enter medical school. Organize and commit to studying from the beginning so that when the time comes to prepare for the USMLE, you will be ready with a strong foundation.

Make a Schedule

After you have defined your goals, map out a study schedule that is consistent with your objectives, your vacation time, the difficulty of your ongoing coursework, and your family and social commitments (see Figure 4). Determine whether you want to spread out your study time or concentrate it into 14-hour study days in the final weeks. Then factor in your own history in

FIGURE 4. Typical Timeline for the USMLE Step 1.



► *Customize your schedule. Tackle your weakest section first.*

preparing for standardized examinations (eg, SAT, MCAT). Talk to students at your school who have recently taken Step 1. Ask them for their study schedules, especially those who have study habits and goals similar to yours. Sample schedules can be found at <https://firstaidteam.com/schedules/>.

Typically, US medical schools allot between four and eight weeks for dedicated Step 1 preparation. The time you dedicate to exam preparation will depend on your target score as well as your success in preparing yourself during the first two years of medical school. Some students reserve about a week at the end of their study period for final review; others save just a few days. When you have scheduled your exam date, do your best to adhere to it. Studies show that a later testing date does not translate into a higher score, so avoid pushing back your test date without good reason.¹⁵

Make your schedule realistic, and set achievable goals. Many students make the mistake of studying at a level of detail that requires too much time for a comprehensive review—reading *Gray's Anatomy* in a couple of days is not a realistic goal! Have one catch-up day per week of studying. No matter how well you stick to your schedule, unexpected events happen. But don't let yourself procrastinate because you have catch-up days; stick to your schedule as closely as possible and revise it regularly on the basis of your actual progress. Be careful not to lose focus. Beware of feelings of inadequacy when comparing study schedules and progress with your peers. **Avoid others who stress you out.** Focus on a few top-rated resources that suit your learning style—not on some obscure books your friends may pass down to you. Accept the fact that you cannot learn it all.

You will need time for uninterrupted and focused study. Plan your personal affairs to minimize crisis situations near the date of the test. Allot an adequate number of breaks in your study schedule to avoid burnout. Maintain a healthy lifestyle with proper diet, exercise, and sleep.

Another important aspect of your preparation is your studying environment. **Study where you have always been comfortable studying.** Be sure to include everything you need close by (review books, notes, coffee, snacks, etc). If you're the kind of person who cannot study alone, form a study group with other students taking the exam. The main point here is to create a comfortable environment with minimal distractions.

► *Avoid burnout. Maintain proper diet, exercise, and sleep habits.*

► Buy review books early (first year) and use while studying for courses.

Year(s) Prior

The knowledge you gained during your first two years of medical school and even during your undergraduate years should provide the groundwork on which to base your test preparation. Student scores on NBME subject tests (commonly known as “shelf exams”) have been shown to be highly correlated with subsequent Step 1 scores.¹⁶ Moreover, undergraduate science GPAs as well as MCAT scores are strong predictors of performance on the Step 1 exam.¹⁷

We also recommend that you buy highly rated review books early in your first year of medical school and use them as you study throughout the two years. When Step 1 comes along, these books will be familiar and personalized to the way in which you learn. It is risky and intimidating to use unfamiliar review books in the final two or three weeks preceding the exam. Some students find it helpful to personalize and annotate *First Aid* throughout the curriculum.

Months Prior

Review test dates and the application procedure. Testing for the USMLE Step 1 is done on a year-round basis. If you have disabilities or special circumstances, contact the NBME as early as possible to discuss test accommodations (see the Section I Supplement at www.firstaidteam.com/bonus).

► Simulate the USMLE Step 1 under “real” conditions before beginning your studies.

Use this time to finalize your ideal schedule. Consider upcoming breaks and whether you want to relax or study. Work backward from your test date to make sure you finish at least one question bank. Also add time to redo missed or flagged questions (which may be half the bank). This is the time to build a structured plan with enough flexibility for the realities of life.

Begin doing blocks of questions from reputable question banks under “real” conditions. Don’t use tutor mode until you’re sure you can finish blocks in the allotted time. It is important to continue balancing success in your normal studies with the Step 1 test preparation process.

Weeks Prior (Dedicated Preparation)

► In the final two weeks, focus on review, practice questions, and endurance. Stay confident!

Your dedicated prep time may be one week or two months. You should have a working plan as you go into this period. Finish your schoolwork strong, take a day off, and then get to work. Start by simulating a full-length USMLE Step 1 if you haven’t yet done so. Consider doing one NBME CBSSA and the free questions from the NBME website. Alternatively, you could choose 7 blocks of randomized questions from a commercial question bank. Make sure you get feedback on your strengths and weaknesses and adjust your studying accordingly. Many students study from review sources or comprehensive programs for part of the day, then do question blocks. Also, keep in mind that reviewing a question block can take upward of two hours. Feedback from CBSSA exams and question banks will help you focus on your weaknesses.

One Week Prior

Make sure you have your CIN (found on your scheduling permit) as well as other items necessary for the day of the examination, including a current driver's license or another form of photo ID with your signature (make sure the name on your ID **exactly** matches that on your scheduling permit). Confirm the Prometric testing center location and test time. Work out how you will get to the testing center and what parking and traffic problems you might encounter. Drive separately from other students taking the test on the same day, and exchange cell phone numbers in case of emergencies. If possible, visit the testing site to get a better idea of the testing conditions you will face. Determine what you will do for lunch. Make sure you have everything you need to ensure that you will be comfortable and alert at the test site. It may be beneficial to adjust your schedule to start waking up at the same time that you will on your test day. And of course, make sure to maintain a healthy lifestyle and get enough sleep.

► *One week before the test:*

- *Sleep according to the same schedule you'll use on test day*
- *Review the CBT tutorial one last time*
- *Call Prometric to confirm test date and time*

One Day Prior

Try your best to relax and rest the night before the test. Double-check your admissions and test-taking materials as well as the comfort measures discussed earlier so that you will not have to deal with such details on the morning of the exam. At this point it will be more effective to review short-term memory material that you're already familiar with than to try to learn new material. The Rapid Review section at the end of this book is high yield for last-minute studying. Remember that regardless of how hard you have studied, you cannot know everything. There will be things on the exam that you have never even seen before, so do not panic. Do not underestimate your abilities.

Many students report difficulty sleeping the night prior to the exam. This is often exacerbated by going to bed much earlier than usual. Do whatever it takes to ensure a good night's sleep (eg, massage, exercise, warm milk, no back-lit screens at night). Do not change your daily routine prior to the exam. Exam day is not the day for a caffeine-withdrawal headache.

Morning of the Exam

On the morning of the Step 1 exam, wake up at your regular time and eat a normal breakfast. If you think it will help you, have a close friend or family member check to make sure you get out of bed. Make sure you have your scheduling permit admission ticket, test-taking materials, and comfort measures as discussed earlier. Wear loose, comfortable clothing. Plan for a variable temperature in the testing center. Arrive at the test site 30 minutes before the time designated on the admission ticket; however, do not come too early, as doing so may intensify your anxiety. When you arrive at the test site, the proctor should give you a USMLE information sheet that will explain critical factors such as the proper use of break time. Seating may be assigned, but ask to be reseated if necessary; you need to be seated in an area

- *No notes, books, calculators, pagers, cell phones, recording devices, or watches of any kind are allowed in the testing area, but they are allowed in lockers.*

► *Arrive at the testing center 30 minutes before your scheduled exam time. If you arrive more than half an hour late, you will not be allowed to take the test.*

that will allow you to remain comfortable and to concentrate. Get to know your testing station, especially if you have never been in a Prometric testing center before. Listen to your proctors regarding any changes in instructions or testing procedures that may apply to your test site.

Finally, remember that it is natural (and even beneficial) to be a little nervous. Focus on being mentally clear and alert. Avoid panic. When you are asked to begin the exam, take a deep breath, focus on the screen, and then begin. Keep an eye on the timer. Take advantage of breaks between blocks to stretch, maybe do some jumping jacks, and relax for a moment with deep breathing or stretching.

After the Test

After you have completed the exam, be sure to have fun and relax regardless of how you may feel. Taking the test is an achievement in itself. Remember, you are much more likely to have passed than not. Enjoy the free time you have before your clerkships. Expect to experience some “reentry” phenomena as you try to regain a real life. Once you have recovered sufficiently from the test (or from partying), we invite you to send us your feedback, corrections, and suggestions for entries, facts, mnemonics, strategies, resource ratings, and the like (see p. xvii, How to Contribute). Sharing your experience will benefit fellow medical students and IMGs.

► STUDY MATERIALS

Quality Considerations

Although an ever-increasing number of review books and software are now available on the market, the quality of such material is highly variable. Some common problems are as follows:

- Certain review books are too detailed to allow for review in a reasonable amount of time or cover subtopics that are not emphasized on the exam.
- Many sample question books were originally written years ago and have not been adequately updated to reflect recent trends.
- Some question banks test to a level of detail that you will not find on the exam.

► *If a given review book is not working for you, stop using it no matter how highly rated it may be or how much it costs.*

Review Books

In selecting review books, be sure to weigh different opinions against each other, read the reviews and ratings in Section IV of this guide, examine the books closely in the bookstore, and choose carefully. You are investing not only money but also your limited study time. Do not worry about finding the “perfect” book, as many subjects simply do not have one, and different students prefer different formats. Supplement your chosen books with personal notes from other sources, including what you learn from question banks.

There are two types of review books: those that are stand-alone titles and those that are part of a series. Books in a series generally have the same style, and you must decide if that style works for you. However, a given style is not optimal for every subject.

You should also find out which books are up to date. Some recent editions reflect major improvements, whereas others contain only cursory changes. Take into consideration how a book reflects the format of the USMLE Step 1.

Apps

With the explosion of smartphones and tablets, apps are an increasingly popular way to review for the Step 1 exam. The majority of apps are compatible with both iOS and Android. Many popular Step 1 review resources (eg, UWorld, USMLE-Rx) have apps that are compatible with their software. Many popular web references (eg, UpToDate) also now offer app versions. All of these apps offer flexibility, allowing you to study while away from a computer (eg, while traveling).

Practice Tests

Taking practice tests provides valuable information about potential strengths and weaknesses in your fund of knowledge and test-taking skills. Some students use practice examinations simply as a means of breaking up the monotony of studying and adding variety to their study schedule, whereas other students rely almost solely on practice. You should also subscribe to one or more high-quality question banks. In addition, students report that many current practice-exam books have questions that are, on average, shorter and less clinically oriented than those on the current USMLE Step 1.

Additionally, some students preparing for the Step 1 exam have started to incorporate case-based books intended primarily for clinical students on the wards or studying for the Step 2 CK exam. *First Aid Cases for the USMLE Step 1* aims to directly address this need.

After taking a practice test, spend time on each question and each answer choice whether you were right or wrong. There are important teaching points in each explanation. Knowing why a wrong answer choice is incorrect is just as important as knowing why the right answer is correct. Do not panic if your practice scores are low as many questions try to trick or distract you to highlight a certain point. Use the questions you missed or were unsure about to develop focused plans during your scheduled catch-up time.

Textbooks and Course Syllabi

Limit your use of textbooks and course syllabi for Step 1 review. Many textbooks are too detailed for high-yield review and include material that is generally not tested on the USMLE Step 1 (eg, drug dosages, complex chemical structures). Syllabi, although familiar, are inconsistent across

► *Charts and diagrams may be the best approach for physiology and biochemistry, whereas tables and outlines may be preferable for microbiology.*

► *Most practice exams are shorter and less clinical than the real thing.*

► *Use practice tests to identify concepts and areas of weakness, not just facts that you missed.*

medical schools and frequently reflect the emphasis of individual faculty, which often does not correspond to that of the USMLE Step 1. Syllabi also tend to be less organized than top-rated books and generally contain fewer diagrams and study questions.

▶ TEST-TAKING STRATEGIES

▶ *Practice! Develop your test-taking skills and strategies well before the test date.*

Your test performance will be influenced by both your knowledge and your test-taking skills. You can strengthen your performance by considering each of these factors. Test-taking skills and strategies should be developed and perfected well in advance of the test date so that you can concentrate on the test itself. We suggest that you try the following strategies to see if they might work for you.

Pacing

You have seven hours to complete up to 280 questions. Note that each one-hour block contains up to 40 questions. This works out to approximately 90 seconds per question. We recommend following the “1 minute rule” to pace yourself. Spend no more than 1 minute on each question. If you are still unsure about the answer after this time, mark the question, make an educated guess, and move on. Following this rule, you should have approximately 20 minutes left after all questions are answered, which you can use to revisit all of your marked questions. Remember that some questions may be experimental and do not count for points (and reassure yourself that these experimental questions are the ones that are stumping you). In the past, pacing errors have been detrimental to the performance of even highly prepared examinees. The bottom line is to keep one eye on the clock at all times!

▶ *Time management is an important skill for exam success.*

Dealing with Each Question

There are several established techniques for efficiently approaching multiple choice questions; find what works for you. One technique begins with identifying each question as easy, workable, or impossible. Your goal should be to answer all easy questions, resolve all workable questions in a reasonable amount of time, and make quick and intelligent guesses on all impossible questions. Most students read the stem, think of the answer, and turn immediately to the choices. A second technique is to first skim the answer choices to get a context, then read the last sentence of the question (the lead-in), and then read through the passage quickly, extracting only information relevant to answering the question. This can be particularly helpful for questions with long clinical vignettes. Try a variety of techniques on practice exams and see what works best for you. If you get overwhelmed, remember that a 30-second time out to refocus may get you back on track.

Guessing

There is **no penalty** for wrong answers. Thus, **no test block should be left with unanswered questions**. A hunch is probably better than a random guess. If you have to guess, we suggest selecting an answer you recognize over one with which you are totally unfamiliar.

Changing Your Answer

The conventional wisdom is not to change answers that you have already marked unless there is a convincing and logical reason to do so—in other words, go with your “first hunch.” Many question banks tell you how many questions you changed from right to wrong, wrong to wrong, and wrong to right. Use this feedback to judge how good a second-guesser you are. If you have extra time, reread the question stem and make sure you didn’t misinterpret the question.

▶ *Go with your first hunch, unless you are certain that you are a good second-guesser.*

▶ CLINICAL VIGNETTE STRATEGIES

In recent years, the USMLE Step 1 has become increasingly clinically oriented. This change mirrors the trend in medical education toward introducing students to clinical problem solving during the basic science years. The increasing clinical emphasis on Step 1 may be challenging to those students who attend schools with a more traditional curriculum.

▶ *Be prepared to read fast and think on your feet!*

What Is a Clinical Vignette?

A clinical vignette is a short (usually paragraph-long) description of a patient, including demographics, presenting symptoms, signs, and other information concerning the patient. Sometimes this paragraph is followed by a brief listing of important physical findings and/or laboratory results. The task of assimilating all this information and answering the associated question in the span of one minute can be intimidating. So be prepared to read quickly and think on your feet. Remember that the question is often indirectly asking something you already know.

▶ *Practice questions that include case histories or descriptive vignettes are critical for Step 1 preparation.*

Strategy

Remember that Step 1 vignettes usually describe diseases or disorders in their most classic presentation. So look for cardinal signs (eg, malar rash for SLE or nuchal rigidity for meningitis) in the narrative history. Be aware that the question will contain classic signs and symptoms instead of buzzwords. Sometimes the data from labs and the physical exam will help you confirm or reject possible diagnoses, thereby helping you rule answer choices in or out. In some cases, they will be a dead giveaway for the diagnosis.

▶ *Step 1 vignettes usually describe diseases or disorders in their most classic presentation.*

Making a diagnosis from the history and data is often not the final answer. Not infrequently, the diagnosis is divulged at the end of the vignette, after you have just struggled through the narrative to come up with a diagnosis of your own. The question might then ask about a related aspect of the diagnosed disease. Consider skimming the answer choices and lead-in before diving into a long stem. However, be careful with skimming the answer choices; going too fast may warp your perception of what the vignette is asking.

▶ IF YOU THINK YOU FAILED

After the test, many examinees feel that they have failed, and most are at the very least unsure of their pass/fail status. There are several sensible steps you can take to plan for the future in the event that you do not achieve a passing score. First, save and organize all your study materials, including review books, practice tests, and notes. Familiarize yourself with the reapplication procedures for Step 1, including application deadlines and upcoming test dates.

▶ *If you pass Step 1 (score of 194 or above), you are not allowed to retake the exam.*

Make sure you know both your school's and the NBME's policies regarding retakes. The NBME allows a maximum of six attempts to pass each Step examination.¹⁸ You may take Step 1 no more than three times within a 12-month period. Your fourth and subsequent attempts must be at least 12 months after your first attempt at that exam and at least six months after your most recent attempt at that exam.

The performance profiles on the back of the USMLE Step 1 score report provide valuable feedback concerning your relative strengths and weaknesses. Study these profiles closely. Set up a study timeline to strengthen gaps in your knowledge as well as to maintain and improve what you already know. Do not neglect high-yield subjects. It is normal to feel somewhat anxious about retaking the test, but if anxiety becomes a problem, seek appropriate counseling.

▶ TESTING AGENCIES

- **National Board of Medical Examiners (NBME) / USMLE Secretariat**
Department of Licensing Examination Services
3750 Market Street
Philadelphia, PA 19104-3102
(215) 590-9500 (operator) or
(215) 590-9700 (automated information line)
Email: webmail@nbme.org
www.nbme.org

- **Educational Commission for Foreign Medical Graduates (ECFMG)**
3624 Market Street
Philadelphia, PA 19104-2685
(215) 386-5900
Email: info@ecfm.org
www.ecfm.org

▶ REFERENCES

1. United States Medical Licensing Examination. Available from: https://www.usmle.org/pdfs/step-1/content_step1.pdf. Accessed October 17, 2019.
2. United States Medical Licensing Examination. 2018 Performance Data. Available from: https://www.usmle.org/performance-data/default.aspx#2018_step-1. Accessed October 17, 2019.
3. Prober CG, Kolars JC, First LR, et al. A plea to reassess the role of United States Medical Licensing Examination Step 1 scores in residency selection. *Acad Med*. 2016;91(1):12–15.
4. Roediger HL, Butler AC. The critical role of retrieval practice in long-term retention. *Trends Cogn Sci*. 2011;15(1):20–27.
5. Dunlosky J, Rawson KA, Marsh EJ, et al. Improving students' learning with effective learning techniques: promising directions from cognitive and educational psychology. *Psychol Sci Publ Int*. 2013;14(1):4–58.
6. Larsen DP, Butler AC, Lawson AL, et al. The importance of seeing the patient: test-enhanced learning with standardized patients and written tests improves clinical application of knowledge. *Adv Health Sci Educ*. 2013;18(3):409–425.
7. Panus PC, Stewart DW, Hagemeyer NE, et al. A subgroup analysis of the impact of self-testing frequency on examination scores in a pathophysiology course. *Am J Pharm Educ*. 2014;78(9):165.
8. Deng F, Gluckstein JA, Larsen DP. Student-directed retrieval practice is a predictor of medical licensing examination performance. *Perspect Med Educ*. 2015;4(6):308–313.
9. McAndrew M, Morrow CS, Atiyeh L, et al. Dental student study strategies: are self-testing and scheduling related to academic performance? *J Dent Educ*. 2016;80(5):542–552.
10. Augustin M. How to learn effectively in medical school: test yourself, learn actively, and repeat in intervals. *Yale J Biol Med*. 2014;87(2):207–212.
11. Bellezza FS. Mnemonic devices: classification, characteristics, and criteria. *Rev Educ Res*. 1981;51(2):247–275.
12. Dyer J-O, Hudon A, Montpetit-Tourangeau K, et al. Example-based learning: comparing the effects of additionally providing three different integrative learning activities on physiotherapy intervention knowledge. *BMC Med Educ*. 2015;15:37.
13. Chamberland M, Mamede S, St-Onge C, et al. Self-explanation in learning clinical reasoning: the added value of examples and prompts. *Med Educ*. 2015;49(2):193–202.
14. Nesbit JC, Adesope OO. Learning with concept and knowledge maps: a meta-analysis. *Rev Educ Res*. 2006;76(3):413–448.

15. Pohl CA, Robeson MR, Hojat M, et al. Sooner or later? USMLE Step 1 performance and test administration date at the end of the second year. *Acad Med.* 2002;77(10):S17–S19.
16. Holtman MC, Swanson DB, Ripkey DR, et al. Using basic science subject tests to identify students at risk for failing Step 1. *Acad Med.* 2001;76(10):S48–S51.
17. Basco WT, Way DP, Gilbert GE, et al. Undergraduate institutional MCAT scores as predictors of USMLE Step 1 performance. *Acad Med.* 2002;77(10):S13–S16.
18. United States Medical Licensing Examination. 2019 USMLE Bulletin of Information. Available from: <https://www.usmle.org/pdfs/bulletin/2019bulletin.pdf>. Accessed July 23, 2018.

SECTION I SUPPLEMENT

Special Situations

Please visit www.firstaidteam.com/bonus/ to view this section.

- ▶ First Aid for the International Medical Graduate 2
- ▶ First Aid for the Osteopathic Medical Student 13
- ▶ First Aid for the Podiatric Medical Student 17
- ▶ First Aid for the Student Requiring Test Accommodations 20

SECTION I SUPPLEMENT

Special Situations

- ▶ First Aid for the International Medical Graduate 2
- ▶ First Aid for the Osteopathic Medical Student 13
- ▶ First Aid for the Podiatric Medical Student 17
- ▶ First Aid for the Student Requiring Test Accommodations 20

▶ *IMGs make up approximately 25% of the US physician population.*

▶ *More detailed information can be found in the ECFMG Information Booklet, available at www.ecfm.org/pubshome.html.*

▶ *Applicants may apply online for USMLE Step 1, Step 2 CK, or Step 2 CS at www.ecfm.org.*

▶ FIRST AID FOR THE INTERNATIONAL MEDICAL GRADUATE

“International medical graduate” (IMG) is the term used to describe any student or graduate of a non-US, non-Canadian, non-Puerto Rican medical school, regardless of whether he or she is a US citizen/resident or not.

IMG’s Steps to Licensure in the United States

To be eligible to take the USMLE Steps, you (the applicant) must be officially enrolled in a medical school located outside the United States and Canada that is listed in the World Directory of Medical Schools (WDOMS; www.wdoms.org) and meet the ECFMG eligibility requirements, both at the time you apply for examination and on your test day. In addition, your “Graduation Year” must be listed as “Current” at the time you apply and on your test day.

If you are an IMG, you must go through the following steps (not necessarily in this order) to apply for residency programs and become licensed to practice in the United States. You must complete these steps even if you are already a practicing physician and have completed a residency program in your own country.

- Pass USMLE Step 1, Step 2 CK, and Step 2 CS, as well as obtain a medical school diploma (not necessarily in this order). All three exams can be taken during medical school. If you have already graduated prior to taking any of the Steps, then you will need to verify your academic credentials (confirmation of enrollment and medical degree) prior to applying for any Step exam.
- You will be certified electronically by the Educational Commission for Foreign Medical Graduates (ECFMG) after above steps are successfully completed. You should receive your formal ECFMG certificate in the mail within the next 1–2 weeks. The ECFMG will not issue a certificate (even if all the USMLE scores are submitted) until it verifies your medical diploma with your medical school.
- You must have a valid ECFMG certificate before entering an accredited residency program in the United States, although you can begin the Electronic Residency Application Service (ERAS) application and interviews before you receive the certificate.
- Apply for residency positions in your fields of interest, either directly or through the ERAS and the National Residency Matching Program (NRMP), otherwise known as “the Match.” To be entered into the Match, you need to have passed all the examinations necessary for ECFMG certification (ie, Step 1, Step 2 CK, and Step 2 CS) by the rank order list deadline (usually in late February before the Match). If you do not pass these exams by the deadline, you will be withdrawn from the Match.

- If you are not a US citizen or green-card holder (permanent resident), you will need to obtain a visa that will allow you to enter and work in the United States after you have matched successfully.
- Sign up to receive the ECFMG and ERAS email newsletter to keep up to date with their most current policies and deadlines.
- If required by the state in which your residency program is located, obtain an educational/training/limited medical license. Your residency program may assist you with this application. Note that medical licensing is the prerogative of each individual state, not of the federal government, and that states vary with respect to their laws about licensing.
- Once you have the ECFMG certification, take the USMLE Step 3 during your residency, and then obtain a full medical license. Once you have a state-issued license, you are permitted to practice in federal institutions such as Veterans Affairs (VA) hospitals and Indian Health Service facilities in any state. This can open the door to “moonlighting” opportunities and possibilities for an H1B visa application if relevant. For details on individual state rules, write to the licensing board in the state in question or contact the Federation of State Medical Boards (FSMB). If you need to apply for an H1B visa for starting residency, you need to first take and pass the USMLE Step 3 exam, preferably before you Match. However, you will be able to apply for and take the USMLE Step 3 exam only after you graduate from medical school.
- Complete your residency and then take the appropriate specialty board exams if you wish to become board certified (eg, in internal medicine or surgery). If you already have a specialty certification in another country, some specialty boards may grant you six months’ or one year’s credit toward your total residency time.
- Currently, most residency programs are accepting applications through ERAS. For more information, see *First Aid for the Match* or contact:

ECFMG/ERAS Program

3624 Market Street
 Philadelphia, PA 19104-2685 USA
 (215) 386-5900
 Email: eras-support@ecfm.org
www.ecfm.org/eras

- For detailed information on the USMLE Steps, visit the USMLE website at <http://www.usmle.org>.

The USMLE and the IMG

The USMLE is a series of standardized exams that give IMGs and US medical graduates a level playing field. The passing marks for IMGs for Step 1, Step 2 CK, and Step 2 CS are determined by a statistical distribution that is based on the scores of US medical school students. For example, to pass Step 1, you will probably have to score higher than the bottom 8–10% of US and Canadian graduates.

► Keep informed by signing up for the ECFMG email newsletter at www.ecfm.org/resources.

► *IMGs have a maximum of six attempts to pass any USMLE Step, and must pass the USMLE Steps required for ECFMG certification within a seven-year period.*

Under USMLE program rules, a maximum of six attempts will be permitted to pass any USMLE Step or component exam. There is a limit of three attempts within a 12-month period for any of the USMLE Steps.

Timing of the USMLE

For an IMG, the timing of a complete application is critical. It is extremely important that you send in your application early if you are to obtain the maximum number of interviews. Complete all exam requirements by August of the year in which you wish to apply. Check the ECFMG website for deadlines to take and pass the various Step exams to be eligible for the NRMP Match.

IMG applicants must pass the USMLE Steps required for ECFMG certification (Step 1, Steps 2 CK and 2 CS) within a seven-year period. The USMLE program recommends, although not all jurisdictions impose, a seven-year limit for completion of the three-step USMLE program.

In terms of USMLE exam order, arguments can be made for taking the Step 1 or the Step 2 CK exam first. For example, you may consider taking the Step 2 CK exam first if you have just graduated from medical school and the clinical topics are still fresh in your mind. However, keep in mind that there is substantial overlap between Step 1 and Step 2 CK topics in areas such as pharmacology, pathophysiology, and biostatistics. You might therefore consider taking the Step 1 and Step 2 CK exams close together to take advantage of this overlap in your test preparation.

USMLE Step 1 and the IMG

Significance of the Test. Step 1 is one of the three exams required for the ECFMG certification. Since most US graduates apply to residency with their Step 1 scores only, it may be the only objective tool available with which to compare IMGs with US graduates.

Signing Up. We advise that you read the FAQ section on the ECFMG website carefully. Most of the services you will need to use involve either IWA or OASIS. If you have not yet completed medical school, follow these steps to sign up for Step 1:

- Apply and pay for an ECFMG/USMLE ID number on the ECFMG website.
- After receiving an email with your ID number, log in to IWA/OASIS, enter your details, and complete the “On-Line part of your USMLE Step 1 application.” Choose your test center location and 3-month eligibility period. Additional fees apply if you need to change your eligibility period.
- Pay the Step 1 fee plus any international test surcharges that may apply.
- Access and complete Form 186 (Certification of Identity Form) from IWA as part of the Application for ECFMG Certification.

- Follow the instructions on the form to notarize Form 186 using the online service NotaryCam.com. The fee for this service is included in the ECFMG application fee.
- Once notarized by NotaryCam.com and submitted, Form 186 will remain valid indefinitely. A valid, previously completed Form 186 will remain valid for five years from the date it was accepted.
- After receiving a confirmation email from the ECFMG, you may book an exam date and location on www.prometric.com.

Eligibility Period. A three-month period of your choice.

Fee. The fee for Step 1 is \$940 plus an international test delivery surcharge (if you choose a testing region other than the United States or Canada).

Statistics. In 2018–2019, 80% of IMG examinees passed Step 1 on their first attempt, compared with 96% of MD degree examinees from the United States and Canada.

Tips. Although few if any students feel totally prepared to take Step 1, IMGs in particular require serious study and preparation in order to reach their full potential on this exam. It is also imperative that IMGs do their best on Step 1, as a poor score on Step 1 is a distinct disadvantage in applying for most residencies. Remember that if you pass Step 1, you cannot retake it in an attempt to improve your score. Your goal should thus be to beat the mean, because you can then assert with confidence that you have done better than average for US students (see Table 1). Higher Step 1 scores will also

► *A higher Step 1 score will improve your chances of getting into a highly competitive specialty.*

TABLE 1. USMLE Step 1 Mean Score of Matched Applicants in 2018.

Specialty	US Graduates	US IMGs	Non-US IMGs
All specialties	233	222	234
Anesthesiology	232	231	240
Dermatology	249	—	238
Diagnostic radiology	240	239	241
Emergency medicine	233	232	229
Family medicine	220	211	220
General surgery	236	237	242
Internal medicine	233	225	236
Neurology	231	227	236
Obstetrics and gynecology	230	229	231
Pathology	233	226	230
Pediatrics	227	221	230
Physical medicine and rehabilitation	225	226	238
Psychiatry	226	214	222

Source: www.nrmp.org.

lend credibility to your residency application and help you get into highly competitive specialties such as radiology, orthopedics, and dermatology.

Commercial Review Courses. Do commercial review courses help improve your scores? Reports vary, and such courses can be expensive. For some students these programs can provide a more structured learning environment with professional support. However, review courses consume a significant chunk of time away from independent study. Many IMGs participate in review courses as they typically need higher scores to compete effectively with US and Canadian candidates for residency positions. (For more information on review courses, see Section IV in the book.)

USMLE Step 2 CK and the IMG

What Is the Step 2 CK? It is a computerized test of the clinical sciences consisting of up to 318 multiple-choice questions divided into eight blocks. Each block contains a maximum of 40 questions and needs to be completed within 60 minutes. It can be taken at Prometric centers in the United States and several other countries.

► *The areas tested on the Step 2 CK relate to the clerkships provided at US medical schools.*

Content. The Step 2 CK includes test items in the following content areas:

- Internal medicine
- Obstetrics and gynecology
- Pediatrics
- Preventive medicine
- Psychiatry
- Surgery
- Other areas relevant to the provision of care under supervision

Significance of the Test. The Step 2 CK is required for the ECFMG certificate. It reflects the level of clinical knowledge of the applicant. It tests clinical subjects, primarily internal medicine. Other areas tested are orthopedics, ENT, ophthalmology, safety science, epidemiology, professionalism, and ethics.

Eligibility. Students and graduates from medical schools that are listed in WDOMS and meet the ECFMG eligibility requirement to take the Step 2 CK. Students must have completed at least two years of medical school. This means that students must have completed the basic medical science component of the medical school curriculum by the beginning of the eligibility period selected.

Eligibility Period. A three-month period of your choice.

Fee. The fee for the Step 2 CK is \$940 plus an international test delivery surcharge (if you choose a testing region other than the United States or Canada).

Statistics. In 2017–2018, 83% of ECFMG candidates passed the Step 2 CK on their first attempt, compared with 97% of MD degree examinees from US and Canadian schools.

Tips. It's better to take the Step 2 CK after your internal medicine rotation because most of the questions on the exam give clinical scenarios and ask you to make medical diagnoses and clinical decisions. In addition, because this is a clinical sciences exam, cultural and geographic considerations play a greater role than is the case with Step 1. For example, if your medical education gave you ample exposure to malaria, brucellosis, and malnutrition but little to alcohol withdrawal, child abuse, and cholesterol screening, you must work to familiarize yourself with topics that are more heavily emphasized in US medicine. You must also have a basic understanding of the legal and social aspects of US medicine, because you will be asked questions about communicating with and advising patients.

► *Be familiar with topics that are heavily emphasized in US medicine, such as cholesterol screening.*

USMLE Step 2 CS and the IMG

What Is the Step 2 CS? The Step 2 CS is a test of clinical and communication skills administered as a one-day, eight-hour exam. It includes 12 encounters with standardized patients (15 minutes each, with 10 minutes to write a note after each encounter).

Content. The Step 2 CS tests the ability to communicate in English as well as interpersonal skills, data-gathering skills, the ability to perform a physical exam, and the ability to formulate a brief note, a differential diagnosis, and a list of diagnostic tests. The areas that are covered in the exam are as follows:

- Internal medicine
- Surgery
- Obstetrics and gynecology
- Pediatrics
- Psychiatry
- Family medicine

Unlike the USMLE Step 1, Step 2 CK, or Step 3, **there are no numerical grades for the Step 2 CS**—it's simply either a “pass” or a “fail.” To pass, a candidate must attain a passing performance in **each** of the following three components:

- Integrated Clinical Encounter (ICE): includes Data Gathering, Physical Exam, and the electronic Patient Note
- Spoken English Proficiency (SEP)
- Communication and Interpersonal Skills (CIS)

According to the NBME, the most commonly failed component for IMGs is the CIS.

► *The Step 2 CS is graded as pass/fail.*

Significance of the Test. The Step 2 CS assesses spoken English language proficiency and is required for the ECFMG certificate. The Test of English as a Foreign Language (TOEFL) is no longer required.

Eligibility. Students must have completed at least two years of medical school in order to take the test. That means students must have completed the basic medical science component of the medical school curriculum at the time they apply for the exam.

Fee. The fee for the Step 2 CS is \$1580.

Statistics. In 2017–2018, 75% of ECFMG candidates passed the Step 2 CS on their first attempt, compared with 95% of MD degree examinees from US and Canadian schools.

► Try to take the Step 2 CS the year before you plan to Match.

Scheduling. You must schedule the Step 2 CS within **four months** of the date indicated on your notification of registration. You must take the exam within 12 months of the date indicated on your notification of registration. It is generally advisable to take the Step 2 CS as soon as possible in the year before your Match, as often the results either come in late or arrive too late to allow you to retake the test and pass it before the Match.

Test Site Locations. The Step 2 CS is currently administered at the following five locations:

- Philadelphia, PA
- Atlanta, GA
- Los Angeles, CA
- Chicago, IL
- Houston, TX

For more information about the Step 2 CS exam, please refer to *First Aid for the Step 2 CS*.

USMLE Step 3 and the IMG

What Is the USMLE Step 3? It is a two-day computerized test in clinical medicine consisting of 413 multiple-choice questions and 13 computer-based case simulations (CCS). The exam aims to test your knowledge and its application to patient care and clinical decision making (ie, this exam tests if you can safely practice medicine independently and without supervision). Please go to the USMLE website to learn more about recent changes to the exam.

► Complete the Step 3 exam before you apply for an H1B visa.

Significance of the Test. Taking Step 3 before residency is critical for IMGs seeking an H1B visa and is also a bonus that can be added to the residency application. Step 3 is also required to obtain a full medical license in the United States and can be taken during residency for this purpose.

Fee. The fee for Step 3 is \$895.

Eligibility. Examinees are no longer required to apply to the Step 3 exam under the eligibility requirements of a specific medical licensing authority. Those wishing to sit for the Step 3 exam, independent of the place of residence, must meet the following requirements:

- Have completed an MD or DO degree from an LCME- or AOA-accredited US or Canadian medical school, or from a medical school outside the US and Canada listed in the World Directory of Medical Schools.
- Have taken and passed the Step 1, Step 2 CK, and Step 2 CS exams.
- If an IMG, be certified by the ECFMG.

The Step 3 exam is not available outside the United States. Applications can be found online at www.fsmb.org and must be submitted to the FSMB.

Statistics. In 2018, 90% of IMG candidates passed the Step 3 on their first attempt, compared with 98% of MD degree examinees from US and Canadian schools.

Residencies and the IMG

In the Match, the number of US-citizen IMG applications has grown over the past few years, while the percentage accepted has remained constant (see Table 2). More information about residency programs can be obtained at www.ama-assn.org.

The Match and the IMG

Given the growing number of IMG candidates with strong applications, you should bear in mind that good USMLE scores are not the only way to gain a competitive edge. However, USMLE Step 1 and Step 2 CK scores continue to be used as the initial screening mechanism when candidates are being considered for interviews.

TABLE 2. IMGs in the Match.

Applicants	2016	2017	2018	2019
US-citizen IMGs	5,323	5,069	5,075	5,080
% US-citizen IMGs accepted	53.9	54.8	57.1	59
Non-US-citizen IMGs	7,460	7,284	7,067	6,869
% non-US-citizen IMGs accepted	50.5	52.4	56.1	58.6
US seniors (non-IMGs)	18,187	18,539	18,818	18,925
% US seniors accepted	93.8	94.3	94.3	93.9
DO graduates		3,590	4,617	6,001
% DO graduates accepted		81.7	81.7	84.6


Source: www.nrmp.org.

Based on accumulated IMG Match experiences over recent years, here are a few pointers to help IMGs maximize their chances for a residency interview:

► All US hospitals allow externship only when the applicant is actively enrolled in a medical school, so plan ahead.

- **Apply early.** Programs offer a limited number of interviews and often select candidates on a first-come, first-served basis. Because of this, you should aim to complete the entire process of applying for the ERAS token, registering with the Association of American Medical Colleges (AAMC), mailing necessary documents to ERAS, and completing the ERAS application by mid-September (see Figure 1). Community programs usually send out interview offers earlier than do university and university-affiliated programs.
- **US clinical experience helps.** Externships and observerships in a US hospital setting have emerged as an important credential on an IMG application. Externships are like short-term medical school internships and offer hands-on clinical experience. Observerships, also called “shadowing,” involve following a physician and observing how he or she manages patients. Some programs require students to have participated in an externship or observership before applying. It is best to gain such an experience before or at the time you apply to various programs so that you can mention it on your

FIGURE 1. IMG Timeline for Application.



ERAS application. If such an experience or opportunity comes up after you apply, be sure to inform the programs accordingly.

- **Clinical research helps.** University programs are attracted to candidates who show a strong interest in clinical research and academics. They may even relax their application criteria for individuals with unique backgrounds and strong research experience. Publications in well-known journals are an added bonus.
- **Time the Step 2 CS well.** ECFMG has published the new Step 2 CS score-reporting schedule for 2019–2020 at <http://www.ecfmg.org>. Most program directors would like to see a passing score on the Step 1, Step 2 CK, and Step 2 CS exams before they rank an IMG on their rank order list in mid-February. There have been many instances in which candidates have lost a potential Match—either because of delayed CS results or because they have been unable to retake the exam on time following a failure. It is difficult to predict a result on the Step 2 CS, since the grading process is not very transparent. Therefore, it is advisable to take the Step 2 CS as early as possible in the application year.
- **US letters of recommendation help.** Letters of recommendation from clinicians practicing in the United States carry more weight than recommendations from home countries.
- **Step up the Step 3.** If H1B visa sponsorship is desired, aim to have Step 3 results by January of the Match year. In addition to the visa advantage you will gain, an early and good Step 3 score may benefit IMGs who have been away from clinical medicine for a while as well as those who have low scores on Step 1 and the Step 2 CK. Note that the Step 3 can be taken only after medical school graduation.
- **Verify medical credentials in a timely manner.** Do not overlook the medical school credential verification process. The ECFMG certificate arrives only after credentials have been verified and after you have passed Step 1, the Step 2 CK, and the Step 2 CS, so you should keep track of the process and check their application status online using IWA/OASIS.
- **Don't count on a pre-Match.** Programs participating in NRMP Match can no longer offer a pre-Match.

► *A good score on the Step 3 may help offset poorer scores on the Step 1 or 2 CK exams.*

What if You Do Not Match?

For applicants who do not Match into a residency program, there's SOAP (Supplemental Offer and Acceptance Program). Under SOAP, unmatched applicants will have access to the list of unfilled programs at noon Eastern time on the Monday of Match week. The unfilled programs electing to participate in SOAP will offer positions to unmatched applicants through the Registration, Ranking, and Results (R3) system. A series of "rounds" will begin at noon Eastern time on Wednesday of Match week until 5:00 PM Eastern time on Friday of Match week. Detailed information about SOAP can be found at the NRMP website at <http://www.nrmp.org>.

Resources for the IMG

- **Educational Commission for Foreign Medical Graduates (ECFMG)**

3624 Market Street
 Philadelphia, PA 19104-2685
 (215) 386-5900
 Fax: (215) 386-9196
 Email: info@ecfm.org
www.ecfm.org

The ECFMG telephone number is answered only between 9:00 AM–5:00 PM Monday through Friday EST. The ECFMG often takes a long time to answer the phone, which is frequently busy at peak times of the year, and then gives you a long voice-mail message—so it is better to email early than to rely on a last-minute phone call. When contacting the ECFMG by email, include your USMLE/ECFMG Identification Number and use the email address that you registered with the ECFMG. Do not contact the NBME, as all IMG exam matters are conducted by the ECFMG. The ECFMG also publishes an information booklet on ECFMG certification and the USMLE program, which gives details on the dates and locations of forthcoming Step tests for IMGs together with application forms. The *Information Booklet* is available to view and download on the ECFMG’s website at www.ecfm.org, where they also have a complete list of fees for certification posted (see Table 3).

TABLE 3. Estimated Costs for IMGs (as of 2019).

Exams and Services	Fee(s)
USMLE Step 1	\$940 + international surcharge (eg, \$195 in all European countries offering the exam)
USMLE Step 2 CK	\$940 + international surcharge (eg, \$220 in all European countries offering the exam)
USMLE Step 2 CS	\$1580
USMLE Step 3	\$895
ERAS	\$130 registration fee (ECFMG token fee) \$80 USMLE transcript assessment \$99 for programs 1–10 \$15 each for programs 11–20 \$19 each for programs 21–30 \$26 each for programs 31+
NRMP	\$85 registration fee (for ranking 20 programs) \$30 per additional program ranked \$35 per partner (couples match only) \$50 late registration fee (sign up before November 30 to avoid paying this fee)
J-1 visa application fee	\$160 visa application fee \$340 annual ECFMG application fee \$220 payable to Homeland Security (SEVIS fee)

- **Federation of State Medical Boards (FSMB)**

400 Fuller Wisser Road, Suite 300
Euless, TX 76039-3856
(817) 868-4041
Email: usmle@fsmb.org
www.fsmb.org

The FSMB has a number of publications available, including free policy documents. All of these documents are available to view and download for free on the FSMB's website at www.fsmb.org. For Step 3 inquiries, the telephone number is (817) 868-4041.

The AMA has dedicated a portion of its website to information on IMG demographics, residencies, immigration, and the like. This information can be found at www.ama-assn.org.

► FIRST AID FOR THE OSTEOPATHIC MEDICAL STUDENT

What Is the COMLEX-USA Level 1?

The National Board of Osteopathic Medical Examiners (NBOME) administers the Comprehensive Osteopathic Medical Licensing Examination, or COMLEX-USA. Like the USMLE, the COMLEX-USA is administered over three levels.

The COMLEX-USA series assesses osteopathic medical knowledge and clinical skills using clinical presentations and physician tasks. A description of the COMLEX-USA Written Examination Blueprints for each level, which outline the various clinical presentations and physician tasks that examinees will encounter, is given on the NBOME website. Another stated goal of the COMLEX-USA Level 1 is to create a more primary care-oriented exam that integrates osteopathic principles into clinical situations.

To be eligible to take the COMLEX-USA Level 1, you must be on track to satisfactorily complete your first two years in an AOA-accredited medical school. The office of the dean at each school informs the NBOME that the student will complete the first two years of medical school and is in good standing. At this point, the NBOME sends out an email with detailed instructions on how to register for the exam.

For all three levels of the COMLEX-USA, raw scores are converted to a percentile score and a score ranging from 5 to 800. For Levels 1 and 2, a score of 400 is required to pass; for Level 3, a score of 350 is needed. COMLEX-USA scores are posted at the NBOME website 4–6 weeks after the test and usually mailed within 8 weeks after the test. The mean score is always 500.

If you pass a COMLEX-USA examination, you are not allowed to retake it to improve your grade. Currently, if you fail, there is no specific limit to the number of times you can retake it in order to pass. However, a student may not take the exam more than four times in one year. Levels 2 and 3 exams must be passed in sequential order within seven years of passing Level 1.

Note that candidates taking COMLEX-USA examinations will be limited to a total of six attempts for each examination.

What Is the Structure of the COMLEX-USA Level 1?

The COMLEX-USA Level 1 is a computer-based examination consisting of 400 questions over an eight-hour period in a single day (nine hours counting breaks). Most of the questions are in one-best-answer format, but a small number are matching-type questions. Some one-best-answer questions are bundled together around a common question stem that usually takes the form of a clinical scenario. Every section of the COMLEX-USA Level 1 ends with either matching questions, multiple questions around a single stem, or both. New question formats may gradually be introduced, but candidates will be notified if this occurs. Multimedia questions are also included on the exam.

Questions are grouped into eight subsections of 50 questions each in a manner similar to that of the USMLE. The individual subsections are not timed, but the exam is divided into two blocks consisting of four subsections. Each subsection consists of 200 questions to be completed within four hours. Reviewing and changing answers may be done only in the current subsection. A “review page” is presented for each subsection in order to advise test takers of questions completed, questions marked for further review, and incomplete questions for which no answer has been given.

Breaks are even more structured with COMLEX-USA than they are with the USMLE. Students are allowed to take an optional 10-minute break at the end of the second and sixth subsections. After subsection 4, students are given a 40-minute lunch break. These are the only times a student is permitted a break. Any unused break time will not be added to the time allotted for taking the examination. More information about the computer-based COMLEX-USA examinations can be obtained from www.nbome.org.

What Is the Difference Between the USMLE and the COMLEX-USA?

According to the NBOME, the COMLEX-USA Level 1 focuses broadly on the following categories, with osteopathic principles and practices integrated into each section:

- Health promotion and disease prevention
- The history and physical
- Diagnostic technologies

- Management
- Scientific understanding of mechanisms
- Health care delivery

Although the COMLEX-USA and the USMLE are similar in scope, content, and emphasis, some differences are worth noting. For example, the interface is different; you cannot search for lab values. Instead, lab values and reference ranges (where appropriate) are included directly in the clinical vignette or test question. Fewer details are given about a patient's condition, so a savvy student needs to know how to differentiate between similar pathologies. Also, age, gender, and race are key factors for diagnosis on the COMLEX-USA. Images or videos are embedded in the question stem and the examinee has to click an attachment button to see the image. If you don't read the question carefully, the attachment buttons are very easy to miss. A standard calculator feature is embedded in the examination interface.

COMLEX-USA Level 1 tests osteopathic principles in addition to basic science materials but does not emphasize lab techniques. Although both exams often require that you apply and integrate knowledge over several areas of basic science to answer a given question, many students who took both tests reported that the questions differed somewhat in style. Students reported, for example, that USMLE questions generally required that the test taker reason and draw from the information given (often a two-step process), whereas those on the COMLEX-USA exam tended to be more straightforward and that multiple different questions are asked pertaining to one question stem.

COMLEX-USA test takers can expect to have only a few questions on biochemistry, molecular biology, or lab technique. On the other hand, microbiology is very heavily tested by clinical presentation and by lab identification. The COMLEX-USA exam also focuses more on disease management, specific legal principles (eg, Tarasoff case and the Emergency Treatment Act) and more detailed ethical principles (eg, *res ipsa loquitur*) than the USMLE Step 1. Another main difference is that the COMLEX-USA exam stresses osteopathic manipulative medicine. Therefore, question banks specific to the USMLE will not be adequate, and supplementation with a question bank specific to the COMLEX-USA is highly recommended. The most commonly used are COMBANK or COMQUEST.

Students also commented that the COMLEX-USA utilized “buzzwords,” although limited in their use (eg, “rose spots” in typhoid fever), whereas the USMLE avoided buzzwords in favor of descriptions of clinical findings or symptoms (eg, rose-colored papules on the abdomen rather than rose spots). Finally, USMLE appeared to have more photographs than did the COMLEX-USA. In general, the overall impression was that the USMLE was a more “thought-provoking” exam, while the COMLEX-USA was more of a “knowledge-based” exam.

► *The test interface for the COMLEX-USA Level 1 is not the same as the USMLE Step 1 interface.*

Who Should Take Both the USMLE and the COMLEX-USA?

Aside from facing the COMLEX-USA Level 1, you must decide if you will also take the USMLE Step 1. We recommend that you consider taking both the USMLE and the COMLEX-USA under the following circumstances:

► If you're not sure whether you need to take either the COMLEX-USA Level 1 or the USMLE Step 1, consider taking both to keep your Match options open.

- **If you are applying to allopathic residencies.** Although there is growing acceptance of COMLEX-USA certification on the part of allopathic residencies, some allopathic programs prefer or even require passage of the USMLE Step 1. These include many academic programs, programs in competitive specialties (eg, orthopedics, ophthalmology, or dermatology), and programs in competitive geographic areas (eg, Vermont, Utah, and California). Fourth-year osteopathic medical students who have already Matched may be a good source of information about which programs and specialties look for USMLE scores. It is also a good idea to contact program directors at the institutions you are interested in to ask about their policy regarding the COMLEX-USA versus the USMLE.
- **If you are unsure about your postgraduate training plans.** Successful passage of both the COMLEX-USA Level 1 and the USMLE Step 1 is certain to provide you with the greatest possible range of options when you are applying for internship and residency training.

In addition, the COMLEX-USA Level 1 has in recent years placed increasing emphasis on questions related to primary care medicine and prevention. Having a strong background in family or primary care medicine can help test takers when they face questions on prevention.

How Do I Prepare for the COMLEX-USA Level 1?

Student experience suggests that you should start studying for the COMLEX-USA four to six months before the test is given, as an early start will allow you to spend up to a month on each subject. The recommendations made in Section I regarding study and testing methods, strategies, and resources, as well as the books suggested in Section IV for the USMLE Step 1, hold true for the COMLEX-USA as well.

Another important source of information is in the *Examination Guidelines and Sample Exam*, a booklet that discusses the breakdown of each subject while also providing sample questions and corresponding answers. Many students, however, felt that this breakdown provided only a general guideline and was not representative of the level of difficulty of the actual COMLEX-USA. The sample questions did not provide examples of clinical vignettes, which made up approximately 25% of the exam. You will receive this publication with registration materials for the COMLEX-USA Level 1, but you can also receive a copy and additional information by writing:

NBOME

8765 W. Higgins Road, Suite 200
Chicago, IL 60631-4174
(773) 714-0622
www.nbome.org

The NBOME developed the Comprehensive Osteopathic Medical Self-Assessment Examination (COMSAE) series to fill the need for self-assessment on the part of osteopathic medical students. Many students take the COMSAE exam before the COMLEX-USA in addition to using test-bank questions and board review books. Students can purchase a copy of this exam at www.nbome.org/comsae.asp.

In recent years, students have reported an emphasis in certain areas. For example:

- There was an increased emphasis on upper limb anatomy/brachial plexus.
- Specific topics were repeatedly tested on the exam. These included cardiovascular physiology and pathology, acid-base physiology, diabetes, benign prostatic hyperplasia, sexually transmitted diseases, measles, and rubella. Thyroid and adrenal function, neurology (head injury), specific drug treatments for bacterial infection, migraines/cluster headaches, and drug mechanisms also received heavy emphasis.
- Behavioral science questions were based on psychiatry.
- High-yield osteopathic manipulative technique (OMT) topics included an emphasis on the sympathetic and parasympathetic innervations of viscera and nerve roots, rib mechanics/diagnosis, and basic craniosacral theory. Students who spend time reviewing basic anatomy, studying nerve and dermatome innervations, and understanding how to perform basic OMT techniques (eg, muscle energy or counterstrain) can improve their scores.

The COMLEX-USA Level 1 also includes multimedia-based questions. Such questions test the student's ability to perform a good physical exam and to elicit various physical diagnostic signs (eg, Murphy sign).

▶ *You must know the Chapman reflex points and the obscure names of physical exam signs.*

▶ *COMLEX is heavy on "bugs and drugs."*

▶ FIRST AID FOR THE PODIATRIC MEDICAL STUDENT

The National Board of Podiatric Medical Examiners (NBPME) offers the American Podiatric Medical Licensing Examinations (APMLE), which are designed to assess whether a candidate possesses the knowledge required to practice as a minimally competent entry-level podiatric surgeon. The APMLE is used as part of the licensing process governing the practice of podiatric medicine and surgery. The APMLE is recognized by all 50 states and the District of Columbia, the US Army, the US Navy, and the Canadian provinces of Alberta, British Columbia, and Ontario. Individual states use the examination scores differently; therefore, doctor of podiatric medicine (DPM) candidates should refer to the *NBPME Part I and Part II Information Bulletin 2019*.

► Areas tested on the NBPME Part I:

- General anatomy
- Lower extremity anatomy
- Biochemistry
- Physiology
- Medical microbiology & immunology
- Pathology
- Pharmacology

The APMLE Part I is generally taken after the completion of the second year of podiatric medical education. Unlike the USMLE Step 1, there is no behavioral science section, nor is biomechanics tested. The exam samples seven basic science disciplines: general anatomy (13%); lower extremity anatomy (25%); biochemistry (10%); physiology (13%); microbiology and immunology (13%); pathology (13%); and pharmacology (13%). A detailed outline of topics and subtopics covered on the exam can be found in the *Candidate Information Bulletin Part I Examination*, available at www.apmle.org.

Your APMLE Appointment

Applicants have to register for the exam online at www.prometric.com/NBPME. Once registration is completed, you will receive an Authorization to Test (ATT) email notification that allows you to schedule your exam online. This should be done promptly to secure the testing location and exam date of your choice. The exam will be offered at an independent Prometric testing facility. Test centers within a 50-mile radius of a podiatric medicine school specifically reserve a number of seats on each APMLE Part I exam date. You may take the exam at any Prometric site regardless of which school you attend. Specific instructions about exam dates and registration deadlines can be found in the *Candidate Information Bulletin*.

Exam Format

The APMLE Part I is a written exam consisting of 205 questions. The test consists exclusively of one-best-answer multiple choice questions with four options per question. A review screen showing all answered, unanswered, and marked questions will be available at the end. Students are encouraged to mark questions and return to these for review at the end of the exam if time allows. Examinees have four hours in which to complete the exam and are given scratch paper that must be turned in at the end of the exam. Some questions on the exam will be “trial questions.” These questions are evaluated as future board questions but are not counted in your score.

Interpreting Your Score

Exam results are emailed to examinees approximately four weeks after the exam date, and are also available online via the Prometric dashboard. APMLE scores are reported as pass/fail, with a scaled score of at least 75 needed to pass. Historically, 85% of first-time test takers pass the APMLE Part I. Failing candidates receive a report with a score between 55 and 74 in addition to diagnostic messages intended to help identify strengths or weaknesses in specific content areas. If you fail the APMLE Part I, you must retake the entire examination at a later date. There is no limit to the number of times you can retake the exam.

Preparation for the APMLE Part I

Begin studying for the APMLE Part I at least three months prior to the test date. The suggestions made in Section I regarding study and testing methods for the USMLE Step 1 can be applied to the APMLE as well. This book should, however, be used as a supplement and not as the sole source of information. Neither you nor your school or future residency will ever see your actual passing numerical score. Competing with colleagues should not be an issue, and study groups are beneficial to many.

A study method that helps many students is to copy the outline of the material to be tested from the *Candidate Information Bulletin*. Check off each topic during your study, because doing so will ensure that you have engaged each topic. If you are pressed for time, prioritize subjects on the basis of their weight on the exam. A full 25% of the APMLE Part I focuses on lower extremity anatomy. In this area, students should rely on the notes and material that they received from their class. Remember, lower extremity anatomy is the podiatric physician's specialty—so everything about it is important. Do not forget to study osteology. Keep your old tests and look through old lower extremity class exams, since each of the podiatric colleges submits questions from its faculty. This strategy will give you an understanding of the types of questions that may be asked. On the APMLE Part I, you will see some of the same classic lower extremity anatomy questions you were tested on in school.

The APMLE, like the USMLE, requires that you apply and integrate knowledge over several areas of basic science in order to answer exam questions. Students report that many questions emphasize clinical presentations; however, the facts in this book are very useful in helping students recall the various diseases and organisms. DPM candidates should expand on the high-yield pharmacology section and study antifungal drugs and treatments for *Pseudomonas*, methicillin-resistant *S aureus*, candidiasis, and erythrasma. The high-yield section focusing on pathology is very useful; however, additional emphasis on diabetes mellitus and all its secondary manifestations, particularly peripheral neuropathy, should not be overlooked. Students should also focus on renal physiology and drug elimination, the biochemistry of gout, and neurophysiology, all of which have been noted to be important topics on the APMLE Part I exam.

A sample set of questions is found on the APMLE website www.apmle.org. These samples are somewhat similar in difficulty to actual board questions. If you have any questions regarding registration, fees, test centers, authorization forms, or score reports, please contact your college registrar or:

Prometric

877-302-8952

Email: nbpmeinquiry@prometric.com

www.prometric.com

► *Know the anatomy of the lower extremity!*

► FIRST AID FOR THE STUDENT REQUIRING TEST ACCOMMODATIONS

The USMLE provides accommodations for students with documented disabilities. The basis for such accommodations is the Americans with Disabilities Act (ADA) of 1990. The ADA defines a disability as “a significant limitation in one or more major life activities.” This includes both “observable/physical” disabilities (eg, blindness, hearing loss, narcolepsy) and “hidden/mental disabilities” (eg, attention-deficit hyperactivity disorder, chronic fatigue syndrome, learning disabilities).

► *US students seeking ADA-compliant accommodations must contact the NBME directly; IMGs, contact the ECFMG.*

To provide appropriate support, the administrators of the USMLE must be informed of both the nature and the severity of an examinee’s disability. Such documentation is required for an examinee to receive testing accommodations. Accommodations include extra time on tests, low-stimulation environments, extra or extended breaks, and zoom text.

Who Can Apply for Accommodations?

Students or graduates of a school in the United States or Canada that is accredited by the Liaison Committee on Medical Education (LCME) or the AOA may apply for test accommodations directly from the NBME. Requests are granted only if they meet the ADA definition of a disability. If you are a disabled student or a disabled graduate of a foreign medical school, you must contact the ECFMG (see the following page).

Who Is Not Eligible for Accommodations?

Individuals who do not meet the ADA definition of disabled are not eligible for test accommodations. Difficulties not eligible for test accommodations include test anxiety, slow reading without an identified underlying cognitive deficit, English as a second language, and learning difficulties that have not been diagnosed as a medically recognized disability.

Understanding the Need for Documentation

Although most learning-disabled medical students are all too familiar with the often exhausting process of providing documentation of their disability, you should realize that **applying for USMLE accommodation is different from these previous experiences**. This is because the NBME determines whether an individual is disabled solely on the basis of the guidelines set by the ADA. **Previous accommodation does not in itself justify provision of an accommodation for the USMLE**, so be sure to review the NBME guidelines carefully.

Getting the Information

The first step in applying for USMLE special accommodations is to contact the NBME and obtain a guidelines and questionnaire booklet. For the Step 1, Step 2 CK, and Step 2 CS exams, this can be obtained by calling or writing to:

Disability Services

National Board of Medical Examiners
3750 Market Street
Philadelphia, PA 19104-3102
(215) 590-9509
Email: disabilityservices@nbme.org
www.usmle.org/test-accommodations

Internet access to this information is also available at www.nbme.org. This information is also relevant for IMGs, since the information is the same as that sent by the ECFMG.

Foreign graduates should contact the ECFMG to obtain information on special accommodations by calling or writing to:

ECFMG

3624 Market Street
Philadelphia, PA 19104-2685
(215) 386-5900
www.ecfmg.org

When you get this information, take some time to read it carefully. The guidelines are clear and explicit about what you need to do to obtain accommodations.

SECTION II

High-Yield General Principles

“There comes a time when for every addition of knowledge you forget something that you knew before. It is of the highest importance, therefore, not to have useless facts elbowing out the useful ones.”

—Sir Arthur Conan Doyle, *A Study in Scarlet*

“Never regard study as a duty, but as the enviable opportunity to learn.”

—Albert Einstein

“Live as if you were to die tomorrow. Learn as if you were to live forever.”

—Gandhi

▶ How to Use the Database	30
▶ Biochemistry	33
▶ Immunology	95
▶ Microbiology	123
▶ Pathology	205
▶ Pharmacology	229
▶ Public Health Sciences	255

▶ HOW TO USE THE DATABASE

The 2020 edition of *First Aid for the USMLE Step 1* contains a revised and expanded database of basic science material that students, student authors, and faculty authors have identified as high yield for board review. The information is presented in a partially organ-based format. Hence, Section II is devoted to the foundational principles of biochemistry, microbiology, immunology, basic pathology, basic pharmacology, and public health sciences. Section III focuses on organ systems, with subsections covering the embryology, anatomy and histology, physiology, clinical pathology, and clinical pharmacology relevant to each. Each subsection is then divided into smaller topic areas containing related facts. Individual facts are generally presented in a three-column format, with the **Title** of the fact in the first column, the **Description** of the fact in the second column, and the **Mnemonic** or **Special Note** in the third column. Some facts do not have a mnemonic and are presented in a two-column format. Others are presented in list or tabular form in order to emphasize key associations.




The database structure used in Sections II and III is useful for reviewing material already learned. These sections are **not** ideal for learning complex or highly conceptual material for the first time.

The database of high-yield facts is not comprehensive. Use it to complement your core study material and not as your primary study source. The facts and notes have been condensed and edited to emphasize the essential material, and as a result, each entry is “incomplete” and arguably “over-simplified.” Often, the more you research a topic, the more complex it becomes, with certain topics resisting simplification. Work with the material, add your own notes and mnemonics, and recognize that not all memory techniques work for all students.

We update the database of high-yield facts annually to keep current with new trends in boards emphasis, including clinical relevance. However, we must note that inevitably many other high-yield topics are not yet included in our database.

We actively encourage medical students and faculty to submit high-yield topics, well-written entries, diagrams, clinical images, and useful mnemonics so that we may enhance the database for future students. We also solicit recommendations of alternate tools for study that may be useful in preparing for the examination, such as charts, flash cards, apps, and online resources (see How to Contribute, p. xix).

Image Acknowledgments

All images and diagrams marked with  are © USMLE-Rx.com (MedIQ Learning, LLC) and reproduced here by special permission. All images marked with  are © Dr. Richard P. Usatine, author of *The Color Atlas of Family Medicine*, *The Color Atlas of Internal Medicine*, and *The Color Atlas of Pediatrics*, and are reproduced here by special permission (www.usatinemedia.com). Images and diagrams marked with  are adapted or reproduced with permission of other sources as listed on page 727. Images and diagrams with no acknowledgment are part of this book.

Disclaimer

The entries in this section reflect student opinions of what is high yield. Because of the diverse sources of material, no attempt has been made to trace or reference the origins of entries individually. We have regarded mnemonics as essentially in the public domain. Errata will gladly be corrected if brought to the attention of the authors, either through our online errata submission form at www.firstaidteam.com or directly by email to firstaid@scholarrx.com.

HIGH-YIELD PRINCIPLES IN

Biochemistry

“Biochemistry is the study of carbon compounds that crawl.”

—Mike Adams

“We think we have found the basic mechanism by which life comes from life.”

—Francis H. C. Crick

“The biochemistry and biophysics are the notes required for life; they conspire, collectively, to generate the real unit of life, the organism.”


—Ursula Goodenough

This high-yield material includes molecular biology, genetics, cell biology, and principles of metabolism (especially vitamins, cofactors, minerals, and single-enzyme-deficiency diseases). When studying metabolic pathways, emphasize important regulatory steps and enzyme deficiencies that result in disease, as well as reactions targeted by pharmacologic interventions. For example, understanding the defect in Lesch-Nyhan syndrome and its clinical consequences is higher yield than memorizing every intermediate in the purine salvage pathway.

Do not spend time learning details of organic chemistry, mechanisms, or physical chemistry. Detailed chemical structures are infrequently tested; however, many structures have been included here to help students learn reactions and the important enzymes involved. Familiarity with the biochemical techniques that have medical relevance—such as ELISA, immunoelectrophoresis, Southern blotting, and PCR—is useful. Review the related biochemistry when studying pharmacology or genetic diseases as a way to reinforce and integrate the material.

▶ Molecular	34
▶ Cellular	46
▶ Laboratory Techniques	52
▶ Genetics	56
▶ Nutrition	65
▶ Metabolism	72

► BIOCHEMISTRY—MOLECULAR


Chromatin structure

DNA exists in the condensed, chromatin form to fit into the nucleus. DNA loops twice around a histone octamer to form a nucleosome (“**beads on a string**”). H1 binds to the nucleosome and to “linker DNA,” thereby stabilizing the chromatin fiber.

Phosphate groups give DNA a \ominus charge. Lysine and arginine give histones a \oplus charge.

In mitosis, DNA condenses to form chromosomes. DNA and histone synthesis occurs during S phase.

Mitochondria have their own DNA, which is circular and does not utilize histones.

Heterochromatin

Condensed, appears darker on EM (labeled H in **A**; Nu, nucleolus). Sterically inaccessible, thus transcriptionally inactive. \uparrow methylation, \downarrow acetylation.

HeteroChromatin = **H**ighly **C**ondensed.

Barr bodies (inactive X chromosomes) may be visible on the periphery of nucleus.

Euchromatin

Less condensed, appears lighter on EM (labeled E in **A**). Transcriptionally active, sterically accessible.

Eu = true, “truly transcribed.”

Euchromatin is **E**xpressed.

DNA methylation

Changes the expression of a DNA segment without changing the sequence. Involved with aging, carcinogenesis, genomic imprinting, transposable element repression, and inactivation of the X chromosome.

DNA is methylated in imprinting.

Methylation within gene promoter (CpG islands) typically represses (silences) gene transcription. CpG **M**ethylation **M**akes DNA **M**ute.

Histone methylation

Usually causes reversible transcriptional suppression, but can also cause activation depending on location of methyl groups.

Histone **M**ethylation **M**ostly **M**akes DNA **M**ute.

Histone acetylation

Removal of histone’s \oplus charge \rightarrow relaxed DNA coiling \rightarrow \uparrow transcription.

Histone **A**cetylation makes DNA **A**ctive.

Histone deacetylation

Removal of acetyl groups \rightarrow tightened DNA coiling \rightarrow \downarrow transcription.

Nucleotides

Nucleo**S**ide = base + (deoxy)ribose (**S**ugar).

Nucleo**T**ide = base + (deoxy)ribose + phospho**T**e;
linked by 3'-5' phosphodiester bond.

5' end of incoming nucleotide bears the triphosphate (energy source for the bond).
Triphosphate bond is target of 3' hydroxyl attack.

PURines (**A,G**)—2 rings.

PYrimidines (**C,U,T**)—1 ring.

Deamination reactions:

Cytosine → uracil

Adenine → hypoxanthine

Guanine → xanthine

5-methylcytosine → thymine

Uracil found in RNA; thymine in DNA.

Methylation of uracil makes thymine.

PURe **A**s **G**old.

CUT the **PY** (pie).

Thymine has a **meth**yl.

C-G bond (3 H bonds) stronger than A-T bond (2 H bonds). ↑ C-G content → ↑ melting temperature of DNA. "**C-G** bonds are like **Crazy G**lue."


Amino acids necessary for **pur**ine synthesis (cats

purr until they **GAG**):

Glycine


Aspartate

Glutamine



De novo pyrimidine and purine synthesis

Various immunosuppressive, antineoplastic, and antibiotic drugs function by interfering with nucleotide synthesis:



Pyrimidine synthesis:

- **Leflunomide:** inhibits dihydroorotate dehydrogenase
- **5-fluorouracil (5-FU)** and its prodrug **capecitabine:** form 5-F-dUMP, which inhibits thymidylate synthase (↓ dTMP)

Purine synthesis:

- **6-mercaptopurine (6-MP)** and its prodrug **azathioprine:** inhibit de novo purine synthesis
- **Mycophenolate** and **ribavirin:** inhibit inosine monophosphate dehydrogenase

Purine and pyrimidine synthesis:


- **Hydroxyurea:** inhibits ribonucleotide reductase
- **Methotrexate (MTX), trimethoprim (TMP),** and **pyrimethamine:** inhibit dihydrofolate reductase (↓ deoxythymidine monophosphate [dTMP]) in humans, bacteria, and protozoa, respectively

CPS1 = mItochondria (urea cycle)

CPS2 = cyTWOsol

Ex

Purine salvage deficiencies



ADA, adenosine deaminase; APRT, adenine phosphoribosyltransferase; HGPRT, hypoxanthine guanine phosphoribosyltransferase; XO, xanthine oxidase.



Adenosine deaminase deficiency

ADA is required for degradation of adenosine and deoxyadenosine. \downarrow ADA \rightarrow \uparrow dATP
 \rightarrow \downarrow ribonucleotide reductase activity
 \rightarrow lymphotoxicity.

One of the major causes of autosomal recessive SCID.

Lesch-Nyhan syndrome

Defective purine salvage due to absent **HGPRT**, which converts hypoxanthine to IMP and guanine to GMP. Results in excess uric acid production and de novo purine synthesis. X-linked recessive.

Findings: intellectual disability, self-mutilation, aggression, hyperuricemia (orange "sand" [sodium urate crystals] in diaper), gout, dystonia, macrocytosis.

Treatment: allopurinol or febuxostat (2nd line).

HGPRT:

Hyperuricemia

Gout

Pissed off (aggression, self-mutilation)

Retardation (intellectual disability)

DysTonia

Genetic code features

Unambiguous

Each codon specifies only 1 amino acid.

Degenerate/redundant

Most amino acids are coded by multiple codons. **Wobble**—codons that differ in 3rd ("wobble") position may code for the same tRNA/amino acid. Specific base pairing is usually required only in the first 2 nucleotide positions of mRNA codon.

Exceptions: methionine (AUG) and tryptophan (UGG) encoded by only 1 codon.

Commaless, nonoverlapping

Read from a fixed starting point as a continuous sequence of bases.


Exceptions: some viruses.

Universal

Genetic code is conserved throughout evolution.

Exception in humans: mitochondria.

DNA replication	Eukaryotic DNA replication is more complex than in prokaryotes but uses many enzymes analogous to those listed below. In both prokaryotes and eukaryotes, DNA replication is semiconservative, involves continuous and discontinuous (Okazaki fragment) synthesis, and occurs in the 5' → 3' direction.	
Origin of replication A	Particular consensus sequence in genome where DNA replication begins. May be single (prokaryotes) or multiple (eukaryotes).	AT-rich sequences (such as TATA box regions) are found in promoters and origins of replication.
Replication fork B	Y-shaped region along DNA template where leading and lagging strands are synthesized.	
Helicase C	Unwinds DNA template at replication fork.	Helicase Halves DNA. Deficient in Bloom syndrome (<i>BLM</i> gene mutation).
Single-stranded binding proteins D	Prevent strands from reannealing.	
DNA topoisomerases E	Create a single- or double-stranded break in the helix to add or remove supercoils.	In eukaryotes: irinotecan/topotecan inhibit topoisomerase (TOP) I, etoposide/teniposide inhibit TOP II. In prokaryotes: fluoroquinolones inhibit TOP II (DNA gyrase) and TOP IV.
Primase F	Makes an RNA primer on which DNA polymerase III can initiate replication.	
DNA polymerase III G	Prokaryotes only. Elongates leading strand by adding deoxynucleotides to the 3' end. Elongates lagging strand until it reaches primer of preceding fragment.	DNA polymerase III has 5' → 3' synthesis and proofreads with 3' → 5' exonuclease. Drugs blocking DNA replication often have a modified 3' OH, thereby preventing addition of the next nucleotide ("chain termination").
DNA polymerase I H	Prokaryotes only. Degrades RNA primer; replaces it with DNA.	Same functions as DNA polymerase III, also excises RNA primer with 5' → 3' exonuclease.
DNA ligase I	Catalyzes the formation of a phosphodiester bond within a strand of double-stranded DNA.	Joins Okazaki fragments. Ligase Links DNA.
Telomerase	Eukaryotes only. A reverse transcriptase (RNA-dependent DNA polymerase) that adds DNA (TTAGGG) to 3' ends of chromosomes to avoid loss of genetic material with every duplication.	Often dysregulated in cancer cells, allowing unlimited replication. Telomerase TAGs for Greatness and Glory.



Mutations in DNA

Severity of damage: silent \ll missense $<$ nonsense $<$ frameshift.

Types of single nucleotide (point) mutations:

- **Transition**—purine to purine (eg, A to G) or pyrimidine to pyrimidine (eg, C to T).
- **Transversion**—purine to pyrimidine (eg, A to T) or pyrimidine to purine (eg, C to G).

Single nucleotide substitutions**Silent mutation**

Nucleotide substitution codes for same (synonymous) amino acid; often base change in 3rd position of codon (tRNA wobble).

Missense mutation

Nucleotide substitution results in changed amino acid (called conservative if new amino acid has similar chemical structure).

Examples include sickle cell disease (substitution of glutamic acid with valine).

Nonsense mutation

Nucleotide substitution results in early **stop** codon (UGA, UAA, UAG). Usually results in nonfunctional protein. **Stop the nonsense!**

Other mutations**Frameshift mutation**


Deletion or insertion of a number of nucleotides not divisible by 3 \rightarrow misreading of all nucleotides downstream. Protein may be shorter or longer, and its function may be disrupted or altered.

Examples include Duchenne muscular dystrophy, Tay-Sachs disease.

Splice site mutation

Retained intron in mRNA \rightarrow protein with impaired or altered function.



Examples include rare causes of cancers, dementia, epilepsy, some types of β -thalassemia, Gaucher disease, Marfan syndrome.



Lac operon

Classic example of a genetic response to an environmental change. Glucose is the preferred metabolic substrate in *E. coli*, but when glucose is absent and lactose is available, the *lac* operon is activated to switch to lactose metabolism. Mechanism of shift:

- Low glucose → ↑ adenylate cyclase activity → ↑ generation of cAMP from ATP → activation of catabolite activator protein (CAP) → ↑ transcription.
- High lactose → unbinds repressor protein from repressor/operator site → ↑ transcription.

**DNA repair****Single strand****Nucleotide excision repair**

Specific endonucleases release the oligonucleotides containing damaged bases; DNA polymerase and ligase fill and reseal the gap, respectively. Repairs bulky helix-distorting lesions. Occurs in G₁ phase of cell cycle.

Defective in xeroderma pigmentosum (inability to repair DNA pyrimidine dimers caused by UV exposure). Findings: dry skin, extreme light sensitivity, skin cancer.

Base excision repair

Base-specific **G**lycosylase removes altered base and creates AP site (apurinic/aprimidinic). One or more nucleotides are removed by **AP-Endonuclease**, which cleaves 5' end. **AP-Lyase** cleaves 3' end. **DNA Polymerase-β** fills the gap and **DNA Ligase** seals it. Occurs throughout cell cycle.

Important in repair of spontaneous/toxic deamination. "**GEL P**lease"

Mismatch repair

Mismatched nucleotides in newly synthesized (unmethylated) strand are removed and gap is filled and resealed. Occurs predominantly in S phase of cell cycle.

Defective in Lynch syndrome (hereditary nonpolyposis colorectal cancer [HNPCC]).

Double strand**Nonhomologous end joining**

Brings together 2 ends of DNA fragments to repair double-stranded breaks.


Defective in ataxia-telangiectasia. No requirement for homology. Some DNA may be lost.

Homologous recombination

Requires 2 homologous DNA duplexes. A strand from damaged dsDNA is repaired using a complementary strand from intact homologous dsDNA as a template.

Defective in breast/ovarian cancers with *BRCA1* mutation and in Fanconi anemia. Restores duplexes accurately without loss of nucleotides.

Functional organization of a eukaryotic gene



Regulation of gene expression

Promoter

Site where RNA polymerase II and multiple other transcription factors bind to DNA upstream from gene locus (AT-rich upstream sequence with TATA and CAAT boxes).

Promoter mutation commonly results in dramatic ↓ in level of gene transcription.

Enhancer


DNA locus where regulatory proteins (“**activators**”) bind, **increasing** expression of a gene on the same chromosome.

Enhancers and silencers may be located close to, far from, or even within (in an intron) the gene whose expression they regulate.

Silencer

DNA locus where regulatory proteins (“**repressors**”) bind, **decreasing** expression of a gene on the same chromosome.

RNA processing (eukaryotes)



Initial transcript is called heterogeneous nuclear RNA (hnRNA). hnRNA is then modified and becomes mRNA.

The following processes occur in the nucleus:

- Capping of 5' end (addition of 7-methylguanosine cap)
- Polyadenylation of 3' end (≈ 200 As)
- Splicing out of introns

Capped, tailed, and spliced transcript is called mRNA.

mRNA is transported out of nucleus to be translated in cytosol.

mRNA quality control occurs at cytoplasmic processing bodies (P-bodies), which contain exonucleases, decapping enzymes, and microRNAs; mRNAs may be degraded or stored in P-bodies for future translation.

Poly-A polymerase does not require a template. AAUAAA = polyadenylation signal.

RNA polymerases**Eukaryotes**

RNA polymerase I makes **r**rRNA, the most common (**r**ampant) type; present only in nucleolus.

RNA polymerase II makes **m**rRNA (**m**assive), **mi**croRNA (**mi**rRNA), and **s**mall **n**uclear RNA (**sn**rRNA).

RNA polymerase III makes 5S rRNA, **t**rRNA (**t**iny).

No proofreading function, but can initiate chains. RNA polymerase II opens DNA at promoter site.

I, II, and III are numbered in the same order that their products are used in protein synthesis: rRNA, mRNA, then tRNA.

α -amanitin, found in *Amanita phalloides* (death cap mushrooms), inhibits RNA polymerase II. Causes severe hepatotoxicity if ingested.

Actinomycin D, also called dactinomycin, inhibits RNA polymerase in both prokaryotes and eukaryotes.

Prokaryotes


1 RNA polymerase (multisubunit complex) makes all 3 kinds of RNA.

Rifampin inhibits DNA-dependent RNA polymerase in prokaryotes.

Splicing of pre-mRNA

Part of process by which precursor mRNA (pre-mRNA) is transformed into mature mRNA.

Alterations in snRNP assembly can cause clinical disease; eg, in spinal muscular atrophy, snRNP assembly is affected due to \downarrow SMN protein \rightarrow congenital degeneration of anterior horns of spinal cord \rightarrow symmetric weakness (hypotonia, or “floppy baby syndrome”).



Introns vs exons


Exons contain the actual genetic information coding for protein.

Introns do not code for protein, but are important in regulation of gene expression.

Different exons are frequently combined by alternative splicing to produce a larger number of unique proteins.

Alternative splicing can produce a variety of protein products from a single hnRNA sequence (eg, transmembrane vs secreted Ig, tropomyosin variants in muscle, dopamine receptors in the brain).

Introns are **int**ervening sequences and stay **in** the nucleus, whereas **exons exit** and are **exp**ressed.



tRNA**Structure**

75–90 nucleotides, 2° structure, cloverleaf form, anticodon end is opposite 3' aminoacyl end. All tRNAs, both eukaryotic and prokaryotic, have CCA at 3' end along with a high percentage of chemically modified bases. The amino acid is covalently bound to the 3' end of the tRNA. **CCA Can Carry Amino acids.**


T-arm: contains the TΨC (ribothymidine, pseudouridine, cytidine) sequence necessary for tRNA-ribosome binding. **T**-arm **T**ethers tRNA molecule to ribosome.

D-arm: contains **D**ihydrouridine residues necessary for tRNA recognition by the correct aminoacyl-tRNA synthetase. **D**-arm allows **D**etection of the tRNA by aminoacyl-tRNA synthetase.

Attachment site: the 5'-CCA-3' is the amino acid acceptor site.

Charging

Aminoacyl-tRNA synthetase (uses ATP; 1 unique enzyme per respective amino acid) and binding of charged tRNA to the codon are responsible for the accuracy of amino acid selection. Aminoacyl-tRNA synthetase matches an amino acid to the tRNA by scrutinizing the amino acid before and after it binds to tRNA. If an incorrect amino acid is attached, the bond is hydrolyzed. A mischarged tRNA reads the usual codon but inserts the wrong amino acid.

**Start and stop codons**

mRNA start codons	AUG (or rarely GUG).	AUG in AUG urates protein synthesis.
Eukaryotes	Codes for methionine, which may be removed before translation is completed.	
Prokaryotes	Codes for <i>N</i> -formylmethionine (fMet).	fMet stimulates neutrophil chemotaxis.
mRNA stop codons	UGA, UAA, UAG.	UGA = U Go Away. UAA = U Are Away. UAG = U Are Gone.

Protein synthesis**Initiation**

1. Eukaryotic initiation factors (eIFs) identify the 5' cap.
2. eIFs help assemble the 40S ribosomal subunit with the initiator tRNA.
3. eIFs released when the mRNA and the ribosomal 60S subunit assemble with the complex. Requires GTP.

Eukaryotes: 40S + 60S → 80S (**E**ven).

Prokaryotes: 30S + 50S → 70S (**P**rime).

Synthesis occurs from N-terminus to C-terminus.

ATP—tRNA **A**ctivation (charging).

GTP—tRNA **G**ripping and **G**oing places (translocation).

Elongation

- 1 Aminoacyl-tRNA binds to A site (except for initiator methionine, which binds the P site), requires an elongation factor and GTP.
- 2 rRNA (“ribozyme”) catalyzes peptide bond formation, transfers growing polypeptide to amino acid in A site.
- 3 Ribosome advances 3 nucleotides toward 3' end of mRNA, moving peptidyl tRNA to P site (translocation).

Think of “going **APE**”:


A site = incoming **A**minoacyl-tRNA.

P site = accommodates growing **P**eptide.

E site = holds **E**mpy tRNA as it **E**xits.

Termination

Eukaryotic release factors (eRFs) recognize the stop codon and halt translation → completed polypeptide is released from ribosome. Requires GTP.

**Posttranslational modifications****Trimming**

Removal of N- or C-terminal propeptides from zymogen to generate mature protein (eg, trypsinogen to trypsin).

Covalent alterations

Phosphorylation, glycosylation, hydroxylation, methylation, acetylation, and ubiquitination.

Chaperone protein

Intracellular protein involved in facilitating and maintaining protein folding. In yeast, heat shock proteins (eg, HSP60) are expressed at high temperatures to prevent protein denaturing/misfolding.

► BIOCHEMISTRY—CELLULAR

Cell cycle phases Checkpoints control transitions between phases of cell cycle. This process is regulated by cyclins, cyclin-dependent kinases (CDKs), and tumor suppressors. M phase (shortest phase of cell cycle) includes mitosis (prophase, prometaphase, metaphase, anaphase, telophase) and cytokinesis (cytoplasm splits in two). G₁ and G₀ are of variable duration.


REGULATION OF CELL CYCLE

Cyclin-dependent kinases Constitutively expressed but inactive when not bound to cyclin.

Cyclins Regulatory proteins that control cell cycle events; phase specific; activate CDKs.

Cyclin-CDK complexes Phosphorylate other proteins to coordinate cell cycle progression; must be activated and inactivated at appropriate times for cell cycle to progress.

Tumor suppressors p53 → p21 induction → CDK inhibition → Rb hypophosphorylation (activation) → G₁-S progression inhibition. Mutations in tumor suppressor genes can result in unrestrained cell division (eg, Li-Fraumeni syndrome). Growth factors (eg, insulin, PDGF, EPO, EGF) bind tyrosine kinase receptors to transition the cell from G₁ to S phase.



CELL TYPES

Permanent	Remain in G ₀ , regenerate from stem cells.	Neurons, skeletal and cardiac muscle, RBCs.
Stable (quiescent)	Enter G ₁ from G ₀ when stimulated.	Hepatocytes, lymphocytes, PCT, periosteal cells.
Labile	Never go to G ₀ , divide rapidly with a short G ₁ . Most affected by chemotherapy.	Bone marrow, gut epithelium, skin, hair follicles, germ cells.

Rough endoplasmic reticulum Site of synthesis of secretory (exported) proteins and of N-linked oligosaccharide addition to lysosomal and other proteins. Nissl bodies (RER in neurons)—synthesize peptide neurotransmitters for secretion. Free ribosomes—unattached to any membrane; site of synthesis of cytosolic, peroxisomal, and mitochondrial proteins. Mucus-secreting goblet cells of the small intestine and antibody-secreting plasma cells are rich in RER. Proteins within organelles (eg, ER, Golgi bodies, lysosomes) are formed in RER.


Smooth endoplasmic reticulum Site of steroid synthesis and detoxification of drugs and poisons. Lacks surface ribosomes. Location of glucose-6-phosphatase (last step of glycogenolysis). Liver hepatocytes and steroid hormone-producing cells of the adrenal cortex and gonads are rich in SER.

Cell trafficking

Golgi is distribution center for proteins and lipids from ER to vesicles and plasma membrane. Posttranslational events in Golgi include modifying N-oligosaccharides on asparagine, adding O-oligosaccharides on serine and threonine, and adding mannose-6-phosphate to proteins for lysosomal trafficking.

Endosomes are sorting centers for material from outside the cell or from the Golgi, sending it to lysosomes for destruction or back to the membrane/Golgi for further use.

I-cell disease (inclusion cell disease/mucopolipidosis type II)—inherited lysosomal storage disorder (autosomal recessive); defect in *N*-acetylglucosaminyl-1-phosphotransferase → failure of the Golgi to phosphorylate mannose residues (↓ mannose-6-phosphate) on glycoproteins → proteins are secreted extracellularly rather than delivered to lysosomes. Results in coarse facial features, gingival hyperplasia, clouded corneas, restricted joint movements, claw hand deformities, kyphoscoliosis, and high plasma levels of lysosomal enzymes. Often fatal in childhood.



Signal recognition particle (SRP)

Abundant, cytosolic ribonucleoprotein that traffics polypeptide-ribosome complex from the cytosol to the RER. Absent or dysfunctional SRP → accumulation of protein in cytosol.

Vesicular trafficking proteins

COPI: Golgi → Golgi (retrograde); *cis*-Golgi → ER.

COPII: ER → *cis*-Golgi (anterograde).

“Two (COPII) steps forward (anterograde); one (COPI) step back (retrograde).”

Clathrin: *trans*-Golgi → lysosomes; plasma membrane → endosomes (receptor-mediated endocytosis [eg, LDL receptor activity]).

Peroxisome

Membrane-enclosed organelle involved in:

- β -oxidation of very-long-chain fatty acids (VLCFA) (strictly peroxisomal process)
- α -oxidation of branched-chain fatty acids (strictly peroxisomal process)
- Catabolism of amino acids and ethanol
- Synthesis of cholesterol, bile acids, and plasmalogens (important membrane phospholipid, especially in white matter of brain)

Zellweger syndrome—autosomal recessive disorder of peroxisome biogenesis due to mutated *PEX* genes. Hypotonia, seizures, hepatomegaly, early death.

Refsum disease—autosomal recessive disorder of α -oxidation → phytanic acid not metabolized to pristanic acid. Scaly skin, ataxia, cataracts/night blindness, shortening of 4th toe, epiphyseal dysplasia. Treatment: diet, plasmapheresis.

Adrenoleukodystrophy—X-linked recessive disorder of β -oxidation due to mutation in *ABCD1* gene → VLCFA buildup in **adrenal** glands, white (**leuko**) matter of brain, testes. Progressive disease that can lead to adrenal gland crisis, coma, and death.


Proteasome

Barrel-shaped protein complex that degrades damaged or ubiquitin-tagged proteins. Defects in the ubiquitin-proteasome system have been implicated in some cases of Parkinson disease.

Cytoskeletal elements

A network of protein fibers within the cytoplasm that supports cell structure, cell and organelle movement, and cell division.

TYPE OF FILAMENT	PREDOMINANT FUNCTION	EXAMPLES
Microfilaments	Muscle contraction, cytokinesis	Actin, microvilli.
Intermediate filaments	Maintain cell structure	Vimentin, desmin, cytokeratin, lamins, glial fibrillary acidic protein (GFAP), neurofilaments.
Microtubules	Movement, cell division	Cilia, flagella, mitotic spindle, axonal trafficking, centrioles.

Microtubule

Cylindrical outer structure composed of a helical array of polymerized heterodimers of α - and β -tubulin. Each dimer has 2 GTP bound. Incorporated into flagella, cilia, mitotic spindles. Grows slowly, collapses quickly. Also involved in slow axoplasmic transport in neurons.

Molecular motor proteins—transport cellular cargo toward opposite ends of microtubule.

- **RE**trograde to microtubule (+ \rightarrow -)—**DY**nein.
- **A**nterograde to microtubule (- \rightarrow +)—**K**inesin.

Clostridium tetani, herpes simplex virus, poliovirus, and rabies virus use dynein for retrograde transport to the neuronal cell body.

Drugs that act on microtubules (**M**icrotubules

Get **C**onstructed **V**ery **P**oorly):

- **M**ebendazole (antihelminthic)
- **G**riseofulvin (antifungal)
- **C**olchicine (antigout)
- **V**incristine/**V**inblastine (anticancer)
- **P**aclitaxel (anticancer)

Negative end **N**ear **N**ucleus.


Positive end **P**oints to **P**eriphery.

REaDY? AttacK!

Cilia structure

9 doublet + 2 singlet arrangement of microtubules **A**.
 Basal body (base of cilium below cell membrane) consists of 9 microtubule triplets **B** with no central microtubules.
 Axonemal dynein—ATPase that links peripheral 9 doublets and causes bending of cilium by differential sliding of doublets.
 Gap junctions enable coordinated ciliary movement.


Kartagener syndrome (1° ciliary dyskinesia)—immotile cilia due to a dynein arm defect. Autosomal recessive. Results in ↓ male and female fertility due to immotile sperm and dysfunctional fallopian tube cilia, respectively; ↑ risk of ectopic pregnancy. Can cause bronchiectasis, recurrent sinusitis, chronic ear infections, conductive hearing loss, and situs inversus (eg, dextrocardia on CXR **C**). ↓ nasal nitric oxide (used as screening test). (Kartagener's restaurant: take-out only; there's no **dynein** "dine-in".)

**Sodium-potassium pump**

Na^+ - K^+ ATPase is located in the plasma membrane with ATP site on cytosolic side. For each ATP consumed, 3 Na^+ leave the cell (pump phosphorylated) and 2 K^+ enter the cell (pump dephosphorylated). Plasma membrane is an asymmetric lipid bilayer containing cholesterol, phospholipids, sphingolipids, glycolipids, and proteins.


Pumpkin = pump K^+ in.

Ouabain (a cardiac glycoside) inhibits by binding to K^+ site. Cardiac glycosides (digoxin and digitoxin) directly inhibit the Na^+ - K^+ ATPase, which leads to indirect inhibition of Na^+ / Ca^{2+} exchange → ↑ $[\text{Ca}^{2+}]_i$ → ↑ cardiac contractility.



Collagen	Most abundant protein in the human body. Extensively modified by posttranslational modification. Organizes and strengthens extracellular matrix.	Be So Totally Cool, Read Books.
Type I	Most common (90%)— B one (made by osteoblasts), S kin, T endon, dentin, fascia, cornea, l ate wound repair.	Type I : b one. ↓ production in osteogenesis imperfecta type I.
Type II	C artilage (including hyaline), vitreous body, nucleus pulposus.	Type II : c artw o lage.
Type III	R eticulin—skin, b lood v essels, uterus, fetal tissue, e arly wound repair.	Type III : deficient in the uncommon, v ascular type of E hlers- D anlos syndrome (ThreE D).
Type IV	B asement membrane (basal lamina), lens.	Type IV : under the f loor (basement membrane). Defective in Alport syndrome; targeted by autoantibodies in Goodpasture syndrome.

Collagen synthesis and structure



- Synthesis**—translation of collagen α chains (preprocollagen)—usually Gly-X-Y (X and Y are proline or lysine). Collagen is $\frac{1}{3}$ glycine; glycine content of collagen is less variable than that of lysine and proline. Hydroxyproline is used for lab quantification of collagen.
- Hydroxylation**—hydroxylation of specific proline and lysine residues. Requires vitamin C; deficiency \rightarrow scurvy.
- Glycosylation**—glycosylation of pro- α -chain hydroxylysine residues and formation of procollagen via hydrogen and disulfide bonds (triple helix of 3 collagen α chains). Problems forming triple helix \rightarrow osteogenesis imperfecta.
- Exocytosis**—exocytosis of procollagen into extracellular space.
- Proteolytic processing**—cleavage of disulfide-rich terminal regions of procollagen \rightarrow insoluble tropocollagen.
- Cross-linking**—reinforcement of many staggered tropocollagen molecules by covalent lysine-hydroxylysine cross-linkage (by copper-containing lysyl oxidase) to make collagen fibrils. Problems with cross-linking \rightarrow Menkes disease.



Osteogenesis imperfecta



Genetic bone disorder (brittle bone disease) caused by a variety of gene defects (most commonly *COL1A1* and *COL1A2*). Most common form is autosomal dominant with ↓ production of otherwise normal type I collagen. Manifestations include:

- Multiple fractures and bone deformities after minimal trauma (eg, during birth)
- Blue sclerae **B** due to the translucent connective tissue over choroidal veins
- Some forms have tooth abnormalities, including opalescent teeth that wear easily due to lack of dentin (dentinogenesis imperfecta)
- Conductive hearing loss (abnormal ossicles)

May be confused with child abuse.
Treat with bisphosphonates to ↓ fracture risk.
Patients can't **BITE**:

Bones = multiple fractures

I (eye) = blue sclerae

Teeth = dental imperfections

Ear = hearing loss



Ehlers-Danlos syndrome


Faulty collagen synthesis causing hyperextensible skin **A**, hypermobile joints **B**, and tendency to bleed (easy bruising).

Multiple types. Inheritance and severity vary. Can be autosomal dominant or recessive. May be associated with joint dislocation, berry and aortic aneurysms, organ rupture.

Hypermobility type (joint instability): most common type.


Classical type (joint and skin symptoms): caused by a mutation in type V collagen (eg, *COL5A1*, *COL5A2*).

Vascular type (fragile tissues including vessels [eg, aorta], muscles, and organs that are prone to rupture [eg, gravid uterus]): mutations in type III procollagen (eg, *COL3A1*).



Menkes disease

X-linked recessive connective tissue disease caused by impaired copper absorption and transport due to defective Menkes protein (*ATP7A*, vs *ATP7B* in Wilson disease). Low copper levels (vs high levels in Wilson disease). Leads to ↓ activity of lysyl oxidase (copper is a necessary cofactor) → defective collagen. Results in brittle, “kinky” hair, growth retardation, hypotonia, ↑ risk of cerebral aneurysms.

Elastin

Stretchy protein within skin, lungs, large arteries, elastic ligaments, vocal cords, ligamenta flava (connect vertebrae → relaxed and stretched conformations).

Rich in nonhydroxylated proline, glycine, and lysine residues, vs the hydroxylated residues of collagen.


Tropoelastin with fibrillin scaffolding.

Cross-linking takes place extracellularly and gives elastin its elastic properties.

Broken down by elastase, which is normally inhibited by α_1 -antitrypsin.

α_1 -Antitrypsin deficiency results in unopposed elastase activity, which can cause COPD.


Changes with aging: ↓ dermal collagen and elastin, ↓ synthesis of collagen fibrils; cross-linking remains normal.



Marfan syndrome—autosomal dominant (with variable expression) connective tissue disorder affecting skeleton, heart, and eyes. *FBN1* gene mutation on chromosome 15 (fifteen) results in defective fibrillin, a glycoprotein that forms a sheath around elastin. Findings: tall with long extremities; pectus carinatum (more specific) or pectus excavatum **A**; hypermobile joints; long, tapering fingers and toes (arachnodactyly); cystic medial necrosis of aorta; aortic root aneurysm rupture or dissection (most common cause of death); mitral valve prolapse. Subluxation of lenses, typically upward and temporally (vs downward and medially in homocystinuria).

► BIOCHEMISTRY—LABORATORY TECHNIQUES**Polymerase chain reaction**

Molecular biology lab procedure used to amplify a desired fragment of DNA. Useful as a diagnostic tool (eg, neonatal HIV, herpes encephalitis).



- ① **Denaturation**—DNA is heated to $\sim 95^\circ\text{C}$ to separate the strands.
- ② **Annealing**—Sample is cooled to $\sim 55^\circ\text{C}$. DNA primers, a heat-stable DNA polymerase (*Taq*), and deoxynucleotide triphosphates (dNTPs) are added. DNA primers anneal to the specific sequence to be amplified on each strand.
- ③ **Elongation**—Temperature is increased to $\sim 72^\circ\text{C}$. DNA polymerase attaches dNTPs to the strand to replicate the sequence after each primer.

Heating and cooling cycles continue until the DNA sample size is sufficient.


Reverse transcriptase polymerase chain reaction

Detects and quantifies mRNA levels in a sample. Uses reverse transcription to create a complementary DNA template that is amplified via standard PCR procedure.


CRISPR/Cas9

A genome editing tool derived from bacteria. Consists of a guide RNA (gRNA) **1**, which is complementary to a target DNA sequence, and an endonuclease (Cas9), which makes a single- or double-strand break at the target site **2**. Break imperfectly repaired by nonhomologous end joining (NHEJ) → accidental frameshift mutations (“knock-out”) **3A**, or a donor DNA sequence can be added to fill in the gap using homology-directed repair (HDR) **3B**.

Not used clinically. Potential applications include removing virulence factors from pathogens, replacing disease-causing alleles of genes with healthy variants, and specifically targeting tumor cells.

**Blotting procedures****Southern blot**

1. DNA sample is enzymatically cleaved into smaller pieces, which are separated on a gel by electrophoresis, and then transferred to a filter.
2. Filter is exposed to radiolabeled DNA probe that recognizes and anneals to its complementary strand.
3. Resulting double-stranded, labeled piece of DNA is visualized when filter is exposed to film.

**Northern blot**

Similar to Southern blot, except that an RNA sample is electrophoresed. Useful for studying mRNA levels, which are reflective of gene expression.

SNoW**DRoP:**

Southern = DNA
Northern = RNA
Western = Protein

Western blot

Sample protein is separated via gel electrophoresis and transferred to a membrane. Labeled antibody is used to bind to relevant **protein**.

Southwestern blot

Identifies **DNA-binding proteins** (eg, c-Jun, c-Fos [leucine zipper motif]) using labeled double-stranded DNA probes.

Flow cytometry



Laboratory technique to assess size, granularity, and protein expression (immunophenotype) of individual cells in a sample.

Cells are tagged with antibodies specific to surface or intracellular proteins. Antibodies are then tagged with a unique fluorescent dye. Sample is analyzed one cell at a time by focusing a laser on the cell and measuring light scatter and intensity of fluorescence.

Data are plotted either as histogram (one measure) or scatter plot (any two measures, as shown). In illustration:

- Cells in left lower quadrant ⊖ for both CD8 and CD3.
- Cells in right lower quadrant ⊕ for CD8 and ⊖ for CD3. In this example, right lower quadrant is empty because all CD8-expressing cells also express CD3.
- Cells in left upper quadrant ⊕ for CD3 and ⊖ for CD8.
- Cells in right upper quadrant ⊕ for both CD8 and CD3.

Commonly used in workup of hematologic abnormalities (eg, leukemia, paroxysmal nocturnal hemoglobinuria, fetal RBCs in mother's blood) and immunodeficiencies (eg, CD4⁺ cell count in HIV).

**Microarrays**

Thousands of nucleic acid sequences are arranged in grids on glass or silicon. DNA or RNA probes are hybridized to the chip, and a scanner detects the relative amounts of complementary binding. Used to profile gene expression levels of thousands of genes simultaneously to study certain diseases and treatments. Able to detect single nucleotide polymorphisms (SNPs) and copy number variations (CNVs) for a variety of applications including genotyping, clinical genetic testing, forensic analysis, cancer mutations, and genetic linkage analysis.

Enzyme-linked immunosorbent assay

Immunologic test used to detect the presence of either a specific antigen or antibody in a patient's blood sample. Detection involves the use of an antibody linked to an enzyme. Added substrate reacts with enzyme, producing a detectable signal. Can have high sensitivity and specificity, but is less specific than Western blot.

Karyotyping

Colchicine is added to cultured cells to halt chromosomes in metaphase. Chromosomes are stained, ordered, and numbered according to morphology, size, arm-length ratio, and banding pattern (arrows in **A** point to extensive abnormalities in a cancer cell).

Can be performed on a sample of blood, bone marrow, amniotic fluid, or placental tissue. Used to diagnose chromosomal imbalances (eg, autosomal trisomies, sex chromosome disorders).

**Fluorescence in situ hybridization**

Fluorescent DNA or RNA probe binds to specific gene site of interest on chromosomes (arrows in **A** point to abnormalities in a cancer cell, whose karyotype is seen above; each fluorescent color represents a chromosome-specific probe).

Used for specific localization of genes and direct visualization of chromosomal anomalies at the molecular level.

- Microdeletion—no fluorescence on a chromosome compared to fluorescence at the same locus on the second copy of that chromosome.
- Translocation—fluorescence signal that corresponds to one chromosome is found in a different chromosome (two white arrows in **A** show fragments of chromosome 17 that have translocated to chromosome 19).
- Duplication—a second copy of a chromosome, resulting in a trisomy or tetrasomy (two blue arrows show duplicated chromosomes 8, resulting in a tetrasomy).

**Molecular cloning**

Production of a recombinant DNA molecule in a bacterial host.

Steps:

1. Isolate eukaryotic mRNA (post-RNA processing) of interest.
2. Add reverse transcriptase (an RNA-dependent DNA polymerase) to produce complementary DNA (cDNA, lacks introns).
3. Insert cDNA fragments into bacterial plasmids containing antibiotic resistance genes.
4. Transform (insert) recombinant plasmid into bacteria.
5. Surviving bacteria on antibiotic medium produce cloned DNA (copies of cDNA).


Gene expression modifications	Transgenic strategies in mice involve: <ul style="list-style-type: none"> ▪ Random insertion of gene into mouse genome ▪ Targeted insertion or deletion of gene through homologous recombination with mouse gene 	Knock- out = removing a gene, taking it out . Knock- in = inserting a gene.
Cre-lox system	Can inducibly manipulate genes at specific developmental points (eg, to study a gene whose deletion causes embryonic death).	Random insertion—constitutive expression. Targeted insertion—conditional expression.
RNA interference	Process whereby small non-coding RNA molecules target mRNAs to inhibit gene expression.	
MicroRNA (miRNA)	Naturally produced by the cell as hairpin structures. Loose nucleotide pairing allows broader targeting of related mRNAs, blocking translation and accelerating mRNA degradation.	Abnormal expression of miRNAs contributes to certain malignancies (eg, by silencing an mRNA from a tumor suppressor gene).
Small interfering RNA (siRNA)	Usually derived from exogenous dsRNA source (eg, virus). Once inside a cell, siRNA requires complete nucleotide pairing, leading to highly specific mRNA targeting. Results in mRNA cleavage prior to translation.	Can be produced by in vitro transcription for gene “knockdown” experiments.

► BIOCHEMISTRY—GENETICS

Genetic terms

TERM	DEFINITION	EXAMPLE
Codominance	Both alleles contribute to the phenotype of the heterozygote.	Blood groups A, B, AB; α_1 -antitrypsin deficiency; HLA groups.
Variable expressivity	Patients with the same genotype have varying phenotypes.	2 patients with neurofibromatosis type 1 (NF1) may have varying disease severity.
Incomplete penetrance	Not all individuals with a mutant genotype show the mutant phenotype. % penetrance \times probability of inheriting genotype = risk of expressing phenotype.	<i>BRCA1</i> gene mutations do not always result in breast or ovarian cancer.
Pleiotropy	One gene contributes to multiple phenotypic effects.	Untreated phenylketonuria (PKU) manifests with light skin, intellectual disability, and musty body odor.
Anticipation	Increased severity or earlier onset of disease in succeeding generations.	Trinucleotide repeat diseases (eg, Huntington disease).
Loss of heterozygosity	If a patient inherits or develops a mutation in a tumor suppressor gene, the complementary allele must be deleted/mutated before cancer develops. This is not true of oncogenes.	Retinoblastoma and the “two-hit hypothesis,” Lynch syndrome (HNPCC), Li-Fraumeni syndrome.

Genetic terms (continued)

TERM	DEFINITION	EXAMPLE
Dominant negative mutation	Exerts a dominant effect. A heterozygote produces a nonfunctional altered protein that also prevents the normal gene product from functioning.	A single mutated <i>p53</i> tumor suppressor gene results in a protein that is able to bind DNA and block the nonmutated <i>p53</i> from binding to the promoter.
Linkage disequilibrium	Tendency for certain alleles at 2 linked loci to occur together more or less often than expected by chance. Measured in a population, not in a family, and often varies in different populations.	
Mosaicism	Presence of genetically distinct cell lines in the same individual. Somatic mosaicism—mutation arises from mitotic errors after fertilization and propagates through multiple tissues or organs. Gonadal mosaicism—mutation only in egg or sperm cells. If parents and relatives do not have the disease, suspect gonadal (or germline) mosaicism.	McCune-Albright syndrome —due to G_s -protein activating mutation. Presents with unilateral café-au-lait spots A with ragged edges, polyostotic fibrous dysplasia (bone is replaced by collagen and fibroblasts), and at least one endocrinopathy (eg, precocious puberty). Lethal if mutation occurs before fertilization (affecting all cells), but survivable in patients with mosaicism.
		
Locus heterogeneity	Mutations at different loci can produce a similar phenotype.	Albinism.
Allelic heterogeneity	Different mutations in the same locus produce the same phenotype.	β -thalassemia.
Heteroplasmy	Presence of both normal and mutated mtDNA, resulting in variable expression in mitochondrially inherited disease.	mtDNA passed from mother to all children.
Uniparental disomy	Offspring receives 2 copies of a chromosome from 1 parent and no copies from the other parent. Heterodisomy (heterozygous) indicates a meiosis I error. Isodisomy (homozygous) indicates a meiosis II error or postzygotic chromosomal duplication of one of a pair of chromosomes, and loss of the other of the original pair.	Uniparental is euploid (correct number of chromosomes). Most occurrences of uniparental disomy (UPD) → normal phenotype. Consider isodisomy in an individual manifesting a recessive disorder when only one parent is a carrier. Examples: Prader-Willi and Angelman syndromes.

Hardy-Weinberg population genetics

	A (p)	a (q)
A (p)	AA (p^2)	Aa (pq)
a (q)	Aa (pq)	aa (q^2)

If **p** and **q** represent the frequencies of alleles A and a, respectively, in a population, then $p + q = 1$:

- p^2 = frequency of homozygosity for allele A
- q^2 = frequency of homozygosity for allele a
- $2pq$ = frequency of heterozygosity (carrier frequency, if an autosomal recessive disease)

Therefore, the sum of the frequencies of these genotypes is $p^2 + 2pq + q^2 = 1$.
The frequency of an X-linked recessive disease in males = q and in females = q^2 .

Hardy-Weinberg law assumptions include:

- No mutation occurring at the locus
- Natural selection is not occurring
- Completely random mating
- No net migration
- Large population


If a population is in Hardy-Weinberg equilibrium, then the values of **p** and **q** remain constant from generation to generation.

Disorders of imprinting Imprinting—one gene copy is silenced by methylation, and only the other copy is expressed
→ parent-of-origin effects.

	Prader-Willi syndrome	Angelman syndrome
WHICH GENE IS SILENT?	Maternally derived genes are silenced Disease occurs when the P aternal allele is deleted or mutated	Paternally derived <i>UBE3A</i> is silenced Disease occurs when the M aternal allele is deleted or mutated
SIGNS AND SYMPTOMS	Hyperphagia, obesity, intellectual disability, hypogonadism, hypotonia	S eizures, A taxia, severe I ntellectual disability, inappropriate L aughter (“happy puppet”) Set SAIL for Angel Island
CHROMOSOMES INVOLVED	Chromosome 15 of paternal origin	<i>UBE3A</i> on maternal copy of chromosome 15
NOTES	25% of cases are due to maternal uniparental disomy P rader has no D ad (P aternal D eletion)	5% of cases are due to paternal uniparental disomy M Ds are angels (M aternal D eletion)

Modes of inheritance

Autosomal dominant




Often due to defects in structural genes. Many generations, both males and females are affected.

	A	a
a	Aa	aa
a	Aa	aa

Often pleiotropic (multiple apparently unrelated effects) and variably expressive (different between individuals). Family history crucial to diagnosis. With one affected (heterozygous) parent, on average, ½ of children affected.

Autosomal recessive




With 2 carrier (heterozygous) parents, on average: ¼ of children will be affected (homozygous), ½ of children will be carriers, and ¼ of children will be neither affected nor carriers.

	A	a
A	AA	Aa
a	Aa	aa

Often due to enzyme deficiencies. Usually seen in only 1 generation. Commonly more severe than dominant disorders; patients often present in childhood.

↑ risk in consanguineous families. Unaffected individual with affected sibling has 2/3 probability of being a carrier.

X-linked recessive




Sons of heterozygous mothers have a 50% chance of being affected. No male-to-male transmission. Skips generations.

	X	X	X	X
X	XX	XX	X	XX
Y	XY	XY	Y	XY

Commonly more severe in males. Females usually must be homozygous to be affected.

X-linked dominant




Transmitted through both parents. Mothers transmit to 50% of daughters and sons; fathers transmit to all daughters but no sons.

	X	X	X	X
X	XX	XX	X	XX
Y	XY	XY	Y	XY

Examples: fragile X syndrome, Alport syndrome, hypophosphatemic rickets (also called X-linked hypophosphatemia)—phosphate wasting at proximal tubule → rickets-like presentation.

Mitochondrial inheritance



Transmitted only through the mother. All offspring of affected females may show signs of disease.

Variable expression in a population or even within a family due to heteroplasmy.

Mitochondrial myopathies—rare disorders; often present with myopathy, lactic acidosis, and CNS disease, eg, MELAS syndrome (mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes). 2° to failure in oxidative phosphorylation. Muscle biopsy often shows “ragged red fibers” (due to accumulation of diseased mitochondria in the subsarcolemma of the muscle fiber).

Leber hereditary optic neuropathy—cell death in optic nerve neurons → subacute bilateral vision loss in teens/young adults, 90% males. Usually permanent.

□ = unaffected male; ■ = affected male; ○ = unaffected female; ● = affected female.

Autosomal dominant diseases


Achondroplasia, autosomal dominant polycystic kidney disease, familial adenomatous polyposis, familial hypercholesterolemia, hereditary hemorrhagic telangiectasia (Osler-Weber-Rendu syndrome), hereditary spherocytosis, Huntington disease, Li-Fraumeni syndrome, Marfan syndrome, multiple endocrine neoplasias, myotonic muscular dystrophy, neurofibromatosis type 1 (von Recklinghausen disease), neurofibromatosis type 2, tuberous sclerosis, von Hippel-Lindau disease.

Autosomal recessive diseases

Oculocutaneous albinism, autosomal recessive polycystic kidney disease (ARPKD), cystic fibrosis, Friedreich ataxia, glycogen storage diseases, hemochromatosis, Kartagener syndrome, mucopolysaccharidoses (except Hunter syndrome), phenylketonuria, sickle cell anemia, sphingolipidoses (except Fabry disease), thalassemias, Wilson disease.

Cystic fibrosis

GENETICS	Autosomal recessive; defect in <i>CFTR</i> gene on chromosome 7; commonly a deletion of Phe508. Most common lethal genetic disease in Caucasian population.
PATHOPHYSIOLOGY	<i>CFTR</i> encodes an ATP-gated Cl^- channel that secretes Cl^- in lungs and GI tract, and reabsorbs Cl^- in sweat glands. Most common mutation \rightarrow misfolded protein \rightarrow protein retained in RER and not transported to cell membrane, causing \downarrow Cl^- (and H_2O) secretion; \uparrow intracellular Cl^- results in compensatory \uparrow Na^+ reabsorption via epithelial Na^+ channels (ENaC) \rightarrow \uparrow H_2O reabsorption \rightarrow abnormally thick mucus secreted into lungs and GI tract. \uparrow Na^+ reabsorption also causes more negative transepithelial potential difference.
DIAGNOSIS	\uparrow Cl^- concentration in pilocarpine-induced sweat test is diagnostic. Can present with contraction alkalosis and hypokalemia (ECF effects analogous to a patient taking a loop diuretic) because of ECF $\text{H}_2\text{O}/\text{Na}^+$ losses via sweating and concomitant renal K^+/H^+ wasting. \uparrow immunoreactive trypsinogen (newborn screening).
COMPLICATIONS	Recurrent pulmonary infections (eg, <i>S aureus</i> [infancy and early childhood], <i>P aeruginosa</i> [adulthood], allergic bronchopulmonary aspergillosis [ABPA]), chronic bronchitis and bronchiectasis \rightarrow reticulonodular pattern on CXR, opacification of sinuses. Pancreatic insufficiency, malabsorption with steatorrhea, fat-soluble vitamin deficiencies (A, D, E, K), biliary cirrhosis, liver disease. Meconium ileus in newborns. Infertility in men (absence of vas deferens, spermatogenesis may be unaffected) and subfertility in women (amenorrhea, abnormally thick cervical mucus). Nasal polyps, clubbing of nails.
TREATMENT	Multifactorial: chest physiotherapy, albuterol, aerosolized dornase alfa (DNase), and hypertonic saline facilitate mucus clearance. Azithromycin used as anti-inflammatory agent. Ibuprofen slows disease progression. Pancreatic enzyme replacement therapy for pancreatic insufficiency. In patients with Phe508 deletion: combination of lumacaftor (corrects misfolded proteins and improves their transport to cell surface) and ivacaftor (opens Cl^- channels \rightarrow improved chloride transport).



X-linked recessive disorders

Ornithine transcarbamylase deficiency, Fabry disease, Wiskott-Aldrich syndrome, Ocular albinism, G6PD deficiency, Hunter syndrome, Bruton agammaglobulinemia, Hemophilia A and B, Lesch-Nyhan syndrome, Duchenne (and Becker) muscular dystrophy.

X-inactivation (lyonization)—one copy of female X chromosome forms a transcriptionally inactive Barr body. Female carriers variably affected depending on the pattern of inactivation of the X chromosome carrying the mutant vs normal gene.

Oblivious Female Will Often Give Her Boys Her x-Linked Disorders

Females with Turner syndrome (45,XO) are more likely to have an X-linked recessive disorder.

Muscular dystrophies**Duchenne**

X-linked disorder typically due to **frameshift** deletions or nonsense mutations → truncated or absent dystrophin protein → progressive myofiber damage. Weakness begins in pelvic girdle muscles and progresses superiorly. Pseudohypertrophy of calf muscles due to fibrofatty replacement of muscle **A**. Waddling gait.


Onset before 5 years of age. Dilated cardiomyopathy is common cause of death.

Gowers sign—patient uses upper extremities to help stand up. Classically seen in Duchenne muscular dystrophy, but also seen in other muscular dystrophies and inflammatory myopathies (eg, polymyositis).

Duchenne = deleted dystrophin.

Dystrophin gene (*DMD*) is the largest protein-coding human gene → ↑ chance of spontaneous mutation. Dystrophin helps anchor muscle fibers, primarily in skeletal and cardiac muscle. It connects the intracellular cytoskeleton (actin) to the transmembrane proteins α - and β -dystroglycan, which are connected to the extracellular matrix (ECM). Loss of dystrophin → myonecrosis.

↑ CK and aldolase; genetic testing confirms diagnosis.

**Becker**

X-linked disorder typically due to **non-frameshift** deletions in dystrophin gene (partially functional instead of truncated). Less severe than Duchenne (**Becker is better**). Onset in adolescence or early adulthood.

Deletions can cause both Duchenne and Becker muscular dystrophies. $\frac{2}{3}$ of cases have large deletions spanning one or more exons.

Myotonic dystrophy

Autosomal dominant. **CTG** trinucleotide repeat expansion in the *DMPK* gene → abnormal expression of myotonin protein kinase → myotonia (eg, difficulty releasing hand from handshake), muscle wasting, cataracts, testicular atrophy, frontal balding, arrhythmia.

Cataracts, Toupee (early balding in men), **Gonadal atrophy**.

Rett syndrome Sporadic disorder seen almost exclusively in girls (affected males die in utero or shortly after birth). Most cases are caused by de novo mutation of *MECP2* on X chromosome. Symptoms of **Rett** syndrome usually appear between ages 1–4 and are characterized by regression (**Ret**turn) in motor, verbal, and cognitive abilities; ataxia; seizures; growth failure; and stereotyped hand-wringing.

Fragile X syndrome X-linked dominant inheritance. Trinucleotide repeat in *FMRI* gene → hypermethylation → ↓ expression. Most common inherited cause of intellectual disability (Down syndrome is the most common genetic cause, but most cases occur sporadically).
Findings: post-pubertal macroorchidism (enlarged testes), long face with a large jaw, large everted ears, autism, mitral valve prolapse, hypermobile joints.


Trinucleotide repeat expansion [(CGG)_n] occurs during oogenesis.

Trinucleotide repeat expansion diseases **Huntington** disease, **my**otonic dystrophy, **fragile X** syndrome, and **Friedreich** ataxia. May show genetic anticipation (disease severity ↑ and age of onset ↓ in successive generations).
Try (trinucleotide) hunting for my fragile cage-free eggs (X).

DISEASE	TRINUCLEOTIDE REPEAT	MODE OF INHERITANCE	MNEMONIC
Huntington disease	(CAG) _n	AD	Caudate has ↓ ACh and GABA
Myotonic dystrophy	(CTG) _n	AD	Cataracts, Toupee (early balding in men), Gonadal atrophy in men, reduced fertility in women
Fragile X syndrome	(CGG) _n	XD	Chin (protruding), Giant Gonads
Friedreich ataxia	(GAA) _n	AR	Ataxic GAAit

Autosomal trisomies

Down syndrome (trisomy 21)



Single palmar crease

Findings: intellectual disability, flat facies, prominent epicanthal folds, single palmar crease, incurved 5th finger, gap between 1st 2 toes, duodenal atresia, Hirschsprung disease, congenital heart disease (eg, ASD), Brushfield spots. Associated with early-onset Alzheimer disease (chromosome 21 codes for amyloid precursor protein), ↑ risk of AML/ALL. 95% of cases due to meiotic nondisjunction (↑ with advanced maternal age; from 1:1500 in women < 20 to 1:25 in women > 45 years old). 4% of cases due to unbalanced Robertsonian translocation, most typically between chromosomes 14 and 21. Only 1% of cases are due to postfertilization mitotic error.

Incidence 1:700.
Drinking age (21).
 Most common viable chromosomal disorder and most common cause of genetic intellectual disability.
 First-trimester ultrasound commonly shows ↑ nuchal translucency and hypoplastic nasal bone. Markers for Down syndrome are **HI** up: ↑ hCG, ↑ inhibin.

- The **5 A's** of Down syndrome:
- **A**dvanced maternal age
 - **A**tresia (duodenal)
 - **A**trioventricular septal defect
 - **A**lzheimer disease (early onset)
 - **A**ML/ALL

Edwards syndrome (trisomy 18)




Overlapping fingers

Findings: **PRINCE** Edward—**P**rominent occiput, **R**ocker-bottom feet, **I**ntellectual disability, **N**ondisjunction, **C**lenched fists with overlapping fingers, low-set **E**ars, micrognathia (small jaw), congenital heart disease, omphalocele, myelomeningocele. Death usually occurs by age 1 year.

Incidence 1:8000.
Election age (18).
 2nd most common autosomal trisomy resulting in live birth (most common is Down syndrome).


Patau syndrome (trisomy 13)



Cutis aplasia

Findings: severe intellectual disability, rocker-bottom feet, microphthalmia, microcephaly, cleft lip/**P**alate, holo**P**rosencephaly, **P**olydactyly, cutis **a**Plasia, congenital heart (**P**ump) disease, **P**olycystic kidney disease, omphalocele. Death usually occurs by age 1.

Incidence 1:15,000.
Puberty (13).
 Defect in fusion of prechordal mesoderm → midline defects.



1st trimester screening		
Trisomy	β-hCG	PAPP-A
21	↑	↓
18	↓	↓
13	↓	↓

2nd trimester screening				
Trisomy	β-hCG	Inhibin A	Estriol	AFP
21	↑	↑	↓	↓
18	↓	— or ↓	↓	↓
13	—	—	—	—

Genetic disorders by chromosome

CHROMOSOME	SELECTED EXAMPLES
3	von Hippel-Lindau disease, renal cell carcinoma
4	ADPKD (<i>PKD2</i>), achondroplasia, Huntington disease
5	Cri-du-chat syndrome, familial adenomatous polyposis
6	Hemochromatosis (<i>HFE</i>)
7	Williams syndrome, cystic fibrosis
9	Friedreich ataxia, tuberous sclerosis (<i>TSC1</i>)
11	Wilms tumor, β -globin gene defects (eg, sickle cell disease, β -thalassemia), MEN1
13	Patau syndrome, Wilson disease, retinoblastoma (<i>RBI</i>), <i>BRCA2</i>
15	Prader-Willi syndrome, Angelman syndrome, Marfan syndrome
16	ADPKD (<i>PKD1</i>), α -globin gene defects (eg, α -thalassemia), tuberous sclerosis (<i>TSC2</i>)
17	Neurofibromatosis type 1, <i>BRCA1</i> , <i>TP53</i>
18	Edwards syndrome
21	Down syndrome
22	Neurofibromatosis type 2, DiGeorge syndrome (22q11)
X	Fragile X syndrome, X-linked agammaglobulinemia, Klinefelter syndrome (XXY)

Robertsonian translocation

Chromosomal translocation that commonly involves chromosome pairs 21, 22, 13, 14, and 15. One of the most common types of translocation. Occurs when the long arms of 2 acrocentric chromosomes (chromosomes with centromeres near their ends) fuse at the centromere and the 2 short arms are lost. Balanced translocations normally do not cause any abnormal phenotype. Unbalanced translocations can result in miscarriage, stillbirth, and chromosomal imbalance (eg, Down syndrome, Patau syndrome).

Cri-du-chat syndrome

Cri du chat = **cry** of the **cat**. Congenital deletion on short arm of chromosome 5 (46,XX or XY, 5p-). Findings: microcephaly, moderate to severe intellectual disability, high-pitched **crying/meowing**, epicanthal folds, cardiac abnormalities (VSD).

Williams syndrome

A



Congenital microdeletion of long arm of chromosome 7 (deleted region includes elastin gene). Findings: distinctive “**e**lf**i**n” facies **A**, intellectual disability, hypercalcemia, well-developed verbal skills, extreme friendliness with strangers, cardiovascular problems (eg, supravalvular aortic stenosis, renal artery stenosis). Think **Will** Ferrell in **Elf**.

▶ BIOCHEMISTRY—NUTRITION

Vitamins: fat soluble	A, D, E, K. Absorption dependent on ileum and pancreas. Toxicity more common than for water-soluble vitamins because fat-soluble vitamins accumulate in fat.	Malabsorption syndromes with steatorrhea (eg, cystic fibrosis and celiac disease) or mineral oil intake can cause fat-soluble vitamin deficiencies.
Vitamins: water soluble	B ₁ (thiamine: TPP) B ₂ (riboflavin: FAD, FMN) B ₃ (niacin: NAD ⁺) B ₅ (pantothenic acid: CoA) B ₆ (pyridoxine: PLP) B ₇ (biotin) B ₉ (folate) B ₁₂ (cobalamin) C (ascorbic acid)	All wash out easily from body except B ₁₂ and B ₉ (folate). B ₁₂ stored in liver for ~ 3–4 years. B ₉ stored in liver for ~ 3–4 months. B-complex deficiencies often result in dermatitis, glossitis, and diarrhea. Can be coenzymes (eg, ascorbic acid) or precursors to coenzymes (eg, FAD, NAD ⁺).

Vitamin A


Includes retinal, retinol, retinoic acid.

FUNCTION

Antioxidant; constituent of visual pigments (**retinal**); essential for normal differentiation of epithelial cells into specialized tissue (pancreatic cells, mucus-secreting cells); prevents squamous metaplasia. Used to treat measles and acute promyelocytic leukemia (APL).

Retinol is vitamin **A**, so think **retin-A** (used topically for wrinkles and **Acne**).
Found in liver and leafy vegetables.
Supplementation in vitamin A-deficient measles patients may improve outcomes.
Use oral isotretinoin to treat severe cystic acne.
Use *all-trans* retinoic acid to treat acute promyelocytic leukemia.

DEFICIENCY



Night blindness (nyctalopia); dry, scaly skin (xerosis cutis); corneal squamous metaplasia → Bitot spots (keratin debris; foamy appearance on conjunctiva **A**); corneal degeneration (keratomalacia); immunosuppression.

EXCESS

Acute toxicity—nausea, vomiting, vertigo, and blurred vision.
Chronic toxicity—alopecia, dry skin (eg, scaliness), hepatic toxicity and enlargement, arthralgias, and idiopathic intracranial hypertension.
Teratogenic (cleft palate, cardiac abnormalities), therefore a ⊖ pregnancy test and two forms of contraception are required before isotretinoin (vitamin A derivative) is prescribed.

Isotretinoin is teratogenic.

Vitamin B₁

Also called thiamine.

FUNCTION

In thiamine pyrophosphate (TPP), a cofactor for several dehydrogenase enzyme reactions:

- **B**ranched-chain ketoacid dehydrogenase
- **α**-ketoglutarate dehydrogenase (TCA cycle)
- **P**yruvate dehydrogenase (links glycolysis to TCA cycle)
- **T**ransketolase (HMP shunt)

Be APT.

Spell beriberi as **BerlBerl** to remember vitamin **B₁**.

Wernicke encephalopathy—acute, life-threatening, neurologic condition; classic triad of confusion, ophthalmoplegia, ataxia.

Korsakoff syndrome—amnesic disorder due to chronic alcohol consumption; presents with confabulation, personality changes, memory loss (permanent).


Wernicke-Korsakoff syndrome—damage to medial dorsal nucleus of thalamus, mammillary bodies. Presentation is combination of Wernicke encephalopathy and Korsakoff syndrome.

Dry beriberi—polyneuropathy, symmetric muscle wasting.

Wet beriberi—high-output cardiac failure (dilated cardiomyopathy), edema.

DEFICIENCY

Impaired glucose breakdown → ATP depletion worsened by glucose infusion; highly aerobic tissues (eg, brain, heart) are affected first.
In alcoholic or malnourished patients, give thiamine before dextrose to ↓ risk of precipitating Wernicke encephalopathy.
Diagnosis made by ↑ in RBC transketolase activity following vitamin B₁ administration.

Vitamin B₂ Also called riboflavin.		
FUNCTION	Component of flavins FAD and FMN, used as cofactors in redox reactions, eg, the succinate dehydrogenase reaction in the TCA cycle.	FAD and FMN are derived from ribo F lavin (B ₂ ≈ 2 ATP).
DEFICIENCY	Cheilosis (inflammation of lips, scaling and fissures at the corners of the mouth), Corneal vascularization.	The 2 C 's of B ₂ .
Vitamin B₃ Also called niacin, nicotinic acid.		
FUNCTION	Constituent of NAD ⁺ , NADP ⁺ (used in redox reactions). Derived from tryptophan. Synthesis requires vitamins B ₂ and B ₆ . Used to treat dyslipidemia; lowers levels of VLDL and raises levels of HDL.	NAD derived from N iacin (B ₃ ≈ 3 ATP).
DEFICIENCY	Glossitis. Severe deficiency leads to pellagra, which can also be caused by Hartnup disease, malignant carcinoid syndrome (↑ tryptophan metabolism), and isoniazid (↓ vitamin B ₆). Symptoms of pellagra: D iarrhea, D ementia (also hallucinations), D ermatitis (C3/C4 dermatome circumferential “broad collar” rash [Casal necklace], hyperpigmentation of sun-exposed limbs A).	The 3 D 's of B ₃ . Hartnup disease —autosomal recessive. Deficiency of neutral amino acid (eg, tryptophan) transporters in proximal renal tubular cells and on enterocytes → neutral aminoaciduria and ↓ absorption from the gut → ↓ tryptophan for conversion to niacin → pellagra-like symptoms. Treat with high-protein diet and nicotinic acid. Deficiency of vitamin B ₃ → pellagra .
		
EXCESS	Facial flushing (induced by prostaglandin, not histamine; can avoid by taking aspirin with niacin), hyperglycemia, hyperuricemia.	Excess of vitamin B ₃ → podagra .
Vitamin B₅ Also called pantothenic acid.		
FUNCTION	Essential component of coenzyme A (CoA, a cofactor for acyl transfers) and fatty acid synthase.	B ₅ is “ pento ”thenic acid.
DEFICIENCY	Dermatitis, enteritis, alopecia, adrenal insufficiency.	
Vitamin B₆ Also called pyridoxine.		
FUNCTION	Converted to pyridoxal phosphate (PLP), a cofactor used in transamination (eg, ALT and AST), decarboxylation reactions, glycogen phosphorylase. Synthesis of glutathione, cystathionine, heme, niacin, histamine, and neurotransmitters including serotonin, epinephrine, norepinephrine (NE), dopamine, and GABA.	
DEFICIENCY	Convulsions, hyperirritability, peripheral neuropathy (deficiency inducible by isoniazid and oral contraceptives), sideroblastic anemia (due to impaired hemoglobin synthesis and iron excess).	

Vitamin B₇

Also called biotin.

FUNCTION

Cofactor for carboxylation enzymes (which add a 1-carbon group):

- Pyruvate carboxylase: pyruvate (3C)
→ oxaloacetate (4C)
- Acetyl-CoA carboxylase: acetyl-CoA (2C)
→ malonyl-CoA (3C)
- Propionyl-CoA carboxylase: propionyl-CoA (3C) → methylmalonyl-CoA (4C)

DEFICIENCY

Relatively rare. Dermatitis, enteritis, alopecia.
Caused by long-term antibiotic use or
excessive ingestion of raw egg whites.

“**A**vidin in egg whites **avidly** binds biotin.”

Vitamin B₉

Also called folate.

FUNCTION

Converted to tetrahydrofolic acid (THF), a
coenzyme for 1-carbon transfer/methylation
reactions.

Important for the synthesis of nitrogenous bases
in DNA and RNA.

Found in leafy green vegetables. Absorbed in
jejunum. **F**olate from **f**oliage.
Small reserve pool stored primarily in the liver.

DEFICIENCY

Macrocytic, megaloblastic anemia;
hypersegmented polymorphonuclear cells
(PMNs); glossitis; no neurologic symptoms (as
opposed to vitamin B₁₂ deficiency).
Labs: ↑ homocysteine, normal methylmalonic
acid levels. Seen in alcoholism and pregnancy.

Deficiency can be caused by several drugs (eg,
phenytoin, sulfonamides, methotrexate).
Supplemental maternal folic acid at least 1 month
prior to conception and during early pregnancy
to ↓ risk of neural tube defects. Give vitamin B₉
for the **9** months of pregnancy.

Vitamin B₁₂

Also called cobalamin.

FUNCTION

Cofactor for methionine synthase (transfers CH₃ groups as methylcobalamin) and methylmalonyl-CoA mutase. Important for DNA synthesis.

Found in animal products.


Synthesized only by microorganisms. Very large reserve pool (several years) stored primarily in the liver. Deficiency caused by malabsorption (eg, sprue, enteritis, *Diphyllobothrium latum*, achlorhydria, bacterial overgrowth, alcohol excess), lack of intrinsic factor (eg, pernicious anemia, gastric bypass surgery), absence of terminal ileum (surgical resection, eg, for Crohn disease), certain drugs (eg, metformin), or insufficient intake (eg, veganism).

DEFICIENCY

Macrocytic, megaloblastic anemia; hypersegmented PMNs; paresthesias and subacute combined degeneration (degeneration of dorsal columns, lateral corticospinal tracts, and spinocerebellar tracts) due to abnormal myelin. Associated with ↑ serum homocysteine and methylmalonic acid levels, along with 2° folate deficiency. Prolonged deficiency → irreversible nerve damage.

Anti-intrinsic factor antibodies diagnostic for pernicious anemia.

Folate supplementation can mask the hematologic symptoms of B₁₂ deficiency, but not the neurologic symptoms.

**Vitamin C**

Also called ascorbic acid.

FUNCTION

Antioxidant; also facilitates iron absorption by reducing it to Fe²⁺ state. Necessary for hydroxylation of proline and lysine in collagen synthesis. Necessary for dopamine β-hydroxylase, which converts dopamine to NE.

Found in fruits and vegetables.

Pronounce “**absorbic**” acid.

Ancillary treatment for methemoglobinemia by reducing Fe³⁺ to Fe²⁺.

DEFICIENCY

Scurvy—swollen gums, easy bruising, petechiae, hemarthrosis, anemia, poor wound healing, perifollicular and subperiosteal hemorrhages, “corkscrew” hair. Weakened immune response.

Vitamin **C** deficiency causes **sCurvy** due to a **C**ollagen synthesis defect.

EXCESS

Nausea, vomiting, diarrhea, fatigue, calcium oxalate nephrolithiasis. Can ↑ iron toxicity in predisposed individuals by increasing dietary iron absorption (ie, can worsen hereditary hemochromatosis or transfusion-related iron overload).

Vitamin D

D₃ (cholecalciferol) from exposure of skin (stratum basale) to sun, ingestion of fish, milk, plants.
 D₂ (ergocalciferol) from ingestion of plants, fungi, yeasts.
 Both converted to 25-OH D₃ (storage form) in liver and to the active form 1,25-(OH)₂ D₃ (calcitriol) in kidney.

FUNCTION

↑ intestinal absorption of Ca²⁺ and PO₄³⁻.
 ↑ bone mineralization at low levels.
 ↑ bone resorption at higher levels.

REGULATION

↑ PTH, ↓ Ca²⁺, ↓ PO₄³⁻ → ↑ 1,25-(OH)₂D₃ production.
 1,25-(OH)₂D₃ feedback inhibits its own production.
 ↑ PTH → ↑ Ca²⁺ reabsorption and ↓ PO₄³⁻ reabsorption in the kidney.

DEFICIENCY

Rickets in children (deformity, such as genu varum “bowlegs” **A**), osteomalacia in adults (bone pain and muscle weakness), hypocalcemic tetany.
 Caused by malabsorption, ↓ sun exposure, poor diet, chronic kidney disease (CKD), advanced liver disease.
 Give oral vitamin D to breastfed infants.
 Deficiency is exacerbated by pigmented skin, premature birth.

EXCESS

Hypercalcemia, hypercalciuria, loss of appetite, stupor. Seen in granulomatous diseases (↑ activation of vitamin D by epithelioid macrophages).

Vitamin E

Includes tocopherol, tocotrienol.

FUNCTION

Antioxidant (protects RBCs and membranes from free radical damage).

DEFICIENCY

Hemolytic anemia, acanthocytosis, muscle weakness, demyelination of posterior columns (↓ position and vibration sensation) and spinocerebellar tract (ataxia).

Neurologic presentation may appear similar to vitamin B₁₂ deficiency, but without megaloblastic anemia, hypersegmented neutrophils, or ↑ serum methylmalonic acid levels.

EXCESS

Risk of enterocolitis in infants.

High-dose supplementation may alter metabolism of vitamin K → enhanced anticoagulant effects of warfarin.

Vitamin K

Includes phytymenadione, phylloquinone, phytonadione, menaquinone.

FUNCTION

Activated by epoxide reductase to the reduced form, which is a cofactor for the γ -carboxylation of glutamic acid residues on various proteins required for blood clotting. Synthesized by intestinal flora.

K is for **K**oagulation. Necessary for the maturation of clotting factors II, VII, IX, X, and proteins C and S. Warfarin inhibits vitamin K–dependent synthesis of these factors and proteins.

DEFICIENCY

Neonatal hemorrhage with \uparrow PT and \uparrow aPTT but normal bleeding time (neonates have sterile intestines and are unable to synthesize vitamin K). Can also occur after prolonged use of broad-spectrum antibiotics.

Not in breast milk; neonates are given vitamin K injection at birth to prevent hemorrhagic disease of the newborn.

Zinc**FUNCTION**

Mineral essential for the activity of 100+ enzymes. Important in the formation of zinc fingers (transcription factor motif).

DEFICIENCY

Delayed wound healing, suppressed immunity, male hypogonadism, \downarrow adult hair (axillary, facial, pubic), dysgeusia, anosmia. Associated with acrodermatitis enteropathica (**A**, defect in intestinal zinc absorption). May predispose to alcoholic cirrhosis.

**Protein-energy malnutrition****Kwashiorkor**

Protein malnutrition resulting in skin lesions, edema due to \downarrow plasma oncotic pressure, liver malfunction (fatty change due to \downarrow apolipoprotein synthesis). Clinical picture is small child with swollen abdomen **A**.

Kwashiorkor results from protein-

deficient **MEALS**:

Malnutrition

Edema

Anemia

Liver (fatty)


Skin lesions (eg, hyperkeratosis, dyspigmentation)

**Marasmus**


Malnutrition not causing edema. Diet is deficient in calories but no nutrients are entirely absent.

Marasmus results in **M**uscle wasting **B**.

Ethanol metabolism



Fomepizole—blocks alcohol DH; antidote **F**or **O**verdoses of **M**ethanol or **E**thylene glycol.
Disulfiram— blocks acetaldehyde dehydrogenase → ↑ acetaldehyde → ↑ hangover symptoms → **d**iscouraging drinking.
 NAD⁺ is the limiting reagent.
 Alcohol dehydrogenase operates via zero-order kinetics.



Ethanol metabolism ↑ NADH/NAD⁺ ratio in liver, causing:

- 1 Lactic acidosis—↑ pyruvate conversion to lactate
- 2 Fasting hypoglycemia—↓ gluconeogenesis due to ↑ conversion of OAA to malate
- 3 Ketoacidosis—diversion of acetyl-CoA into ketogenesis rather than TCA cycle
- 4 Hepatosteatois— ↑ conversion of DHAP to glycerol-3-P
 - 4A; acetyl-CoA diverges into fatty acid synthesis 4B, which combines with glycerol-3-P to synthesize triglycerides

↑ NADH/NAD⁺ ratio inhibits TCA cycle → ↑ acetyl-CoA used in ketogenesis (→ ketoacidosis), lipogenesis (→ hepatosteatois).

► BIOCHEMISTRY—METABOLISM

Metabolism sites

Mitochondria	Fatty acid oxidation (β-oxidation), acetyl-CoA production, TCA cycle, oxidative phosphorylation, ketogenesis.
Cytoplasm	Glycolysis, HMP shunt, and synthesis of cholesterol (SER), proteins (ribosomes, RER), fatty acids, and nucleotides.
Both	Heme synthesis, Urea cycle, Gluconeogenesis. HUGs take two (both).

Enzyme terminology An enzyme's name often describes its function. For example, glucokinase is an enzyme that catalyzes the phosphorylation of glucose using a molecule of ATP. The following are commonly used enzyme descriptors.


Kinase	Catalyzes transfer of a phosphate group from a high-energy molecule (usually ATP) to a substrate (eg, phosphofructokinase).
Phosphorylase	Adds inorganic phosphate onto substrate without using ATP (eg, glycogen phosphorylase).
Phosphatase	Removes phosphate group from substrate (eg, fructose-1,6-bisphosphatase).
Dehydrogenase	Catalyzes oxidation-reduction reactions (eg, pyruvate dehydrogenase).
Hydroxylase	Adds hydroxyl group (–OH) onto substrate (eg, tyrosine hydroxylase).
Carboxylase	Transfers CO ₂ groups with the help of biotin (eg, pyruvate carboxylase).
Mutase	Relocates a functional group within a molecule (eg, vitamin B ₁₂ –dependent methylmalonyl-CoA mutase).
Synthase/synthetase	Joins two molecules together using a source of energy (eg, ATP, acetyl-CoA, nucleotide sugar).

Rate-determining enzymes of metabolic processes

PROCESS	ENZYME	REGULATORS
Glycolysis	Phosphofructokinase-1 (PFK-1)	AMP ⊕, fructose-2,6-bisphosphate ⊕ ATP ⊖, citrate ⊖
Gluconeogenesis	Fructose-1,6-bisphosphatase	AMP ⊖, fructose-2,6-bisphosphate ⊖
TCA cycle	Isocitrate dehydrogenase	ADP ⊕ ATP ⊖, NADH ⊖
Glycogenesis	Glycogen synthase	Glucose-6-phosphate ⊕, insulin ⊕, cortisol ⊕ Epinephrine ⊖, glucagon ⊖
Glycogenolysis	Glycogen phosphorylase	Epinephrine ⊕, glucagon ⊕, AMP ⊕ Glucose-6-phosphate ⊖, insulin ⊖, ATP ⊖
HMP shunt	Glucose-6-phosphate dehydrogenase (G6PD)	NADP ⁺ ⊕ NADPH ⊖
De novo pyrimidine synthesis	Carbamoyl phosphate synthetase II	ATP ⊕, PRPP ⊕ UTP ⊖
De novo purine synthesis	Glutamine-phosphoribosylpyrophosphate (PRPP) amidotransferase	AMP ⊖, inosine monophosphate (IMP) ⊖, GMP ⊖
Urea cycle	Carbamoyl phosphate synthetase I	N-acetylglutamate ⊕
Fatty acid synthesis	Acetyl-CoA carboxylase (ACC)	Insulin ⊕, citrate ⊕ Glucagon ⊖, palmitoyl-CoA ⊖
Fatty acid oxidation	Carnitine acyltransferase I	Malonyl-CoA ⊖
Ketogenesis	HMG-CoA synthase	
Cholesterol synthesis	HMG-CoA reductase	Insulin ⊕, thyroxine ⊕, estrogen ⊕ Glucagon ⊖, cholesterol ⊖

Summary of pathways

- 1 Galactokinase (*mild galactosemia*)
- 2 Galactose-1-phosphate uridylyltransferase (*severe galactosemia*)
- 3 Hexokinase/glucokinase
- 4 Glucose-6-phosphatase (*von Gierke disease*)
- 5 Glucose-6-phosphate dehydrogenase
- 6 Transketolase
- 7 Phosphofructokinase-1
- 8 Fructose-1,6-bisphosphatase
- 9 Fructokinase (*essential fructosuria*)
- 10 Aldolase B (*fructose intolerance*)
- 11 Aldolase B (*liver*), A (*muscle*)
- 12 Triose phosphate isomerase
- 13 Pyruvate kinase
- 14 Pyruvate dehydrogenase
- 15 Pyruvate carboxylase
- 16 PEP carboxykinase
- 17 Citrate synthase
- 18 Isocitrate dehydrogenase
- 19 α -ketoglutarate dehydrogenase
- 20 Carbamoyl phosphate synthetase I
- 21 Ornithine transcarbamylase
- 22 Propionyl-CoA carboxylase
- 23 HMG-CoA reductase



ATP production

Aerobic metabolism of one glucose molecule produces 32 net ATP via malate-aspartate shuttle (heart and liver), 30 net ATP via glycerol-3-phosphate shuttle (muscle). Anaerobic glycolysis produces only 2 net ATP per glucose molecule. ATP hydrolysis can be coupled to energetically unfavorable reactions.

Arsenic causes glycolysis to produce zero net ATP.

Activated carriers

CARRIER MOLECULE	CARRIED IN ACTIVATED FORM
ATP	Phosphoryl groups
NADH, NADPH, FADH ₂	Electrons
CoA, lipoamide	Acyl groups
Biotin	CO ₂
Tetrahydrofolates	1-carbon units
S-adenosylmethionine (SAM)	CH ₃ groups
TPP	Aldehydes

Universal electron acceptors

Nicotinamides (NAD⁺, NADP⁺ from vitamin B₃) and flavin nucleotides (FAD from vitamin B₂). NADPH is a product of the HMP shunt. NADPH is used in:

- Anabolic processes
- Respiratory burst
- Cytochrome P-450 system
- Glutathione reductase

NAD⁺ is generally used in **catabolic** processes to carry reducing equivalents away as NADH. NADPH is used in **anabolic** processes (eg, steroid and fatty acid synthesis) as a supply of reducing equivalents.

Hexokinase vs glucokinase

Phosphorylation of glucose to yield glucose-6-phosphate is catalyzed by glucokinase in the liver and hexokinase in other tissues. Hexokinase sequesters glucose in tissues, where it is used even when glucose concentrations are low. At high glucose concentrations, glucokinase helps to store glucose in liver.

	Hexokinase	Glucokinase
Location	Most tissues, except liver and pancreatic β cells	Liver, β cells of pancreas
K _m	Lower (\uparrow affinity)	Higher (\downarrow affinity)
V _{max}	Lower (\downarrow capacity)	Higher (\uparrow capacity)
Induced by insulin	No	Yes
Feedback inhibition by	Glucose-6-phosphate	Fructose-6-phosphate

Glycolysis regulation, key enzymes

Net glycolysis (cytoplasm):




Equation not balanced chemically, and exact balanced equation depends on ionization state of reactants and products.

REQUIRE ATP	Glucose $\xrightarrow{\text{Hexokinase/glucokinase}}$ Glucose-6-P	Glucose-6-P \ominus hexokinase. Fructose-6-P \ominus glucokinase.
	Fructose-6-P $\xrightarrow{\text{Phosphofructokinase-1 (rate-limiting step)}}$ Fructose-1,6-BP	AMP \oplus , fructose-2,6-bisphosphate \oplus . ATP \ominus , citrate \ominus .
PRODUCE ATP	1,3-BPG $\xleftarrow{\text{Phosphoglycerate kinase}}$ 3-PG	
	Phosphoenolpyruvate $\xrightarrow{\text{Pyruvate kinase}}$ Pyruvate	Fructose-1,6-bisphosphate \oplus . ATP \ominus , alanine \ominus .

Regulation by fructose-2,6-bisphosphate

Fructose bispophatase-2 (FBPase-2) and **phosphofructokinase-2 (PFK-2)** are the same bifunctional enzyme whose function is reversed by phosphorylation by protein kinase A.



Fasting state: \uparrow glucagon \rightarrow \uparrow cAMP \rightarrow \uparrow protein kinase A \rightarrow \uparrow FBPase-2, \downarrow PFK-2, less glycolysis, more gluconeogenesis.


FaBian the Peasant (FBP) has to work hard when starving.

Fed state: \uparrow insulin \rightarrow \downarrow cAMP \rightarrow \downarrow protein kinase A \rightarrow \downarrow FBPase-2, \uparrow PFK-2, more glycolysis, less gluconeogenesis.

Prince FrederickK (PFK) works only when fed.

Pyruvate dehydrogenase complex

Mitochondrial enzyme complex linking glycolysis and TCA cycle. Differentially regulated in fed (active)/fasting (inactive) states.



Contains 3 enzymes requiring 5 cofactors:

1. **T**hiamine pyrophosphate (B_1)
2. **L**ipoic acid
3. **C**oA (B_5 , pantothenic acid)
4. **F**AD (B_2 , riboflavin)
5. **N**AD $^+$ (B_3 , niacin)

Activated by: \uparrow NAD $^+$ /NADH ratio, \uparrow ADP \uparrow Ca $^{2+}$.

The complex is similar to the α -ketoglutarate dehydrogenase complex (same cofactors, similar substrate and action), which converts α -ketoglutarate \rightarrow succinyl-CoA (TCA cycle).

The Lovely Coenzymes For Nerds.

Arsenic inhibits lipoic acid. Arsenic poisoning clinical findings: imagine a vampire (pigmentary skin changes, skin cancer), vomiting and having diarrhea, running away from a cutie (QT prolongation) with garlic breath.

Pyruvate dehydrogenase complex deficiency

Causes a buildup of pyruvate that gets shunted to lactate (via LDH) and alanine (via ALT).
X-linked.


FINDINGS

Neurologic defects, lactic acidosis, ↑ serum alanine starting in infancy.

TREATMENT

↑ intake of ketogenic nutrients (eg, high fat content or ↑ lysine and leucine).


Pyruvate metabolism



Functions of different pyruvate metabolic pathways (and their associated cofactors):

- 1 Alanine aminotransferase (B_6): alanine carries amino groups to the liver from muscle
- 2 Pyruvate carboxylase (biotin): oxaloacetate can replenish TCA cycle or be used in gluconeogenesis
- 3 Pyruvate dehydrogenase (B_1, B_2, B_3, B_5 , lipoic acid): transition from glycolysis to the TCA cycle
- 4 Lactic acid dehydrogenase (B_3): end of anaerobic glycolysis (major pathway in RBCs, WBCs, kidney medulla, lens, testes, and cornea)

TCA cycle



Also called Krebs cycle. Pyruvate → acetyl-CoA produces 1 NADH, 1 CO₂.


The TCA cycle produces 3 NADH, 1 FADH₂, 2 CO₂, 1 GTP per acetyl-CoA = 10 ATP/ acetyl-CoA (2× everything per glucose). TCA cycle reactions occur in the mitochondria.

α-ketoglutarate dehydrogenase complex requires the same cofactors as the pyruvate dehydrogenase complex (vitamins B₁, B₂, B₃, B₅, lipoic acid).

Citrate Is Krebs' Starting Substrate For Making Oxaloacetate.

Electron transport chain and oxidative phosphorylation

NADH electrons from glycolysis enter mitochondria via the malate-aspartate or glycerol-3-phosphate shuttle. FADH₂ electrons are transferred to complex II (at a lower energy level than NADH). The passage of electrons results in the formation of a proton gradient that, coupled to oxidative phosphorylation, drives the production of ATP.



ATP PRODUCED VIA ATP SYNTHASE

1 NADH → 2.5 ATP; 1 FADH₂ → 1.5 ATP.

OXIDATIVE PHOSPHORYLATION POISONS

Electron transport inhibitors

Directly inhibit electron transport, causing a ↓ proton gradient and block of ATP synthesis.

Rotenone: complex **one** inhibitor.
“An-3-mycin” (antimycin) A: complex **3** inhibitor.
Cyanide, carbon **monoxide**, **azide** (the **-ides**, 4 letters) inhibit complex **IV**.

ATP synthase inhibitors

Directly inhibit mitochondrial ATP synthase, causing an ↑ proton gradient. No ATP is produced because electron transport stops.

Oligomycin.

Uncoupling agents

↑ permeability of membrane, causing a ↓ proton gradient and ↑ O₂ consumption. ATP synthesis stops, but electron transport continues. Produces heat.

2,4-Dinitrophenol (used illicitly for weight loss), aspirin (fevers often occur after overdose), thermogenin in brown fat (has more mitochondria than white fat).

Gluconeogenesis, irreversible enzymes

Pathway **P**roduces **F**resh **G**lucose.

Pyruvate carboxylase

In mitochondria. Pyruvate → oxaloacetate.

Requires biotin, ATP. Activated by acetyl-CoA.

Phosphoenolpyruvate carboxykinase

In cytosol. Oxaloacetate → phosphoenolpyruvate.

Requires GTP.

Fructose-1,6-bisphosphatase

In cytosol. Fructose-1,6-bisphosphate → fructose-6-phosphate.

Citrate ⊕, AMP ⊖, fructose 2,6-bisphosphate ⊖.

Glucose-6-phosphatase

In ER. Glucose-6-phosphate → glucose.

Occurs primarily in liver; serves to maintain euglycemia during fasting. Enzymes also found in kidney, intestinal epithelium. Deficiency of the key gluconeogenic enzymes causes hypoglycemia. (Muscle cannot participate in gluconeogenesis because it lacks glucose-6-phosphatase). Odd-chain fatty acids yield 1 propionyl-CoA during metabolism, which can enter the TCA cycle (as succinyl-CoA), undergo gluconeogenesis, and serve as a glucose source. Even-chain fatty acids cannot produce new glucose, since they yield only acetyl-CoA equivalents.


Pentose phosphate pathway

Also called HMP shunt. Provides a source of NADPH from abundantly available glucose-6-P (NADPH is required for reductive reactions, eg, glutathione reduction inside RBCs, fatty acid and cholesterol biosynthesis). Additionally, this pathway yields ribose for nucleotide synthesis. Two distinct phases (oxidative and nonoxidative), both of which occur in the cytoplasm. No ATP is used or produced.


Sites: lactating mammary glands, liver, adrenal cortex (sites of fatty acid or steroid synthesis), RBCs.

REACTIONS

Oxidative (irreversible)



Nonoxidative (reversible)



Glucose-6-phosphate dehydrogenase deficiency


NADPH is necessary to keep glutathione reduced, which in turn detoxifies free radicals and peroxides. ↓ NADPH in RBCs leads to hemolytic anemia due to poor RBC defense against oxidizing agents (eg, fava beans, sulfonamides, nitrofurantoin, primaquine/chloroquine, antituberculosis drugs). Infection (most common cause) can also precipitate hemolysis; inflammatory response produces free radicals that diffuse into RBCs, causing oxidative damage.

X-linked recessive disorder; most common human enzyme deficiency; more prevalent among African Americans. ↑ malarial resistance.

Heinz bodies—denatured globin chains precipitate within RBCs due to oxidative stress.

Bite cells—result from the phagocytic removal of **Heinz** bodies by splenic macrophages.

Think, “**Bite** into some **Heinz** ketchup.”



Disorders of fructose metabolism**Essential fructosuria**

Involves a defect in **fructokinase**. Autosomal recessive. A benign, asymptomatic condition (fructokinase deficiency is **kin**der), since fructose is not trapped in cells. Hexokinase becomes 1^o pathway for converting fructose to fructose-6-phosphate.

Symptoms: fructose appears in blood and urine.


Disorders of fructose metabolism cause milder symptoms than analogous disorders of galactose metabolism.

Hereditary fructose intolerance

Hereditary deficiency of **aldolase B**. Autosomal recessive. Fructose-1-phosphate accumulates, causing a ↓ in available phosphate, which results in inhibition of glycogenolysis and gluconeogenesis. Symptoms present following consumption of fruit, juice, or honey. Urine dipstick will be ⊖ (tests for glucose only); reducing sugar can be detected in the urine (nonspecific test for inborn errors of carbohydrate metabolism).

Symptoms: hypoglycemia, jaundice, cirrhosis, vomiting.

Treatment: ↓ intake of fructose, sucrose (glucose + fructose), and sorbitol (metabolized to fructose).

**Disorders of galactose metabolism****Galactokinase deficiency**

Hereditary deficiency of **galactokinase**. Galactitol accumulates if galactose is present in diet. Relatively mild condition. Autosomal recessive.

Symptoms: galactose appears in blood (galactosemia) and urine (galactosuria); infantile cataracts.


May present as failure to track objects or to develop a social smile. Galactokinase deficiency is **kin**der (benign condition).

Classic galactosemia

Absence of **galactose-1-phosphate uridylyltransferase**. Autosomal recessive. Damage is caused by accumulation of toxic substances (including galactitol, which accumulates in the lens of the eye).

Symptoms develop when infant begins feeding (lactose present in breast milk and routine formula) and include failure to thrive, jaundice, hepatomegaly, infantile cataracts, intellectual disability. Can predispose to *E coli* sepsis in neonates.

Treatment: exclude galactose and lactose (galactose + glucose) from diet.



Fructose is to **Aldolase B** as Galactose is to **UridylTransferase (FAB GUT)**.


The more serious defects lead to PO_4^{3-} depletion.

Sorbitol

An alternative method of trapping glucose in the cell is to convert it to its alcohol counterpart, sorbitol, via aldose reductase. Some tissues then convert sorbitol to fructose using sorbitol dehydrogenase; tissues with an insufficient amount/activity of this enzyme are at risk of intracellular sorbitol accumulation, causing osmotic damage (eg, cataracts, retinopathy, and peripheral neuropathy seen with chronic hyperglycemia in diabetes).

High blood levels of galactose also result in conversion to the osmotically active galactitol via aldose reductase.

Liver, Ovaries, and Seminal vesicles have both enzymes (they **LOSE** sorbitol).



Lens has primarily aldose reductase. Retina, Kidneys, and Schwann cells have only aldose reductase (**LuRKS**).

Lactase deficiency

Insufficient lactase enzyme → dietary lactose intolerance. Lactase functions on the intestinal brush border to digest lactose (in milk and milk products) into glucose and galactose.

Primary: age-dependent decline after childhood (absence of lactase-persistent allele), common in people of Asian, African, or Native American descent.

Secondary: loss of intestinal brush border due to gastroenteritis (eg, rotavirus), autoimmune disease.

Congenital lactase deficiency: rare, due to defective gene.

Stool demonstrates ↓ pH and breath shows ↑ hydrogen content with lactose hydrogen breath test.

Intestinal biopsy reveals normal mucosa in patients with hereditary lactose intolerance.

FINDINGS

Bloating, cramps, flatulence, osmotic diarrhea.

TREATMENT

Avoid dairy products or add lactase pills to diet; lactose-free milk.

Amino acids

Only L-amino acids are found in proteins.

Essential

PVT TIM HaLL: Phenylalanine, Valine, Tryptophan, Threonine, Isoleucine, Methionine, Histidine, Leucine, Lysine.

Glucogenic: Methionine, histidine, valine. We **met his valentine**, she is so **sweet** (glucogenic).

Glucogenic/ketogenic: Isoleucine, phenylalanine, threonine, tryptophan.

Ketogenic: Leucine, Lysine. The **onLy pureLy** ketogenic amino acids.

Acidic

Aspartic acid, glutamic acid.

Negatively charged at body pH.

Basic

Arginine, histidine, lysine.

Arginine is most **basic**. Histidine has no charge at body pH.

Arginine and histidine are required during periods of growth.


Arginine and lysine are ↑ in histones which bind negatively charged DNA.

His lys (lies) **are basic**.


Urea cycle

Amino acid catabolism results in the formation of common metabolites (eg, pyruvate, acetyl-CoA), which serve as metabolic fuels. Excess nitrogen generated by this process is converted to urea and excreted by the kidneys.


Ordinarily, Careless Crappers Are Also Frivolous About Urination.



Transport of ammonia by alanine



Hyperammonemia




Can be acquired (eg, liver disease) or hereditary (eg, urea cycle enzyme deficiencies). Presents with flapping tremor (eg, asterixis), slurring of speech, somnolence, vomiting, cerebral edema, blurring of vision.

↑ NH₃ changes relative amounts of α-ketoglutarate, glutamate, GABA, and glutamine to favor ↑ glutamine. CNS toxicity may involve ↓ GABA, ↓ α-ketoglutarate, TCA cycle inhibition, and cerebral edema due to glutamine-induced osmotic shifts.

Treatment: limit protein in diet.

May be given to ↓ ammonia levels:


- Lactulose to acidify GI tract and trap NH₄⁺ for excretion.
- Antibiotics (eg, rifaximin, neomycin) to ↓ ammoniagenic bacteria.
- Benzoate, phenylacetate, or phenylbutyrate react with glycine or glutamine, forming products that are excreted renally.




Ornithine transcarbamylase deficiency

Most common urea cycle disorder. X-linked recessive (vs other urea cycle enzyme deficiencies, which are autosomal recessive). Interferes with the body's ability to eliminate ammonia. Often evident in the first few days of life, but may present later. Excess carbamoyl phosphate is converted to orotic acid (part of the pyrimidine synthesis pathway). Findings: ↑ orotic acid in blood and urine, ↓ BUN, symptoms of hyperammonemia. No megaloblastic anemia (vs orotic aciduria).

Amino acid derivatives



Catecholamine synthesis/tyrosine catabolism



Phenylketonuria

Due to ↓ phenylalanine hydroxylase or ↓ tetrahydrobiopterin (BH₄) cofactor (malignant PKU). Tyrosine becomes essential. ↑ phenylalanine → ↑ phenyl ketones in urine. Findings: intellectual disability, growth retardation, seizures, fair complexion, eczema, musty body odor. Treatment: ↓ phenylalanine and ↑ tyrosine in diet, tetrahydrobiopterin supplementation.

Maternal PKU—lack of proper dietary therapy during pregnancy. Findings in infant: microcephaly, intellectual disability, growth retardation, congenital heart defects.

Autosomal recessive. Incidence ≈ 1:10,000. Screening occurs 2–3 days after birth (normal at birth because of maternal enzyme during fetal life).

Phenyl ketones—phenylacetate, phenyllactate, and phenylpyruvate.

Disorder of **aromatic** amino acid metabolism → musty body **odor**.

PKU patients must avoid the artificial sweetener aspartame, which contains phenylalanine.


Maple syrup urine disease

Blocked degradation of **branched** amino acids (**I**soleucine, **L**eucine, **V**aline) due to ↓ branched-chain α-ketoacid dehydrogenase (B₁). Causes ↑ α-ketoacids in the blood, especially those of leucine. Treatment: restriction of isoleucine, leucine, valine in diet, and thiamine supplementation.

Autosomal recessive.

Presentation: vomiting, poor feeding, urine smells like maple syrup/burnt sugar. Causes severe CNS defects, intellectual disability, death.

I Love **V**ermont **m**aple **s**yrup from maple trees (with **B**₁**r**anches).

Alkaptonuria

Congenital deficiency of homogentisate oxidase in the degradative pathway of tyrosine to fumarate → pigment-forming homogentisic acid builds up in tissue **A**. Autosomal recessive. Usually benign. Findings: bluish-black connective tissue, ear cartilage, and sclerae (ochronosis); urine turns black on prolonged exposure to air. May have debilitating arthralgias (homogentisic acid toxic to cartilage).


Homocystinuria


Causes (all autosomal recessive):

- Cystathionine synthase deficiency (treatment: ↓ methionine, ↑ cysteine, ↑ B₆, B₁₂, and folate in diet)
- ↓ affinity of cystathionine synthase for pyridoxal phosphate (treatment: ↑↑ B₆ and ↑ cysteine in diet)
- Methionine synthase (homocysteine methyltransferase) deficiency (treatment: ↑ methionine in diet)
- Methylene tetrahydrofolate reductase (MTHFR) deficiency (treatment: ↑ folate in diet)

All forms result in excess homocysteine.

HOMOCYstinuria: ↑↑ **H**omocysteine in urine, **O**steoporosis, **M**arfanoid habitus, **O**cular changes (downward and inward lens subluxation), **C**ardiovascular effects (thrombosis and atherosclerosis → stroke and MI), **kY**phosis, intellectual disability, fair complexion. In homocystinuria, lens subluxes “down and in” (vs Marfan, “up and fans out”).



Cystinuria

Hereditary defect of renal PCT and intestinal amino acid transporter that prevents reabsorption of **C**ystine, **O**rnithine, **L**ysine, and **A**rginine (**COLA**).

Excess cystine in the urine can lead to recurrent precipitation of hexagonal cystine stones **A**.

Treatment: urinary alkalization (eg, potassium citrate, acetazolamide) and chelating agents (eg, penicillamine) ↑ solubility of cystine stones; good hydration.

Autosomal recessive. Common (1:7000).
Urinary cyanide-nitroprusside test is diagnostic.

Cystine is made of 2 cysteines connected by a disulfide bond.

Organic acidemias

Most commonly present in infancy with poor feeding, vomiting, hypotonia, high anion gap metabolic acidosis, hepatomegaly, seizures. Organic acid accumulation:

- Inhibits gluconeogenesis → ↓ fasting blood glucose levels, ↑ ketoacidosis → high anion gap metabolic acidosis
- Inhibits urea cycle → hyperammonemia


Propionic acidemia

Deficiency of propionyl-CoA carboxylase
→ ↑ propionyl-CoA, ↓ methylmalonic acid.

Treatment: low-protein diet limited in substances that metabolize into propionyl-CoA: **V**aline, **O**dd-chain fatty acids, **M**ethionine, **I**soleucine, **T**hreonine (**VOMIT**).

Methylmalonic acidemia

Deficiency of methylmalonyl-CoA mutase or vitamin B₁₂.

Glycogen regulation by insulin and glucagon/epinephrine

Glycogen


Branches have α -(1,6) bonds; linkages have α -(1,4) bonds.

Skeletal muscle

Glycogen undergoes glycogenolysis \rightarrow glucose-1-phosphate \rightarrow glucose-6-phosphate, which is rapidly metabolized during exercise.

Hepatocytes

Glycogen is stored and undergoes glycogenolysis to maintain blood sugar at appropriate levels. Glycogen phosphorylase **4** liberates glucose-1-phosphate residues off branched glycogen until 4 glucose units remain on a branch. Then 4- α -D-glucanotransferase (debranching enzyme **5**) moves 3 of the 4 glucose units from the branch to the linkage. Then α -1,6-glucosidase (debranching enzyme **6**) cleaves off the last residue, liberating glucose. "Limit dextrin" refers to the two to four residues remaining on a branch after glycogen phosphorylase has already shortened it.



Note: A small amount of glycogen is degraded in lysosomes by **7** α -1,4-glucosidase (acid maltase).

Glycogen storage diseases





At least 15 types have been identified, all resulting in abnormal glycogen metabolism and an accumulation of glycogen within cells. Periodic acid–Schiff stain identifies glycogen and is useful in identifying these diseases.


Very Poor Carbohydrate Metabolism.
Types I, II, III, and V are autosomal recessive.

DISEASE	FINDINGS	DEFICIENT ENZYME	COMMENTS
Von Gierke disease (type I)	Severe fasting hypoglycemia, ↑↑ Glycogen in liver and kidneys, ↑ blood lactate, ↑ triglycerides, ↑ uric acid (Gout), and hepatomegaly, renomegaly. Liver does not regulate blood glucose.	Glucose-6-phosphatase	Treatment: frequent oral glucose/cornstarch; avoidance of fructose and galactose Impaired gluconeogenesis and glycogenolysis
Pompe disease (type II)	Cardiomegaly, hypertrophic cardiomyopathy, hypotonia, exercise intolerance, and systemic findings lead to early death.	Lysosomal acid α-1,4-glucosidase (acid maltase) with α-1,6-glucosidase activity	PomPe trashes the PumP (1st and 4th letter; heart, liver, and muscle)
Cori disease (type III)	Similar to von Gierke disease, but milder symptoms and normal blood lactate levels. Can lead to cardiomyopathy. Limit dextrin–like structures accumulate in cytosol.	Debranching enzymes (α-1,6-glucosidase and 4-α-D-glucanotransferase)	Gluconeogenesis is intact
McArdle disease (type V)	↑ glycogen in muscle, but muscle cannot break it down → painful Muscle cramps, Myoglobinuria (red urine) with strenuous exercise, and arrhythmia from electrolyte abnormalities. Second-wind phenomenon noted during exercise due to ↑ muscular blood flow.	Skeletal muscle glycogen phosphorylase (Myophosphorylase) Characterized by a flat venous lactate curve with normal rise in ammonia levels during exercise	Blood glucose levels typically unaffected McArdle = Muscle

Lysosomal storage diseases

Each is caused by a deficiency in one of the many lysosomal enzymes. Results in an accumulation of abnormal metabolic products.


DISEASE	FINDINGS	DEFICIENT ENZYME	ACCUMULATED SUBSTRATE	INHERITANCE
Sphingolipidoses				
Tay-Sachs disease 	Progressive neurodegeneration, developmental delay, hyperreflexia, hyperacusis, “cherry-red” spot on macula A , lysosomes with onion skin, no hepatosplenomegaly (vs Niemann-Pick).	1 HeXosaminidase A (“T Ay-SaX ”)	GM ₂ ganglioside	AR
Fabry disease 	Early: triad of episodic peripheral neuropathy, angiokeratomas B , hypohidrosis. Late: progressive renal failure, cardiovascular disease.	2 α-galactosidase A	Ceramide trihexoside (globotriaosylceramide)	XR
Metachromatic leukodystrophy	Central and peripheral demyelination with ataxia, dementia.	3 Arylsulfatase A	Cerebroside sulfate	AR
Krabbe disease	Peripheral neuropathy, destruction of oligodendrocytes, developmental delay, optic atrophy, globoid cells.	4 Galactocerebrosidase (galactosylceramidase)	Galactocerebroside, psychosine	AR
Gaucher disease 	Most common. Hepatosplenomegaly, pancytopenia, osteoporosis, avascular necrosis of femur, bone crises, Gaucher cells C (lipid-laden macrophages resembling crumpled tissue paper).	5 Glucocerebrosidase (β-glucosidase); treat with recombinant glucocerebrosidase	Glucocerebroside	AR
Niemann-Pick disease 	Progressive neurodegeneration, hepatosplenomegaly, foam cells (lipid-laden macrophages) D , “cherry-red” spot on macula A .	6 Sphingomyelinase	Sphingomyelin	AR
Mucopolysaccharidoses				
Hurler syndrome	Developmental delay, gargoylism, airway obstruction, corneal clouding, hepatosplenomegaly.	α-L-iduronidase	Heparan sulfate, dermatan sulfate	AR
Hunter syndrome	Mild Hurler + aggressive behavior, no corneal clouding.	Iduronate-2-sulfatase	Heparan sulfate, dermatan sulfate	XR



No man picks (Niemann-Pick) his nose with his **sphinger** (sphingomyelinase).

Hunters see clearly (no corneal clouding) and aggressively aim for the **X** (X-linked recessive).
↑ incidence of Tay-Sachs, Niemann-Pick, some forms of Gaucher disease in Ashkenazi Jews.

Fatty acid metabolism



Fatty acid synthesis requires transport of citrate from mitochondria to cytosol. Predominantly occurs in liver, lactating mammary glands, and adipose tissue.

Long-chain fatty acid (LCFA) degradation requires carnitine-dependent transport into the mitochondrial matrix.

“**SY**trate” = **SY**nthesis.

CARnitine = **CAR**nage of fatty acids.

Systemic 1° carnitine deficiency—no cellular uptake of carnitine → no transport of LCFAs into mitochondria → toxic accumulation of LCFAs in the cytosol. Causes weakness, hypotonia, hypoketotic hypoglycemia, dilated cardiomyopathy.

Medium-chain acyl-CoA dehydrogenase deficiency—↓ ability to break down fatty acids into acetyl-CoA → accumulation of fatty acyl carnitines in the blood with hypoketotic hypoglycemia. Causes vomiting, lethargy, seizures, coma, liver dysfunction, hyperammonemia. Can lead to sudden death in infants or children. Treat by avoiding fasting.

Ketone bodies

In the liver, fatty acids and amino acids are metabolized to acetoacetate and β -hydroxybutyrate (to be used in muscle and brain).

In prolonged starvation and diabetic ketoacidosis, oxaloacetate is depleted for gluconeogenesis. In alcoholism, excess NADH shunts oxaloacetate to malate. All of these processes lead to a buildup of acetyl-CoA, which is shunted into ketone body synthesis.

Ketone bodies: acetone, acetoacetate, β -hydroxybutyrate.


Breath smells like acetone (fruity odor).

Urine test for ketones can detect acetoacetate, but not β -hydroxybutyrate.


RBCs cannot utilize ketones; they strictly use glucose.

HMG-CoA lyase for ketone production.

HMG-CoA reductase for cholesterol synthesis.




Metabolic fuel use




1g carb/protein = 4 kcal
 1g alcohol = 7 kcal
 1g fatty acid = 9 kcal
 (# letters = # kcal)

Fasting and starvation	Priorities are to supply sufficient glucose to the brain and RBCs and to preserve protein.	
Fed state (after a meal)	Glycolysis and aerobic respiration.	Insulin stimulates storage of lipids, proteins, and glycogen.
Fasting (between meals)	Hepatic glycogenolysis (major); hepatic gluconeogenesis, adipose release of FFA (minor).	Glucagon and epinephrine stimulate use of fuel reserves.
Starvation days 1–3	Blood glucose levels maintained by: <ul style="list-style-type: none"> ▪ Hepatic glycogenolysis ▪ Adipose release of FFA ▪ Muscle and liver, which shift fuel use from glucose to FFA ▪ Hepatic gluconeogenesis from peripheral tissue lactate and alanine, and from adipose tissue glycerol and propionyl-CoA (from odd-chain FFA—the only triacylglycerol components that contribute to gluconeogenesis) 	Glycogen reserves depleted after day 1. RBCs lack mitochondria and therefore cannot use ketones.
Starvation after day 3	Adipose stores (ketone bodies become the main source of energy for the brain). After these are depleted, vital protein degradation accelerates, leading to organ failure and death. Amount of excess stores determines survival time.	




Lipid transport



Key enzymes in lipid transport

Cholesteryl ester transfer protein	Mediates transfer of cholesteryl esters to other lipoprotein particles.
Hepatic lipase	Degrades TGs remaining in IDL.
Hormone-sensitive lipase	Degrades TGs stored in adipocytes.
Lecithin-cholesterol acyltransferase	Catalyzes esterification of $\frac{2}{3}$ of plasma cholesterol.
Lipoprotein lipase	Degrades TGs in circulating chylomicrons.
Pancreatic lipase	Degrades dietary TGs in small intestine.
PCSK9	Degrades LDL receptor → ↑ serum LDL. Inhibition → ↑ LDL receptor recycling → ↓ serum LDL.

**Major apolipoproteins**

Apolipoprotein	Function	Chylomicron					
		Chylomicron	remnant	VLDL	IDL	LDL	HDL
E	Mediates remnant uptake (everything except LDL)	✓	✓	✓	✓		✓
A-I	Found only on alpha-lipoproteins (HDL), activates LCAT						✓
C-II	Lipoprotein lipase cofactor that catalyzes cleavage.	✓		✓	✓		✓
B-48	Mediates chylomicron secretion into lymphatics Only on particles originating from the intestines	✓	✓				
B-100	Binds LDL receptor Only on particles originating from the liver			✓	✓	✓	

Lipoprotein functions	Lipoproteins are composed of varying proportions of cholesterol, TGs, and phospholipids. LDL and HDL carry the most cholesterol. Cholesterol is needed to maintain cell membrane integrity and synthesize bile acids, steroids, and vitamin D.
Chylomicron	Delivers dietary TGs to peripheral tissues. Delivers cholesterol to liver in the form of chylomicron remnants, which are mostly depleted of their TGs. Secreted by intestinal epithelial cells.
VLDL	Delivers hepatic TGs to peripheral tissue. Secreted by liver.
IDL	Delivers TGs and cholesterol to liver. Formed from degradation of VLDL.
LDL	Delivers hepatic cholesterol to peripheral tissues. Formed by hepatic lipase modification of IDL in the liver and peripheral tissue. Taken up by target cells via receptor-mediated endocytosis. LDL is Lethal.
HDL	Mediates reverse cholesterol transport from peripheral tissues to liver. Acts as a repository for apolipoproteins C and E (which are needed for chylomicron and VLDL metabolism). Secreted from both liver and intestine. Alcohol ↑ synthesis. HDL is Healthy.

Abetalipoproteinemia Autosomal recessive. Mutation in gene that encodes microsomal transfer protein (*MTP*). Chylomicrons, VLDL, LDL absent. Deficiency in ApoB-48, ApoB-100. Affected infants present with severe fat malabsorption, steatorrhea, failure to thrive. Later manifestations include retinitis pigmentosa, spinocerebellar degeneration due to vitamin E deficiency, progressive ataxia, acanthocytosis. Intestinal biopsy shows lipid-laden enterocytes. Treatment: restriction of long-chain fatty acids, large doses of oral vitamin E.

Familial dyslipidemias

TYPE	INHERITANCE	PATHOGENESIS	↑ BLOOD LEVEL	CLINICAL
I—Hyperchylomicronemia	AR	Lipoprotein lipase or apolipoprotein C-II deficiency	Chylomicrons, TG, cholesterol	Pancreatitis, hepatosplenomegaly, and eruptive/pruritic xanthomas (no ↑ risk for atherosclerosis). Creamy layer in supernatant.
II—Familial hypercholesterolemia	AD	Absent or defective LDL receptors, or defective ApoB-100	IIa: LDL, cholesterol IIb: LDL, cholesterol, VLDL	Heterozygotes (1:500) have cholesterol ≈ 300 mg/dL; homozygotes (very rare) have cholesterol ≥ 700 mg/dL. Accelerated atherosclerosis (may have MI before age 20), tendon (Achilles) xanthomas, and corneal arcus.
III—Dysbeta-lipoproteinemia	AR	Defective ApoE	Chylomicrons, VLDL	Premature atherosclerosis, tuberoeruptive and palmar xanthomas.
IV—Hypertriglyceridemia	AD	Hepatic overproduction of VLDL	VLDL, TG	Hypertriglyceridemia (> 1000 mg/dL) can cause acute pancreatitis. Related to insulin resistance.

HIGH-YIELD PRINCIPLES IN

Immunology

“I hate to disappoint you, but my rubber lips are immune to your charms.”

—Batman & Robin

“The fully engaged heart is the antibody for the infection of violence.”

—Mark Nepo

Learning the components of the immune system and their roles in host defense at the cellular level is essential for both the understanding of disease pathophysiology and clinical practice. Know the immune mechanisms of responses to vaccines. Both congenital and acquired immunodeficiencies are very testable. Cell surface markers are high yield for understanding immune cell interactions and for laboratory diagnosis. Know the roles and functions of major cytokines and chemokines.

▶ Lymphoid Structures 96

▶ Cellular Components 99

▶ Immune Responses 104

▶ Immunosuppressants 120

▶ IMMUNOLOGY—LYMPHOID STRUCTURES

Immune system organs

- 1° organs:
 - Bone marrow—immune cell production, B cell maturation
 - Thymus—T cell maturation
- 2° organs:
 - Spleen, lymph nodes, tonsils, Peyer patches
 - Allow immune cells to interact with antigen

Lymph node

A 2° lymphoid organ that has many afferents, 1 or more efferents. Encapsulated, with trabeculae **A B**. Functions are nonspecific filtration by macrophages, circulation of B and T cells, and immune response activation.

Follicle

Site of B-cell localization and proliferation. In outer cortex. 1° follicles are dense and quiescent. 2° follicles have pale central germinal centers and are active.


Medulla

Consists of medullary cords (closely packed lymphocytes and plasma cells) and medullary sinuses. Medullary sinuses communicate with efferent lymphatics and contain reticular cells and macrophages.


Paracortex

Contains T cells. Region of cortex between follicles and medulla. Contains high endothelial venules through which T and B cells enter from blood. Not well developed in patients with DiGeorge syndrome.


Paracortex enlarges in an extreme cellular immune response (eg, EBV and other viral infections → paracortical hyperplasia → lymphadenopathy).



Lymphatic drainage associations



Spleen



Located in LUQ of abdomen, anterolateral to left kidney, protected by 9th-11th ribs. Sinusoids are long, vascular channels in red pulp (red arrows in **A**) with fenestrated “barrel hoop” basement membrane.

- T cells are found in the periarteriolar lymphatic sheath (PALS) within the white pulp (white arrows in **A**).
- B cells are found in follicles within the white pulp.
- The marginal zone, in between the red pulp and white pulp, contains macrophages and specialized B cells, and is where antigen-presenting cells (APCs) capture blood-borne antigens for recognition by lymphocytes.


Splenic macrophages remove encapsulated bacteria.

Splenic dysfunction (eg, postsplenectomy state, sickle cell disease autosplenectomy):
 ↓ IgM → ↓ complement activation → ↓ C3b opsonization → ↑ susceptibility to encapsulated organisms.



Postsplenectomy blood findings:

- Howell-Jolly bodies (nuclear remnants)
- Target cells
- Thrombocytosis (loss of sequestration and removal)
- Lymphocytosis (loss of sequestration)

Vaccinate patients undergoing splenectomy or with splenic dysfunction against encapsulated organisms (pneumococci, Hib, meningococci).



Thymus



Located in the anterosuperior mediastinum. Site of T-cell differentiation and maturation. Encapsulated. **T**hymus epithelium is derived from **T**hird pharyngeal pouch (endoderm), whereas thymic lymphocytes are of mesodermal origin. Cortex is dense with immature T cells; **M**edulla is pale with **M**ature T cells and Hassall corpuscles **A** containing epithelial reticular cells. Normal neonatal thymus “sail-shaped” on CXR **B**, involutes by age 3 years.

T cells = **T**hymus

B cells = **B**one marrow

Absent thymic shadow or hypoplastic thymus seen in some immunodeficiencies (eg, SCID, DiGeorge syndrome).

Thymoma—neoplasm of thymus. Associated with myasthenia gravis, superior vena cava syndrome, pure red cell aplasia, Good syndrome.

▶ IMMUNOLOGY—CELLULAR COMPONENTS

Innate vs adaptive immunity



	Innate immunity	Adaptive immunity
COMPONENTS	Neutrophils, macrophages, monocytes, dendritic cells, natural killer (NK) cells (lymphoid origin), complement, physical epithelial barriers, secreted enzymes	T cells, B cells, circulating antibodies
MECHANISM	Germline encoded	Variation through V(D)J recombination during lymphocyte development
RESISTANCE	Resistance persists through generations; does not change within an organism's lifetime	Microbial resistance not heritable
RESPONSE TO PATHOGENS	Nonspecific Occurs rapidly (minutes to hours) No memory response	Highly specific, refined over time Develops over long periods; memory response is faster and more robust
SECRETED PROTEINS	Lysozyme, complement, C-reactive protein (CRP), defensins, cytokines	Immunoglobulins
KEY FEATURES IN PATHOGEN RECOGNITION	Toll-like receptors (TLRs): pattern recognition receptors that recognize pathogen-associated molecular patterns (PAMPs) and lead to activation of NF- κ B. Examples of PAMPs include LPS (gram \ominus bacteria), flagellin (bacteria), nucleic acids (viruses)	Memory cells: activated B and T cells; subsequent exposure to a previously encountered antigen \rightarrow stronger, quicker immune response

Major histocompatibility complex I and II

MHC encoded by HLA genes. Present antigen fragments to T cells and bind T-cell receptors (TCRs).

	MHC I	MHC II
LOCI	HLA- A , HLA- B , HLA- C MHC I loci have 1 letter	HLA- DP , HLA- DQ , HLA- DR MHC II loci have 2 letters
BINDING	TCR and CD8	TCR and CD4
STRUCTURE	1 long chain, 1 short chain	2 equal-length chains (2 α , 2 β)
EXPRESSION	All nucleated cells, APCs, platelets (except RBCs)	APCs
FUNCTION	Present endogenous antigens (eg, viral or cytosolic proteins) to CD8+ cytotoxic T cells	Present exogenous antigens (eg, bacterial proteins) to CD4+ helper T cells
ANTIGEN LOADING	Antigen peptides loaded onto MHC I in RER after delivery via TAP (transporter associated with antigen processing)	Antigen loaded following release of invariant chain in an acidified endosome
ASSOCIATED PROTEINS	β_2 -microglobulin	Invariant chain

STRUCTURE



HLA subtypes associated with diseases

HLA SUBTYPE	DISEASE	MNEMONIC
A3	Hemochromatosis	HA3 mochromatosis
B8	Addison disease, my asthenia gravis, Graves disease	Don't Be late(8), Dr. Addison , or else you'll send my patient to the grave
B27	P soriatic arthritis, A nkylosing spondylitis, I BD-associated arthritis, R eactive arthritis	PAIR . Also called seronegative arthropathies
C	Psoriasis	
DQ2/DQ8	Celiac disease	I ate (8) too (2) much gluten at Dairy Queen
DR2	M ultiple sclerosis, hay fever, SLE, Goodpasture syndrome	DR ive 2 multiple hay pastures
DR3	DM type 1, SLE , Graves disease, Hashimoto thyroiditis, Addison disease	2-3, S-L-E
DR4	R heumatoid arthritis, DM type 1 , Addison disease	There are 4 walls in 1 " rheum " (room)
DR5	H ashimoto thyroiditis	H ashimoto is an odd Dr (DR3, DR5)

Functions of natural killer cells

Lymphocyte member of innate immune system.

Use perforin and granzymes to induce apoptosis of virally infected cells and tumor cells.

Activity enhanced by IL-2, IL-12, IFN- α , and IFN- β .

Induced to kill when exposed to a nonspecific activation signal on target cell and/or to an absence of MHC I on target cell surface.

Also kills via antibody-dependent cell-mediated cytotoxicity (CD16 binds Fc region of bound IgG, activating the NK cell).

Major functions of B and T cells**B cells**

Humoral immunity.

Recognize and present antigen—undergo somatic hypermutation to optimize antigen specificity.

Produce antibody—differentiate into plasma cells to secrete specific immunoglobulins.

Maintain immunologic memory—memory B cells persist and accelerate future response to antigen.

T cells

Cell-mediated immunity.

CD4+ T cells help B cells make antibodies and produce cytokines to recruit phagocytes and activate other leukocytes.


CD8+ T cells directly kill virus-infected and tumor cells via perforin and granzymes (similar to NK cells).

Delayed cell-mediated hypersensitivity (type IV).

Acute and chronic cellular organ rejection.

Rule of 8: MHC II \times CD4 = 8; MHC I \times CD8 = 8.

Differentiation of T cells



Positive selection

Thymic cortex. T cells expressing TCRs capable of binding self-MHC on cortical epithelial cells survive.

Negative selection

Thymic medulla. T cells expressing TCRs with high affinity for self antigens undergo apoptosis or become regulatory T cells. Tissue-restricted self-antigens are expressed in the thymus due to the action of autoimmune regulator (AIRE); deficiency leads to autoimmune polyendocrine syndrome-1 (Chronic mucocutaneous candidiasis, Hypoparathyroidism, Adrenal insufficiency, Recurrent *Candida* infections). “Without AIRE, your body will CHAR”.

Macrophage-lymphocyte interaction

Th1 cells secrete IFN- γ , which enhances the ability of monocytes and macrophages to kill microbes they ingest. This function is also enhanced by interaction of T cell CD40L with CD40 on macrophages. Macrophages also activate lymphocytes via antigen presentation.

Cytotoxic T cells

Kill virus-infected, neoplastic, and donor graft cells by inducing apoptosis. Release cytotoxic granules containing preformed proteins (eg, perforin, granzyme B). Cytotoxic T cells have CD8, which binds to MHC I on virus-infected cells.

Regulatory T cells

Help maintain specific immune tolerance by suppressing CD4⁺ and CD8⁺ T-cell effector functions. Identified by expression of CD3, CD4, CD25, and FOXP3. Activated regulatory T cells (Tregs) produce anti-inflammatory cytokines (eg, IL-10, TGF- β).


IPEX (Immune dysregulation, Polyendocrinopathy, Enteropathy, X-linked) syndrome—genetic deficiency of FOXP3 \rightarrow autoimmunity. Characterized by enteropathy, endocrinopathy, nail dystrophy, dermatitis, and/or other autoimmune dermatologic conditions. Associated with diabetes in male infants.

T- and B-cell activation APCs: B cells, dendritic cells, Langerhans cells, macrophages.

Two signals are required for T-cell activation, B-cell activation, and class switching.


T-cell activation

- ❶ Dendritic cell (specialized APC) samples and processes antigen, then migrates to the draining lymph node.
- ❷ T-cell activation (signal 1): exogenous antigen is presented on MHC II and recognized by TCR on Th (CD4+) cell. Endogenous or cross-presented antigen is presented on MHC I to Tc (CD8+) cell.
- ❸ Proliferation and survival (signal 2): costimulatory signal via interaction of B7 protein (CD80/86) on dendritic cell and CD28 on naïve T cell.
- ❹ Activated Th cell produces cytokines. Tc cell able to recognize and kill virus-infected cell.



B-cell activation and class switching


- ❶ Th-cell activation as above.
- ❷ B-cell receptor-mediated endocytosis.
- ❸ Exogenous antigen is presented on MHC II and recognized by TCR on Th cell.
- ❹ CD40 receptor on B cell binds CD40 ligand (CD40L) on Th cell.
- ❺ Th cells secrete cytokines that determine Ig class switching of B cells.
- ❻ B cells are activated, undergo class switching and affinity maturation, and begin producing antibodies.



▶ IMMUNOLOGY—IMMUNE RESPONSES

Antibody structure and function

Fab (containing the variable/hypervariable regions) consisting of light (L) and heavy (H) chains recognizes antigens. Fc region of IgM and IgG fixes complement. Heavy chain contributes to Fc and Fab regions. Light chain contributes only to Fab region.



Fab:

- Fragment, antigen binding
- Determines idiotype: unique antigen-binding pocket; only 1 antigenic specificity expressed per B cell

Fc (5 C's):


- Constant
- Carboxy terminal
- Complement binding
- Carbohydrate side chains
- Confers (determines) isotype (IgM, IgD, etc)

Generation of antibody diversity (antigen independent)

1. Random recombination of VJ (light-chain) or V(D)J (heavy-chain) genes
2. Random addition of nucleotides to DNA during recombination by terminal deoxynucleotidyl transferase (TdT)
3. Random combination of heavy chains with light chains

Generation of antibody specificity (antigen dependent)

4. Somatic hypermutation and affinity maturation (variable region)
5. Isotype switching (constant region)



Immunoglobulin isotypes

All isotypes can exist as monomers. Mature, naïve B cells prior to activation express IgM and IgD on their surfaces. They may differentiate in germinal centers of lymph nodes by isotype switching (gene rearrangement; induced by cytokines and CD40L) into plasma cells that secrete IgA, IgE, or IgG.


Affinity refers to the individual antibody-antigen interaction, while avidity describes the cumulative binding strength of all antibody-antigen interactions in a multivalent molecule.

IgG




Main antibody in 2° response to an antigen. Most abundant isotype in serum. Fixes complement, opsonizes bacteria, neutralizes bacterial toxins and viruses. Only isotype that crosses the placenta (provides infants with passive immunity that starts to wane after birth). “IgG Greets the Growing fetus.”

IgA



Prevents attachment of bacteria and viruses to mucous membranes; does not fix complement. Monomer (in circulation) or dimer (with J chain when secreted). Crosses epithelial cells by transcytosis. Produced in GI tract (eg, by Peyer patches) and protects against gut infections (eg, *Giardia*). Most produced antibody overall, but has lower serum concentrations. Released into secretions (tears, saliva, mucus) and breast milk. Picks up secretory component from epithelial cells, which protects the Fc portion from luminal proteases.

IgM



Produced in the 1° (immediate) response to an antigen. Fixes complement. Antigen receptor on the surface of B cells. Monomer on B cell, pentamer with J chain when secreted. Pentamer enables avid binding to antigen while humoral response evolves.

IgD



Unclear function. Found on surface of many B cells and in serum.

IgE



Binds mast cells and basophils; cross-links when exposed to allergen, mediating immediate (type I) hypersensitivity through release of inflammatory mediators such as histamine. Contributes to immunity to parasites by activating eosinophils.

Antigen type and memory

Thymus-independent antigens

Antigens lacking a peptide component (eg, lipopolysaccharides from gram \ominus bacteria); cannot be presented by MHC to T cells. Weakly immunogenic; vaccines often require boosters and adjuvants (eg, capsular polysaccharide subunit of *Streptococcus pneumoniae* PPSV23 vaccine).

Thymus-dependent antigens

Antigens containing a protein component (eg, *Streptococcus pneumoniae* PCV13 vaccine, polysaccharides conjugated to diphtheria toxin-like protein). Class switching and immunologic memory occur as a result of direct contact of B cells with Th cells.

Complement

System of hepatically synthesized plasma proteins that play a role in innate immunity and inflammation. Membrane attack complex (MAC) defends against gram \ominus bacteria. The CH₅₀ test is used to screen for activation of the classical complement pathway.

ACTIVATION PATHWAYS

Classic—IgG or IgM mediated.

GM makes **classic** cars.

Alternative—microbe surface molecules.

Lectin—mannose or other sugars on microbe surface.

FUNCTIONS

C3b—opsonization.

C3b binds to lipopolysaccharides on bacteria.

C3a, C4a, C5a—**anaphylaxis**.


C5a—neutrophil chemotaxis.

C5b-9 (MAC)—**cytolysis**.

Opsonins—C3b and IgG are the two 1^o opsonins in bacterial defense; enhance phagocytosis. C3b also helps clear immune complexes.

Opsonin (Greek) = to prepare for eating.

Inhibitors—decay-accelerating factor (DAF, aka CD55) and C1 esterase inhibitor help prevent complement activation on self cells (eg, RBCs).



Complement disorders**Complement protein deficiencies**


Early complement deficiencies (C1-C4) Increased risk of severe, recurrent pyogenic sinus and respiratory tract infections. Increased risk of SLE.

Terminal complement deficiencies (C5-C9) Increased susceptibility to recurrent *Neisseria* bacteremia.

Complement regulatory protein deficiencies

C1 esterase inhibitor deficiency Causes hereditary angioedema due to unregulated activation of kallikrein → ↑ bradykinin. Characterized by ↓ C4 levels. ACE inhibitors are contraindicated (also ↑ bradykinin).


Paroxysmal nocturnal hemoglobinuria A defect in the *PIGA* gene preventing the formation of glycosylphosphatidylinositol (GPI) anchors for complement inhibitors, such as decay-accelerating factor (DAF/CD55) and membrane inhibitor of reactive lysis (MIRL/CD59). Causes complement-mediated intravascular hemolysis → ↓ haptoglobin, dark urine **A**.



Important cytokines		
Acute (IL-1, IL-6, TNF- α), then recruit (IL-8, IL-12).		
SECRETED BY MACROPHAGES		
Interleukin-1	Causes fever, acute inflammation. Activates endothelium to express adhesion molecules. Induces chemokine secretion to recruit WBCs. Also called osteoclast-activating factor.	“Hot T-bone stEAK”: IL-1: fever (hot). IL-2: stimulates T cells. IL-3: stimulates bone marrow. IL-4: stimulates Ig E production. IL-5: stimulates Ig A production. IL-6: stimulates a K ute-phase protein production.
Interleukin-6	Causes fever and stimulates production of acute-phase proteins.	
Tumor necrosis factor-α	Activates endothelium. Causes WBC recruitment, vascular leak.	Causes cachexia in malignancy. Maintains granulomas in TB. IL-1, IL-6, TNF- α can mediate fever and sepsis.
Interleukin-8	Major chemotactic factor for neutrophils.	“Clean up on aisle 8.” Neutrophils are recruited by IL-8 to clear infections.
Interleukin-12	Induces differentiation of T cells into Th1 cells. Activates NK cells.	
SECRETED BY ALL T CELLS		
Interleukin-2	Stimulates growth of helper, cytotoxic, and regulatory T cells, and NK cells.	
Interleukin-3	Supports growth and differentiation of bone marrow stem cells. Functions like GM-CSF.	
FROM Th1 CELLS		
Interferon-γ	Secreted by NK cells and T cells in response to antigen or IL-12 from macrophages; stimulates macrophages to kill phagocytosed pathogens. Inhibits differentiation of Th2 cells.	Also activates NK cells to kill virus-infected cells. Increases MHC expression and antigen presentation by all cells. Activates macrophages to induce granuloma formation.
FROM Th2 CELLS		
Interleukin-4	Induces differentiation of T cells into Th (helper) 2 cells. Promotes growth of B cells. Enhances class switching to Ig E and Ig G .	Ain't too proud 2 BEG 4 help .
Interleukin-5	Promotes growth and differentiation of B cells. Enhances class switching to IgA. Stimulates growth and differentiation of eosinophils.	
Interleukin-10	Attenuates inflammatory response. Decreases expression of MHC class II and Th1 cytokines. Inhibits activated macrophages and dendritic cells. Also secreted by regulatory T cells.	TGF- β and IL-10 both attenuate the immune response.

Respiratory burst

Also called oxidative burst. Involves the activation of the phagocyte NADPH oxidase complex (eg, in neutrophils, monocytes), which utilizes O_2 as a substrate. Plays an important role in the immune response → rapid release of reactive oxygen species (ROS). NADPH plays a role in both the creation and neutralization of ROS. Myeloperoxidase contains a blue-green, heme-containing pigment that gives sputum its color.



Phagocytes of patients with CGD can utilize H_2O_2 generated by invading organisms and convert it to ROS. Patients are at ↑ risk for infection by catalase ⊕ species (eg, *S aureus*, *Aspergillus*) capable of neutralizing their own H_2O_2 , leaving phagocytes without ROS for fighting infections. Pyocyanin of *P aeruginosa* generates ROS to kill competing pathogens. Oxidative burst also leads to K^+ influx, which releases lysosomal enzymes. Lactoferrin is a protein found in secretory fluids and neutrophils that inhibits microbial growth via iron chelation.

Interferons

IFN- α , IFN- β , IFN- γ

MECHANISM

A part of innate host defense, **interferons interfere** with both RNA and DNA viruses. Cells infected with a virus synthesize these glycoproteins, which act on local cells, priming them for viral defense by downregulating protein synthesis to resist potential viral replication and by upregulating MHC expression to facilitate recognition of infected cells. Also play a major role in activating antitumor immunity.

CLINICAL USE

Chronic HBV, Kaposi sarcoma, hairy cell leukemia, condyloma acuminatum, renal cell carcinoma, malignant melanoma, multiple sclerosis, chronic granulomatous disease.

ADVERSE EFFECTS

Flu-like symptoms, depression, neutropenia, myopathy, interferon-induced autoimmunity.

Cell surface proteins

T cells	TCR (binds antigen-MHC complex) CD3 (associated with TCR for signal transduction) CD28 (binds B7 on APC)	
Helper T cells	CD4, CD40L, CXCR4/CCR5 (co-receptors for HIV)	
Cytotoxic T cells	CD8	
Regulatory T cells	CD4, CD25	
B cells	Ig (binds antigen) CD19, CD20, CD21 (receptor for Epstein-Barr virus), CD40 MHC II, B7	Must be 21 to drink Beer in a Barr
Macrophages	CD14 (receptor for PAMPs, eg, LPS), CD40 CCR5 MHC II, B7 (CD80/86) Fc and C3b receptors (enhanced phagocytosis)	
NK cells	CD16 (binds Fc of IgG), CD56 (suggestive marker for NK)	
Hematopoietic stem cells	CD34	

Anergy

State during which a cell cannot become activated by exposure to its antigen. T and B cells become anergic when exposed to their antigen without costimulatory signal (signal 2). Another mechanism of self-tolerance.

Passive vs active immunity

	Passive	Active
MEANS OF ACQUISITION	Receiving preformed antibodies	Exposure to exogenous antigens
ONSET	Rapid	Slow
DURATION	Short span of antibodies (half-life = 3 weeks)	Long-lasting protection (memory)
EXAMPLES	IgA in breast milk, maternal IgG crossing placenta, antitoxin, humanized monoclonal antibody	Natural infection, vaccines, toxoid
NOTES	After exposure to T etanus toxin, B otulinum toxin, H BV, V aricella, R abies virus, or D iphtheria toxin, unvaccinated patients are given preformed antibodies (passive)—“ T o B e H ealed V ery R apidly before D ying”	Combined passive and active immunizations can be given for hepatitis B or rabies exposure


Vaccination

Induces an active immune response (humoral and/or cellular) to specific pathogens.

VACCINE TYPE	DESCRIPTION	PROS/CONS	EXAMPLES
Live attenuated vaccine	Microorganism loses its pathogenicity but retains capacity for transient growth within inoculated host. Induces cellular and humoral responses . MMR and varicella vaccines can be given to HIV ⊕ patients without evidence of immunity if CD4 cell count ≥ 200 cells/mm ³ .	Pros: induces strong, often lifelong immunity. Cons: may revert to virulent form. Often contraindicated in pregnancy and immunodeficiency.	A denovirus (nonattenuated, given to military recruits), T yphoid (Ty21a, oral), P olio (Sabin), V aricella (chickenpox), S mallpox, B CG, Y ellow fever, I nfluenza (intranasal), M MR, R otavirus “ A ttention T eachers! P lease V accinate S mall, B eautiful Y oung I nfants with M MR R egularly!”
Killed or inactivated vaccine	Pathogen is inactivated by heat or chemicals. Maintaining epitope structure on surface antigens is important for immune response. Mainly induces a humoral response .	Pros: safer than live vaccines. Cons: weaker immune response; booster shots usually required.	Hepatitis A , T yphoid (Vi polysaccharide, intramuscular), R abies, I nfluenza, P olio (SalK) A TRIP could Kill you
Subunit	Includes only the antigens that best stimulate the immune system.	Pros: lower chance of adverse reactions. Cons: expensive, weaker immune response.	HBV (antigen = HBsAg), HPV (types 6, 11, 16, and 18), acellular pertussis (aP), <i>Neisseria meningitidis</i> (various strains), <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> type b.
Toxoid	Denatured bacterial toxin with an intact receptor binding site. Stimulates the immune system to make antibodies without potential for causing disease.	Pros: protects against the bacterial toxins. Cons: antitoxin levels decrease with time, may require a booster.	<i>Clostridium tetani</i> , <i>Corynebacterium diphtheriae</i>

Hypersensitivity types Four types (**ABCD**): **A**naphylactic and **A**topical (type I), **A**nti**B**ody-mediated (type II), **I**mmune **C**omplex (type III), **D**elayed (cell-mediated, type IV). Types I, II, and III are all antibody-mediated.

Type I hypersensitivity




Anaphylactic and atopic—two phases:

- Immediate (minutes): antigen crosslinks preformed IgE on presensitized mast cells → immediate degranulation → release of histamine (a vasoactive amine) and tryptase (a marker of mast cell activation).
- Late (hours): chemokines (attract inflammatory cells, eg, eosinophils) and other mediators (eg, leukotrienes) from mast cells → inflammation and tissue damage.

First (type) and **F**ast (anaphylaxis).
 Test: skin test or blood test (ELISA) for allergen-specific IgE.
 Example:

- Anaphylaxis (eg, food, drug, or bee sting allergies)
- Allergic asthma

Type II hypersensitivity



Antibodies bind to cell-surface antigens → cellular destruction, inflammation, and cellular dysfunction.

Cellular destruction—cell is opsonized (coated) by antibodies, leading to either:

- Phagocytosis and/or activation of complement system.
- NK cell killing (antibody-dependent cellular cytotoxicity).

Inflammation—binding of antibodies to cell surfaces → activation of complement system and Fc receptor-mediated inflammation.

Cellular dysfunction—antibodies bind to cell surface receptors → abnormal blockade or activation of downstream process.

Direct Coombs test—detects antibodies attached **directly** to the RBC surface.
 Indirect Coombs test—detects presence of unbound antibodies in the serum

Examples:


- Autoimmune-hemolytic anemia
- Immune thrombocytopenia
- Transfusion reactions
- Hemolytic disease of the newborn

Examples:

- Goodpasture syndrome
- Rheumatic fever
- Hyperacute transplant rejection

Examples:

- Myasthenia gravis
- Graves disease
- Pemphigus vulgaris

Hypersensitivity types (continued)**Type III hypersensitivity**

Immune complex—antigen-antibody (mostly IgG) complexes activate complement, which attracts neutrophils; neutrophils release lysosomal enzymes.

Can be associated with vasculitis and systemic manifestations.

Serum sickness—the prototypic immune complex disease. Antibodies to foreign proteins are produced and 1–2 weeks later, antibody-antigen complexes form and deposit in tissues → complement activation → inflammation and tissue damage.


Arthus reaction—a local subacute immune complex-mediated hypersensitivity reaction. Intradermal injection of antigen into a presensitized (has circulating IgG) individual leads to immune complex formation in the skin (eg, enhanced local reaction to a booster vaccination). Characterized by edema, necrosis, and activation of complement.

In type **III** reaction, imagine an immune complex as **3** things stuck together: antigen-antibody-complement.

Examples:

- SLE
- Polyarteritis nodosa
- Poststreptococcal glomerulonephritis

Fever, urticaria, arthralgia, proteinuria, lymphadenopathy occur 1–2 weeks after antigen exposure. Serum sickness-like reactions are associated with some drugs (may act as haptens, eg, penicillin) and infections (eg, hepatitis B).

Type IV hypersensitivity

Two mechanisms, each involving T cells:

1. Direct cell cytotoxicity: CD8+ cytotoxic T cells kill targeted cells.
2. Inflammatory reaction: effector CD4+ T cells recognize antigen and release inflammation-inducing cytokines (shown in illustration).

Response does not involve antibodies (vs types I, II, and III).










Examples: contact dermatitis (eg, poison ivy, nickel allergy) and graft-versus-host disease.

Tests: PPD for TB infection; patch test for contact dermatitis; *Candida* skin test for T cell immune function.

4T's: **T** cells, **T**ransplant rejections, **T**B skin tests, **T**ouching (contact dermatitis).

Fourth (type) and **last** (delayed).

Blood transfusion reactions

TYPE	PATHOGENESIS	TIMING	CLINICAL PRESENTATION	DONOR BLOOD	HOST BLOOD
Allergic/ anaphylactic reaction	Type I hypersensitivity reaction against plasma proteins in transfused blood IgA-deficient individuals should receive blood products without IgA	Within minutes to 2-3 hr (due to release of preformed inflammatory mediators in degranulating mast cells)	Allergies: urticaria, pruritus Anaphylaxis: wheezing, hypotension, respiratory arrest, shock	 Donor plasma proteins, including IgA	 Host mast cell
Acute hemolytic transfusion reaction	Type II hypersensitivity reaction Typically causes intravascular hemolysis (ABO blood group incompatibility)	During transfusion or within 24 hr (due to preformed antibodies)	Fever, hypotension, tachypnea, tachycardia, flank pain, hemoglobinuria (intravascular), jaundice (extravascular)	 Donor RBC with A and/or B group antigens	 Host anti-A, anti-B IgG, IgM
Febrile nonhemolytic transfusion reaction	Cytokines created by donor WBCs accumulate during storage of blood products Reactions prevented by leukoreduction of blood products	Within 1-6 hr (due to preformed cytokines)	Fever, headaches, chills, flushing More common in children	 Donor WBC releases preformed cytokines	
Transfusion-related acute lung injury	Two-hit mechanism: ▪ Neutrophils are sequestered and primed in pulmonary vasculature due to recipient risk factors ▪ Neutrophils are activated by a product (eg, antileukocyte antibodies) in the transfused blood and release inflammatory mediators → ↑ capillary permeability → pulmonary edema	Within minutes to 6 hr	Respiratory distress, noncardiogenic pulmonary edema	 Donor antileukocyte IgG	 Host neutrophils
Delayed hemolytic transfusion reaction	Anamnestic response to a foreign antigen on donor RBCs (most commonly Rh or other minor blood group antigens) previously encountered by recipient Typically causes extravascular hemolysis	Onset over 24 hr Usually presents within 1-2 wk (due to slow destruction by reticuloendothelial system)	Generally self limited and clinically silent Mild fever, hyperbilirubinemia	 Donor RBC with foreign antigens	 Host IgG



Autoantibodies

AUTOANTIBODY	ASSOCIATED DISORDER
Anti-postsynaptic ACh receptor	Myasthenia gravis
Anti-presynaptic voltage-gated calcium channel	Lambert-Eaton myasthenic syndrome
Anti- β_2 glycoprotein I	Antiphospholipid syndrome
Antinuclear (ANA)	Nonspecific screening antibody, often associated with SLE
Anticardiolipin, lupus anticoagulant	SLE, antiphospholipid syndrome
Anti-dsDNA, anti-Smith	SLE
Antihistone	Drug-induced lupus
Anti-U1 RNP (ribonucleoprotein)	Mixed connective tissue disease
Rheumatoid factor (IgM antibody against IgG Fc region), anti-CCP (more specific)	Rheumatoid arthritis
Anti-Ro/SSA, anti-La/SSB	Sjögren syndrome
Anti-Scl-70 (anti-DNA topoisomerase I)	Scleroderma (diffuse)
Anticentromere	Limited scleroderma (CREST syndrome)
Antisynthetase (eg, anti-Jo-1), anti-SRP, anti-helicase (anti-Mi-2)	Polymyositis, dermatomyositis
Antimitochondrial	1° biliary cholangitis
Anti-smooth muscle	Autoimmune hepatitis type 1
MPO-ANCA/p-ANCA	Microscopic polyangiitis, eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome), ulcerative colitis
PR3-ANCA/c-ANCA	Granulomatosis with polyangiitis (Wegener)
Anti-phospholipase A ₂ receptor	1° membranous nephropathy
Anti-hemidesmosome	Bullous pemphigoid
Anti-desmoglein (anti-desmosome)	Pemphigus vulgaris
Antithyroglobulin, antithyroid peroxidase (antimicrosomal)	Hashimoto thyroiditis
Anti-TSH receptor	Graves disease
IgA anti-endomysial, IgA anti-tissue transglutaminase, IgA and IgG deamidated gliadin peptide	Celiac disease
Anti-glutamic acid decarboxylase, islet cell cytoplasmic antibodies	Type 1 diabetes mellitus
Antiparietal cell, anti-intrinsic factor	Pernicious anemia
Anti-glomerular basement membrane	Goodpasture syndrome

Immunodeficiencies

DISEASE	DEFECT	PRESENTATION	FINDINGS
B-cell disorders			
X-linked (Bruton) agammaglobulinemia	Defect in BTK , a tyrosine kinase gene → no B -cell maturation; X-linked recessive (↑ in B oys)	Recurrent bacterial and enteroviral infections after 6 months (↓ maternal IgG)	Absent B cells in peripheral blood, ↓ Ig of all classes. Absent/scanty lymph nodes and tonsils (1° follicles and germinal centers absent) → live vaccines contraindicated
Selective IgA deficiency	Cause unknown Most common 1° immunodeficiency	Majority A symptomatic Can see A irway and GI infections, A utoimmune disease, A topy, A naphylaxis to Ig A -containing products	↓ IgA with normal IgG, IgM levels ↑ susceptibility to giardiasis Can cause false-positive β-hCG test
Common variable immunodeficiency	Defect in B-cell differentiation. Cause unknown in most cases	May present in childhood but usually diagnosed after puberty ↑ risk of autoimmune disease, bronchiectasis, lymphoma, sinopulmonary infections	↓ plasma cells, ↓ immunoglobulins
T-cell disorders			
Thymic aplasia	22q11 microdeletion; failure to develop 3rd and 4th pharyngeal pouches → absent thymus and parathyroids DiGeorge syndrome —thymic, parathyroid, cardiac defects Velocardiofacial syndrome —palate, facial, cardiac defects	CATCH-22 : Cardiac defects (conotruncal abnormalities [eg, tetralogy of Fallot, truncus arteriosus]), A bnormal facies, T hymic hypoplasia → T-cell deficiency (recurrent viral/fungal infections), C left palate, H ypocalcemia 2° to parathyroid aplasia → tetany	↓ T cells, ↓ PTH, ↓ Ca ²⁺ Thymic shadow absent on CXR
IL-12 receptor deficiency	↓ Th1 response; autosomal recessive	Disseminated mycobacterial and fungal infections; may present after administration of BCG vaccine	↓ IFN-γ Most common cause of Mendelian susceptibility to mycobacterial diseases (MSMD)
Autosomal dominant hyper-IgE syndrome (Job syndrome)	Deficiency of Th17 cells due to STAT3 mutation → impaired recruitment of neutrophils to sites of infection	Cold (noninflamed) staphylococcal A bscesses, retained B aby teeth, C oarse facies, D ermatologic problems (eczema), ↑ IgE , bone F ractures from minor trauma	↑ IgE ↑ eosinophils Learn the ABCDEF 's to get a J ob!
Chronic mucocutaneous candidiasis	T-cell dysfunction Impaired cell-mediated immunity against <i>Candida</i> sp Classic form caused by defects in AIRE	Persistent noninvasive <i>Candida albicans</i> infections of skin and mucous membranes	Absent in vitro T-cell proliferation in response to <i>Candida</i> antigens Absent cutaneous reaction to <i>Candida</i> antigens

Immunodeficiencies (continued)




DISEASE	DEFECT	PRESENTATION	FINDINGS
B- and T-cell disorders			
Severe combined immunodeficiency	Several types including defective IL-2R gamma chain (most common, X-linked recessive); adenosine deaminase deficiency (autosomal recessive); RAG mutation → VDJ recombination defect	Failure to thrive, chronic diarrhea, thrush Recurrent viral, bacterial, fungal, and protozoal infections	↓ T-cell receptor excision circles (TRECs) Absence of thymic shadow (CXR), germinal centers (lymph node biopsy), and T cells (flow cytometry)
Ataxia-telangiectasia 	Defects in ATM gene → failure to detect DNA damage → failure to halt progression of cell cycle → mutations accumulate; autosomal recessive	Triad: cerebellar defects (A taxia), spider A ngiomas (telangiectasia A), IgA deficiency ↑↑ sensitivity to radiation (limit x-ray exposure)	↑ A FP ↓ IgA, IgG, and IgE Lymphopenia, cerebellar atrophy ↑ risk of lymphoma and leukemia
Hyper-IgM syndrome	Most commonly due to defective CD40L on Th cells → class switching defect; X-linked recessive	Severe pyogenic infections early in life; opportunistic infection with <i>Pneumocystis</i> , <i>Cryptosporidium</i> , CMV	Normal or ↑ IgM ↓↓ IgG, IgA, IgE Failure to make germinal centers
Wiskott-Aldrich syndrome	Mutation in WAS gene; leukocytes and platelets unable to reorganize actin cytoskeleton → defective antigen presentation; X-linked recessive	WATER : Wiskott-Aldrich: T hrombocytopenia, E czema, R ecurrent (pyogenic) infections ↑ risk of autoimmune disease and malignancy	↓ to normal IgG, IgM ↑ IgE, IgA Fewer and smaller platelets
Phagocyte dysfunction			
Leukocyte adhesion deficiency (type 1)	Defect in LFA-1 integrin (CD18) protein on phagocytes; impaired migration and chemotaxis; autosomal recessive	L ate separation (>30 days) of umbilical cord, a bsent pus, d ysfunctional neutrophils → recurrent skin and mucosal bacterial infections	↑ neutrophils in blood Absence of neutrophils at infection sites → impaired wound healing
Chédiak-Higashi syndrome 	Defect in lysosomal trafficking regulator gene (<i>LYST</i>) Microtubule dysfunction in phagosome-lysosome fusion; autosomal recessive	PLAIN : Progressive neurodegeneration, L ymphohistiocytosis, A lbinism (partial), recurrent pyogenic I nfections, peripheral N europathy	Giant granules (B , arrows) in granulocytes and platelets. Pancytopenia Mild coagulation defects
Chronic granulomatous disease	Defect of NADPH oxidase → ↓ reactive oxygen species (eg, superoxide) and ↓ respiratory burst in neutrophils; X-linked form most common	↑ susceptibility to catalase ⊕ organisms	Abnormal dihydrorhodamine (flow cytometry) test (↓ green fluorescence) Nitroblue tetrazolium dye reduction test (obsolete) fails to turn blue

Infections in immunodeficiency

PATHOGEN	↓ T CELLS	↓ B CELLS	↓ GRANULOCYTES	↓ COMPLEMENT
Bacteria	Sepsis	Encapsulated (P lease S HINE my S KiS): <i>Pseudomonas aeruginosa</i> , <i>Streptococcus pneumoniae</i> , <i>Haemophilus Influenzae</i> type b, <i>Neisseria meningitidis</i> , <i>Escherichia coli</i> , <i>Salmonella</i> , <i>Klebsiella pneumoniae</i> , Group B <i>Streptococcus</i>	Some B acteria Produce N o Serious granules: <i>Staphylococcus</i> , <i>Burkholderia cepacia</i> , <i>Pseudomonas aeruginosa</i> , <i>Nocardia</i> , <i>Serratia</i>	Encapsulated species with early complement deficiencies <i>Neisseria</i> with late complement (C5–C9) deficiencies
Viruses	CMV, EBV, JC virus, VZV, chronic infection with respiratory/GI viruses	Enteroviral encephalitis, poliovirus (live vaccine contraindicated)	N/A	N/A
Fungi/parasites	<i>Candida</i> (local), PCP, <i>Cryptococcus</i>	GI giardiasis (no IgA)	<i>Candida</i> (systemic), <i>Aspergillus</i> , <i>Mucor</i>	N/A

Note: **B**-cell deficiencies tend to produce recurrent **b**acterial infections, whereas T-cell deficiencies produce more fungal and viral infections.

Transplant rejection


TYPE OF REJECTION	ONSET	PATHOGENESIS	FEATURES
Hyperacute 	Within minutes	Pre-existing recipient antibodies react to donor antigen (type II hypersensitivity reaction), activate complement	Widespread thrombosis of graft vessels (arrows within glomerulus A) → ischemia/necrosis Graft must be removed
Acute 	Weeks to months	Cellular: CD8+ T cells and/or CD4+ T cells activated against donor MHCs (type IV hypersensitivity reaction) Humoral: similar to hyperacute, except antibodies develop after transplant	Vasculitis of graft vessels with dense interstitial lymphocytic infiltrate B Prevent/reverse with immunosuppressants
Chronic 	Months to years	CD4+ T cells respond to recipient APCs presenting donor peptides, including allogeneic MHC Both cellular and humoral components (type II and IV hypersensitivity reactions)	Recipient T cells react and secrete cytokines → proliferation of vascular smooth muscle, parenchymal atrophy, interstitial fibrosis Dominated by arteriosclerosis C Organ-specific examples: <ul style="list-style-type: none"> ▪ Chronic allograft nephropathy ▪ Bronchiolitis obliterans ▪ Accelerated atherosclerosis (heart) ▪ Vanishing bile duct syndrome
Graft-versus-host disease	Varies	Grafted immunocompetent T cells proliferate in the immunocompromised host and reject host cells with “foreign” proteins → severe organ dysfunction Type IV hypersensitivity reaction	Maculopapular rash, jaundice, diarrhea, hepatosplenomegaly Usually in bone marrow and liver transplants (rich in lymphocytes) Potentially beneficial in bone marrow transplant for leukemia (graft-versus-tumor effect) For immunocompromised patients, irradiate blood products prior to transfusion to prevent GVHD

► IMMUNOLOGY—IMMUNOSUPPRESSANTS

Immunosuppressants Agents that block lymphocyte activation and proliferation. Reduce acute transplant rejection by suppressing cellular immunity (used as prophylaxis). Frequently combined to achieve greater efficacy with ↓ toxicity. Chronic suppression ↑ risk of infection and malignancy.

DRUG	MECHANISM	INDICATIONS	TOXICITY	NOTES
Cyclosporine	Calcineurin inhibitor; binds cyclophilin Blocks T-cell activation by preventing IL-2 transcription	Psoriasis, rheumatoid arthritis	Nephrotoxicity , hypertension, hyperlipidemia, neurotoxicity, gingival hyperplasia, hirsutism	Both calcineurin inhibitors are highly nephrotoxic, especially in higher doses or in patients with decreased renal function
Tacrolimus (FK506)	Calcineurin inhibitor; binds FK506 binding protein (FKBP) Blocks T-cell activation by preventing IL-2 transcription		Similar to cyclosporine, ↑ risk of diabetes and neurotoxicity; no gingival hyperplasia or hirsutism	
Sirolimus (Rapamycin)	mTOR inhibitor; binds FKBP Blocks T-cell activation and B-cell differentiation by preventing response to IL-2	Kidney transplant rejection prophylaxis specifically Sir Basil's kidney transplant	“Pan Sirt openia” (pancytopenia), insulin resistance, hyperlipidemia; not nephrotoxic	Kidney “ sir -vives.” Synergistic with cyclosporine Also used in drug-eluting stents
Basiliximab	Monoclonal antibody; blocks IL-2R		Edema, hypertension, tremor	
Azathioprine	Antimetabolite precursor of 6-mercaptopurine Inhibits lymphocyte proliferation by blocking nucleotide synthesis	Rheumatoid arthritis, Crohn disease, glomerulonephritis, other autoimmune conditions	Pancytopenia	6-MP degraded by xanthine oxidase; toxicity ↑ by allopurinol Pronounce “azathio- purine ”
Mycophenolate Mofetil	Reversibly inhibits IMP dehydrogenase, preventing purine synthesis of B and T cells	Lupus nephritis	GI upset, pancytopenia, hypertension, hyperglycemia Less nephrotoxic and neurotoxic	Associated with invasive CMV infection
Glucocorticoids	Inhibit NF-κB Suppress both B- and T-cell function by ↓ transcription of many cytokines Induce T cell apoptosis	Many autoimmune and inflammatory disorders, adrenal insufficiency, asthma, CLL, non-Hodgkin lymphoma	Cushing syndrome, osteoporosis, hyperglycemia, diabetes, amenorrhea, adrenocortical atrophy, peptic ulcers, psychosis, cataracts, avascular necrosis (femoral head)	Demargination of WBCs causes artificial leukocytosis Adrenal insufficiency may develop if drug is stopped abruptly after chronic use

Immunosuppression targets



Recombinant cytokines and clinical uses

CYTOKINE	AGENT	CLINICAL USES
Bone marrow stimulation		
Erythropoietin	Epoetin alfa (EPO analog)	Anemias (especially in renal failure)
Colony stimulating factors	Filgrastim (G -CSF), Sargramostim (GM -CSF)	Leukopenia; recovery of granulocyte and monocyte counts
Thrombopoietin	Romiplostim (TPO analog), eltrombopag (TPO receptor agonist)	Autoimmune thrombocytopenia Platelet stimulator
Immunotherapy		
Interleukin-2	Aldesleukin	Renal cell carcinoma, metastatic melanoma
Interferons	IFN- α	Chronic hepatitis C (not preferred) and B, renal cell carcinoma
	IFN- β	Multiple sclerosis
	IFN- γ	Chronic granulomatous disease

Therapeutic antibodies

AGENT	TARGET	CLINICAL USE	NOTES
Cancer therapy			
Alemtuzumab	CD52	CLL, multiple sclerosis	“ Alymtuzumab ”—chronic lymphocytic leukemia
Bevacizumab	VEGF	Colorectal cancer, renal cell carcinoma, non-small cell lung cancer	Also used for neovascular age-related macular degeneration, proliferative diabetic retinopathy, and macular edema
Rituximab	CD20	B-cell non-Hodgkin lymphoma, CLL, rheumatoid arthritis, ITP, multiple sclerosis	Risk of PML in patients with JC virus CD20—“ ri2ximab ”
Trastuzumab	HER2	Breast cancer, gastric cancer	HER2—“ tras2zumab ”
Autoimmune disease therapy			
Adalimumab, infliximab	Soluble TNF- α	IBD, rheumatoid arthritis, ankylosing spondylitis, psoriasis	Etanercept is a decoy TNF- α receptor and not a monoclonal antibody
Eculizumab	Complement protein C5	Paroxysmal nocturnal hemoglobinuria	
Ixekizumab, secukinumab	IL-17A	Psoriasis, psoriatic arthritis	
Natalizumab	α 4-integrin	Multiple sclerosis, Crohn disease	α 4-integrin: WBC adhesion Risk of PML in patients with JC virus
Ustekinumab	IL-12/IL-23	Psoriasis, psoriatic arthritis	
Other applications			
Abciximab	Platelet glycoproteins IIb/IIIa	Antiplatelet agent for prevention of ischemic complications in patients undergoing percutaneous coronary intervention	ABC is as easy as 123
Denosumab	RANKL	Osteoporosis; inhibits osteoclast maturation (mimics osteoprotegerin)	Denosumab helps make dense bones
Omalizumab	IgE	Refractory allergic asthma; prevents IgE binding to Fc ϵ RI	
Palivizumab	RSV F protein	RSV prophylaxis for high-risk infants	Pali VI zumab— VI rus

HIGH-YIELD PRINCIPLES IN

Microbiology

“Support bacteria. They’re the only culture some people have.”

—Steven Wright

“What lies behind us and what lies ahead of us are tiny matters compared to what lies within us.”

—Henry S. Haskins

“Infectious disease is merely a disagreeable instance of a widely prevalent tendency of all living creatures to save themselves the bother of building, by their own efforts, the things they require.”

—Hans Zinsser

Microbiology questions on the Step 1 exam often require two (or more) steps: Given a certain clinical presentation, you will first need to identify the most likely causative organism, and you will then need to provide an answer regarding some features of that organism or relevant antimicrobial agents. For example, a description of a child with fever and a petechial rash will be followed by a question that reads, “From what site does the responsible organism usually enter the blood?”


This section therefore presents organisms in two major ways: in individual microbial “profiles” and in the context of the systems they infect and the clinical presentations they produce. You should become familiar with both formats. When reviewing the systems approach, remind yourself of the features of each microbe by returning to the individual profiles. Also be sure to memorize the laboratory characteristics that allow you to identify microbes.

▶ Basic Bacteriology	124
▶ Clinical Bacteriology	134
▶ Mycology	151
▶ Parasitology	155
▶ Virology	162
▶ Systems	178
▶ Antimicrobials	187

▶ MICROBIOLOGY—BASIC BACTERIOLOGY


Bacterial structures

STRUCTURE	CHEMICAL COMPOSITION	FUNCTION
Appendages		
Flagellum	Proteins	Motility
Pilus/fimbria	Glycoprotein	Mediate adherence of bacteria to cell surface; sex pilus forms during conjugation
Specialized structures		
Spore	Keratin-like coat; dipicolinic acid; peptidoglycan, DNA	Gram ⊕ only Survival: resist dehydration, heat, chemicals
Cell envelope		
Capsule	Discrete layer usually made of polysaccharides (and rarely proteins)	Protects against phagocytosis
Slime (S) layer	Loose network of polysaccharides	Mediates adherence to surfaces, especially foreign surfaces (eg, indwelling catheters)
Outer membrane	Outer leaflet: contains endotoxin (LPS/LOS) Embedded proteins: porins and other outer membrane proteins (OMPs) Inner leaflet: phospholipids	Gram ⊖ only Endotoxin: lipid A induces TNF and IL-1; antigenic O polysaccharide component Most OMPs are antigenic Porins: transport across outer membrane
Periplasm	Space between cytoplasmic membrane and outer membrane in gram ⊖ bacterial (peptidoglycan in middle)	Accumulates components exiting gram ⊖ cells, including hydrolytic enzymes (eg, β-lactamases)
Cell wall	Peptidoglycan is a sugar backbone with peptide side chains cross-linked by transpeptidase	Net-like structure gives rigid support, protects against osmotic pressure damage
Cytoplasmic membrane	Phospholipid bilayer sac with embedded proteins (eg, penicillin-binding proteins [PBPs]) and other enzymes Lipoteichoic acids (gram positive) only extend from membrane to exterior	Site of oxidative and transport enzymes; PBPs involved in cell wall synthesis Lipoteichoic acids induce TNF-α and IL-1

Cell envelope

Stains

Gram stain	First-line lab test in bacterial identification. Bacteria with thick peptidoglycan layer retain crystal violet dye (gram ⊕); bacteria with thin peptidoglycan layer turn red or pink (gram ⊖) with counterstain. These bugs do not Gram stain well (T hese L ittle M icrobes M ay U nfortunately L ack R eal C olor B ut A re E verywhere):	
	<i>Treponema</i> , <i>Leptospira</i>	Too thin to be visualized
	<i>Mycobacteria</i>	Cell wall has high lipid content
	<i>Mycoplasma</i> , <i>Ureaplasma</i>	No cell wall
	<i>Legionella</i> , <i>Rickettsia</i> , <i>Chlamydia</i> , <i>Bartonella</i> , <i>Anaplasma</i> , <i>Ehrlichia</i>	Primarily intracellular; also, <i>Chlamydia</i> lack classic peptidoglycan because of ↓ muramic acid
Giemsa stain	<i>Rickettsia</i> , <i>Chlamydia</i> , Trypanosomes A , <i>Plasmodium</i> , <i>Borrelia</i> , <i>Helicobacter pylori</i>	R icky got <i>Chlamydia</i> as he T ried to P lease the B ored H ot “ G eisha”
Periodic acid–Schiff stain	Stains glycogen , mucopolysaccharides; used to diagnose Whipple disease (<i>Tropheryma whipplei</i> B)	P aSs the s ugar
Ziehl-Neelsen stain (carbol fuchsin)	Acid-fast bacteria (eg, <i>Mycobacteria</i> C , <i>Nocardia</i> ; stains mycolic acid in cell wall); protozoa (eg, <i>Cryptosporidium</i> oocysts)	Auramine-rhodamine stain is more often used for screening (inexpensive, more sensitive)
India ink stain	<i>Cryptococcus neoformans</i> D ; mucicarmine can also be used to stain thick polysaccharide capsule red	
Silver stain	Fungi (eg, <i>Coccidioides</i> E , <i>Pneumocystis jirovecii</i>), <i>Legionella</i> , <i>Helicobacter pylori</i>	
Fluorescent antibody stain	Used to identify many bacteria, viruses, <i>Pneumocystis jirovecii</i> , <i>Giardia</i> , and <i>Cryptosporidium</i>	Example is FTA-ABS for syphilis



Properties of growth media

The same type of media can possess both (or neither) of these properties.

Selective media

Favors the growth of particular organism while preventing growth of other organisms. Example: Thayer-Martin agar contains antibiotics that allow the selective growth of *Neisseria* by inhibiting the growth of other sensitive organisms.

Indicator (differential) media

Yields a color change in response to the metabolism of certain organisms. Example: MacConkey agar contains a pH indicator; a lactose fermenter like *E coli* will convert lactose to acidic metabolites → color change to pink.

Special culture requirements

BUG	MEDIA USED FOR ISOLATION	MEDIA CONTENTS/OTHER
<i>H influenzae</i>	Chocolate agar	Factors V (NAD ⁺) and X (hematin)
<i>N gonorrhoeae</i> , <i>N meningitidis</i>	Thayer-Martin agar	Selectively favors growth of <i>Neisseria</i> by inhibiting growth of gram ⊕ organisms with V ancomycin, gram ⊖ organisms except <i>Neisseria</i> with T rimethoprim and C olistin, and fungi with N ystatin Very Typically Cultures Neisseria
<i>B pertussis</i>	Bordet-Gengou agar (Bordet for <i>Bordetella</i>) Regan-Lowe medium	Potato extract Charcoal, blood, and antibiotic
<i>C diphtheriae</i>	Tellurite agar, Löffler medium	
<i>M tuberculosis</i>	Löwenstein-Jensen medium, Middlebrook medium, rapid automated broth cultures	
<i>M pneumoniae</i>	Eaton agar	Requires cholesterol
Lactose-fermenting enterics	MacConkey agar	Fermentation produces acid, causing colonies to turn pink
<i>E coli</i>	Eosin–methylene blue (EMB) agar	Colonies with green metallic sheen
<i>Brucella</i> , <i>Francisella</i> , <i>Legionella</i> , <i>Pasteurella</i>	Charcoal yeast extract agar buffered with cysteine and iron	The Ella siblings, Bruce , Francis , a legionnaire , and a pasteur (pastor), built the Sistine (cysteine) chapel out of charcoal and iron .
Fungi	Sabouraud agar	“ Sab ’s a fun guy!”

Aerobes

Use an O₂-dependent system to generate ATP.


Examples include *Nocardia*, *Pseudomonas aeruginosa*, *Mycobacterium tuberculosis*, and *Bordetella pertussis*.

Reactivation of *M tuberculosis* (eg, after immunocompromise or TNF- α inhibitor use) has a predilection for the apices of the lung.

Anaerobes	Examples include <i>Clostridium</i> , <i>Bacteroides</i> , <i>Fusobacterium</i> , and <i>Actinomyces israelii</i> . They lack catalase and/or superoxide dismutase and are thus susceptible to oxidative damage. Generally foul smelling (short-chain fatty acids), are difficult to culture, and produce gas in tissue (CO ₂ and H ₂).	Anaerobes Can't Breathe Fresh Air . Anaerobes are normal flora in GI tract, typically pathogenic elsewhere. AminO ₂ glycosides are ineffective against anaerobes because these antibiotics require O ₂ to enter into bacterial cell.
Facultative anaerobes	May use O ₂ as a terminal electron acceptor to generate ATP, but can also use fermentation and other O ₂ -independent pathways.	Streptococci, staphylococci, and enteric gram ⊖ bacteria.

Intracellular bacteria

Obligate intracellular	<i>Rickettsia</i> , <i>Chlamydia</i> , <i>Coxiella</i> Rely on host ATP	Stay inside (cells) when it is Really Chilly and Cold
Facultative intracellular	<i>Salmonella</i> , <i>Neisseria</i> , <i>Brucella</i> , <i>Mycobacterium</i> , <i>Listeria</i> , <i>Francisella</i> , <i>Legionella</i> , <i>Yersinia pestis</i>	Some Nasty Bugs May Live FacultativeLY

Encapsulated bacteria

Examples are *Pseudomonas aeruginosa*, *Streptococcus pneumoniae* **A**, *Haemophilus influenzae* type b, *Neisseria meningitidis*, *Escherichia coli*, *Salmonella*, *Klebsiella pneumoniae*, and group B **Strep**. Their capsules serve as an antiphagocytic virulence factor.
Capsular polysaccharide + protein conjugate serves as an antigen in vaccines.

Please **SHiNE** my **SKiS**.

Are opsonized, and then cleared by spleen. Asplenic (**No Spleen Here**) have ↓ opsonizing ability and thus ↑ risk for severe infections; need vaccines to protect against:

- *N meningitidis*
- *S pneumoniae*
- *H influenzae*

Encapsulated bacteria vaccines


Some vaccines containing polysaccharide capsule antigens are conjugated to a carrier protein, enhancing immunogenicity by promoting T-cell activation and subsequent class switching. A polysaccharide antigen alone cannot be presented to T cells.

Pneumococcal vaccines: PCV13 (pneumococcal conjugate vaccine), PPSV23 (pneumococcal polysaccharide vaccine with no conjugated protein).
H influenzae type b (conjugate vaccine).
Meningococcal vaccine (conjugate vaccine).

Urease-positive organisms

Proteus, *Cryptococcus*, *H pylori*, *Ureaplasma*, *Nocardia*, *Klebsiella*, *S epidermidis*, *S saprophyticus*. Urease hydrolyzes urea to release ammonia and CO₂ → ↑ pH. Predisposes to struvite (ammonium magnesium phosphate) stones, particularly *Proteus*.

Pee **CHUNKSS**.

Catalase-positive organisms

Catalase degrades H_2O_2 into H_2O and bubbles of O_2 **A** before it can be converted to microbicidal products by the enzyme myeloperoxidase. People with chronic granulomatous disease (NADPH oxidase deficiency) have recurrent infections with certain catalase \oplus organisms.

Examples: *Nocardia*, *Staphylococci*, *Serratia*, *Candida*, *Listeria*, *E coli*, *Burkholderia cepacia*, *Pseudomonas*, *Aspergillus*, *Helicobacter pylori*, *Bordetella pertussis*.

Pigment-producing bacteria

Actinomyces israelii—**yellow** “sulfur” **granules**, which are composed of filaments of bacteria

Israel has **yellow sand**

S aureus—**yellow** pigment

Aureus (Latin) = **gold**

P aeruginosa—**blue-green** pigment (pyocyanin and pyoverdin)

Aerugula is **green**

Serratia marcescens—**red** pigment

Think **red Sriracha** hot sauce

In vivo biofilm-producing bacteria

S epidermidis

Catheter and prosthetic device infections

Viridans streptococci (*S mutans*, *S sanguinis*)


Dental plaques, infective endocarditis

P aeruginosa

Respiratory tree colonization in patients with cystic fibrosis, ventilator-associated pneumonia
Contact lens-associated keratitis

Nontypeable (unencapsulated) *H influenzae*

Otitis media

Spore-forming bacteria

Some gram \oplus bacteria can form spores **A** when nutrients are limited. Spores lack metabolic activity and are highly resistant to heat and chemicals. Core contains dipicolinic acid. Must autoclave to kill spores (as is done to surgical equipment) by steaming at 121°C for 15 minutes.

Examples: *B anthracis* (anthrax), *B cereus* (food poisoning), *C botulinum* (botulism), *C difficile* (pseudomembranous colitis), *C perfringens* (gas gangrene), *C tetani* (tetanus).

Bacterial virulence factors

These promote evasion of host immune response.

Protein A

Binds Fc region of IgG. Prevents opsonization and phagocytosis. Expressed by *S aureus*.

IgA protease

Enzyme that cleaves IgA, allowing bacteria to adhere to and colonize mucous membranes. Secreted by *S pneumoniae*, *H influenzae* type b, and *Neisseria* (**SHiN**).


M protein

Helps prevent phagocytosis. Expressed by group A streptococci. Shares similar epitopes to human cellular proteins (**m**olecular **m**imicry); possibly underlies the autoimmune response seen in acute rheumatic fever.


Bacterial genetics**Transformation**

Competent bacteria can bind and import short pieces of environmental naked bacterial chromosomal DNA (from bacterial cell lysis). The transfer and expression of newly transferred genes is called transformation. A feature of many bacteria, especially *S pneumoniae*, *H influenzae* type b, and *Neisseria* (SHiN).


Adding deoxyribonuclease degrades naked DNA, preventing transformation.

**Conjugation** $F^+ \times F^-$


F^+ plasmid contains genes required for sex pilus and conjugation. Bacteria without this plasmid are termed F^- . Sex pilus on F^+ bacterium contacts F^- bacterium. A single strand of plasmid DNA is transferred across the conjugal bridge (“mating bridge”). No transfer of chromosomal DNA.

 $Hfr \times F^-$


F^+ plasmid can become incorporated into bacterial chromosomal DNA, termed high-frequency recombination (Hfr) cell. Transfer of leading part of plasmid and a few flanking chromosomal genes. High-frequency recombination may integrate some of those bacterial genes. Recipient cell remains F^- but now may have new bacterial genes.

**Transduction****Generalized**

A packaging “error.” Lytic phage infects bacterium, leading to cleavage of bacterial DNA. Parts of bacterial chromosomal DNA may become packaged in phage capsid. Phage infects another bacterium, transferring these genes.

**Specialized**


An “excision” event. Lysogenic phage infects bacterium; viral DNA incorporates into bacterial chromosome. When phage DNA is excised, flanking bacterial genes may be excised with it. DNA is packaged into phage capsid and can infect another bacterium. Genes for the following 5 bacterial toxins are encoded in a lysogenic phage (ABCD'S): Group A strep erythrogenic toxin, Botulinum toxin, Cholera toxin, Diphtheria toxin, Shiga toxin.



Bacterial genetics (continued)


Transposition

A “jumping” process involving a transposon (specialized segment of DNA), which can copy and excise itself and then insert into the same DNA molecule or an unrelated DNA (eg, plasmid or chromosome). Critical in creating plasmids with multiple drug resistance and transfer across species lines (eg, Tn1546 with *vanA* from *Enterococcus* to *S aureus*).



Main features of exotoxins and endotoxins

	Exotoxins	Endotoxins
SOURCE	Certain species of gram ⊕ and gram ⊖ bacteria	Outer cell membrane of most gram ⊖ bacteria
SECRETED FROM CELL	Yes	No
CHEMISTRY	Polypeptide	Lipid A component of LPS (structural part of bacteria; released when lysed)
LOCATION OF GENES	Plasmid or bacteriophage	Bacterial chromosome
ADVERSE EFFECTS	High (fatal dose on the order of 1 μg)	Low (fatal dose on the order of hundreds of micrograms)
CLINICAL EFFECTS	Various effects (see following pages)	Fever, shock (hypotension), DIC
MODE OF ACTION	Various modes (see following pages)	Induces TNF, IL-1, and IL-6
ANTIGENICITY	Induces high-titer antibodies called antitoxins	Poorly antigenic
VACCINES	Toxoids used as vaccines	No toxoids formed and no vaccine available
HEAT STABILITY	Destroyed rapidly at 60°C (except staphylococcal enterotoxin and <i>E coli</i> heat-stable toxin)	Stable at 100°C for 1 hr
TYPICAL DISEASES	Tetanus, botulism, diphtheria, cholera	Meningococemia; sepsis by gram ⊖ rods



Bacteria with exotoxins

BACTERIA	TOXIN	MECHANISM	MANIFESTATION
Inhibit protein synthesis			
<i>Corynebacterium diphtheriae</i>	Diphtheria toxin ^a	Inactivate elongation factor (EF-2)	Pharyngitis with pseudomembranes in throat and severe lymphadenopathy (bull neck), myocarditis
<i>Pseudomonas aeruginosa</i>	Exotoxin A ^a		Host cell death
<i>Shigella</i> spp	Shiga toxin (ST) ^a	Inactivate 60S ribosome by removing adenine from rRNA	GI mucosal damage → dysentery; ST also enhances cytokine release, causing hemolytic-uremic syndrome (HUS)
Enterohemorrhagic <i>E coli</i>	Shiga-like toxin (SLT) ^a		SLT enhances cytokine release, causing HUS (prototypically in EHEC serotype O157:H7) Unlike <i>Shigella</i> , EHEC does not invade host cells
Increase fluid secretion			
Enterotoxigenic <i>E coli</i>	Heat- labile toxin (LT) ^a	Overactivates adenylate cyclase (↑ cAMP) → ↑ Cl ⁻ secretion in gut and H ₂ O efflux	Watery diarrhea: “ labile in the A ir (A denylate cyclase), stable on the G round (G uanylate cyclase)”
	Heat- stable toxin (ST)	Overactivates guanylate cyclase (↑ cGMP) → ↓ resorption of NaCl and H ₂ O in gut	
<i>Bacillus anthracis</i>	Anthrax toxin ^a	Mimics adenylate cyclase (↑ cAMP)	Likely responsible for characteristic edematous borders of black eschar in cutaneous anthrax
<i>Vibrio cholerae</i>	Cholera toxin ^a	Overactivates adenylate cyclase (↑ cAMP) by permanently activating G _s → ↑ Cl ⁻ secretion in gut and H ₂ O efflux	Voluminous “rice-water” diarrhea
Inhibit phagocytic ability			
<i>Bordetella pertussis</i>	Pertussis toxin ^a	Inactivates inhibitory G subunit (G _i) → activation of adenylate cyclase → ↑ cAMP	Whooping cough —child coughs on expiration and “whoops” on inspiration (toxin may not actually be a cause of cough; can cause “100-day cough” in adults)
Inhibit release of neurotransmitter			
<i>Clostridium tetani</i>	Tetanospasmin ^a	Both are proteases that cleave SNARE (soluble NSF attachment protein receptor), a set of proteins required for neurotransmitter release via vesicular fusion	Toxin prevents release of inhibitory (GABA and glycine) neurotransmitters from Renshaw cells in spinal cord → spastic paralysis, risus sardonicus, trismus (lockjaw)
<i>Clostridium botulinum</i>	Botulinum toxin ^a		Toxin prevents release of stimulatory (ACh) signals at neuromuscular junction → flaccid paralysis (floppy baby)

^aAn AB toxin (aka, two-component toxin [or three for anthrax]) with **B** enabling **b**inding and triggering uptake (endocytosis) of the **a**ctive **A** component. The A components are usually ADP ribosyltransferases; others have enzymatic activities as listed in chart.

Bacteria with exotoxins (continued)

BACTERIA	TOXIN	MECHANISM	MANIFESTATION
Lyse cell membranes			
<i>Clostridium perfringens</i>	Alpha toxin	Phospholipase (lecithinase) that degrades tissue and cell membranes	Degradation of phospholipids → myonecrosis (“gas gangrene”) and hemolysis (“double zone” of hemolysis on blood agar)
<i>Streptococcus pyogenes</i>	Streptolysin O	Protein that degrades cell membrane	Lyses RBCs; contributes to β-hemolysis; host antibodies against toxin (ASO) used to diagnose rheumatic fever (do not confuse with immune complexes of poststreptococcal glomerulonephritis)
Superantigens causing shock			
<i>Staphylococcus aureus</i>	Toxic shock syndrome toxin (TSST-1)	Cross-links β region of TCR to MHC class II on APCs outside of the antigen binding site	Toxic shock syndrome: fever, rash, shock; other toxins cause scalded skin syndrome (exfoliative toxin) and food poisoning (heat-stable enterotoxin)
<i>Streptococcus pyogenes</i>	Erythrogenic exotoxin A	→ overwhelming release of IL-1, IL-2, IFN-γ, and TNF-α → shock	Toxic shock–like syndrome: fever, rash, shock; scarlet fever

Endotoxin


LPS found in outer membrane of gram ⊖ bacteria (both cocci and rods). Composed of O antigen + core polysaccharide + lipid A (the toxic component).

Released upon cell lysis or by living cells by blebs detaching from outer surface membrane (vs exotoxin, which is actively secreted).

Three main effects: macrophage activation (TLR4/CD14), complement activation, and tissue factor activation.


ENDOTOXINS:

Edema
Nitric oxide
DIC/Death
Outer membrane
TNF-α
O-antigen + core polysaccharide + lipid A
eXtremely heat stable
IL-1 and **IL-6**
Neutrophil chemotaxis
Shock



▶ MICROBIOLOGY—CLINICAL BACTERIOLOGY

Gram-positive lab algorithm




Important **tests** are in **bold**. Important **pathogens** are in **bold italics**.
 Note: Enterococcus is either α - or γ -hemolytic.



Gram-positive cocci antibiotic tests

Staphylococci	Novobiocin— <i>Saprophyticus</i> is Resistant ; <i>Epidermidis</i> is Sensitive	On the office’s “ staph ” retreat, there was no stress
Streptococci	Optochin— <i>Viridans</i> is Resistant ; <i>Pneumoniae</i> is Sensitive	OVRPS (overpass)
	Bacitracin—group B strep are Resistant ; group A strep are Sensitive	B-BRAS

α-hemolytic bacteria


Gram ⊕ cocci. Partial oxidation of hemoglobin causes greenish or brownish color without clearing around growth on blood agar **A**. Include the following organisms:

- *Streptococcus pneumoniae* (catalase ⊖ and optochin sensitive)
- Viridans streptococci (catalase ⊖ and optochin resistant)

β-hemolytic bacteria

Gram ⊕ cocci. Complete lysis of RBCs → pale/clear area surrounding colony on blood agar **A**. Include the following organisms:

- *Staphylococcus aureus* (catalase and coagulase ⊕)
- *Streptococcus pyogenes*—group A strep (catalase ⊖ and bacitracin sensitive)
- *Streptococcus agalactiae*—group B strep (catalase ⊖ and bacitracin resistant)

Staphylococcus aureus

Gram ⊕, β-hemolytic, catalase ⊕, coagulase ⊕ cocci in clusters **A**. Protein A (virulence factor) binds Fc-IgG, inhibiting complement activation and phagocytosis. Commonly colonizes the nares, ears, axilla, and groin.

Causes:

- Inflammatory disease—skin infections, organ abscesses, pneumonia (often after influenza virus infection), endocarditis, septic arthritis, and osteomyelitis.
- Toxin-mediated disease—toxic shock syndrome (TSST-1), scalded skin syndrome (exfoliative toxin), rapid-onset food poisoning (enterotoxins).

MRSA (methicillin-resistant *S aureus*)—important cause of serious nosocomial and community-acquired infections; resistance due to altered penicillin-binding protein. *mecA* gene from staphylococcal chromosomal cassette involved in penicillin resistance.

TSST-1 is a superantigen that binds to MHC II and T-cell receptor, resulting in polyclonal T-cell activation and cytokine release.

Staphylococcal toxic shock syndrome (TSS)—fever, vomiting, rash, desquamation, shock, end-organ failure. TSS results in ↑ AST, ↑ ALT, ↑ bilirubin. Associated with prolonged use of vaginal tampons or nasal packing.

Compare with *Streptococcus pyogenes* TSS (a toxic shock-like syndrome associated with painful skin infection).

S aureus food poisoning due to ingestion of preformed toxin → short incubation period (2–6 hr) followed by nonbloody diarrhea and emesis. Enterotoxin is heat stable → not destroyed by cooking.

S aureus makes coagulase and toxins. Forms fibrin clot around itself → abscess.

Staphylococcus epidermidis


Gram ⊕, catalase ⊕, coagulase ⊖, urease ⊕ cocci in clusters. Novobiocin sensitive. Does not ferment mannitol (vs *S aureus*).

Normal flora of skin; contaminates blood cultures.

Infects prosthetic devices (eg, hip implant, heart valve) and IV catheters by producing adherent biofilms.

Staphylococcus saprophyticus

Gram \oplus , catalase \oplus , coagulase \ominus , urease \oplus cocci in clusters. Novobiocin resistant. Normal flora of female genital tract and perineum. Second most common cause of uncomplicated UTI in young women (most common is *E coli*).

Streptococcus pneumoniae

Gram \oplus , α -hemolytic, lancet-shaped diplococci **A**. Encapsulated. IgA protease. Optochin sensitive and bile soluble. Most commonly causes:

- Meningitis
- Otitis media (in children)
- Pneumonia
- Sinusitis

Pneumococcus is associated with “rusty” sputum, sepsis in patients with sickle cell disease, and asplenic patients. No virulence without capsule.

Viridans group streptococci


Gram \oplus , α -hemolytic cocci. Optochin resistant and bile insoluble. Normal flora of the oropharynx.

Streptococcus mutans and *S mitis* cause dental caries.

S sanguinis makes dextrans that bind to fibrin-platelet aggregates on damaged **heart** valves, causing subacute bacterial endocarditis.

Viridans group strep live in the mouth, because they are not afraid **of-the-chin** (**op-to-chin** resistant).

Sanguinis = **blood**. Think, “there is lots of **blood** in the **heart**” (endocarditis).

***Streptococcus pyogenes* (group A streptococci)**

Gram \oplus cocci in chains **A**. Group A strep cause:

- Pyogenic—pharyngitis, cellulitis, impetigo (“honey-crusted” lesions), erysipelas
- Toxigenic—scarlet fever, toxic shock-like syndrome, necrotizing fasciitis
- Immunologic—rheumatic fever, glomerulonephritis

Bacitracin sensitive, β -hemolytic, pyrrolidonyl arylamidase (PYR) \oplus . Hyaluronic acid capsule and M protein inhibit phagocytosis. Antibodies to M protein enhance host defenses against *S pyogenes* but can give rise to rheumatic fever. ASO titer or anti-DNase B antibodies indicate recent *S pyogenes* infection.

“**Ph**”yogenes **ph**aryngitis can result in rheumatic “**p**hever” and glomerulone**ph**ritis. Strains causing impetigo can induce glomerulonephritis.

Scarlet fever—blanching, sandpaper-like body rash, strawberry tongue, and circumoral pallor in the setting of group A streptococcal pharyngitis (erythrogenic toxin \oplus).

***Streptococcus agalactiae* (group B streptococci)**

Gram ⊕ cocci, bacitracin resistant, β-hemolytic, Group **B** for **B**abies!
 colonizes vagina; causes pneumonia, meningitis, and sepsis, mainly in **babies**.
 Produces CAMP factor, which enlarges the area of hemolysis formed by *S aureus*. (Note: CAMP stands for the authors of the test, not cyclic AMP.) Hippurate test ⊕. PYR ⊖.
 Screen pregnant women at 35–37 weeks of gestation with rectal and vaginal swabs.
 Patients with ⊕ culture receive intrapartum penicillin/ampicillin prophylaxis.

Streptococcus bovis

Gram ⊕ cocci, colonizes the gut. *S gallolyticus* (*S bovis* biotype 1) can cause bacteremia and subacute endocarditis and is associated with colon cancer. **B**ovis in the **b**lood = **c**ancer in the **c**olon.

Enterococci

Gram ⊕ cocci. Enterococci (*E faecalis* and *E faecium*) are normal colonic flora that are penicillin G resistant and cause UTI, biliary tract infections, and subacute endocarditis (following GI/GU procedures). Catalase ⊖, PYR ⊕, variable hemolysis.
 VRE (vancomycin-resistant enterococci) are an important cause of nosocomial infection.

Enterococci are more resilient than streptococci, can grow in 6.5% NaCl and bile (lab test).

Entero = intestine, *faecalis* = feces, *strepto* = twisted (chains), *coccus* = berry.

Bacillus anthracis

Gram ⊕, spore-forming rod that produces anthrax toxin (an exotoxin consisting of protective antigen, lethal factor, and edema factor). Has a polypeptide capsule (poly D-glutamate). Colonies show a halo of projections, sometimes referred to as “medusa head” appearance.

Cutaneous anthrax

Painless papule surrounded by vesicles → ulcer with black eschar **A** (painless, necrotic) → uncommonly progresses to bacteremia and death.

**Pulmonary anthrax**

Inhalation of spores, most commonly from contaminated animals or animal products, although also a potential bioweapon → flu-like symptoms that rapidly progress to fever, pulmonary hemorrhage, mediastinitis (CXR may show widened mediastinum), and shock. Also called woolsorter's disease.

Bacillus cereus

Gram ⊕ rod. Causes food poisoning. Spores survive cooking rice (reheated rice syndrome).

Keeping rice warm results in germination of spores and enterotoxin formation.

Emetic type causes nausea and vomiting within 1–5 hours. Caused by cereulide, a preformed toxin.

Diarrheal type causes watery, nonbloody diarrhea and GI pain within 8–18 hours.

Management: supportive care (antibiotics are ineffective against toxins).

Clostridia

Gram ⊕, spore-forming, obligate anaerobic rods. Tetanus toxin and botulinum toxin are proteases that cleave SNARE proteins involved in neurotransmission.

Clostridium tetani

Produces tetanospasmin, an exotoxin causing tetanus. Tetanospasmin blocks release of GABA and glycine from Renshaw cells in spinal cord.

Causes **spastic** paralysis, trismus (lockjaw), risus sardonicus (raised eyebrows and open grin), opisthotonos (spasms of spinal extensors).

Prevent with tetanus vaccine. Treat with antitoxin +/- vaccine booster, antibiotics, diazepam (for muscle spasms), and wound debridement.

Tetanus is **tetanic** paralysis.

Clostridium botulinum

Produces a heat-labile toxin that inhibits ACh release at the neuromuscular junction, causing botulism. In adults, disease is caused by ingestion of preformed toxin. In babies, ingestion of spores (eg, in honey) leads to disease (**floppy** baby syndrome). Treat with human botulinum immunoglobulin.


Symptoms of botulism (the **4 D's**): **D**iplopia, **D**ysarthria, **D**ysphagia, **D**yspnea.

Botulinum is from bad **bott**les of food, juice, and honey (causes a descending **flaccid** paralysis). Local botulinum toxin A (Botox) injections used to treat focal dystonia, hyperhidrosis, muscle spasms, and cosmetic reduction of facial wrinkles.

Clostridium perfringens

Produces α-toxin (lecithinase, a phospholipase) that can cause myonecrosis (gas gangrene **A**); presents as soft tissue crepitus) and hemolysis. If heavily spore-contaminated food is cooked but left standing too long at < 60°C, spores germinate → vegetative bacteria → heat-labile enterotoxin → food poisoning symptoms in 10-12 hours, resolution in 24 hours.

Perfringens perforates a gangrenous leg.

Clostridioides difficile


Produces toxins A and B, which damage enterocytes. Both toxins lead to watery diarrhea → pseudomembranous colitis **B**. Often 2° to antibiotic use, especially clindamycin or ampicillin; associated with PPIs.

Diagnosed by PCR or antigen detection of one or both toxins in stool.

Complications: toxic megacolon.

Difficile causes **diarrhea**.

Treatment: oral vancomycin, metronidazole, or fidaxomicin. For recurrent cases, consider repeating prior regimen or fecal microbiota transplant.

Corynebacterium diphtheriae

Gram \oplus rods occurring in angular arrangements; transmitted via respiratory droplets. Causes diphtheria via exotoxin encoded by β -prophage. Potent exotoxin inhibits protein synthesis via ADP-ribosylation of EF-2, leading to possible necrosis in pharynx, cardiac, and CNS tissue.

Symptoms include pseudomembranous pharyngitis (grayish-white membrane **A**) with lymphadenopathy. Toxin dissemination may cause myocarditis, arrhythmias, neuropathies.

Lab diagnosis based on gram \oplus rods with metachromatic (blue and red) granules and \oplus Elek test for toxin.

Toxoid vaccine prevents diphtheria.

Coryne = club shaped (metachromatic granules on Löffler media).

Black colonies on cystine-tellurite agar.

ABCDEFG:

ADP-ribosylation

β -prophage


Corynebacterium

Diphtheriae

Elongation **F**actor 2

Granules

Treatment: antibiotic therapy +/- diphtheria antitoxin.



Listeria monocytogenes

Gram \oplus , facultative intracellular rod; acquired by ingestion of unpasteurized dairy products and cold deli meats, transplacental transmission, by vaginal transmission during birth. Grows well at refrigeration temperatures ("cold enrichment").

Forms "rocket tails" (red in **A**) via actin polymerization that allow intracellular movement and cell-to-cell spread across cell membranes, thereby avoiding antibody. Characteristic tumbling motility in broth.

Can cause amnionitis, septicemia, and spontaneous abortion in pregnant women; granulomatous infantiseptica; meningitis in immunocompromised patients, neonates, and older adults; mild, self-limited gastroenteritis in healthy individuals.

Treatment: ampicillin.

Nocardia vs Actinomyces

Both are gram \oplus and form long, branching filaments resembling fungi.

Nocardia

Aerobe

Acid fast (weak) **A**

Found in soil

Causes pulmonary infections in immunocompromised (can mimic TB but with \ominus PPD); cutaneous infections after trauma in immunocompetent; can spread to CNS

Treat with sulfonamides (TMP-SMX)

Treatment is a **SNAP**: Sulfonamides—*Nocardia*; *Actinomyces*—Penicillin

Actinomyces

Anaerobe


Not acid fast **B**

Normal oral, reproductive, and GI flora

Causes oral/facial abscesses that drain through sinus tracts; often associated with dental caries/extraction and other maxillofacial trauma; forms yellow "sulfur granules"; can also cause PID with IUDs

Treat with penicillin


Mycobacteria




Gram ⊕ acid fast rods (pink rods, arrows in **A**).
Mycobacterium tuberculosis (TB, often resistant to multiple drugs).
M avium–intracellulare (causes disseminated, non-TB disease in AIDS; often resistant to multiple drugs). Prophylaxis with azithromycin when CD4+ count < 50 cells/mm³.
M scrofulaceum (cervical lymphadenitis in children).
M marinum (hand infection in aquarium handlers).

TB symptoms include fever, night sweats, weight loss, cough (nonproductive or productive), hemoptysis.
 Cord factor creates a “serpentine cord” appearance in virulent *M tuberculosis* strains; activates macrophages (promoting granuloma formation) and induces release of TNF-α.
 Sulfatides (surface glycolipids) inhibit phagolysosomal fusion.

Tuberculosis



Interferon-γ release assay (IGRA) has fewer false positives from BCG vaccination.
 PPD ⊕ if current infection or past exposure.
 PPD ⊖ if no infection and in sarcoidosis or HIV infection (especially with low CD4+ cell count).
 Caseating granulomas with central necrosis and Langhans giant cell (single example in **A**) are characteristic of 2° tuberculosis.




Leprosy

Also called Hansen disease. Caused by *Mycobacterium leprae*, an acid-fast bacillus that likes cool temperatures (infects skin and superficial nerves—“glove and stocking” loss of sensation **A**) and cannot be grown in vitro. Diagnosed via skin biopsy or tissue PCR. Reservoir in United States: armadillos.

Leprosy has 2 forms (many cases fall temporarily between two extremes):


- **Lepromatous**—presents diffusely over the skin, with **L**ionine (**L**ion-like) facies **B**, and is communicable (high bacterial load); characterized by low cell-mediated immunity with a largely Th2 response. **L**epromatous form can be **L**ethal.
- **Tuberculoid**—limited to a few hypoesthetic, hairless skin plaques; characterized by high cell-mediated immunity with a largely Th1-type response and low bacterial load.

Treatment: dapsone and rifampin for tuberculoid form; clofazimine is added for lepromatous form.

Gram-negative lab algorithm

Important **tests** are in **bold**. Important **pathogens** are in **bold italics**.

^aPleomorphic rod/coccobacillus

Neisseria

Gram \ominus diplococci. Metabolize glucose and produce IgA proteases. Contain lipooligosaccharides (LOS) with strong endotoxin activity. *N gonorrhoeae* is often intracellular (within neutrophils) **A**.

Acid production: **MeninG**ococci—**M**altose and **G**lucose; **Gonococci**—**G**lucose.

Gonococci

No polysaccharide capsule

No maltose acid detection

No vaccine due to antigenic variation of pilus proteins

Sexually or perinatally transmitted

Causes gonorrhea, septic arthritis, neonatal conjunctivitis (2–5 days after birth), pelvic inflammatory disease (PID), and Fitz-Hugh–Curtis syndrome

Diagnosed with NAT

Condoms \downarrow sexual transmission, erythromycin eye ointment prevents neonatal blindness

Treatment: ceftriaxone (+ azithromycin or doxycycline, for possible chlamydial coinfection)

Meningococci

Polysaccharide capsule

Maltose acid detection

Vaccine (type B vaccine available for at-risk individuals)



Transmitted via respiratory and oral secretions

Causes meningococemia with petechial hemorrhages and gangrene of toes **B**, meningitis, Waterhouse-Friderichsen syndrome (adrenal insufficiency, fever, DIC, shock)

Diagnosed via culture-based tests or PCR

Rifampin, ciprofloxacin, or ceftriaxone prophylaxis in close contacts

Treatment: ceftriaxone or penicillin G

Haemophilus influenzae

Small gram \ominus (coccobacillary) rod. Aerosol transmission. Nontypeable (unencapsulated) strains are the most common cause of mucosal infections (otitis media, conjunctivitis, bronchitis) as well as invasive infections since the vaccine for capsular type b was introduced. Produces IgA protease.

Culture on chocolate agar, which contains factors V (NAD^+) and X (hematin) for growth; can also be grown with *S aureus*, which provides factor V via RBC hemolysis.

HaEMOPhilus causes **E**piglottitis (endoscopic appearance in **A**), can be “cherry red” in children; “thumb sign” on lateral neck x-ray **B**), **M**eningitis, **O**titis media, and **P**neumonia.

Vaccine contains type b capsular polysaccharide (polyribosylribitol phosphate) conjugated to diphtheria toxoid or other protein. Given between 2 and 18 months of age.

Does not cause the flu (influenza virus does).

Treatment: amoxicillin +/- clavulanate for mucosal infections; ceftriaxone for meningitis; rifampin prophylaxis for close contacts.

Acinetobacter baumannii

Gram \ominus , strictly aerobic, oxidase \ominus coccobacillus. Commensal opportunist but increasingly associated with resistant hospital-acquired infections, especially in ICU. Can cause ventilator-associated pneumonia and septicemia in immunocompromised patients.

Bordetella pertussis

Gram \ominus , aerobic coccobacillus. Virulence factors include pertussis toxin (disables G_i), adenylate cyclase toxin (\uparrow cAMP), and tracheal cytotoxin. Three clinical stages:

- **C**atarrhal—low-grade fevers, **C**oryza.
- **P**aroxysmal—paroxysms of intense cough followed by inspiratory “whoop” (“whooping cough”), posttussive vomiting.
- **C**onvalescent—gradual recovery of chronic cough.


Prevented by Tdap, DTaP vaccines. May be mistaken as viral infection due to lymphocytic infiltrate resulting from immune response.

Treatment: macrolides; if allergic use TMP-SMX.

Brucella

Gram \ominus , aerobic coccobacillus. Transmitted via ingestion of contaminated animal products (eg, **un**pasteurized milk). Survives in macrophages in the reticuloendothelial system. Can form non-caseating granulomas. Typically presents with **undulant** fever, night sweats, and arthralgia.

Treatment: doxycycline + rifampin or streptomycin.

Legionella pneumophila

Gram \ominus rod. Gram stains poorly—use **silver** stain. Grow on **charcoal** yeast extract medium with **iron** and **cysteine**. Detected by presence of antigen in urine. Labs may show hyponatremia.


Aerosol transmission from environmental water source habitat (eg, air conditioning systems, hot water tanks). No person-to-person transmission.

Treatment: macrolide or quinolone.

Think of a French **legionnaire** (soldier) with his **silver** helmet, sitting around a campfire (**charcoal**) with his **iron** dagger—he is no **sissy** (cysteine).

Legionnaires' disease—severe pneumonia (often unilateral and lobar **A**), fever, GI and CNS symptoms. Common in smokers and in chronic lung disease.

Pontiac fever—mild flu-like syndrome.

Pseudomonas aeruginosa

Aeruginosa—**aerobic**; motile, catalase \oplus , gram \ominus rod. Non-lactose fermenting.

Oxidase \oplus . Frequently found in water. Has a grape-like odor.

PSEUDOMONAS is associated with:

Pneumonia, **S**epsis, **E**cthyma gangrenosum, **U**TI, **D**iabetes, **O**steomyelitis, **M**ucoid polysaccharide capsule, **O**tis externa (swimmer's ear), **N**osocomial infections (eg, catheters, equipment), **A**ddicts (drug abusers), **S**kin infections (eg, hot tub folliculitis, wound infection in burn victims).

Mucoid polysaccharide capsule may contribute to chronic pneumonia in cystic fibrosis patients due to biofilm formation.

Produces **PEEP**: **P**hospholipase C (degrades cell membranes); **E**ndotoxin (fever, shock); **E**xotoxin A (inactivates EF-2); **P**igments: pyoverdine and pyocyanin (blue-green pigment **A**; also generates reactive oxygen species).

Corneal ulcers/keratitis in contact lens wearers/minor eye trauma.

Ecthyma gangrenosum—rapidly progressive, necrotic cutaneous lesion **B** caused by *Pseudomonas* bacteremia. Typically seen in immunocompromised patients.

Treatments include “**CAMPFIRE**” drugs:

- **C**arbapenems
- **A**minoglycosides
- **M**onobactams
- **P**olymyxins (eg, polymyxin B, colistin)
- **F**luoroquinolones (eg, ciprofloxacin, levofloxacin)
- **T**HIRd- and fourth-generation cephalosporins (eg, ceftazidime, cefepime)
- **E**xtended-spectrum penicillins (eg, piperacillin, ticarcillin)

Salmonella vs Shigella Both *Salmonella* and *Shigella* are gram \ominus rods, non-lactose fermenters, oxidase \ominus , and can invade the GI tract via M cells of Peyer patches.

	<i>Salmonella typhi</i> (ty-Vi)	<i>Salmonella</i> spp. (except <i>S typhi</i>)	<i>Shigella</i>
RESERVOIRS	Humans only	Humans and animals	Humans only
SPREAD	Can disseminate hematogenously	Can disseminate hematogenously	Cell to cell; no hematogenous spread
H ₂ S PRODUCTION	Yes	Yes	No
FLAGELLA	Yes (salmon swim)	Yes (salmon swim)	No
VIRULENCE FACTORS	Endotoxin; Vi capsule	Endotoxin	Endotoxin; Shiga toxin (enterotoxin)
INFECTIOUS DOSE (ID ₅₀)	High—large inoculum required; acid-labile (inactivated by gastric acids)	High	Low—very small inoculum required; acid stable (resistant to gastric acids)
EFFECT OF ANTIBIOTICS ON FECAL EXCRETION	Prolongs duration	Prolongs duration	Shortens duration
IMMUNE RESPONSE	Primarily monocytes	PMNs in disseminated disease	Primarily PMN infiltration
GI MANIFESTATIONS	Constipation, followed by diarrhea	Diarrhea (possibly bloody)	Crampy abdominal pain → tenesmus, bloody mucoid stools (bacillary dysentery)
VACCINE	Oral vaccine contains live attenuated <i>S typhi</i> IM vaccine contains Vi capsular polysaccharide	No vaccine	No vaccine
UNIQUE PROPERTIES	<ul style="list-style-type: none"> Causes typhoid fever (rose spots on abdomen, constipation, abdominal pain, fever; later GI ulceration and hemorrhage); treat with ceftriaxone or fluoroquinolone Carrier state with gallbladder colonization 	<ul style="list-style-type: none"> Poultry, eggs, pets, and turtles are common sources Antibiotics not indicated Gastroenteritis is usually caused by non-typhoidal <i>Salmonella</i> 	<ul style="list-style-type: none"> 4 F's: Fingers, Flies, Food, Feces In order of decreasing severity (less toxin produced): <i>S dysenteriae</i>, <i>S flexneri</i>, <i>S boydii</i>, <i>S sonnei</i> Invasion of M cells is key to pathogenicity: organisms that produce little toxin can cause disease

Yersinia enterocolitica Gram \ominus pleomorphic rod/coccobacillus. Usually transmitted from pet feces (eg, puppies), contaminated milk, or pork. Can cause acute bloody diarrhea, pseudoappendicitis (right lower abdominal pain due to mesenteric adenitis and/or terminal ileitis), reactive arthritis in adults.

Lactose-fermenting enteric bacteria


Fermentation of **lactose** → pink colonies on MacCon**key** agar. Examples include *E coli*, *Enterobacter*, *Klebsiella*. *E coli* produces β -galactosidase, which breaks down lactose into glucose and galactose.

Lactose is key.
EMB agar—lactose fermenters grow as purple/black colonies. *E coli* grows colonies with a green sheen.

Escherichia coli


Gram \ominus , indole \oplus rod. *E coli* virulence factors: fimbriae—cystitis and pyelonephritis (P pili); K capsule—pneumonia, neonatal meningitis; LPS endotoxin—septic shock.

STRAIN	TOXIN AND MECHANISM	PRESENTATION
Enteroinvasive <i>E coli</i>	Microbe invades intestinal mucosa and causes necrosis and inflammation.	EIEC is I nvasive; dysentery. Clinical manifestations similar to <i>Shigella</i> .
Enterotoxigenic <i>E coli</i>	Produces heat-labile and heat-stable enterotoxins. No inflammation or invasion.	ETEC; T raveler's diarrhea (watery).
Enteropathogenic <i>E coli</i>	No toxin produced. Adheres to apical surface, flattens villi, prevents absorption.	Diarrhea, usually in children (think EPEC and P ediatrics).
Enterohemorrhagic <i>E coli</i>	O157:H7 is most common serotype in US. Often transmitted via undercooked meat, raw leafy vegetables. Shiga-like toxin causes hemolytic-uremic syndrome : triad of anemia, thrombocytopenia, and acute kidney injury due to microthrombi forming on damaged endothelium → mechanical hemolysis (with schistocytes on peripheral blood smear), platelet consumption, and ↓ renal blood flow.	Dysentery (toxin alone causes necrosis and inflammation). Does not ferment sorbitol (vs other <i>E coli</i>). H emorrhagic, H amburgers, H emolytic-uremic syndrome.

Klebsiella


Gram \ominus rod; intestinal flora that causes lobar pneumonia in alcoholics and diabetics when aspirated. Very mucoid colonies **A** caused by abundant polysaccharide capsules. Dark red “currant jelly” sputum (blood/mucus).
Also cause of nosocomial UTIs. Associated with evolution of multidrug resistance (MDR).

ABCDE's of *Klebsiella*:
Aspiration pneumonia
aBscess in lungs and liver
“Currant jelly” sputum
Diabetes
EtOH abuse

Campylobacter jejuni


Gram \ominus , comma or S shaped (with polar flagella) **A**, oxidase \oplus , grows at **42°C** (“*Campylobacter* likes the **hot campfire**”).
Major cause of bloody diarrhea, especially in children. Fecal-oral transmission through person-to-person contact or via ingestion of undercooked contaminated poultry or meat, unpasteurized milk. Contact with infected animals (dogs, cats, pigs) is also a risk factor.
Common antecedent to Guillain-Barré syndrome and reactive arthritis.

Vibrio cholerae



Gram \ominus , flagellated, comma shaped **A**, oxidase \oplus , grows in alkaline media. Endemic to developing countries. Produces profuse rice-water diarrhea via enterotoxin that permanently activates G_s, \uparrow cAMP. Sensitive to stomach acid (acid labile); requires large inoculum (high ID₅₀) unless host has \downarrow gastric acidity. Transmitted via ingestion of contaminated water or uncooked food (eg, raw shellfish). Treat promptly with oral rehydration solution.


Helicobacter pylori



Curved, flagellated (motile), gram \ominus rod **A** that is **triple** \oplus : catalase \oplus , oxidase \oplus , and urease \oplus (can use urea breath test or fecal antigen test for diagnosis). Urease produces ammonia, creating an alkaline environment, which helps *H pylori* survive in acidic mucosa. Colonizes mainly antrum of stomach; causes gastritis and peptic ulcers (especially duodenal). Risk factor for peptic ulcer disease, gastric adenocarcinoma, and MALT lymphoma.

Most common initial treatment is **triple** therapy: Amoxicillin (metronidazole if penicillin allergy) + Clarithromycin + Proton pump inhibitor; Antibiotics Cure *Pylori*. Bismuth-based quadruple therapy if concerned about macrolide resistance.

Spirochetes



Spiral-shaped bacteria **A** with axial filaments. Includes *Borrelia* (big size), *Leptospira*, and *Treponema*. Only *Borrelia* can be visualized using aniline dyes (Wright or Giemsa stain) in light microscopy due to size. *Treponema* is visualized by dark-field microscopy or direct fluorescent antibody (DFA) microscopy.

BLT:
Borrelia is Big.

Lyme disease



Caused by *Borrelia burgdorferi*, which is transmitted by the *Ixodes* deer tick **A** (also vector for *Anaplasma* spp. and protozoa *Babesia*). Natural reservoir is the mouse; deer are essential to tick life cycle but do not harbor *Borrelia*.

A Key **Lyme** pie to the **FACE**:
Facial nerve palsy (typically bilateral)
Arthritis
Cardiac block
Erythema migrans

Treatment: doxycycline (1st line); amoxicillin and, if severe illness, CNS signs, or heart block, ceftriaxone



Common in northeastern United States.
Stage 1—early localized: erythema migrans (typical “bulls-eye” configuration **B** is pathognomonic but not always present), flu-like symptoms.
Stage 2—early disseminated: secondary lesions, carditis, AV block, facial nerve (Bell) palsy, migratory myalgias/transient arthritis.
Stage 3—late disseminated: encephalopathy, chronic arthritis.

Leptospira interrogans Spirochete with hook-shaped ends found in water contaminated with animal urine.

Leptospirosis—flu-like symptoms, myalgias (classically of calves), jaundice, photophobia with conjunctival suffusion (erythema without exudate). Prevalent among surfers and in tropics (eg, Hawaii).

Weil disease (icterohemorrhagic leptospirosis)—severe form with jaundice and azotemia from liver and kidney dysfunction, fever, hemorrhage, and anemia.

Syphilis

Caused by spirochete *Treponema pallidum*. Treatment: penicillin G.

Primary syphilis

Localized disease presenting with **painless chancre** **A**. Use fluorescent or dark-field microscopy to visualize treponemes in fluid from chancre **B**. VDRL ⊕ in ~ 80%.

Secondary syphilis

Disseminated disease with constitutional symptoms, maculopapular rash **C** (including palms **D** and soles), condylomata lata **E** (smooth, painless, wart-like white lesions on genitals), lymphadenopathy, patchy hair loss; also confirmable with dark-field microscopy.

Serologic testing: VDRL/RPR (nonspecific), confirm diagnosis with specific test (eg, FTA-ABS).

Secondary syphilis = Systemic. Latent syphilis (⊕ serology without symptoms) may follow.

Tertiary syphilis

Gummas **F** (chronic granulomas), aortitis (vasa vasorum destruction), neurosyphilis (tabes dorsalis, “general paresis”), Argyll Robertson pupil (constricts with accommodation but is not reactive to light; also called “prostitute’s pupil” since it accommodates but does not react).

Signs: broad-based ataxia, ⊕ Romberg, Charcot joint, stroke without hypertension.

For neurosyphilis: test spinal fluid with VDRL, FTA-ABS, and PCR.

Congenital syphilis

Presents with facial abnormalities such as rhagades (linear scars at angle of mouth, black arrow in **G**), snuffles (nasal discharge, red arrow in **G**), saddle nose, notched (Hutchinson) teeth **H**, mulberry molars, and short maxilla; saber shins; CN VIII deafness.

To prevent, treat mother early in pregnancy, as placental transmission typically occurs after first trimester.




VDRL false positives


VDRL detects nonspecific antibody that reacts with beef cardiolipin. Quantitative, inexpensive, and widely available test for syphilis (sensitive but not specific).

False-Positive results on **VDRL** with:

- P**regnancy
- V**iral infection (eg, EBV, hepatitis)
- D**rugs (eg, chlorpromazine, procainamide)
- R**heumatic fever (rare)
- L**upus and leprosy

**Jarisch-Herxheimer reaction**

Flu-like syndrome (fever, chills, headache, myalgia) after antibiotics are started; due to killed bacteria (usually spirochetes) releasing toxins.


Gardnerella vaginalis

A pleomorphic, gram-variable rod involved in bacterial vaginosis. Presents as a gray vaginal discharge with a **fishy** smell; nonpainful (vs vaginitis). Associated with sexual activity, but not sexually transmitted. Bacterial **vaginosis** is also characterized by overgrowth of certain anaerobic bacteria in vagina (due to ↓ lactobacilli). **Clue** cells (vaginal epithelial cells covered with *Gardnerella*) have stippled appearance along outer margin (arrow in **A**).

I don't have a **clue** why I smell **fish** in the **vagina garden!**

Amine whiff test—mixing discharge with 10% KOH enhances fishy odor.

Treatment: metronidazole or clindamycin.

Chlamydiae

Chlamydiae cannot make their own ATP. They are obligate intracellular organisms that cause mucosal infections. 2 forms:

- **E**lementary body (small, dense) is “**E**nfectious” and **E**nters cell via **E**ndocytosis; transforms into reticulate body.
- **R**eticulate body **R**eplicates in cell by fission; **R**eorganizes into elementary bodies.

Chlamydia trachomatis causes neonatal and follicular adult conjunctivitis **A**, nongonococcal urethritis, PID, and reactive arthritis.

Chlamydoiphila pneumoniae and *Chlamydoiphila psittaci* cause atypical pneumonia; transmitted by aerosol.

Chlamydial cell wall lacks classic peptidoglycan (due to reduced muramic acid), rendering β-lactam antibiotics ineffective.

Chlamys = cloak (intracellular).

C psittaci—has an avian reservoir (**p**arrots), causes atypical **p**neumonia.

Lab diagnosis: PCR, nucleic acid amplification test. Cytoplasmic inclusions (reticulate bodies) seen on Giemsa or fluorescent antibody-stained smear.

Treatment: azithromycin (favored because one-time treatment) or doxycycline. Add ceftriaxone for possible concomitant gonorrhea.

***Chlamydia trachomatis* serotypes**

Types A, B, and C	Chronic infection, cause blindness due to follicular conjunctivitis in Africa.	ABC = Africa, Blindness, Chronic infection.
Types D–K	Urethritis/PID, ectopic pregnancy, neonatal pneumonia (staccato cough) with eosinophilia, neonatal conjunctivitis (1–2 weeks after birth).	D–K = everything else. Neonatal disease can be acquired during passage through infected birth canal.
Types L1, L2, and L3	Lymphogranuloma venereum —small, painless ulcers on genitals → swollen, painful inguinal lymph nodes that ulcerate (buboes). Treat with doxycycline.	

Zoonotic bacteria Zoonosis—infectious disease transmitted between animals and humans.

SPECIES	DISEASE	TRANSMISSION AND SOURCE
<i>Anaplasma</i> spp	Anaplasmosis	<i>Ixodes</i> ticks (live on deer and mice)
<i>Bartonella</i> spp	Cat scratch disease, bacillary angiomatosis	Cat scratch
<i>Borrelia burgdorferi</i>	Lyme disease	<i>Ixodes</i> ticks (live on deer and mice)
<i>Borrelia recurrentis</i>	Relapsing fever	Louse (recurrent due to variable surface antigens)
<i>Brucella</i> spp	Brucellosis/ undulant fever	Un pasteurized dairy
<i>Campylobacter</i>	Bloody diarrhea	Feces from infected pets/animals; contaminated meats/foods/hands
<i>Chlamydophila psittaci</i>	Psittacosis	Parrots, other birds
<i>Coxiella burnetii</i>	Q fever	Aerosols of cattle/sheep amniotic fluid
<i>Ehrlichia chaffeensis</i>	Ehrlichiosis	<i>Amblyomma</i> (Lone Star tick)
<i>Francisella tularensis</i>	Tularemia	Ticks, rabbits, deer flies
<i>Leptospira</i> spp	Leptospirosis	Animal urine in water; recreational water use
<i>Mycobacterium leprae</i>	Leprosy	Humans with lepromatous leprosy; armadillo (rare)
<i>Pasteurella multocida</i>	Cellulitis, osteomyelitis	Animal bite, cats, dogs
<i>Rickettsia prowazekii</i>	Epidemic typhus	Human to human via human body louse
<i>Rickettsia rickettsii</i>	Rocky Mountain spotted fever	<i>Dermacentor</i> (dog tick)
<i>Rickettsia typhi</i>	Endemic typhus	Fleas
<i>Salmonella</i> spp (except <i>S typhi</i>)	Diarrhea (which may be bloody), vomiting, fever, abdominal cramps	Reptiles and poultry
<i>Yersinia pestis</i>	Plague	Fleas (rats and prairie dogs are reservoirs)

Rickettsial diseases and vector-borne illnesses

Treatment: doxycycline.

RASH COMMON

Rocky Mountain spotted fever

Rickettsia rickettsii, vector is tick. Despite its name, disease occurs primarily in the South Atlantic states, especially North Carolina. Rash typically starts at wrists **A** and ankles and then spreads to trunk, palms, and soles.

Classic triad—headache, fever, rash (vasculitis). **Palms** and **soles** rash is seen in **Coxsackievirus A** infection (hand, foot, and mouth disease), **Rocky Mountain spotted fever**, and **2° Syphilis** (you drive **CARS** using your **palms** and **soles**).

Typhus

Endemic (fleas)—*R typhi*.
Epidemic (human body louse)—*R prowazekii*.
Rash starts centrally and spreads out, sparing palms and soles.

Rickettsii on the w**R**ists, **T**yphus on the **T**runk.

RASH RARE

Ehrlichiosis

Ehrlichia, vector is tick. **Monocytes** with morulae **B** (mul**berry**-like inclusions) in cytoplasm.

MEGA berry—
Monocytes = **E**hrlichiosis
Granulocytes = **A**naplasmosis


Anaplasmosis

Anaplasma, vector is tick. **Granulocytes** with morulae **C** in cytoplasm.


Q fever

Coxiella burnetii, no arthropod vector. Spores inhaled as aerosols from cattle/sheep amniotic fluid. Presents with headache, cough, influenza-like symptoms, pneumonia, possibly in combination with hepatitis. Common cause of culture ⊖ endocarditis.

Q fever is caused by a **Q**uite **C**omplexed **B**ug because it has no rash or vector and its causative organism can survive outside in its endospore form. Not in the *Rickettsia* genus, but closely related.



Mycoplasma pneumoniae



Classic cause of atypical “walking pneumonia” (insidious onset, headache, nonproductive cough, patchy or diffuse interstitial infiltrate). Occurs frequently in those <30 years old; outbreaks in military recruits, prisons, colleges. X-ray looks worse than patient. High titer of **cold** agglutinins (IgM), which can agglutinate RBCs. Treatment: macrolides, doxycycline, or fluoroquinolone (penicillin ineffective since *Mycoplasma* has no cell wall).

Not seen on Gram stain. Pleomorphic **A**. Bacterial membrane contains sterols for stability. Grown on Eaton agar.

Mycoplasma gets **cold** without a **coat** (no cell wall). Can cause atypical variant of Stevens-Johnson syndrome, typically in children and adolescents.

▶ MICROBIOLOGY—MYCOLOGY

Systemic mycoses





All of the following can cause pneumonia and can disseminate.


All are caused by dimorphic fungi: **cold** (20°C) = **mold**; **heat** (37°C) = **yeast**. Only exception is

Coccidioides, which is a spherule (not yeast) in tissue.

Systemic mycoses can form granulomas (like TB); cannot be transmitted person-to-person (unlike TB).

Treatment: fluconazole or itraconazole for **local** infection; amphotericin B for **systemic** infection.

DISEASE	ENDEMIC LOCATION	PATHOLOGIC FEATURES	UNIQUE SIGNS/SYMPTOMS	NOTES
<p>Histoplasmosis</p> 	Mississippi and Ohio River Valleys	Macrophage filled with <i>Histoplasma</i> (smaller than RBC) A	Palatal/tongue ulcers, splenomegaly, pancytopenia	Histo hides (within macrophages) Associated with bird or bat droppings (eg, spelunking) Diagnosis via urine/serum antigen
<p>Blastomycosis</p> 	Eastern and Central US, Great Lakes	Broad -based budding of <i>Blastomyces</i> (same size as RBC) B	Inflammatory lung disease Disseminates to bone/skin (may mimic SCC) Forms granulomatous nodules	Blasto buds broadly
<p>Coccidioidomycosis</p> 	Southwestern US, California	Spherule (much larger than RBC) filled with endospores of <i>Coccidioides</i> C	Disseminates to skin/bone Erythema nodosum (desert bumps) or multiforme Arthralgias (desert rheumatism) Can cause meningitis	Associated with dust exposure in endemic areas (eg, archeological excavations, earthquakes)
<p>Para-coccidioidomycosis</p> 	Latin America	Budding yeast of <i>Paracoccidioides</i> with “ captain’s wheel ” formation (much larger than RBC) D	Similar to blastomycosis, males > females	Paracoccidio parasails with the captain’s wheel all the way to Latin America



Cutaneous mycoses

Tinea (dermatophytes)	Clinical name for dermatophyte (cutaneous fungal) infections. Dermatophytes include <i>Microsporum</i> , <i>Trichophyton</i> , and <i>Epidermophyton</i> . Branching septate hyphae visible on KOH preparation with blue fungal stain A . Associated with pruritus.
Tinea capitis	Occurs on head, scalp. Associated with lymphadenopathy, alopecia, scaling B .
Tinea corporis	Occurs on body (usually torso). Characterized by enlarging erythematous, scaly rings (“ringworm”) with central clearing C . Can be acquired from contact with infected pets or farm animals.
Tinea cruris	Occurs in inguinal area D . Often does not show the central clearing seen in tinea corporis.
Tinea pedis	Three varieties: <ul style="list-style-type: none"> ▪ Interdigital E; most common ▪ Moccasin distribution F ▪ Vesicular type
Tinea unguium	Onychomycosis; occurs on nails.
Tinea (pityriasis) versicolor	Caused by <i>Malassezia</i> spp. (<i>Pityrosporum</i> spp.), a yeast-like fungus (not a dermatophyte despite being called tinea). Degradation of lipids produces acids that inhibit tyrosinase (involved in melanin synthesis) → hypopigmentation G ; hyperpigmentation and/or pink patches can also occur due to inflammatory response. Less pruritic than dermatophytes. Can occur any time of year, but more common in summer (hot, humid weather). “Spaghetti and meatballs” appearance on microscopy H . Treatment: selenium sulfide, topical and/or oral antifungal medications.



Opportunistic fungal infections***Candida albicans***

alba = white. Dimorphic; forms pseudohyphae and budding yeasts at 20°C **A**, germ tubes at 37°C **B**.

Systemic or superficial fungal infection. Causes oral **C** and esophageal thrush in immunocompromised (neonates, steroids, diabetes, AIDS), vulvovaginitis (diabetes, use of antibiotics), diaper rash, endocarditis (IV drug users), disseminated candidiasis (especially in neutropenic patients), chronic mucocutaneous candidiasis.

Treatment: oral fluconazole/topical azoles for vaginal; nystatin, azoles, or, rarely, echinocandins for oral; fluconazole, echinocandins, or amphotericin B for esophageal or systemic disease.

Aspergillus fumigatus

Septate hyphae that branch at 45° Acute Angle **D E**.

Causes invasive aspergillosis in immunocompromised patients, neutrophil dysfunction (eg, chronic granulomatous disease).

Can cause aspergillomas **F** in pre-existing lung cavities, especially after TB infection.

Some species of *Aspergillus* produce **A**flatoxins (associated with hepatocellular carcinoma).

Treatment: voriconazole or echinocandins (2nd-line).

Allergic bronchopulmonary aspergillosis (ABPA)—hypersensitivity response to *Aspergillus* growing in lung mucus. Associated with asthma and cystic fibrosis; may cause bronchiectasis and eosinophilia.

Cryptococcus neoformans

5–10 μm with narrow budding. Heavily encapsulated yeast. Not dimorphic.

Found in soil, pigeon droppings. Acquired through inhalation with hematogenous dissemination to meninges. Highlighted with India ink (clear halo **G**) and mucicarmine (red inner capsule **H**).

Latex agglutination test detects polysaccharide capsular antigen and is more sensitive and specific. Causes cryptococcosis, cryptococcal meningitis, cryptococcal encephalitis (“soap bubble” lesions in brain), primarily in immunocompromised.

Treatment: amphotericin B + flucytosine followed by fluconazole for cryptococcal meningitis.


***Mucor and Rhizopus* spp**

Irregular, broad, nonseptate hyphae branching at wide angles **I**.

Causes mucormycosis, mostly in ketoacidotic diabetic and/or neutropenic patients (eg, leukemia).

Inhalation of spores → fungi proliferate in blood vessel walls, penetrate cribriform plate, and enter brain. Rhinocerebral, frontal lobe abscess; cavernous sinus thrombosis. Headache, facial pain, black necrotic eschar on face **J**; may have cranial nerve involvement.


Treatment: surgical debridement, amphotericin B or isavuconazole.



Pneumocystis jirovecii

Causes *Pneumocystis* pneumonia (PCP), a diffuse interstitial pneumonia **A**. Yeast-like fungus (originally classified as protozoan). Most infections are asymptomatic. Immunosuppression (eg, AIDS) predisposes to disease. Diffuse, bilateral ground-glass opacities on chest imaging, with pneumatoceles **B**. Diagnosed by bronchoalveolar lavage or lung biopsy. Disc-shaped yeast seen on methenamine silver stain of lung tissue **C** or with fluorescent antibody.

Treatment/prophylaxis: TMP-SMX, pentamidine, dapsone (prophylaxis as single agent, or treatment in combination with TMP), atovaquone. Start prophylaxis when CD4⁺ count drops to < 200 cells/mm³ in HIV patients.

***Sporothrix schenckii***

Causes sporotrichosis. Dimorphic fungus. Exists as a **cigar**-shaped yeast at 37 °C in the human body and as hyphae with spores in soil (conidia). Lives on vegetation. When spores are traumatically introduced into the skin, typically by a thorn (“**rose gardener’s** disease”), causes local pustule or ulcer with nodules along draining lymphatics (ascending lymphangitis **A**).

Disseminated disease possible in immunocompromised host.

Treatment: itraconazole or **pot**assium iodide (only for cutaneous/lymphocutaneous).

Think of a **rose gardener** who smokes a **cigar** and **pot**.

▶ MICROBIOLOGY—PARASITOLOGY


Protozoa—gastrointestinal infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
<i>Giardia lamblia</i>	Giardiasis —bloating, flatulence, foul-smelling, fatty diarrhea (often seen in campers/hikers)—think fat -rich Ghirardelli chocolates for fatty stools of Giardia	Cysts in water	Multinucleated trophozoites A or cysts B in stool, antigen detection	Metronidazole
<i>Entamoeba histolytica</i>	Amebiasis —bloody diarrhea (dysentery), liver abscess (“anchovy paste” exudate), RUQ pain; histology of colon biopsy shows flask-shaped ulcers	Cysts in water	Serology, antigen testing, and/or trophozoites (with engulfed RBCs C in the cytoplasm) or cysts with up to 4 nuclei in stool D ; Entamoeba Eats Erythrocytes	Metronidazole; paromomycin or iodoquinol for asymptomatic cyst passers
<i>Cryptosporidium</i>	Severe diarrhea in AIDS Mild disease (watery diarrhea) in immunocompetent hosts	Oocysts in water	Oocysts on acid-fast stain E , antigen detection	Prevention (by filtering city water supplies); nitazoxanide in immunocompetent hosts





Protozoa—CNS infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
<i>Toxoplasma gondii</i>	Immunocompetent: mononucleosis-like symptoms, ⊖ heterophile antibody test Reactivation in AIDS → brain abscesses usually seen as multiple ring-enhancing lesions on MRI A Congenital toxoplasmosis: classic triad of chorioretinitis, hydrocephalus, and intracranial calcifications	Cysts in meat (most common); oocysts in cat feces; crosses placenta (pregnant women should avoid cats)	Serology, biopsy (tachyzoite) B	Sulfadiazine + pyrimethamine
<i>Naegleria fowleri</i>	Rapidly fatal meningoencephalitis	Swimming in warm freshwater; enters via cribriform plate	Amoebas in CSF C	Amphotericin B has been effective for a few survivors
<i>Trypanosoma brucei</i>	African sleeping sickness —enlarged lymph nodes, recurring fever (due to antigenic variation), somnolence, coma	Tsetse fly, a painful bite	Trypomastigote in blood smear D	Suramin for blood-borne disease or melarsoprol for CNS penetration (“ I sure am mellow when I’m sleeping ”)



Protozoa—hematologic infections

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
<p>Plasmodium <i>P vivax/ovale</i> <i>P falciparum</i> <i>P malariae</i></p> 	<p>Malaria—fever, headache, anemia, splenomegaly <i>P vivax/ovale</i>—48-hr cycle (tertian; includes fever on first day and third day, thus fevers are actually 48 hr apart); dormant form (hypnozoite) in liver <i>P falciparum</i>—severe; irregular fever patterns; parasitized RBCs occlude capillaries in brain (cerebral malaria), kidneys, lungs <i>P malariae</i>—72-hr cycle (quartan)</p>	<p><i>Anopheles</i> mosquito</p>	<p>Blood smear: trophozoite ring form within RBC A, schizont containing merozoites; red granules (Schüffner stippling) B throughout RBC cytoplasm seen with <i>P vivax/ovale</i></p>	<p>Chloroquine (for sensitive species); if resistant, use mefloquine or atovaquone/proguanil If life-threatening, use intravenous quinidine or artesunate (test for G6PD deficiency) For <i>P vivax/ovale</i>, add primaquine for hypnozoite (test for G6PD deficiency)</p>
<p>Babesia</p> 	<p>Babesiosis—fever and hemolytic anemia; predominantly in northeastern United States; asplenia ↑ risk of severe disease</p>	<p><i>Ixodes</i> tick (also vector for <i>Borrelia burgdorferi</i> and <i>Anaplasma</i> spp)</p>	<p>Blood smear: ring form C1, “Maltese cross” C2; PCR</p>	<p>Atovaquone + azithromycin</p>

Protozoa—others

ORGANISM	DISEASE	TRANSMISSION	DIAGNOSIS	TREATMENT
Visceral infections				
<i>Trypanosoma cruzi</i>	Chagas disease —dilated cardiomyopathy with apical atrophy, megacolon, megaesophagus; predominantly in South America Unilateral periorbital swelling (Romaña sign) characteristic of acute stage	Triatomine insect (kissing bug) bites and defecates around the mouth or eyes; fecal transmission into bite site or mucosa	Trypomastigote in blood smear A	Benznidazole or nifurtimox ; crusing in my Benz , with a fur coat on
<i>Leishmania</i> spp	Visceral leishmaniasis (kala-azar) —spiking fevers, hepatosplenomegaly, pancytopenia Cutaneous leishmaniasis —skin ulcers B	Sandfly	Macrophages containing amastigotes C	Amphotericin B, sodium stibogluconate
Sexually transmitted infections				
<i>Trichomonas vaginalis</i>	Vaginitis —foul-smelling, greenish discharge; itching and burning; do not confuse with <i>Gardnerella vaginalis</i> , a gram-variable bacterium associated with bacterial vaginosis	Sexual (cannot exist outside human because it cannot form cysts)	Trophozoites (motile) D on wet mount; punctate cervical hemorrhages (“strawberry cervix”)	Metronidazole for patient and partner (prophylaxis; check for STI)



Nematode routes of infection

Ingested—*Enterobius*, *Ascaris*, *Toxocara*, *Trichinella*, *Trichuris*
Cutaneous—*Strongyloides*, *Ancylostoma*, *Necator*
Bites—*Loa loa*, *Onchocerca volvulus*, *Wuchereria bancrofti*



You'll get sick if you **EATTT** these!

These get into your feet from the **SANd**

Lay **LOW** to avoid getting bitten

Nematodes (roundworms)

ORGANISM	DISEASE	TRANSMISSION	TREATMENT
Intestinal			
<i>Enterobius vermicularis</i> (pinworm)	Causes anal pruritus (diagnosed by seeing egg A via the tape test).	Fecal-oral.	Bendazoles (bendy worms), pyrantel pamoate.
<i>Ascaris lumbricoides</i> (giant roundworm)	May cause obstruction at ileocecal valve, biliary obstruction, intestinal perforation, migrates from nose/mouth.	Fecal-oral; knobby-coated, oval eggs seen in feces under microscope B .	Bendazoles.
<i>Strongyloides stercoralis</i> (threadworm)	GI (eg, duodenitis), pulmonary (eg, dry cough, hemoptysis), and cutaneous (eg, pruritus) symptoms. Hyperinfection syndrome caused by autoinfection (larvae enter bloodstream).	Larvae in soil penetrate skin; rhabditiform larvae seen in feces under microscope.	Ivermectin or bendazoles.
<i>Ancylostoma</i> spp, <i>Necator americanus</i> (hookworms)	Cause microcytic anemia by sucking blood from intestinal wall. Cutaneous larva migrans —pruritic, serpiginous rash C from walking barefoot on contaminated beach.	Larvae penetrate skin.	Bendazoles or pyrantel pamoate.
<i>Trichinella spiralis</i>	Larvae enter bloodstream, encyst in striated muscle D → myositis. Trichinosis —fever, vomiting, nausea, periorbital edema, myalgia.	Undercooked meat (especially pork); fecal-oral (less likely).	Bendazoles.
<i>Trichuris trichiura</i> (whipworm)	Often asymptomatic; loose stools, anemia, rectal prolapse in children.	Fecal-oral.	Bendazoles.
Tissue			
<i>Toxocara canis</i>	Visceral larva migrans —nematodes migrate to blood through intestinal wall → inflammation affecting liver, eyes (visual impairment, blindness), CNS (seizures, coma), heart (myocarditis).	Fecal-oral.	Bendazoles.
<i>Onchocerca volvulus</i>	Skin changes, loss of elastic fibers, river blindness (black skin nodules, “ black sight”); allergic reaction possible.	Female black fly.	Ivermectin (ivermectin for river blindness).
<i>Loa loa</i>	Swelling in skin, worm in conjunctiva.	Deer fly, horse fly, mango fly.	Diethylcarbamazine.
<i>Wuchereria bancrofti</i>	Lymphatic filariasis (elephantiasis) —worms invade lymph nodes. → inflammation → lymphedema E ; symptom onset after 9 mo–1 yr.	Female mosquito.	Diethylcarbamazine.




Cestodes (tapeworms)

ORGANISM	DISEASE	TRANSMISSION	TREATMENT
<i>Taenia solium</i> A	Intestinal tapeworm	Ingestion of larvae encysted in undercooked pork	Praziquantel
	Cysticercosis, neurocysticercosis (cystic CNS lesions, seizures) B	Ingestion of eggs in food contaminated with human feces	Praziquantel; albendazole for neurocysticercosis
<i>Diphyllobothrium latum</i>	Vitamin B ₁₂ deficiency (tapeworm competes for B ₁₂ in intestine) → megaloblastic anemia	Ingestion of larvae in raw freshwater fish	Praziquantel
<i>Echinococcus granulosus</i> C	Hydatid cysts D (“eggshell calcification”) in liver E ; cyst rupture can cause anaphylaxis	Ingestion of eggs in food contaminated with dog feces Sheep are an intermediate host	Albendazole



Trematodes (flukes)


ORGANISM	DISEASE	TRANSMISSION	TREATMENT
<i>Schistosoma</i> A B	Liver and spleen enlargement (<i>S mansoni</i> , egg with lateral spine A), fibrosis, inflammation, portal hypertension Chronic infection with <i>S haematobium</i> (egg with terminal spine B) can lead to squamous cell carcinoma of the bladder (painless hematuria) and pulmonary hypertension	Snails are intermediate host; cercariae penetrate skin of humans in contact with contaminated fresh water (eg, swimming or bathing)	Praziquantel
	<i>Clonorchis sinensis</i>	Biliary tract inflammation → pigmented gallstones Associated with cholangiocarcinoma	Undercooked fish

Ectoparasites***Sarcoptes scabiei***

Mites burrow into stratum corneum and cause **scabies**—pruritus (worse at night) and serpiginous burrows (lines) often between fingers and toes **A**.

Common in children, crowded populations (jails, nursing homes); transmission through skin-to-skin contact (most common) or via fomites.

Treatment: permethrin cream, washing/drying all clothing/bedding, treat close contacts.

Pediculus humanus/Phthirus pubis

Blood-sucking lice that cause intense pruritus with associated excoriations, commonly on scalp and neck (head lice), waistband and axilla (body lice), or pubic and perianal regions (pubic lice).

Body lice can transmit *Rickettsia prowazekii* (epidemic typhus), *Borrelia recurrentis* (relapsing fever), *Bartonella quintana* (trench fever).


Treatment: pyrethroids, malathion, or ivermectin lotion, and nit **B** combing. Children with head lice can be treated at home without interrupting school attendance.

Parasite hints

ASSOCIATIONS	ORGANISM
Biliary tract disease, cholangiocarcinoma	<i>Clonorchis sinensis</i>
Brain cysts, seizures	<i>Taenia solium</i> (neurocysticercosis)
Hematuria, squamous cell bladder cancer	<i>Schistosoma haematobium</i>
Liver (hydatid) cysts	<i>Echinococcus granulosus</i>
Microcytic anemia	<i>Ancylostoma</i> , <i>Necator</i>
Myalgias, periorbital edema	<i>Trichinella spiralis</i>
Perianal pruritus	<i>Enterobius</i>
Portal hypertension	<i>Schistosoma mansoni</i> , <i>Schistosoma japonicum</i>
Vitamin B ₁₂ deficiency	<i>Diphyllobothrium latum</i>

▶ MICROBIOLOGY—VIROLOGY


Viral structure—general features



Viral genetics

Recombination

Exchange of genes between 2 chromosomes by crossing over within regions of significant base sequence homology.




Reassortment

When viruses with segmented genomes (eg, influenza virus) exchange genetic material. For example, the 2009 novel H1N1 influenza A pandemic emerged via complex viral reassortment of genes from human, swine, and avian viruses. Has potential to cause antigenic shift.




Complementation

When 1 of 2 viruses that infect the cell has a mutation that results in a nonfunctional protein, the nonmutated virus “complements” the mutated one by making a functional protein that serves both viruses. For example, hepatitis D virus requires the presence of replicating hepatitis B virus to supply HBsAg, the envelope protein for HDV.



Phenotypic mixing

Occurs with simultaneous infection of a cell with 2 viruses. For progeny 1, genome of virus A can be partially or completely coated (forming pseudovirion) with the surface proteins of virus B. Type B protein coat determines the tropism (infectivity) of the hybrid virus. Progeny from subsequent infection of a cell by progeny 1 will have a type A coat that is encoded by its type A genetic material.



DNA viral genomes All DNA viruses have dsDNA genomes except Parvoviridae (ssDNA). All are linear except papilloma-, polyoma-, and hepadnaviruses (circular). All are dsDNA (like our cells), except “**part-of-a-virus**” (**parvovirus**) is ssDNA. *Parvus* = small.

RNA viral genomes All RNA viruses have ssRNA genomes except Reoviridae (dsRNA).
 ⊕ stranded RNA viruses: I went to a **retro** (**retrovirus**) **toga** (**togavirus**) party, where I drank **flavored** (**flavivirus**) **Corona** (**coronavirus**) and ate **hippie** (**hepevirus**) **California** (**calicivirus**) **pickles** (**picornavirus**). All are ssRNA, except “**repeato-virus**” (**reovirus**) is dsRNA.


Naked viral genome infectivity Purified nucleic acids of most dsDNA viruses (except poxviruses and HBV) and ⊕ strand ssRNA (≈ mRNA) viruses are infectious. Naked nucleic acids of ⊖ strand ssRNA and dsRNA viruses are not infectious. They require polymerases contained in the complete virion.

Viral envelopes Generally, enveloped viruses acquire their envelopes from plasma membrane when they exit from cell. Exceptions include herpesviruses, which acquire envelopes from nuclear membrane. **Naked** (nonenveloped) viruses include **P**apillomavirus, **A**denovirus, **P**arvovirus, **P**olyomavirus, **C**alicivirus, **P**icornavirus, **R**eovirus, and **H**epevirus. DNA = **PAPP**; RNA = **CPR** and **hepevirus**. Give **PAPP** smears and **CPR** to a **naked hippie** (**hepevirus**). Enveloped DNA viruses **Have Helpful Protection** (**Herpesvirus**, **Hepadnavirus**, **Poxvirus**).

DNA virus characteristics

Some general rules—all DNA viruses:

GENERAL RULE	COMMENTS
Are HHAPPPP y viruses	H epadna, H erpes, A deno, P ox, P arvo, P apilloma, P olyoma.
Are double stranded	Except parvo (single stranded).
Have linear genomes	Except papilloma and polyoma (circular, supercoiled) and hepadna (circular, incomplete).
Are icosahedral	Except pox (complex).
Replicate in the nucleus	Except pox (carries own DNA-dependent RNA polymerase).


DNA viruses		All replicate in the nucleus (except poxvirus). “ P ox is out of the box (nucleus).”	
VIRAL FAMILY	ENVELOPE	DNA STRUCTURE	MEDICAL IMPORTANCE
Herpesviruses	Yes	DS and linear	See Herpesviruses entry
Poxvirus	Yes	DS and linear (largest DNA virus)	Smallpox eradicated world wide by use of the live-attenuated vaccine Cowpox (“milkmaid blisters”) Molluscum contagiosum —flesh-colored papule with central umbilication
Hepadnavirus	Yes	Partially DS and circular	HBV: <ul style="list-style-type: none"> Acute or chronic hepatitis Not a retrovirus but has reverse transcriptase
Adenovirus	No	DS and linear	Febrile pharyngitis A —sore throat Acute hemorrhagic cystitis Pneumonia Conjunctivitis—“pink eye” Gastroenteritis Myocarditis
			
Papillomavirus	No	DS and circular	HPV—warts (serotypes 1, 2, 6, 11), CIN, cervical cancer (most commonly 16, 18)
Polyomavirus	No	DS and circular	JC virus—progressive multifocal leukoencephalopathy (PML) in HIV BK virus—transplant patients, commonly targets kidney JC: Junky Cerebrum; BK: Bad Kidney
Parvovirus	No	SS and linear (smallest DNA virus)	B19 virus—aplastic crises in sickle cell disease, “slapped cheek” rash in children (erythema infectiosum, or fifth disease); infects RBC precursors and endothelial cells → RBC destruction → hydrops fetalis and death in fetus, pure RBC aplasia and rheumatoid arthritis–like symptoms in adults

Herpesviruses Enveloped, DS, and linear viruses


VIRUS	ROUTE OF TRANSMISSION	CLINICAL SIGNIFICANCE	NOTES
Herpes simplex virus-1	Respiratory secretions, saliva	Gingivostomatitis, keratoconjunctivitis A , herpes labialis (cold sores) B , herpetic whitlow on finger, temporal lobe encephalitis, esophagitis, erythema multiforme	Most commonly latent in trigeminal ganglia Most common cause of sporadic encephalitis, can present as altered mental status, seizures, and/or aphasia
Herpes simplex virus-2	Sexual contact, perinatal	Herpes genitalis C , neonatal herpes	Most commonly latent in sacral ganglia Viral meningitis more common with HSV-2 than with HSV-1

Herpesviruses (continued)

VIRUS	ROUTE OF TRANSMISSION	CLINICAL SIGNIFICANCE	NOTES
Varicella-Zoster virus (HHV-3)	Respiratory secretions, contact with fluid from vesicles	Varicella-zoster (chickenpox D , shingles E), encephalitis, pneumonia Most common complication of shingles is post-herpetic neuralgia	Latent in dorsal root or trigeminal ganglia; CN V ₁ branch involvement can cause herpes zoster ophthalmicus
Epstein-Barr virus (HHV-4)	Respiratory secretions, saliva; aka “kissing disease,” (common in teens, young adults)	Mononucleosis —fever, hepatosplenomegaly F , pharyngitis, and lymphadenopathy (especially posterior cervical nodes); avoid contact sports until resolution due to risk of splenic rupture Associated with lymphomas (eg, endemic Burkitt lymphoma), nasopharyngeal carcinoma (especially Asian adults), lymphoproliferative disease in transplant patients	Infects B cells through CD21, “Must be 21 to drink B eer in a B arr” Atypical lymphocytes on peripheral blood smear G —not infected B cells but reactive cytotoxic T cells ⊕ Monospot test—heterophile antibodies detected by agglutination of sheep or horse RBCs Use of amoxicillin in mononucleosis can cause characteristic maculopapular rash
Cytomegalovirus (HHV-5)	Congenital, transfusion, sexual contact, saliva, urine, transplant	Mononucleosis (⊖ Monospot) in immunocompetent patients; infection in immunocompromised, especially pneumonia in transplant patients; esophagitis; AIDS retinitis (“ sight omegalovirus”): hemorrhage, cotton-wool exudates, vision loss Congenital CMV	Infected cells have characteristic “owl eye” intranuclear inclusions H Latent in mononuclear cells
Human herpesviruses 6 and 7	Saliva	Roseola infantum (exanthem subitum): high fevers for several days that can cause seizures, followed by diffuse macular rash (starts on trunk then spreads to extremities) I	Roseola : fever first, R osy (rash) l ater. HHV-7—less common cause of roseola
Human herpesvirus 8	Sexual contact	Kaposi sarcoma (neoplasm of endothelial cells). Seen in HIV/AIDS and transplant patients. Dark/violaceous plaques or nodules J representing vascular proliferations	Can also affect GI tract and lungs



HSV identification



Viral culture for skin/genitalia.

CSF PCR for herpes encephalitis.

Tzanck test—a smear of an opened skin vesicle to detect multinucleated giant cells **A** commonly seen in HSV-1, HSV-2, and VZV infection. PCR of skin lesions is test of choice.

Tzanck heavens I do not have herpes.

Intranuclear eosinophilic Cowdry A inclusions also seen with HSV-1, HSV-2, VZV.

Receptors used by viruses

VIRUS	RECEPTORS
CMV	Integrins (heparan sulfate)
EBV	CD21
HIV	CD4, CXCR4, CCR5
Parvovirus B19	P antigen on RBCs
Rabies	Nicotinic AChR
Rhinovirus	ICAM-1 (I came to see the rhino)

RNA viruses		All replicate in the cytoplasm (except retrovirus and influenza virus). “ Retro flu is outta cyt (sight).”		
VIRAL FAMILY	ENVELOPE	RNA STRUCTURE	CAPSID SYMMETRY	MEDICAL IMPORTANCE
Reoviruses	No	DS linear Multisegmented	Icosahedral (double)	Coltivirus ^a — Colorado tick fever Rotavirus—cause of fatal diarrhea in children
Picornaviruses	No	SS ⊕ linear	Icosahedral	Poliovirus —polio-Salk/Sabin vaccines—IPV/OPV Echovirus —aseptic meningitis Rhinovirus —“common cold” Coxsackievirus —aseptic meningitis; herpangina (mouth blisters, fever); hand, foot, and mouth disease; myocarditis; pericarditis HAV —acute viral hepatitis PERCH
Hepevirus	No	SS ⊕ linear	Icosahedral	HEV
Caliciviruses	No	SS ⊕ linear	Icosahedral	Norovirus—viral gastroenteritis
Flaviviruses	Yes	SS ⊕ linear	Icosahedral	HCV Yellow fever ^a Dengue ^a St. Louis encephalitis ^a West Nile virus ^a —meningoencephalitis, flaccid paralysis Zika virus ^a
Togaviruses	Yes	SS ⊕ linear	Icosahedral	Toga CREW — Chikungunya virus ^a (co-infection with dengue virus can occur), Rubella , Eastern and Western equine encephalitis
Retroviruses	Yes	SS ⊕ linear 2 copies	Icosahedral (HTLV), complex and conical (HIV)	Have reverse transcriptase HTLV—T-cell leukemia HIV—AIDS
Coronaviruses	Yes	SS ⊕ linear	Helical	“Common cold,” SARS, MERS
Orthomyxoviruses	Yes	SS ⊖ linear 8 segments	Helical	Influenza virus
Paramyxoviruses	Yes	SS ⊖ linear Nonsegmented	Helical	PaRaMyxovirus : Parainfluenza —croup RSV —bronchiolitis in babies Measles , Mumps
Rhabdoviruses	Yes	SS ⊖ linear	Helical	Rabies
Filoviruses	Yes	SS ⊖ linear	Helical	Ebola/Marburg hemorrhagic fever—often fatal.
Arenaviruses	Yes	SS ⊕ and ⊖ circular 2 segments	Helical	LCMV—lymphocytic choriomeningitis virus Lassa fever encephalitis—spread by rodents
Bunyaviruses	Yes	SS ⊖ circular 3 segments	Helical	California encephalitis ^a Sandfly/Rift Valley fevers ^a Crimean-Congo hemorrhagic fever ^a Hantavirus—hemorrhagic fever, pneumonia
Delta virus	Yes	SS ⊖ circular	Uncertain	HDV is a “defective” virus that requires the presence of HBV to replicate

SS, single-stranded; DS, double-stranded; ⊕, positive sense; ⊖, negative sense; ^a= **arbovirus**, **arthropod borne** (mosquitoes, ticks).

Negative-stranded viruses

Must transcribe \ominus strand to \oplus . Virion brings its own RNA-dependent RNA polymerase. They include **A**renaviruses, **B**unyaviruses, **P**aramyxoviruses, **O**rthomyxoviruses, **F**iloviruses, and **R**habdoviruses.

Always **B**ring **P**olymerase **O**r **F**ail **R**eplication.

Segmented viruses

All are RNA viruses. They include **B**unyaviruses (3 segments), **O**rthomyxoviruses (influenza viruses) (8 segments), **A**renaviruses (2 segments), and **R**eoviruses (10-12 segments).

BOARding flight **382** in **10-12** minutes.

Picornavirus

Includes **P**oliovirus, **E**chovirus, **R**hinovirus, **C**oxsackievirus, and **H**AV. RNA is translated into 1 large polypeptide that is cleaved by virus-encoded proteases into functional viral proteins. Can cause aseptic (viral) meningitis (except rhinovirus and HAV). All are enteroviruses except rhinovirus and HAV.

Pico**RNA**virus = small **RNA** virus.
PERCH on a “**peak**” (pico).

Rhinovirus


A picornavirus. Nonenveloped RNA virus. Cause of common cold; > 100 serologic types. Acid labile—destroyed by stomach acid; therefore, does not infect the GI tract (unlike the other picornaviruses).

Rhino has a runny **nose**.

Yellow fever virus

A flavivirus (also an arbovirus) transmitted by *Aedes* mosquitoes. Virus has a monkey or human reservoir. Symptoms: high fever, black vomitus, and jaundice. May see Councilman bodies (eosinophilic apoptotic globules) on liver biopsy.

Flavi = yellow, jaundice.

Rotavirus

Segmented dsRNA virus (a reovirus) **A**. Most important global cause of infantile gastroenteritis. Major cause of acute diarrhea in the United States during winter, especially in day care centers, kindergartens. Villous destruction with atrophy leads to ↓ absorption of Na^+ and loss of K^+ .

ROTAvirus = **R**ight **O**ut **T**he **A**nus. CDC recommends routine vaccination of all infants except those with a history of intussusception or SCID.

Influenza viruses

Orthomyxoviruses. Enveloped, \ominus ssRNA viruses with 8-segment genome. Contain hemagglutinin (binds sialic acid and promotes viral entry) and neuraminidase (promotes progeny virion release) antigens. Patients at risk for fatal bacterial superinfection, most commonly *S aureus*, *S pneumoniae*, and *H influenzae*.

Reformulated vaccine (“the flu shot”) contains viral strains most likely to appear during the flu season, due to the virus’ rapid genetic change. Killed viral vaccine is most frequently used. Live attenuated vaccine contains temperature-sensitive mutant that replicates in the nose but not in the lung; administered intranasally. Treatment: supportive +/- neuraminidase inhibitor (eg, oseltamivir, zanamivir).


Genetic/antigenic shift

Infection of 1 cell by 2 different segmented viruses (eg, swine influenza and human influenza viruses) → RNA segment reassortment → dramatically different virus (genetic shift) → major global outbreaks (pandemics).

Sudden shift is more deadly than gradual drift.

Genetic/antigenic drift

Random mutation in hemagglutinin or neuraminidase genes → minor changes (antigenic drift) → local outbreaks (epidemics).



Rubella virus




A togavirus. Causes rubella, once known as German (3-day) measles. Fever, postauricular and other lymphadenopathy, arthralgias, and fine, maculopapular rash that starts on face and spreads centrifugally to involve trunk and extremities **A**.

Causes mild disease in children but serious congenital disease (a TORCH infection). Congenital rubella findings include “blueberry muffin” appearance due to dermal extramedullary hematopoiesis.

Paramyxoviruses


Paramyxoviruses cause disease in children. They include those that cause parainfluenza (croup), mumps, measles, RSV, and human metapneumovirus, which causes respiratory tract infection (bronchiolitis, pneumonia) in infants. All contain surface F (fusion) protein, which causes respiratory epithelial cells to fuse and form multinucleated cells. Palivizumab (monoclonal antibody against F protein) prevents pneumonia caused by RSV infection in premature infants. **P**alivizumab for **P**aramyxovirus (RSV) **P**rophylaxis in **P**reemies.

Acute laryngotracheobronchitis



Also called croup. Caused by parainfluenza viruses. Virus membrane contains hemagglutinin (binds sialic acid and promotes viral entry) and neuraminidase (promotes progeny virion release) antigens. Results in a “seal-like” barking cough and inspiratory stridor. Narrowing of upper trachea and subglottis leads to characteristic steeple sign on x-ray **A**. Severe croup can result in pulsus paradoxus 2° to upper airway obstruction.

Measles (rubeola) virus



Usual presentation involves prodromal fever with cough, coryza, and conjunctivitis, then eventually Koplik spots (bright red spots with blue-white center on buccal mucosa **A**), followed 1–2 days later by a maculopapular rash **B** that starts at the head/neck and spreads downward.

Lymphadenitis with Warthin-Finkeldey giant cells (fused lymphocytes) in a background of paracortical hyperplasia. Possible sequelae:

- Subacute sclerosing panencephalitis (SSPE): personality changes, dementia, autonomic dysfunction, death (occurs years later)
- Encephalitis (1:1000): symptoms appear within few days of rash
- Giant cell pneumonia (rare except in immunosuppressed)

4 **C**'s of measles:

- C**ough
- C**oryza
- C**onjunctivitis
- “**C**”oplik spots

Vitamin A supplementation can reduce morbidity and mortality from measles, particularly in malnourished children.

Pneumonia is the most common cause of measles-associated death in children.



Mumps virus



Uncommon due to effectiveness of MMR vaccine.

Symptoms: **P**arotitis **A**, **O**rchitis (inflammation of testes), aseptic **M**eningitis, and **P**ancreatitis. Can cause sterility (especially after puberty).

Mumps makes your parotid glands and testes as big as **POM-Poms**.


Rabies virus

Bullet-shaped virus **A**. Negri bodies (cytoplasmic inclusions **B**) commonly found in Purkinje cells of cerebellum and in hippocampal neurons. Rabies has long incubation period (weeks to months) before symptom onset. Postexposure prophylaxis is wound cleaning plus immunization with killed vaccine and rabies immunoglobulin. Example of passive-active immunity.

Travels to the CNS by migrating in a retrograde fashion (via dynein motors) up nerve axons after binding to ACh receptors.

Progression of disease: fever, malaise
→ agitation, photophobia, hydrophobia, hypersalivation → paralysis, coma → death.

Infection more commonly from bat, raccoon, and skunk bites than from dog bites in the United States; aerosol transmission (eg, bat caves) also possible.

Ebola virus

A filovirus **A** that targets endothelial cells, phagocytes, hepatocytes. Following an incubation period of up to 21 days, presents with abrupt onset of flu-like symptoms, diarrhea/vomiting, high fever, myalgia. Can progress to DIC, diffuse hemorrhage, shock. Diagnosed with RT-PCR within 48 hr of symptom onset. High mortality rate.

Transmission requires direct contact with bodily fluids, fomites (including dead bodies), infected bats or primates (apes/monkeys); high incidence of nosocomial infection.

Supportive care, no definitive treatment. Strict isolation of infected individuals and barrier practices for health care workers are key to preventing transmission.

Zika virus

A flavivirus most commonly transmitted by *Aedes* mosquito bites. Causes conjunctivitis, low-grade pyrexia, and itchy rash in 20% of cases. Can lead to congenital microcephaly or miscarriage if transmitted in utero. Diagnose with RT-PCR or serology.

Sexual and vertical transmission possible. Outbreaks more common in tropical and subtropical climates. Supportive care, no definitive treatment.

Hepatitis viruses

Signs and symptoms of all hepatitis viruses: episodes of fever, jaundice, ↑ ALT and AST. Naked viruses (HAV and HEV) lack an envelope and are not destroyed by the gut: the **vowels** hit your **bowels**.

HBV DNA polymerase has DNA- and RNA-dependent activities. Upon entry into nucleus, the polymerase completes the partial dsDNA. Host RNA polymerase transcribes mRNA from viral DNA to make viral proteins. The DNA polymerase then reverse transcribes viral RNA to DNA, which is the genome of the progeny virus.

HCV lacks 3'-5' exonuclease activity → no proofreading ability → antigenic variation of HCV envelope proteins. Host antibody production lags behind production of new mutant strains of HCV.


Virus	HAV	HBV	HCV	HDV	HEV
FAMILY	RNA picornavirus	DNA hepadnavirus	RNA flavivirus	RNA deltavirus	RNA hepevirus
TRANSMISSION	Fecal-oral (shellfish, travelers, day care)	Parenteral (B lood), sexual (B aby-making), perinatal (B irthing)	Primarily blood (IVDU, post-transfusion)	Parenteral, sexual, perinatal	Fecal-oral, especially waterborne
INCUBATION	Short (weeks)	Long (months)	Long	Superinfection (HDV after HBV) = short Coinfection (HDV with HBV) = long	Short
CLINICAL COURSE	A cute and self limiting (adults), A symptomatic (children)	Initially like serum sickness (fever, arthralgias, rash); may progress to carcinoma	May progress to C irrhosis or C arcinoma	Similar to HBV	Fulminant hepatitis in E xpectant (pregnant) women
PROGNOSIS	Good	Adults → mostly full resolution; neonates → worse prognosis	Majority develop stable, C hronic hepatitis C	Superinfection → worse prognosis	High mortality in pregnant women
HCC RISK	No	Yes	Yes	Yes	No
LIVER BIOPSY	Hepatocyte swelling, monocyte infiltration, Councilman bodies	Granular eosinophilic "ground glass" appearance; cytotoxic T cells mediate damage	Lymphoid aggregates with focal areas of macrovesicular steatosis	Similar to HBV	Patchy necrosis
NOTES	No carrier state	Carrier state common	C arrier state very common	D efective virus, D epends on HBV HBsAg coat for entry into hepatocytes	E nteric, E pidemic (eg, in parts of Asia, Africa, Middle East), no carrier state

Extrahepatic manifestations of hepatitis B and C

	Hepatitis B	Hepatitis C
HEMATOLOGIC	Aplastic anemia	Essential mixed cryoglobulinemia, ↑ risk B-cell NHL, ITP, autoimmune hemolytic anemia
RENAL	Membranous GN > membranoproliferative GN	Membranoproliferative GN > membranous GN
VASCULAR	Polyarteritis nodosa	Leukocytoclastic vasculitis
DERMATOLOGIC		Sporadic porphyria cutanea tarda, lichen planus
ENDOCRINE		↑ risk of diabetes mellitus, autoimmune hypothyroidism


Hepatitis serologic markers

Anti-HAV (IgM)	IgM antibody to HAV; best test to detect acute hepatitis A.
Anti-HAV (IgG)	IgG antibody indicates prior HAV infection and/or prior vaccination; protects against reinfection.
HBsAg	Antigen found on surface of HBV; indicates hepatitis B infection.
Anti-HBs	Antibody to HBsAg; indicates immunity to hepatitis B due to vaccination or recovery from infection.
HBcAg	Antigen associated with core of HBV.
Anti-HBc	Antibody to HBcAg; IgM = acute/recent infection; IgG = prior exposure or chronic infection. IgM anti-HBc may be the sole ⊕ marker of infection during window period.
HBeAg	Secreted by infected hepatocyte into circulation. Not part of mature HBV virion. Indicates active viral replication and therefore high transmissibility and poorer prognosis.
Anti-HBe	Antibody to HBeAg; indicates low transmissibility.



	HBsAg	Anti-HBs	HBeAg	Anti-HBe	Anti-HBc
Acute HBV	✓		✓		IgM
Window				✓	IgM
Chronic HBV (high infectivity)	✓		✓		IgG
Chronic HBV (low infectivity)	✓			✓	IgG
Recovery		✓		✓	IgG
Immunized		✓			

HIV



Diploid genome (2 molecules of RNA).

The 3 structural genes (protein coded for):

- *env* (gp120 and gp41):
 - Formed from cleavage of gp160 to form envelope glycoproteins.
 - gp120—attachment to host CD4+ T cell.
 - gp41—fusion and entry.
- *gag* (p24 and p17)—capsid and matrix proteins, respectively.
- *pol*—Reverse transcriptase, Integrase, Protease; RIP “Pol” (Paul)

Reverse transcriptase synthesizes dsDNA from genomic RNA; dsDNA integrates into host genome.


Virus binds CD4 as well as a coreceptor, either CCR5 on macrophages (early infection) or CXCR4 on T cells (late infection).

Homozygous CCR5 mutation = immunity.
Heterozygous CCR5 mutation = slower course.


HIV diagnosis

Presumptive diagnosis made with HIV-1/2 Ag/Ab immunoassays. These immunoassays detect viral p24 Ag capsid protein and IgG Abs to HIV-1/2. Very high sensitivity/specificity. Viral load tests determine the amount of viral RNA in the plasma. High viral load associated with poor prognosis. Also use viral load to monitor effect of drug therapy. Use HIV genotyping to determine appropriate therapy. AIDS diagnosis: ≤ 200 CD4+ cells/mm³ (normal: 500–1500 cells/mm³) or HIV ⊕ with AIDS-defining condition (eg, *Pneumocystis pneumonia*).

Western blot tests are no longer recommended by the CDC for confirmatory testing. HIV-1/2 Ag/Ab testing is not recommended in babies with suspected HIV due to maternally transferred antibody. Use HIV viral load instead.



Time course of untreated HIV infection



Dashed lines on CD4+ count axis indicate moderate immunocompromise (< 400 CD4+ cells/mm³) and when AIDS-defining illnesses emerge (< 200 CD4+ cells/mm³).

Most patients who do not receive treatment eventually die of complications of HIV infection.

Four stages of untreated infection:

1. Flu-like (acute)
2. Feeling fine (latent)
3. Falling count
4. Final crisis

During clinical latency phase, virus replicates in lymph nodes

Common diseases of HIV-positive adults

↓ CD4+ cell count → reactivation of past infections (eg, TB, HSV, shingles), dissemination of bacterial infections and fungal infections (eg, coccidioidomycosis), and non-Hodgkin lymphomas.

PATHOGEN	PRESENTATION	FINDINGS
CD4+ cell count < 500/mm³		
<i>Candida albicans</i>	Oral thrush	Scrapable white plaque, pseudohyphae on microscopy
EBV	Oral hairy leukoplakia	Unscrapable white plaque on lateral tongue
HHV-8	Kaposi sarcoma	Biopsy with lymphocytic inflammation
HPV	Squamous cell carcinoma, commonly of anus (men who have sex with men) or cervix	
CD4+ cell count < 200/mm³		
<i>Histoplasma capsulatum</i>	Fever, weight loss, fatigue, cough, dyspnea, nausea, vomiting, diarrhea	Oval yeast cells within macrophages
HIV	Dementia	Cerebral atrophy on neuroimaging
JC virus (reactivation)	Progressive multifocal leukoencephalopathy	Nonenhancing areas of demyelination on MRI
<i>Pneumocystis jirovecii</i>	<i>Pneumocystis</i> pneumonia	“Ground-glass” opacities on chest imaging
CD4+ cell count < 100/mm³		
<i>Aspergillus fumigatus</i>	Hemoptysis, pleuritic pain	Cavitation or infiltrates on chest imaging
<i>Bartonella</i> spp	Bacillary angiomatosis	Biopsy with neutrophilic inflammation
<i>Candida albicans</i>	Esophagitis	White plaques on endoscopy; yeast and pseudohyphae on biopsy
CMV	Colitis, Retinitis, Esophagitis, Encephalitis, Pneumonitis (CREEP)	Linear ulcers on endoscopy, cotton-wool spots on funduscopy Biopsy reveals cells with intranuclear (owl eye) inclusion bodies
<i>Cryptococcus neoformans</i>	Meningitis	Encapsulated yeast on India ink stain or capsular antigen ⊕
<i>Cryptosporidium</i> spp	Chronic, watery diarrhea	Acid-fast oocysts in stool
EBV	B-cell lymphoma (eg, non-Hodgkin lymphoma, CNS lymphoma)	CNS lymphoma—ring enhancing, may be solitary (vs <i>Toxoplasma</i>)
<i>Mycobacterium avium–intracellulare</i> , <i>Mycobacterium avium</i> complex	Nonspecific systemic symptoms (fever, night sweats, weight loss) or focal lymphadenitis	Most common if CD4+ cell count < 50/mm ³
<i>Toxoplasma gondii</i>	Brain abscesses	Multiple ring-enhancing lesions on MRI

Prions

Prion diseases are caused by the conversion of a normal (predominantly α -helical) protein termed prion protein (PrP^c) to a β -pleated form (PrP^{sc}), which is transmissible via CNS-related tissue (iatrogenic CJD) or food contaminated by BSE-infected animal products (variant CJD). PrP^{sc} resists protease degradation and facilitates the conversion of still more PrP^c to PrP^{sc}. Resistant to standard sterilizing procedures, including standard autoclaving. Accumulation of PrP^{sc} results in spongiform encephalopathy and dementia, ataxia, and death.

Creutzfeldt-Jakob disease—rapidly progressive dementia, typically sporadic (some familial forms).

Bovine spongiform encephalopathy—also called “mad cow disease.”

Kuru—acquired prion disease noted in tribal populations practicing human cannibalism.

▶ MICROBIOLOGY—SYSTEMS

**Normal flora:
dominant**

Neonates delivered by C-section have no flora but are rapidly colonized after birth.

LOCATION	MICROORGANISM
Skin	<i>S epidermidis</i>
Nose	<i>S epidermidis</i> ; colonized by <i>S aureus</i>
Oropharynx	Viridans group streptococci
Dental plaque	<i>S mutans</i>
Colon	<i>B fragilis</i> > <i>E coli</i>
Vagina	<i>Lactobacillus</i> ; colonized by <i>E coli</i> and group B strep

**Bugs causing food-
borne illness**

S aureus and *B cereus* food poisoning starts quickly and ends quickly.

MICROORGANISM	SOURCE OF INFECTION
<i>B cereus</i>	Reheated rice. “Food poisoning from reheated rice? Be serious! ” (<i>B cereus</i>)
<i>C botulinum</i>	Improperly canned foods (toxins), raw honey (spores)
<i>C perfringens</i>	Reheated meat
<i>E coli</i> O157:H7	Undercooked meat
<i>L monocytogenes</i>	Deli meats, soft cheeses
<i>Salmonella</i>	Poultry, meat, and eggs
<i>S aureus</i>	Meats, mayonnaise, custard; preformed toxin
<i>V parahaemolyticus</i> and <i>V vulnificus</i> ^a	Raw/undercooked seafood

^a*V vulnificus* can also cause wound infections from contact with contaminated water or shellfish.

Bugs causing diarrhea

Bloody diarrhea	
<i>Campylobacter</i>	Comma- or S-shaped organisms; growth at 42°C
<i>E histolytica</i>	Protozoan; amebic dysentery; liver abscess
Enterohemorrhagic <i>E coli</i>	O157:H7; can cause HUS; makes Shiga-like toxin
Enteroinvasive <i>E coli</i>	Invades colonic mucosa
<i>Salmonella</i> (non-typhoidal)	Lactose ⊖; flagellar motility; has animal reservoir, especially poultry and eggs
<i>Shigella</i>	Lactose ⊖; very low ID ₅₀ ; produces Shiga toxin; human reservoir only; bacillary dysentery
<i>Y enterocolitica</i>	Day care outbreaks; pseudoappendicitis
Watery diarrhea	
<i>C difficile</i>	Pseudomembranous colitis; associated with antibiotics and PPIs; occasionally bloody diarrhea
<i>C perfringens</i>	Also causes gas gangrene
Enterotoxigenic <i>E coli</i>	Travelers' diarrhea; produces heat-labile (LT) and heat-stable (ST) toxins
Protozoa	<i>Giardia</i> , <i>Cryptosporidium</i>
<i>V cholerae</i>	Comma-shaped organisms; rice-water diarrhea; often from infected seafood
Viruses	Rotavirus, norovirus, enteric adenovirus

Common causes of pneumonia

NEONATES (< 4 WK)	CHILDREN (4 WK–18 YR)	ADULTS (18–40 YR)	ADULTS (40–65 YR)	ELDERLY
Group B streptococci	Viruses (RSV)	<i>Mycoplasma</i>	<i>S pneumoniae</i>	<i>S pneumoniae</i>
<i>E coli</i>	Mycoplasma	<i>C pneumoniae</i>	<i>H influenzae</i>	Influenza virus
	C trachomatis	<i>S pneumoniae</i>	Anaerobes	Anaerobes
	(infants–3 yr)	Viruses (eg, influenza)	Viruses	<i>H influenzae</i>
	C pneumoniae		<i>Mycoplasma</i>	Gram ⊖ rods
	(school-aged children)			
	S pneumoniae			
	Runts May Cough			
	Chunky Sputum			

Special groups

Alcoholic	<i>Klebsiella</i> , anaerobes usually due to aspiration (eg, <i>Peptostreptococcus</i> , <i>Fusobacterium</i> , <i>Prevotella</i> , <i>Bacteroides</i>)
IV drug users	<i>S pneumoniae</i> , <i>S aureus</i>
Aspiration	Anaerobes
Atypical	<i>Mycoplasma</i> , <i>Chlamydomphila</i> , <i>Legionella</i> , viruses (RSV, CMV, influenza, adenovirus)
Cystic fibrosis	<i>Pseudomonas</i> , <i>S aureus</i> , <i>S pneumoniae</i> , <i>Burkholderia cepacia</i>
Immunocompromised	<i>S aureus</i> , enteric gram ⊖ rods, fungi, viruses, <i>P jirovecii</i> (with HIV)
Nosocomial (hospital acquired)	<i>S aureus</i> , <i>Pseudomonas</i> , other enteric gram ⊖ rods
Postviral	<i>S pneumoniae</i> , <i>S aureus</i> , <i>H influenzae</i>

Common causes of meningitis

NEWBORN (0–6 MO)	CHILDREN (6 MO–6 YR)	6–60 YR	60 YR +
Group B <i>Streptococcus</i>	<i>S pneumoniae</i>	<i>N meningitidis</i>	<i>S pneumoniae</i>
<i>E coli</i>	<i>N meningitidis</i>	<i>S pneumoniae</i>	<i>N meningitidis</i>
<i>Listeria</i>	<i>H influenzae</i> type b	Enteroviruses	<i>H influenzae</i> type b
	Group B <i>Streptococcus</i>	HSV	Group B <i>Streptococcus</i>
	Enteroviruses		<i>Listeria</i>

Give ceftriaxone and vancomycin empirically (add ampicillin if *Listeria* is suspected).

Viral causes of meningitis: enteroviruses (especially coxsackievirus), HSV-2 (HSV-1 = encephalitis), HIV, West Nile virus (also causes encephalitis), VZV.

In HIV: *Cryptococcus* spp.

Note: Incidence of Group B streptococcal meningitis in neonates has ↓ greatly due to screening and antibiotic prophylaxis in pregnancy. Incidence of *H influenzae* meningitis has ↓ greatly due to conjugate *H influenzae* vaccinations. Today, cases are usually seen in unimmunized children.

Cerebrospinal fluid findings in meningitis

	OPENING PRESSURE	CELL TYPE	PROTEIN	GLUCOSE
Bacterial	↑	↑ PMNs	↑	↓
Fungal/TB	↑	↑ lymphocytes	↑	↓
Viral	Normal/↑	↑ lymphocytes	Normal/↑	Normal

Infections causing brain abscess

Most commonly viridans streptococci and *Staphylococcus aureus*. If dental infection or extraction precedes abscess, oral anaerobes commonly involved.

Multiple abscesses are usually from bacteremia; single lesions from contiguous sites: otitis media and mastoiditis → temporal lobe and cerebellum; sinusitis or dental infection → frontal lobe.

Toxoplasma reactivation in AIDS.

Osteomyelitis



RISK FACTOR	ASSOCIATED INFECTION
Assume if no other information is available	<i>S aureus</i> (most common overall)
Sexually active	<i>Neisseria gonorrhoeae</i> (rare), septic arthritis more common
Sickle cell disease	<i>Salmonella</i> and <i>S aureus</i>
Prosthetic joint replacement	<i>S aureus</i> and <i>S epidermidis</i>
Vertebral involvement	<i>S aureus</i> , <i>M tuberculosis</i> (Pott disease)
Cat and dog bites	<i>Pasteurella multocida</i>
IV drug abuse	<i>S aureus</i> ; also <i>Pseudomonas</i> , <i>Candida</i>

Elevated ESR and CRP sensitive but not specific.

Radiographs are insensitive early but can be useful in chronic osteomyelitis (A, left). MRI is best for detecting acute infection and detailing anatomic involvement (A, right).

Urinary tract infections

Cystitis presents with dysuria, frequency, urgency, suprapubic pain, and WBCs (but not WBC casts) in urine. Primarily caused by ascension of microbes from urethra to bladder. Ascension to kidney results in pyelonephritis, which presents with fever, chills, flank pain, costovertebral angle tenderness, hematuria, and WBC casts.


Ten times more common in women (shorter urethras colonized by fecal flora).

Risk factors: obstruction (eg, kidney stones, enlarged prostate), kidney surgery, catheterization, congenital GU malformation (eg, vesicoureteral reflux), diabetes, pregnancy.

SPECIES	FEATURES	COMMENTS
<i>Escherichia coli</i>	Leading cause of UTI. Colonies show strong pink lactose-fermentation on MacConkey agar.	Diagnostic markers: ⊕ Leukocyte esterase = evidence of WBC activity. ⊕ Nitrite test = reduction of urinary nitrates by gram ⊖ bacterial species (eg, <i>E coli</i>). ⊕ Urease test = urease-producing bugs (eg, <i>S saprophyticus</i> , <i>Proteus</i> , <i>Klebsiella</i>).
<i>Staphylococcus saprophyticus</i>	2nd leading cause of UTI in sexually active women.	
<i>Klebsiella pneumoniae</i>	3rd leading cause of UTI. Large mucoid capsule and viscous colonies.	
<i>Serratia marcescens</i>	Some strains produce a red pigment; often nosocomial and drug resistant.	
<i>Enterococcus</i>	Often nosocomial and drug resistant.	
<i>Proteus mirabilis</i>	Motility causes “swarming” on agar; associated with struvite stones.	
<i>Pseudomonas aeruginosa</i>	Blue-green pigment and fruity odor; usually nosocomial and drug resistant.	

Common vaginal infections

	Bacterial vaginosis	<i>Trichomonas vaginitis</i>	<i>Candida vulvovaginitis</i>
SIGNS AND SYMPTOMS	No inflammation Thin, white discharge A with fishy odor	Inflammation (“strawberry cervix”) Frothy, yellow-green, foul-smelling discharge	Inflammation Thick, white, “cottage cheese” discharge C
LAB FINDINGS	Clue cells pH > 4.5 ⊕ KOH whiff test	Motile pear-shaped trichomonads B pH > 4.5	Pseudohyphae pH normal (4.0–4.5)
TREATMENT	Metronidazole or clindamycin	Metronidazole Treat sexual partner(s)	Azoles



TORCH infections

Microbes that may pass from mother to fetus. Transmission is transplacental in most cases, or via delivery (especially HSV-2). Nonspecific signs common to many **ToRCHHeS** infections include hepatosplenomegaly, jaundice, thrombocytopenia, and growth retardation.

Other important infectious agents include *Streptococcus agalactiae* (group B streptococci), *E coli*, and *Listeria monocytogenes*—all causes of meningitis in neonates. Parvovirus B19 causes hydrops fetalis.

AGENT	MODES OF MATERNAL TRANSMISSION	MATERNAL MANIFESTATIONS	NEONATAL MANIFESTATIONS
Toxoplasma gondii	Cat feces or ingestion of undercooked meat	Usually asymptomatic; lymphadenopathy (rarely)	Classic triad: chorioretinitis, hydrocephalus, and intracranial calcifications, +/- “blueberry muffin” rash A
Rubella	Respiratory droplets	Rash, lymphadenopathy, polyarthrits, polyarthralgia	Classic triad: abnormalities of eye (cataracts B) and ear (deafness) and congenital heart disease (PDA); +/- “blueberry muffin” rash. “ I (eye) ♥ ruby (rubella) earrings ”
Cytomegalovirus	Sexual contact, organ transplants	Usually asymptomatic; mononucleosis-like illness	Hearing loss, seizures, petechial rash, “blueberry muffin” rash, chorioretinitis, periventricular calcifications C
HIV	Sexual contact, needlestick	Variable presentation depending on CD4+ cell count	Recurrent infections, chronic diarrhea
Herpes simplex virus-2	Skin or mucous membrane contact	Usually asymptomatic; herpetic (vesicular) lesions	Meningoencephalitis, herpetic (vesicular) lesions
Syphilis	Sexual contact	Chancre (1°) and disseminated rash (2°) are the two stages likely to result in fetal infection	Often results in stillbirth, hydrops fetalis; if child survives, presents with facial abnormalities (eg, notched teeth, saddle nose, short maxilla), saber shins, CN VIII deafness







Red rashes of childhood

AGENT	ASSOCIATED SYNDROME/DISEASE	CLINICAL PRESENTATION
Coxsackievirus type A	Hand-foot-mouth disease	Oval-shaped vesicles on palms and soles A ; vesicles and ulcers in oral mucosa (herpangina)
Human herpesvirus 6	Roseola (exanthem subitum)	Asymptomatic rose-colored macules appear on body after several days of high fever; can present with febrile seizures; usually affects infants
Measles virus	Measles (rubeola)	Confluent rash beginning at head and moving down; preceded by cough, coryza, conjunctivitis, and blue-white (Koplik) spots on buccal mucosa
Parvovirus B19	Erythema infectiosum (fifth disease)	“Slapped cheek” rash on face B (can cause hydrops fetalis in pregnant women)
Rubella virus	Rubella	Pink macules and papules begin at head and move down, remain discrete → fine desquamating truncal rash; postauricular lymphadenopathy
<i>Streptococcus pyogenes</i>	Scarlet fever	Flushed cheeks and circumoral pallor C on face; erythematous, sandpaper-like rash from neck to trunk and extremities; fever, sore throat, strawberry tongue
Varicella-Zoster virus	Chickenpox	Vesicular rash begins on trunk; spreads to face D and extremities with lesions of different stages



Sexually transmitted infections

DISEASE	CLINICAL FEATURES	ORGANISM
AIDS	Opportunistic infections, Kaposi sarcoma, lymphoma	HIV
Chancroid 	Painful genital ulcer with exudate, inguinal adenopathy A	<i>Haemophilus ducreyi</i> (it's so painful, you "do cry")
Chlamydia	Urethritis, cervicitis, epididymitis, conjunctivitis, reactive arthritis, PID	<i>Chlamydia trachomatis</i> (D–K)
Condylomata acuminata	Genital warts, koilocytes	HPV-6 and -11
Genital herpes	Painful penile, vulvar, or cervical vesicles and ulcers; can cause systemic symptoms such as fever, headache, myalgia	HSV-2, less commonly HSV-1
Gonorrhea	Urethritis, cervicitis, PID, prostatitis, epididymitis, arthritis, creamy purulent discharge	<i>Neisseria gonorrhoeae</i>
Granuloma inguinale (Donovanosis) 	Painless, beefy red ulcer that bleeds readily on contact B Uncommon in US	<i>Klebsiella (Calymmatobacterium) granulomatis</i> ; cytoplasmic Donovan bodies (bipolar staining) seen on microscopy
Hepatitis B	Jaundice	HBV
Lymphogranuloma venereum	Infection of lymphatics; painless genital ulcers, painful lymphadenopathy (ie, buboes)	<i>C trachomatis</i> (L1–L3)
Primary syphilis	Painless chancre	<i>Treponema pallidum</i>
Secondary syphilis	Fever, lymphadenopathy, skin rashes, condylomata lata	
Tertiary syphilis	Gummas, tabes dorsalis, general paresis, aortitis, Argyll Robertson pupil	
Trichomoniasis	Vaginitis, strawberry cervix, motile in wet prep	<i>Trichomonas vaginalis</i>

Pelvic inflammatory disease

Top bugs—*Chlamydia trachomatis* (subacute, often undiagnosed), *Neisseria gonorrhoeae* (acute).

C trachomatis—most common bacterial STI in the United States.

Signs include cervical motion tenderness, adnexal tenderness, purulent cervical discharge **A**.

PID may include salpingitis, endometritis, hydrosalpinx, and tubo-ovarian abscess.

Salpingitis is a risk factor for ectopic pregnancy, infertility, chronic pelvic pain, and adhesions. Can lead to perihepatitis (**Fitz-Hugh–Curtis syndrome**)—infection and inflammation of liver capsule and “violin string” adhesions of peritoneum to liver **B**.

Nosocomial infections *E coli* (UTI) and *S aureus* (wound infection) are the two most common causes.

RISK FACTOR	PATHOGEN	UNIQUE SIGNS/SYMPTOMS
Antibiotic use	<i>Clostridium difficile</i>	Watery diarrhea, leukocytosis
Aspiration (2° to altered mental status, old age)	Polymicrobial, gram \ominus bacteria, often anaerobes	Right lower lobe infiltrate or right upper/middle lobe (patient recumbent); purulent malodorous sputum
Decubitus ulcers, surgical wounds, drains	<i>S aureus</i> (including MRSA), gram \ominus anaerobes (<i>Bacteroides</i> , <i>Prevotella</i> , <i>Fusobacterium</i>)	Erythema, tenderness, induration, drainage from surgical wound sites
Intravascular catheters	<i>S aureus</i> (including MRSA), <i>S epidermidis</i> (long term), <i>Enterobacter</i>	Erythema, induration, tenderness, drainage from access sites
Mechanical ventilation, endotracheal intubation	Late onset: <i>P aeruginosa</i> , <i>Klebsiella</i> , <i>Acinetobacter</i> , <i>S aureus</i>	New infiltrate on CXR, \uparrow sputum production; sweet odor (<i>Pseudomonas</i>)
Renal dialysis unit, needlestick	HBV, HCV	
Urinary catheterization	<i>Proteus</i> spp, <i>E coli</i> , <i>Klebsiella</i> (infections in your PEcKer)	Dysuria, leukocytosis, flank pain or costovertebral angle tenderness
Water aerosols	<i>Legionella</i>	Signs of pneumonia, GI symptoms (diarrhea, nausea, vomiting), neurologic abnormalities

Bugs affecting unvaccinated children


CLINICAL PRESENTATION	FINDINGS/LABS	PATHOGEN
Dermatologic		
Rash	Beginning at head and moving down with postauricular lymphadenopathy	Rubella virus
	Beginning at head and moving down; preceded by cough, coryza, conjunctivitis, and Koplik spots	Measles virus
Neurologic		
Meningitis	Microbe colonizes nasopharynx	<i>H influenzae</i> type b
	Can also lead to myalgia and paralysis	Poliovirus
Tetanus	Muscle spasms and spastic paralysis (eg, lockjaw, opisthotonus)	<i>Clostridium tetani</i>
Respiratory		
Epiglottitis	Fever with dysphagia, drooling, and difficulty breathing due to edema	<i>H influenzae</i> type b (also capable of causing epiglottitis in fully immunized children)
Pertussis	Low-grade fevers, coryza → whooping cough, post-tussive vomiting → gradual recovery	<i>Bordetella pertussis</i>
Pharyngitis	Grayish pseudomembranes (may obstruct airways)	<i>Corynebacterium diphtheriae</i>

Bug hints

CHARACTERISTIC	ORGANISM
Asplenic patients	Encapsulated microbes, especially SHiN (<i>S pneumoniae</i> >> <i>H influenzae</i> type b > <i>N meningitidis</i>)
Branching rods in oral infection, sulfur granules	<i>Actinomyces israelii</i>
Chronic granulomatous disease	Catalase ⊕ microbes, especially <i>S aureus</i>
“Currant jelly” sputum	<i>Klebsiella</i>
Dog or cat bite	<i>Pasteurella multocida</i>
Facial nerve palsy (typically bilateral)	<i>Borrelia burgdorferi</i> (Lyme disease)
Human bite	Human oral flora (eg, <i>Eikenella</i> , <i>Fusobacterium</i>)
Neutropenic patients	<i>Candida albicans</i> (systemic), <i>Aspergillus</i>
Organ transplant recipient	CMV
PAS ⊕	<i>Tropheryma whipplei</i> (Whipple disease)
Pediatric infection	<i>Haemophilus influenzae</i> (including epiglottitis)
Pneumonia in cystic fibrosis, burn infection	<i>Pseudomonas aeruginosa</i>
Puncture wound, lockjaw	<i>Clostridium tetani</i>
Pus, empyema, abscess	<i>S aureus</i>
Rash on hands and feet	Coxsackie A, Rickettsii, Syphilis (CARS)
Sepsis/meningitis in newborn	Group B strep
Sinus/CNS infection in diabetics	<i>Mucor</i> or <i>Rhizopus</i> spp.
Surgical wound	<i>S aureus</i>
Traumatic open wound	<i>Clostridium perfringens</i>

► MICROBIOLOGY—ANTIMICROBIALS

Antimicrobial therapy

**Penicillin G, V**

Penicillin G (IV and IM form), penicillin V (oral). Prototype β -lactam antibiotics.

MECHANISM

D-Ala-D-Ala structural analog. Bind penicillin-binding proteins (transpeptidases). Block transpeptidase cross-linking of peptidoglycan in cell wall. Activate autolytic enzymes.

CLINICAL USE

Mostly used for gram \oplus organisms (*S pneumoniae*, *S pyogenes*, *Actinomyces*). Also used for gram \ominus cocci (mainly *N meningitidis*) and spirochetes (mainly *T pallidum*). Bactericidal for gram \oplus cocci, gram \oplus rods, gram \ominus cocci, and spirochetes. β -lactamase sensitive.

ADVERSE EFFECTS

Hypersensitivity reactions, direct Coombs \oplus hemolytic anemia, drug-induced interstitial nephritis.

RESISTANCE

β -lactamase cleaves the β -lactam ring. Mutations in PBPs.

Penicillinase-sensitive penicillins

	Amoxicillin, ampicillin; aminopenicillins.	
MECHANISM	Same as penicillin. Wider spectrum; penicillinase sensitive. Also combine with clavulanic acid to protect against destruction by β -lactamase.	AM ino P enicillins are AMP ed-up penicillin. Am O xicillin has greater O ral bioavailability than ampicillin.
CLINICAL USE	Extended-spectrum penicillin— H influenzae, H pylori, E coli, L isteria monocytogenes, P roteus mirabilis, S almonella, S higella, enterococci.	Coverage: ampicillin/amoxicillin HHELPSS kill enterococci.
ADVERSE EFFECTS	Hypersensitivity reactions, rash, pseudomembranous colitis.	
MECHANISM OF RESISTANCE	Penicillinase (a type of β -lactamase) cleaves β -lactam ring.	

Penicillinase-resistant penicillins

	Dicloxacillin, nafcillin, oxacillin.	
MECHANISM	Same as penicillin. Narrow spectrum; penicillinase resistant because bulky R group blocks access of β -lactamase to β -lactam ring.	
CLINICAL USE	<i>S aureus</i> (except MRSA).	“Use naf (nafcillin) for staph .”
ADVERSE EFFECTS	Hypersensitivity reactions, interstitial nephritis.	
MECHANISM OF RESISTANCE	MRSA has altered penicillin-binding protein target site.	

Antipseudomonal penicillins

	Piperacillin, ticarcillin.	
MECHANISM	Same as penicillin. Extended spectrum. Penicillinase sensitive; use with β -lactamase inhibitors.	
CLINICAL USE	<i>Pseudomonas</i> spp. and gram \ominus rods.	
ADVERSE EFFECTS	Hypersensitivity reactions.	

Cephalosporins

MECHANISM	β -lactam drugs that inhibit cell wall synthesis but are less susceptible to penicillinases. Bactericidal.	Organisms typically not covered by 1st–4th generation cephalosporins are LAME : L <i>isteria</i> , A typicals (<i>Chlamydia</i> , <i>Mycoplasma</i>), M RSA, and E nterococci.
CLINICAL USE	1st generation (cefazolin, cephalexin)—gram \oplus cocci, <i>Proteus mirabilis</i> , <i>E coli</i> , <i>Klebsiella pneumoniae</i> . Cefazolin used prior to surgery to prevent <i>S aureus</i> wound infections. 2nd generation (cefaclor, cefoxitin, cefuroxime, cefotetan)—gram \oplus cocci, <i>H influenzae</i> , <i>Enterobacter aerogenes</i> , <i>Neisseria</i> spp., <i>Serratia marcescens</i> , <i>Proteus mirabilis</i> , <i>E coli</i> , <i>Klebsiella pneumoniae</i> . 3rd generation (ceftriaxone, cefotaxime, cefpodoxime, ceftazidime)—serious gram \ominus infections resistant to other β -lactams. 4th generation (cefepime)—gram \ominus organisms, with \uparrow activity against <i>Pseudomonas</i> and gram \oplus organisms. 5th generation (ceftaroline)—broad gram \oplus and gram \ominus organism coverage; unlike 1st–4th generation cephalosporins, ceftaroline covers MRSA, and <i>Enterococcus faecalis</i> —does not cover <i>Pseudomonas</i> .	1st generation— \oplus PEcK . 2nd graders wear fake fox fur to tea parties. 2nd generation— \oplus HENS PEcK . Can cross blood-brain barrier. Ceftriaxone—meningitis, gonorrhea, disseminated Lyme disease. Ceftazidime— <i>Pseudomonas</i> .
ADVERSE EFFECTS	Hypersensitivity reactions, autoimmune hemolytic anemia, disulfiram-like reaction, vitamin K deficiency. Low rate of cross-reactivity even in penicillin-allergic patients. \uparrow nephrotoxicity of aminoglycosides.	
MECHANISM OF RESISTANCE	Inactivated by cephalosporinases (a type of β -lactamase). Structural change in penicillin-binding proteins (transpeptidases).	
β-lactamase inhibitors	Include C lavulanic acid, A vibactam, S ulbactam, T azobactam. Often added to penicillin antibiotics to protect the antibiotic from destruction by β -lactamase.	CAST (eg, amoxicillin-clavulanate, ceftazidime-avibactam, ampicillin-sulbactam, piperacillin-tazobactam).

Carbapenems

Doripenem, Imipenem, Meropenem, Ertapenem (**DIME** antibiotics are given when there is a 10/10 [life-threatening] infection).


MECHANISM	Imipenem is a broad-spectrum, β -lactamase-resistant carbapenem. Always administered with cilastatin (inhibitor of renal dehydropeptidase I) to \downarrow inactivation of drug in renal tubules.	With imipenem, “the kill is lastin ’ with cilastatin .” Newer carbapenems include ertapenem (limited <i>Pseudomonas</i> coverage) and doripenem.
CLINICAL USE	Gram \oplus cocci, gram \ominus rods, and anaerobes. Wide spectrum and significant side effects limit use to life-threatening infections or after other drugs have failed. Meropenem has a \downarrow risk of seizures and is stable to dehydropeptidase I.	
ADVERSE EFFECTS	GI distress, rash, and CNS toxicity (seizures) at high plasma levels.	
MECHANISM OF RESISTANCE	Inactivated by carbapenemases produced by, eg, <i>K pneumoniae</i> , <i>E coli</i> , <i>E aerogenes</i> .	

Monobactams


Aztreonam

MECHANISM	Less susceptible to β -lactamases. Prevents peptidoglycan cross-linking by binding to penicillin-binding protein 3. Synergistic with aminoglycosides. No cross-allergenicity with penicillins.
CLINICAL USE	Gram \ominus rods only—no activity against gram \oplus rods or anaerobes. For penicillin-allergic patients and those with renal insufficiency who cannot tolerate aminoglycosides.
ADVERSE EFFECTS	Usually nontoxic; occasional GI upset.

Vancomycin

MECHANISM	Inhibits cell wall peptidoglycan formation by binding D-Ala-D-Ala portion of cell wall precursors. Bactericidal against most bacteria (bacteriostatic against <i>C difficile</i>). Not susceptible to β -lactamases.
CLINICAL USE	Gram \oplus bugs only—for serious, multidrug-resistant organisms, including MRSA, <i>S epidermidis</i> , sensitive <i>Enterococcus</i> species, and <i>Clostridium difficile</i> (oral dose for pseudomembranous colitis).
ADVERSE EFFECTS	Well tolerated in general but NOT trouble Free . N ephrotoxicity, O tototoxicity, T hrombophlebitis, diffuse F lushing (red man syndrome A idiopathic reaction largely preventable by pretreatment with antihistamines), DRESS syndrome.
	
MECHANISM OF RESISTANCE	Occurs in bacteria (eg, <i>Enterococcus</i>) via amino acid modification of D-Ala-D-Ala to D-Ala-D-Lac . “If you Lack a D-Ala (dollar), you can’t ride the van (vancomycin).”

Protein synthesis inhibitors



Specifically target smaller bacterial ribosome (70S, made of 30S and 50S subunits), leaving human ribosome (80S) unaffected.

All are bacteriostatic, except aminoglycosides (bactericidal) and linezolid (variable).

30S inhibitors

- Aminoglycosides
- Tetracyclines

50S inhibitors

- Chloramphenicol, Clindamycin
- Erythromycin (macrolides)
- Linezolid

“Buy AT 30, CCEL (sell) at 50.”

Aminoglycosides

Gentamicin, Neomycin, Amikacin, Tobramycin, Streptomycin.

“Mean” (aminoglycoside) GNATS caNNOT kill anaerobes.

MECHANISM	Bactericidal; irreversible inhibition of initiation complex through binding of the 30S subunit. Can cause misreading of mRNA. Also block translocation. Require O ₂ for uptake; therefore ineffective against anaerobes.
CLINICAL USE	Severe gram ⊖ rod infections. Synergistic with β-lactam antibiotics. Neomycin for bowel surgery.
ADVERSE EFFECTS	Nephrotoxicity, Neuromuscular blockade (absolute contraindication with myasthenia gravis), Ototoxicity (especially with loop diuretics), Teratogenicity.
MECHANISM OF RESISTANCE	Bacterial transferase enzymes inactivate the drug by acetylation, phosphorylation, or adenylation.

Tetracyclines

Tetracycline, doxycycline, minocycline.

MECHANISM	Bacteriostatic; bind to 30S and prevent attachment of aminoacyl-tRNA. Limited CNS penetration. Doxycycline is fecally eliminated and can be used in patients with renal failure. Do not take tetracyclines with milk (Ca^{2+}), antacids (eg, Ca^{2+} or Mg^{2+}), or iron-containing preparations because divalent cations inhibit drugs' absorption in the gut.
CLINICAL USE	<i>Borrelia burgdorferi</i> , <i>M pneumoniae</i> . Drugs' ability to accumulate intracellularly makes them very effective against <i>Rickettsia</i> and <i>Chlamydia</i> . Also used to treat acne. Doxycycline effective against community-acquired MRSA.
ADVERSE EFFECTS	GI distress, discoloration of teeth and inhibition of bone growth in children, photosensitivity. Contraindicated in pregnancy.
MECHANISM OF RESISTANCE	↓ uptake or ↑ efflux out of bacterial cells by plasmid-encoded transport pumps.

Tigecycline

MECHANISM	Tetracycline derivative. Binds to 30S, inhibiting protein synthesis. Generally bacteriostatic.
CLINICAL USE	Broad-spectrum anaerobic, gram \ominus , and gram \oplus coverage. Multidrug-resistant organisms (MRSA, VRE) or infections requiring deep tissue penetration.
ADVERSE EFFECTS	GI symptoms: nausea, vomiting.

Chloramphenicol

MECHANISM	Blocks peptidyltransferase at 50S ribosomal subunit. Bacteriostatic.
CLINICAL USE	Meningitis (<i>Haemophilus influenzae</i> , <i>Neisseria meningitidis</i> , <i>Streptococcus pneumoniae</i>) and rickettsial diseases (eg, Rocky Mountain spotted fever [<i>Rickettsia rickettsii</i>]). Limited use due to toxicity but often still used in developing countries because of low cost.
ADVERSE EFFECTS	Anemia (dose dependent), aplastic anemia (dose independent), gray baby syndrome (in premature infants because they lack liver UDP-glucuronosyltransferase).
MECHANISM OF RESISTANCE	Plasmid-encoded acetyltransferase inactivates the drug.

Clindamycin

MECHANISM	Blocks peptide transfer (translocation) at 50S ribosomal subunit. Bacteriostatic.
CLINICAL USE	Anaerobic infections (eg, <i>Bacteroides</i> spp., <i>Clostridium perfringens</i>) in aspiration pneumonia, lung abscesses, and oral infections. Also effective against invasive group A streptococcal infection. Treats anaerobic infections above the diaphragm vs metronidazole (anaerobic infections below diaphragm).
ADVERSE EFFECTS	Pseudomembranous colitis (<i>C difficile</i> overgrowth), fever, diarrhea.

Linezolid

MECHANISM	Inhibits protein synthesis by binding to 50S subunit and preventing formation of the initiation complex.
CLINICAL USE	Gram ⊕ species including MRSA and VRE.
ADVERSE EFFECTS	Bone marrow suppression (especially thrombocytopenia), peripheral neuropathy, serotonin syndrome (due to partial MAO inhibition).
MECHANISM OF RESISTANCE	Point mutation of ribosomal RNA.

Macrolides

Azithromycin, clarithromycin, erythromycin.

MECHANISM	Inhibit protein synthesis by blocking translocation (“macroslides”); bind to the 23S rRNA of the 50S ribosomal subunit. Bacteriostatic.
CLINICAL USE	Atypical pneumonias (<i>Mycoplasma</i> , <i>Chlamydia</i> , <i>Legionella</i>), STIs (<i>Chlamydia</i>), gram ⊕ cocci (streptococcal infections in patients allergic to penicillin), and <i>B pertussis</i> .
ADVERSE EFFECTS	MACRO: Gastrointestinal Motility issues, Arrhythmia caused by prolonged QT interval, acute Cholestatic hepatitis, Rash, eOsinophilia. Increases serum concentration of theophylline, oral anticoagulants. Clarithromycin and erythromycin inhibit cytochrome P-450.
MECHANISM OF RESISTANCE	Methylation of 23S rRNA-binding site prevents binding of drug.

Polymyxins

Colistin (polymyxin E), polymyxin B.

MECHANISM	Cation polypeptides that bind to phospholipids on cell membrane of gram ⊖ bacteria. Disrupt cell membrane integrity → leakage of cellular components → cell death.
CLINICAL USE	Salvage therapy for multidrug-resistant gram ⊖ bacteria (eg, <i>P aeruginosa</i> , <i>E coli</i> , <i>K pneumoniae</i>). Polymyxin B is a component of a triple antibiotic ointment used for superficial skin infections.
ADVERSE EFFECTS	Nephrotoxicity, neurotoxicity (eg, slurred speech, weakness, paresthesias), respiratory failure.

Sulfonamides

Sulfamethoxazole (SMX), sulfisoxazole, sulfadiazine.


MECHANISM	Inhibit dihydropteroate synthase, thus inhibiting folate synthesis. Bacteriostatic (bactericidal when combined with trimethoprim).
CLINICAL USE	Gram \oplus , gram \ominus , <i>Nocardia</i> . TMP-SMX for simple UTI.
ADVERSE EFFECTS	Hypersensitivity reactions, hemolysis if G6PD deficient, nephrotoxicity (tubulointerstitial nephritis), photosensitivity, Stevens-Johnson syndrome, kernicterus in infants, displace other drugs from albumin (eg, warfarin).
MECHANISM OF RESISTANCE	Altered enzyme (bacterial dihydropteroate synthase), \downarrow uptake, or \uparrow PABA synthesis.

Dapsone

MECHANISM	Similar to sulfonamides, but structurally distinct agent.
CLINICAL USE	Leprosy (lepromatous and tuberculoid), <i>Pneumocystis jirovecii</i> prophylaxis, or treatment when used in combination with TMP.
ADVERSE EFFECTS	Hemolysis if G6PD deficient, methemoglobinemia, agranulocytosis.

Trimethoprim


MECHANISM	Inhibits bacterial dihydrofolate reductase. Bacteriostatic.
CLINICAL USE	Used in combination with sulfonamides (trimethoprim-sulfamethoxazole [TMP-SMX]), causing sequential block of folate synthesis. Combination used for UTIs, <i>Shigella</i> , <i>Salmonella</i> , <i>Pneumocystis jirovecii</i> pneumonia treatment and prophylaxis, toxoplasmosis prophylaxis.
ADVERSE EFFECTS	Hyperkalemia (high doses), megaloblastic anemia, leukopenia, granulocytopenia, which may be avoided with coadministration of leucovorin (folinic acid). TMP Treats Marrow Poorly.



Fluoroquinolones	Ciprofloxacin, enoxacin, norfloxacin, ofloxacin; respiratory fluoroquinolones—gemifloxacin, levofloxacin, moxifloxacin.	
MECHANISM	Inhibit prokaryotic enzymes topoisomerase II (DNA gyrase) and topoisomerase IV. Bactericidal. Must not be taken with antacids.	
CLINICAL USE	Gram \ominus rods of urinary and GI tracts (including <i>Pseudomonas</i>), some gram \oplus organisms, otitis externa.	
ADVERSE EFFECTS	GI upset, superinfections, skin rashes, headache, dizziness. Less commonly, can cause leg cramps and myalgias. Contraindicated in pregnant women, nursing mothers, and children < 18 years old due to possible damage to cartilage. Some may prolong QT interval. May cause tendonitis or tendon rupture in people > 60 years old and in patients taking prednisone. Ciprofloxacin inhibits cytochrome P-450.	Fluoroquinolones hurt attachments to your bones .
MECHANISM OF RESISTANCE	Chromosome-encoded mutation in DNA gyrase, plasmid-mediated resistance, efflux pumps.	
Daptomycin		
MECHANISM	Lipopeptide that disrupts cell membranes of gram \oplus cocci by creating transmembrane channels.	
CLINICAL USE	<i>S aureus</i> skin infections (especially MRSA), bacteremia, endocarditis, VRE.	Not used for pneumonia (avidly binds to and is inactivated by surfactant). “Daptomy skin ” is used for skin infections.
ADVERSE EFFECTS	Myopathy, rhabdomyolysis.	
Metronidazole		
MECHANISM	Forms toxic free radical metabolites in the bacterial cell that damage DNA. Bactericidal, antiprotozoal.	
CLINICAL USE	Treats <i>Giardia</i> , <i>Entamoeba</i> , <i>Trichomonas</i> , <i>Gardnerella vaginalis</i> , Anaerobes (<i>Bacteroides</i> , <i>C difficile</i>). Can be used in place of amoxicillin in <i>H pylori</i> “triple therapy” in case of penicillin allergy.	GET GAP on the Metro with metronidazole ! Treats anaerobic infection below the diaphragm vs clindamycin (anaerobic infections above diaphragm).
ADVERSE EFFECTS	Disulfiram-like reaction (severe flushing, tachycardia, hypotension) with alcohol; headache, metallic taste.	

Antimycobacterial therapy

BACTERIUM	PROPHYLAXIS	TREATMENT
<i>M tuberculosis</i>	Isoniazid	Rifampin, Isoniazid, Pyrazinamide, Ethambutol (RIPE for treatment)
<i>M avium–intracellulare</i>	Azithromycin, rifabutin	Azithromycin or clarithromycin + ethambutol Can add rifabutin or ciprofloxacin
<i>M leprae</i>	N/A	Long-term treatment with dapsone and rifampin for tuberculoid form Add clofazimine for lepromatous form

MYCOBACTERIAL CELL**Rifamycins**

Rifampin, rifabutin.

MECHANISM	Inhibit DNA-dependent RNA polymerase.	Rifampin's 4 R's : RNA polymerase inhibitor Ramps up microsomal cytochrome P-450 Red/orange body fluids Rapid resistance if used alone
CLINICAL USE	<i>Mycobacterium tuberculosis</i> ; delay resistance to dapsone when used for leprosy. Used for meningococcal prophylaxis and chemoprophylaxis in contacts of children with <i>H influenzae</i> type b.	Rifampin ramps up cytochrome P-450, but rifabutin does not.
ADVERSE EFFECTS	Minor hepatotoxicity and drug interactions (↑ cytochrome P-450); orange body fluids (nonhazardous side effect). Rifabutin favored over rifampin in patients with HIV infection due to less cytochrome P-450 stimulation.	
MECHANISM OF RESISTANCE	Mutations reduce drug binding to RNA polymerase. Monotherapy rapidly leads to resistance.	

Isoniazid

MECHANISM	↓ synthesis of mycolic acids. Bacterial catalase-peroxidase (encoded by KatG) needed to convert INH to active metabolite.	
CLINICAL USE	<i>Mycobacterium tuberculosis</i> . The only agent used as solo prophylaxis against TB. Also used as monotherapy for latent TB.	Different INH half-lives in fast vs slow acetylators.
ADVERSE EFFECTS	Hepatotoxicity, cytochrome P-450 inhibition, drug-induced SLE, anion gap metabolic acidosis, vitamin B ₆ deficiency (peripheral neuropathy, sideroblastic anemia), seizures (in high doses, refractory to benzodiazepines). Administer with pyridoxine (B ₆).	INH Injures Neurons and Hepatocytes.
MECHANISM OF RESISTANCE	Mutations leading to underexpression of KatG.	

Pyrazinamide

MECHANISM	Mechanism uncertain. Pyrazinamide is a prodrug that is converted to the active compound pyrazinoic acid. Works best at acidic pH (eg, in host phagolysosomes).	
CLINICAL USE	<i>Mycobacterium tuberculosis</i> .	
ADVERSE EFFECTS	Hyperuricemia, hepatotoxicity.	

Ethambutol

MECHANISM	↓ carbohydrate polymerization of mycobacterium cell wall by blocking arabinosyltransferase.	
CLINICAL USE	<i>Mycobacterium tuberculosis</i> .	
ADVERSE EFFECTS	Optic neuropathy (red-green color blindness, usually reversible). Pronounce “ ey ethambutol.”	

Streptomycin

MECHANISM	Interferes with 30S component of ribosome.	
CLINICAL USE	<i>Mycobacterium tuberculosis</i> (2nd line).	
ADVERSE EFFECTS	Tinnitus, vertigo, ataxia, nephrotoxicity.	

Antimicrobial prophylaxis


CLINICAL SCENARIO	MEDICATION
Exposure to meningococcal infection	Ceftriaxone, ciprofloxacin, or rifampin
High risk for endocarditis and undergoing surgical or dental procedures	Amoxicillin
History of recurrent UTIs	TMP-SMX
Malaria prophylaxis for travelers	Atovaquone-proguanil, mefloquine, doxycycline, primaquine, or chloroquine (for areas with sensitive species)
Pregnant woman carrying group B strep	Intrapartum penicillin G or ampicillin
Prevention of gonococcal conjunctivitis in newborn	Erythromycin ointment on eyes
Prevention of postsurgical infection due to <i>S aureus</i>	Cefazolin
Prophylaxis of strep pharyngitis in child with prior rheumatic fever	Benzathine penicillin G or oral penicillin V

Prophylaxis in HIV/AIDS patients

CELL COUNT	PROPHYLAXIS	INFECTION
CD4 < 200 cells/mm ³	TMP-SMX	<i>Pneumocystis pneumonia</i>
CD4 < 100 cells/mm ³	TMP-SMX	<i>Pneumocystis pneumonia</i> and toxoplasmosis
CD4 < 50 cells/mm ³	Azithromycin or clarithromycin	<i>Mycobacterium avium</i> complex

Treatment of highly resistant bacteria

MRSA: vancomycin, daptomycin, linezolid, tigecycline, ceftaroline, doxycycline.
 VRE: linezolid, tigecycline, and streptogramins (quinupristin, dalbavand).
 Multidrug-resistant *P aeruginosa*, multidrug-resistant *Acinetobacter baumannii*: polymyxins B and E (colistin).

Antifungal therapy

Amphotericin B

MECHANISM	Binds ergosterol (unique to fungi); forms membrane pores that allow leakage of electrolytes.	Amphotericin “tears” holes in the fungal membrane by forming pores.
CLINICAL USE	Serious, systemic mycoses. <i>Cryptococcus</i> (amphotericin B with +/- without flucytosine for cryptococcal meningitis), <i>Blastomyces</i> , <i>Coccidioides</i> , <i>Histoplasma</i> , <i>Candida</i> , <i>Mucor</i> . Intrathecally for coccidioidal meningitis.	Supplement K ⁺ and Mg ²⁺ because of altered renal tubule permeability.
ADVERSE EFFECTS	Fever/chills (“shake and bake”), hypotension, nephrotoxicity, arrhythmias, anemia, IV phlebitis (“amphoterrible”). Hydration ↓ nephrotoxicity. Liposomal amphotericin ↓ toxicity.	

Nystatin

MECHANISM	Same as amphotericin B. Topical use only as too toxic for systemic use.
CLINICAL USE	“Swish and swallow” for oral candidiasis (thrush); topical for diaper rash or vaginal candidiasis.

Flucytosine

MECHANISM	Inhibits DNA and RNA biosynthesis by conversion to 5-fluorouracil by cytosine deaminase.
CLINICAL USE	Systemic fungal infections (especially meningitis caused by <i>Cryptococcus</i>) in combination with amphotericin B.
ADVERSE EFFECTS	Bone marrow suppression.

Azoles

MECHANISM	Clotrimazole, fluconazole, isavuconazole, itraconazole, ketoconazole, miconazole, voriconazole. Inhibit fungal sterol (ergosterol) synthesis by inhibiting the cytochrome P-450 enzyme that converts lanosterol to ergosterol.
CLINICAL USE	Local and less serious systemic mycoses. Fluconazole for chronic suppression of cryptococcal meningitis in AIDS patients and candidal infections of all types. Itraconazole may be used for <i>Blastomyces</i> , <i>Coccidioides</i> , <i>Histoplasma</i> , <i>Sporothrix schenckii</i> . Clotrimazole and miconazole for topical fungal infections. Voriconazole for <i>Aspergillus</i> and some <i>Candida</i> . Isavuconazole for serious <i>Aspergillus</i> and <i>Mucor</i> infections.
ADVERSE EFFECTS	Testosterone synthesis inhibition (gynecomastia, especially with ketoconazole), liver dysfunction (inhibits cytochrome P-450).

Terbinafine

MECHANISM	Inhibits the fungal enzyme squalene epoxidase.
CLINICAL USE	Dermatophytoses (especially onychomycosis—fungal infection of finger or toe nails).
ADVERSE EFFECTS	GI upset, headaches, hepatotoxicity, taste disturbance.

Echinocandins	Anidulafungin, caspofungin, micafungin.
MECHANISM	Inhibit cell wall synthesis by inhibiting synthesis of β -glucan.
CLINICAL USE	Invasive aspergillosis, <i>Candida</i> .
ADVERSE EFFECTS	GI upset, flushing (by histamine release).

Griseofulvin

MECHANISM	Interferes with microtubule function; disrupts mitosis. Deposits in keratin-containing tissues (eg, nails).
CLINICAL USE	Oral treatment of superficial infections; inhibits growth of dermatophytes (tinea, ringworm).
ADVERSE EFFECTS	Teratogenic, carcinogenic, confusion, headaches, disulfiram-like reaction, \uparrow cytochrome P-450 and warfarin metabolism.

Antiprotozoal therapy	Pyrimethamine (toxoplasmosis), suramin and melarsoprol (<i>Trypanosoma brucei</i>), nifurtimox (<i>T cruzi</i>), sodium stibogluconate (leishmaniasis).
------------------------------	---


Anti-mite/louse therapy	Permethrin (inhibits Na^+ channel deactivation \rightarrow neuronal membrane depolarization), malathion (acetylcholinesterase inhibitor), topical +/- oral ivermectin. Used to treat scabies (<i>Sarcoptes scabiei</i>) and lice (<i>Pediculus</i> and <i>Pthirus</i>).
--------------------------------	--

Chloroquine

MECHANISM	Blocks detoxification of heme into hemozoin. Heme accumulates and is toxic to plasmodia.
CLINICAL USE	Treatment of plasmodial species other than <i>P falciparum</i> (frequency of resistance in <i>P falciparum</i> is too high). Resistance due to membrane pump that \downarrow intracellular concentration of drug. Treat <i>P falciparum</i> with artemether/lumefantrine or atovaquone/proguanil. For life-threatening malaria, use quinidine in US (quinine elsewhere) or artesunate.
ADVERSE EFFECTS	Retinopathy; pruritus (especially in dark-skinned individuals).

Anthelmintic therapy	Pyrantel pamoate, Ivermectin, Mebendazole (microtubule inhibitor), Praziquantel ($\uparrow \text{Ca}^{2+}$ permeability, \uparrow vacuolization), Diethylcarbamazine. Helminths get PIMP'D.
-----------------------------	--

Antiviral therapy



Oseltamivir, zanamivir

MECHANISM

Inhibit influenza neuraminidase → ↓ release of progeny virus.

CLINICAL USE

Treatment and prevention of influenza A and B. Beginning therapy within 48 hours of symptom onset may shorten duration of illness.

Acyclovir, famciclovir, valacyclovir

MECHANISM

Guanosine analogs. Monophosphorylated by HSV/VZV thymidine kinase and not phosphorylated in uninfected cells → few adverse effects. Triphosphate formed by cellular enzymes. Preferentially inhibit viral DNA polymerase by chain termination.

CLINICAL USE

HSV and VZV. Weak activity against EBV. No activity against CMV. Used for HSV-induced mucocutaneous and genital lesions as well as for encephalitis. Prophylaxis in immunocompromised patients. No effect on latent forms of HSV and VZV. Valacyclovir, a prodrug of acyclovir, has better oral bioavailability. For herpes zoster, use famciclovir.

ADVERSE EFFECTS

Obstructive crystalline nephropathy and acute kidney injury if not adequately hydrated.

MECHANISM OF RESISTANCE

Mutated viral thymidine kinase.

Ganciclovir

MECHANISM	5'-monophosphate formed by a CMV viral kinase. Guanosine analog. Triphosphate formed by cellular kinases. Preferentially inhibits viral DNA polymerase.
CLINICAL USE	CMV, especially in immunocompromised patients. Valganciclovir, a prodrug of ganciclovir, has better oral bioavailability.
ADVERSE EFFECTS	Bone marrow suppression (leukopenia, neutropenia, thrombocytopenia), renal toxicity. More toxic to host enzymes than acyclovir.
MECHANISM OF RESISTANCE	Mutated viral kinase.

Foscarnet

MECHANISM	Viral DNA/RNA polymerase inhibitor and HIV reverse transcriptase inhibitor. Binds to pyrophosphate-binding site of enzyme. Does not require any kinase activation.	Foscarnet = pyro fos phate analog.
CLINICAL USE	CMV retinitis in immunocompromised patients when ganciclovir fails; acyclovir-resistant HSV.	
ADVERSE EFFECTS	Nephrotoxicity, electrolyte abnormalities (hypo- or hypercalcemia, hypo- or hyperphosphatemia, hypokalemia, hypomagnesemia) can lead to seizures.	
MECHANISM OF RESISTANCE	Mutated DNA polymerase.	

Cidofovir

MECHANISM	Preferentially inhibits viral DNA polymerase. Does not require phosphorylation by viral kinase.
CLINICAL USE	CMV retinitis in immunocompromised patients; acyclovir-resistant HSV. Long half-life.
ADVERSE EFFECTS	Nephrotoxicity (coadminister with probenecid and IV saline to ↓ toxicity).

HIV therapy

Antiretroviral therapy (ART): often initiated at the time of HIV diagnosis.

Strongest indication for use with patients presenting with AIDS-defining illness, low CD4+ cell counts (< 500 cells/mm³), or high viral load. Regimen consists of 3 drugs to prevent resistance: 2 NRTIs and preferably an integrase inhibitor.

All ARTs are active against HIV-1 and HIV-2 with the exception of NNRTIs and enfuvirtide.

DRUG	MECHANISM	TOXICITY
NRTIs		
Abacavir (ABC) Didanosine (ddI) Emtricitabine (FTC) Lamivudine (3TC) Stavudine (d4T) Tenofovir (TDF) Zidovudine (ZDV, formerly AZT)	Competitively inhibit nucleotide binding to reverse transcriptase and terminate the DNA chain (lack a 3' OH group). T enofovir is a nucleo T ide; the others are nucleosides. All need to be phosphorylated to be active. ZDV can be used for general prophylaxis and during pregnancy to ↓ risk of fetal transmission. Have you dined (vudine) with my nuclear (nucleosides) family?	Bone marrow suppression (can be reversed with granulocyte colony-stimulating factor [G-CSF] and erythropoietin), peripheral neuropathy, lactic acidosis (nucleosides), anemia (ZDV), pancreatitis (didanosine). Abacavir contraindicated if patient has HLA-B*5701 mutation due to ↑ risk of hypersensitivity.
NNRTIs		
Delavirdine Efavirenz Nevirapine	Bind to reverse transcriptase at site different from NRTIs. Do not require phosphorylation to be active or compete with nucleotides.	Rash and hepatotoxicity are common to all NNRTIs. Vivid dreams and CNS symptoms are common with efavirenz.
Integrase inhibitors		
Bictegravir Dolutegravir Elvitegravir Raltegravir	Inhibits HIV genome integration into host cell chromosome by reversibly inhibiting HIV integrase.	↑ creatine kinase.
Protease inhibitors		
Atazanavir Darunavir Fosamprenavir Indinavir Lopinavir Ritonavir Saquinavir	Assembly of virions depends on HIV-1 protease (<i>pol</i> gene), which cleaves the polypeptide products of HIV mRNA into their functional parts. Thus, protease inhibitors prevent maturation of new viruses. Ritonavir can “boost” other drug concentrations by inhibiting cytochrome P-450. Navir (never) tease a protease.	Hyperglycemia, GI intolerance (nausea, diarrhea), lipodystrophy (Cushing-like syndrome). Nephropathy, hematuria, thrombocytopenia (indinavir). Rifampin (potent CYP/UGT inducer) reduces protease inhibitor concentrations; use rifabutin instead.
Entry inhibitors		
Enfuvirtide	Binds gp41, inhibiting viral entry.	Skin reaction at injection sites. En f uvirtide inhibits f usion.
Maraviroc	Binds CCR-5 on surface of T cells/monocytes, inhibiting interaction with gp120.	Maraviro c inhibits d ocking.

Hepatitis C therapy Chronic HCV infection treated with multidrug therapy that targets specific steps within HCV replication cycle (HCV-encoded proteins). Examples of drugs are provided.

DRUG	MECHANISM	TOXICITY
NS5A inhibitors		
Ledipasvir	Inhibits NS5A, a viral phosphoprotein that plays a key role in RNA replication	Headache, diarrhea
Ombitasvir		
Velpatasvir	Exact mechanism unknown	
NS5B inhibitors		
Sofosbuvir	Inhibits NS5B, an RNA-dependent RNA polymerase acting as a chain terminator Prevents viral RNA replication	Fatigue, headache
Dasabuvir		
NS3/4A inhibitors		
Grazoprevir	Inhibits NS3/4A, a viral protease, preventing viral replication	Grazoprevir: headache, fatigue Simeprevir: photosensitivity reactions, rash
Simeprevir		
Alternative drugs		
Ribavirin	Inhibits synthesis of guanine nucleotides by competitively inhibiting IMP dehydrogenase Used as adjunct in cases refractory to newer medications	Hemolytic anemia, severe teratogen

Disinfection and sterilization Goals include the reduction of pathogenic organism counts to safe levels (disinfection) and the inactivation of all microbes including spores (sterilization).

Autoclave	Pressurized steam at > 120°C. Sporicidal. May not reliably inactivate prions.
Alcohols	Denature proteins and disrupt cell membranes. Not sporicidal.
Chlorhexidine	Denatures proteins and disrupts cell membranes. Not sporicidal.
Chlorine	Oxidizes and denatures proteins. Sporicidal.
Ethylene oxide	Alkylating agent. Sporicidal.
Hydrogen peroxide	Free radical oxidation. Sporicidal.
Iodine and iodophors	Halogenation of DNA, RNA, and proteins. May be sporicidal.
Quaternary amines	Impair permeability of cell membranes. Not sporicidal.

Antimicrobials to avoid in pregnancy

ANTIMICROBIAL	ADVERSE EFFECT
S ulfonamides	Kernicterus
A minoglycosides	Ototoxicity
F luoroquinolones	Cartilage damage
C larithromycin	Embryotoxic
T etracyclines	Discolored teeth, inhibition of bone growth
R ibavirin	Teratogenic
G riseofulvin	Teratogenic
C hloramphenicol	Gray baby syndrome
SAFe Children Take Really Good Care.	

HIGH-YIELD PRINCIPLES IN

Pathology

“Digressions, objections, delight in mockery, carefree mistrust are signs of health; everything unconditional belongs in pathology.”

—Friedrich Nietzsche

“You cannot separate passion from pathology any more than you can separate a person’s spirit from his body.”


—Richard Selzer

The fundamental principles of pathology are key to understanding diseases in all organ systems. Major topics such as inflammation and neoplasia appear frequently in questions across different organ systems, and such topics are definitely high yield. For example, the concepts of cell injury and inflammation are key to understanding the inflammatory response that follows myocardial infarction, a very common subject of board questions. Similarly, a familiarity with the early cellular changes that culminate in the development of neoplasias—for example, esophageal or colon cancer—is critical. Finally, make sure you recognize the major tumor-associated genes and are comfortable with key cancer concepts such as tumor staging and metastasis.

▶ Cellular Injury	206
▶ Inflammation	213
▶ Neoplasia	219

▶ PATHOLOGY—CELLULAR INJURY

Cellular adaptations	Reversible changes that can be physiologic (eg, uterine enlargement during pregnancy) or pathologic (eg, myocardial hypertrophy 2° to systemic HTN). If stress is excessive or persistent, adaptations can progress to cell injury (eg, significant LV hypertrophy → injury to myofibrils → HF).
Hypertrophy	↑ structural proteins and organelles → ↑ in size of cells. Example: cardiac hypertrophy.
Hyperplasia	Controlled proliferation of stem cells and differentiated cells → ↑ in number of cells. Excessive stimulation → pathologic hyperplasia (eg, endometrial hyperplasia), which may progress to dysplasia and cancer. Example: benign prostatic hyperplasia.
Atrophy	↓ in tissue mass due to ↓ in size (↑ cytoskeleton degradation via ubiquitin-proteasome pathway and autophagy; ↓ protein synthesis) and/or number of cells (apoptosis). Causes include disuse, denervation, loss of blood supply, loss of hormonal stimulation, poor nutrition.
Metaplasia	Reprogramming of stem cells → replacement of one cell type by another that can adapt to a new stress. Usually due to exposure to an irritant, such as gastric acid (→ Barrett esophagus) or cigarette smoke (→ respiratory ciliated columnar epithelium replaced by stratified squamous epithelium). May progress to dysplasia → malignant transformation with persistent insult (eg, Barrett esophagus → esophageal adenocarcinoma). Metaplasia of connective tissue can also occur (eg, myositis ossificans, the formation of bone within muscle after trauma).
Dysplasia	Disordered, precancerous epithelial cell growth; not considered a true adaptive response. Characterized by loss of uniformity of cell size and shape (pleomorphism); loss of tissue orientation; nuclear changes (eg, ↑ nuclear:cytoplasmic ratio and clumped chromatin). Mild and moderate dysplasias (ie, do not involve entire thickness of epithelium) may regress with alleviation of inciting cause. Severe dysplasia often becomes irreversible and progresses to carcinoma in situ. Usually preceded by persistent metaplasia or pathologic hyperplasia.




Cell injury

Reversible cell injury

- ↓ ATP → ↓ activity of Ca²⁺ and Na⁺/K⁺ pumps → cellular swelling (earliest morphologic manifestation), mitochondrial swelling
- Ribosomal/polysomal detachment → ↓ protein synthesis
- Plasma membrane changes (eg, blebbing)
- Nuclear changes (eg, chromatin clumping)
- Rapid loss of function (eg, myocardial cells are noncontractile after 1-2 minutes of ischemia)
- Myelin figures (aggregation of peroxidized lipids)

Irreversible cell injury

- Breakdown of plasma membrane → cytosolic enzymes (eg, troponin) leak outside of cell, influx of Ca²⁺ → activation of degradative enzymes
- Mitochondrial damage/dysfunction → loss of electron transport chain → ↓ ATP
- Cytoplasmic vacuolization accompanies programmed cell death (apoptosis)
- Rupture of lysosomes → autolysis
- Nuclear degradation: pyknosis (nuclear condensation) → karyorrhexis (nuclear fragmentation caused by endonuclease-mediated cleavage) → karyolysis (nuclear dissolution)
- Amorphous densities/inclusions in mitochondria



Apoptosis

ATP-dependent programmed cell death.

Intrinsic and extrinsic pathways; both pathways activate caspases (cytosolic proteases) → cellular breakdown including cell shrinkage, chromatin condensation, membrane blebbing, and formation of apoptotic bodies, which are then phagocytosed.

Characterized by deeply eosinophilic cytoplasm and basophilic nucleus, pyknosis, and karyorrhexis. Cell membrane typically remains intact without significant inflammation (unlike necrosis). DNA laddering (fragments in multiples of 180 bp) is a sensitive indicator of apoptosis.

Intrinsic (mitochondrial) pathway

Involved in tissue remodeling in embryogenesis. Occurs when a regulating factor is withdrawn from a proliferating cell population (eg, ↓ IL-2 after a completed immunologic reaction → apoptosis of proliferating effector cells). Also occurs after exposure to injurious stimuli (eg, radiation, toxins, hypoxia).

Regulated by Bcl-2 family of proteins. BAX and BAK are proapoptotic, while Bcl-2 and Bcl-xL are antiapoptotic.

BAX and BAK form pores in the mitochondrial membrane → release of cytochrome C from inner mitochondrial membrane into the cytoplasm → activation of caspases.

Bcl-2 keeps the mitochondrial membrane impermeable, thereby preventing cytochrome C release. Bcl-2 overexpression (eg, follicular lymphoma t[14;18]) → ↓ caspase activation → tumorigenesis.

Extrinsic (death receptor) pathway


2 pathways:

- Ligand receptor interactions (FasL binding to Fas [CD95] or TNF- α binding to its receptor)
- Immune cell (cytotoxic T-cell release of perforin and granzyme B)

Fas-FasL interaction is necessary in thymic medullary negative selection.

Fas mutations ↑ numbers of circulating self-reacting lymphocytes due to failure of clonal deletion.


Defective Fas-FasL interactions cause autoimmune lymphoproliferative syndrome.




Necrosis

Exogenous injury → plasma membrane damage → cell undergoes enzymatic degradation and protein denaturation, intracellular components leak → local inflammatory reaction (unlike apoptosis).

TYPE	SEEN IN	DUE TO	HISTOLOGY
Coagulative	Ischemia/infarcts in most tissues (except brain)	Ischemia or infarction; injury denatures enzymes → proteolysis blocked	Preserved cellular architecture (cell outlines seen), but nuclei disappear; ↑ cytoplasmic binding of eosin stain (→ ↑ eosinophilia; red/pink color) A
Liquefactive	Bacterial abscesses, brain infarcts	Neutrophils release lysosomal enzymes that digest the tissue B	Early: cellular debris and macrophages Late: cystic spaces and cavitation (brain) Neutrophils and cell debris seen with bacterial infection
Caseous	TB, systemic fungi (eg, <i>Histoplasma capsulatum</i>), <i>Nocardia</i>	Macrophages wall off the infecting microorganism → granular debris C	Fragmented cells and debris surrounded by lymphocytes and macrophages (granuloma)
Fat	Enzymatic: acute pancreatitis (saponification of peripancreatic fat) Nonenzymatic: traumatic (eg, injury to breast tissue)	Damaged pancreatic cells release lipase, which breaks down triglycerides; liberated fatty acids bind calcium → saponification (chalky-white appearance)	Outlines of dead fat cells without peripheral nuclei; saponification of fat (combined with Ca ²⁺) appears dark blue on H&E stain D
Fibrinoid	Immune vascular reactions (eg, PAN) Nonimmune vascular reactions (eg, hypertensive emergency, preeclampsia)	Immune complex deposition (type III hypersensitivity reaction) and/or plasma protein (eg, fibrin) leakage from damaged vessel	Vessel walls are thick and pink E
Gangrenous	Distal extremity and GI tract, after chronic ischemia	Dry: ischemia F Wet: superinfection	Coagulative Liquefactive superimposed on coagulative



Ischemia

Inadequate blood supply to meet demand. Mechanisms include ↓ arterial perfusion (eg, atherosclerosis), ↓ venous drainage (eg, testicular torsion, Budd-Chiari syndrome), shock. Regions most vulnerable to hypoxia/ischemia and subsequent infarction:


ORGAN	REGION
Brain	ACA/MCA/PCA boundary areas ^{a,b}
Heart	Subendocardium (LV) A
Kidney	Straight segment of proximal tubule (medulla) Thick ascending limb (medulla)
Liver	Area around central vein (zone III)
Colon	Splenic flexure (Griffith point), ^a rectosigmoid junction (Sudeck point) ^a

^aWatershed areas (border zones) receive blood supply from most distal branches of 2 arteries with limited collateral vascularity. These areas are susceptible to ischemia from hypoperfusion.

^bNeurons most vulnerable to hypoxic-ischemic insults include Purkinje cells of the cerebellum and pyramidal cells of the hippocampus and neocortex (zones 3, 5, 6).

Types of infarcts**Red infarct**

Occurs in venous occlusion and tissues with multiple blood supplies (eg, liver, lung **A**, intestine, testes), and with reperfusion (eg, after angioplasty). **Re**perfusion injury is due to damage by free radicals.

**Pale infarct**

Occurs in solid organs with a single (end-arterial) blood supply (eg, heart, kidney **B**).

Free radical injury


Free radicals damage cells via membrane lipid peroxidation, protein modification, DNA breakage. Initiated via radiation exposure (eg, cancer therapy), metabolism of drugs (phase I), redox reactions, nitric oxide (eg, inflammation), transition metals, WBC (eg, neutrophils, macrophages) oxidative burst.

Free radicals can be eliminated by scavenging enzymes (eg, catalase, superoxide dismutase, glutathione peroxidase), spontaneous decay, antioxidants (eg, vitamins A, C, E), and certain metal carrier proteins (eg, transferrin, ceruloplasmin).


Examples:

- Oxygen toxicity: retinopathy of prematurity (abnormal vascularization), bronchopulmonary dysplasia, reperfusion injury after thrombolytic therapy
- Drug/chemical toxicity: acetaminophen overdose (hepatotoxicity), carbon tetrachloride (converted by cytochrome P-450 into CCl₃ free radical → fatty liver [cell injury → ↓ apolipoprotein synthesis → fatty change], centrilobular necrosis)
- Metal storage diseases: hemochromatosis (iron) and Wilson disease (copper)

Types of calcification		
	Calcium deposits appear deeply basophilic (arrow in A) on H&E stain.	
	Dystrophic calcification	Metastatic calcification
Ca²⁺ DEPOSITION	In abnormal (D iseased) tissues	In normal tissues
EXTENT	Tends to be localized (eg, calcific aortic stenosis)	Widespread (ie, diffuse, metastatic)
ASSOCIATED CONDITIONS	TB (lung and pericardium) and other granulomatous infections, liquefactive necrosis of chronic abscesses, fat necrosis, infarcts, thrombi, schistosomiasis, congenital CMV, toxoplasmosis, rubella, psammoma bodies, CREST syndrome, atherosclerotic plaques can become calcified	Predominantly in interstitial tissues of kidney, lung, and gastric mucosa (these tissues lose acid quickly; ↑ pH favors Ca ²⁺ deposition) Nephrocalcinosis of collecting ducts may lead to nephrogenic diabetes insipidus and renal failure
ETIOLOGY	2° to injury or necrosis	2° to hypercalcemia (eg, 1° hyperparathyroidism, sarcoidosis, hypervitaminosis D) or high calcium-phosphate product levels (eg, chronic kidney disease with 2° hyperparathyroidism, long-term dialysis, calciphylaxis, multiple myeloma)
SERUM Ca²⁺ LEVELS	Normal	Usually abnormal



Lipofuscin




A yellow-brown “wear and tear” pigment **A** associated with normal aging. Composed of polymers of lipids and phospholipids complexed with protein. May be derived through lipid peroxidation of polyunsaturated lipids of subcellular membranes. Autopsy of elderly person will reveal deposits in heart, colon, liver, kidney, eye, and other organs.

Amyloidosis

Abnormal aggregation of proteins (or their fragments) into β -pleated linear sheets \rightarrow insoluble fibrils \rightarrow cellular damage and apoptosis. Amyloid deposits visualized by Congo red stain (red/orange on nonpolarized light [arrows in **A**]), (apple-green birefringence on polarized light [arrows in **B**]), and H&E stain (shows deposits in glomerular mesangial areas). Tubular basement membranes are enlarged on light microscopy.

COMMON TYPES	FIBRIL PROTEIN	DESCRIPTION	
Systemic			
Primary amyloidosis	AL (from Ig Light chains)	Seen in Plasma cell disorders (eg, multiple myeloma)	Manifestations include: <ul style="list-style-type: none"> ▪ Cardiac (eg, restrictive cardiomyopathy) ▪ GI (eg, macroglossia, hepatomegaly) ▪ Renal (eg, nephrotic syndrome) ▪ Hematologic (eg, easy bruising, splenomegaly) ▪ Neurologic (eg, neuropathy) ▪ Musculoskeletal (eg, carpal tunnel syndrome)
Secondary amyloidosis	Serum Amyloid A (AA)	Seen in chronic inflammatory conditions, (eg, rheumatoid arthritis, IBD, familial Mediterranean fever, protracted infection)	
Dialysis-related amyloidosis	β_2 -microglobulin	Seen in patients with ESRD and/or on long-term dialysis	
Localized			
Alzheimer disease	β -amyloid protein	Cleaved from amyloid precursor protein (APP)	
Type 2 diabetes mellitus	Islet amyloid polypeptide (IAPP)	Caused by deposition of amylin in pancreatic islets	
Medullary thyroid cancer	Calcitonin		
Isolated atrial amyloidosis	ANP	Common in normal aging \uparrow risk of atrial fibrillation	
Systemic senile (age-related) amyloidosis	Normal (wild-type) transthyretin (TTR)	Seen predominantly in cardiac ventricles	Cardiac dysfunction more insidious than in AL amyloidosis
Hereditary			
Familial amyloid cardiomyopathy	Mutated transthyretin (ATTR)	Ventricular endomyocardium deposition \rightarrow restrictive cardiomyopathy, arrhythmias	5% of African Americans are carriers of mutant allele
Familial amyloid polyneuropathies	Mutated transthyretin (ATTR)	Due to transthyretin gene mutation	



▶ PATHOLOGY—INFLAMMATION

Inflammation Response to eliminate initial cause of cell injury, to remove necrotic cells resulting from the original insult, and to initiate tissue repair. Divided into acute and chronic. The inflammatory response itself can be harmful to the host if the reaction is excessive (eg, septic shock), prolonged (eg, persistent infections such as TB), or inappropriate (eg, autoimmune diseases such as SLE).

Cardinal signs

SIGN	MECHANISM	MEDIATORS
Rubor (redness), calor (warmth)	Vasodilation (relaxation of arteriolar smooth muscle) → ↑ blood flow	Histamine, prostaglandins, bradykinin, NO
Tumor (swelling)	Endothelial contraction/disruption (eg, from tissue damage) → ↑ vascular permeability → leakage of protein-rich fluid from postcapillary venules into interstitial space (exudate) → ↑ interstitial oncotic pressure	Endothelial contraction: leukotrienes (C ₄ , D ₄ , E ₄), histamine, serotonin
Dolor (pain)	Sensitization of sensory nerve endings	Bradykinin, PGE ₂ , histamine
Functio laesa (loss of function)	Cardinal signs above impair function (eg, inability to make fist with hand that has cellulitis)	

Systemic manifestations (acute-phase reaction)

Fever	Pyrogens (eg, LPS) induce macrophages to release IL-1 and TNF → ↑ COX activity in perivascular cells of hypothalamus → ↑ PGE ₂ → ↑ temperature set point	
Leukocytosis	Elevation of WBC count; type of cell that is predominantly elevated depends on the inciting agent or injury (eg, bacteria → ↑ neutrophils)	
↑ plasma acute-phase proteins	Factors whose serum concentrations change significantly in response to inflammation Produced by the liver in both acute and chronic inflammatory states	Notably induced by IL-6

Acute phase reactants More **FFiSH** in the **C** (sea).

POSITIVE (UPREGULATED)

Ferritin	Binds and sequesters iron to inhibit microbial iron scavenging.
Fibrinogen	Coagulation factor; promotes endothelial repair; correlates with ESR.
Serum amyloid A	Prolonged elevation can lead to amyloidosis.
Hepcidin	↓ iron absorption (by degrading ferroportin) and ↓ iron release (from macrophages) → anemia of chronic disease.
C-reactive protein	Opsonin; fixes complement and facilitates phagocytosis. Measured clinically as a nonspecific sign of ongoing inflammation.

NEGATIVE (DOWNREGULATED)

Albumin	Reduction conserves amino acids for positive reactants.
Transferrin	Internalized by macrophages to sequester iron.

Erythrocyte sedimentation rate


RBCs normally remain separated via \ominus charges. Products of inflammation (eg, fibrinogen) coat RBCs \rightarrow \downarrow \ominus charge \rightarrow \uparrow RBC aggregation. Denser RBC aggregates fall at a faster rate within a pipette tube \rightarrow \uparrow ESR. Often co-tested with CRP (more specific marker of inflammation).

 \uparrow ESR

Most anemias
Infections
Inflammation (eg, giant cell [temporal] arteritis, polymyalgia rheumatica)
Cancer (eg, metastases, multiple myeloma)
Renal disease (end-stage or nephrotic syndrome)
Pregnancy

 \downarrow ESR

Sickle cell anemia (altered shape)
Polycythemia (\uparrow RBCs “dilute” aggregation factors)
HF
Microcytosis
Hypofibrinogenemia

Acute inflammation

Transient and early response to injury or infection. Characterized by neutrophils in tissue **A**, often with associated edema. Rapid onset (seconds to minutes) and short duration (minutes to days). Represents a reaction of the innate immune system (ie, less specific response than chronic inflammation).

STIMULI

Infections, trauma, necrosis, foreign bodies.

MEDIATORS

Toll-like receptors, arachidonic acid metabolites, neutrophils, eosinophils, antibodies (pre-existing), mast cells, basophils, complement, Hageman factor (factor XII).

Inflammasome—Cytoplasmic protein complex that recognizes products of dead cells, microbial products, and crystals (eg, uric acid crystals) \rightarrow activation of IL-1 and inflammatory response.

COMPONENTS

- Vascular: vasodilation (\rightarrow \uparrow blood flow and stasis) and \uparrow endothelial permeability
- Cellular: extravasation of leukocytes (mainly neutrophils) from postcapillary venules and accumulation in the focus of injury followed by leukocyte activation

To bring cells and proteins to site of injury or infection.

Leukocyte extravasation has 4 steps: margination and rolling, adhesion, transmigration, and migration (chemoattraction).

OUTCOMES


- Resolution and healing (IL-10, TGF- β)
- Persistent acute inflammation (IL-8)
- Abscess (acute inflammation walled off by fibrosis)
- Chronic inflammation (antigen presentation by macrophages and other APCs \rightarrow activation of CD4⁺ Th cells)
- Scarring

Macrophages predominate in the late stages of acute inflammation (peak 2–3 days after onset) and influence outcome by secreting cytokines.

Leukocyte extravasation

Extravasation predominantly occurs at postcapillary venules.

STEP	VASCULATURE/STROMA	LEUKOCYTE
1 Margination and rolling— defective in leukocyte adhesion deficiency type 2 (↓ Sialyl Lewis ^X)	E-selectin (upregulated by TNF and IL-1) P-selectin (released from Weibel- Palade bodies) GlyCAM-1, CD34	Sialyl Lewis ^X Sialyl Lewis ^X L-selectin
2 Tight binding (adhesion)— defective in leukocyte adhesion deficiency type 1 (↓ CD18 integrin subunit)	ICAM-1 (CD54) VCAM-1 (CD106)	CD11/18 integrins (LFA-1, Mac-1) VLA-4 integrin
3 Diapedesis (transmigration)— WBC travels between endothelial cells and exits blood vessel	PECAM-1 (CD31)	PECAM-1 (CD31)
4 Migration—WBC travels through interstitium to site of injury or infection guided by chemotactic signals	Chemotactic factors: C5a, IL-8, LTB ₄ , kallikrein, platelet-activating factor	Various



Chronic inflammation	Prolonged inflammation characterized by mononuclear infiltration (macrophages, lymphocytes, plasma cells), which leads to simultaneous tissue destruction and repair (including angiogenesis and fibrosis). May be preceded by acute inflammation.
STIMULI	Persistent infections (eg, TB, <i>T pallidum</i> , certain fungi and viruses) → type IV hypersensitivity, autoimmune diseases, prolonged exposure to toxic agents (eg, silica) and foreign material.
MEDIATORS	Macrophages are the dominant cells. Interaction of macrophages and T lymphocytes → chronic inflammation. <ul style="list-style-type: none"> ▪ Th1 cells secrete IFN-γ → macrophage classical activation (proinflammatory) ▪ Th2 cells secrete IL-4 and IL-13 → macrophage alternative activation (repair and anti-inflammatory)
OUTCOMES	Scarring, amyloidosis, and neoplastic transformation (eg, chronic HCV infection → chronic inflammation → hepatocellular carcinoma; <i>Helicobacter pylori</i> infection → chronic gastritis → gastric adenocarcinoma).


Wound healing

Tissue mediators	MEDIATOR	ROLE
	FGF	Stimulates angiogenesis
	TGF- β	Angiogenesis, fibrosis
	VEGF	Stimulates angiogenesis
	PDGF	Secreted by activated platelets and macrophages Induces vascular remodeling and smooth muscle cell migration Stimulates fibroblast growth for collagen synthesis
	Metalloproteinases	Tissue remodeling
	EGF	Stimulates cell growth via tyrosine kinases (eg, EGFR/ <i>ErbB1</i>)
PHASE OF WOUND HEALING	EFFECTOR CELLS	CHARACTERISTICS
Inflammatory (up to 3 days after wound)	Platelets, neutrophils, macrophages	Clot formation, \uparrow vessel permeability and neutrophil migration into tissue; macrophages clear debris 2 days later
Proliferative (day 3–weeks after wound)	Fibroblasts, myofibroblasts, endothelial cells, keratinocytes, macrophages	Deposition of granulation tissue and type III collagen, angiogenesis, epithelial cell proliferation, dissolution of clot, and wound contraction (mediated by myofibroblasts) Delayed second phase of wound healing in vitamin C and copper deficiency
Remodeling (1 week–6+ months after wound)	Fibroblasts	Type III collagen replaced by type I collagen, \uparrow tensile strength of tissue Collagenases (require zinc to function) break down type III collagen Zinc deficiency → delayed wound healing

Granulomatous inflammation

A pattern of chronic inflammation. Can be induced by persistent T-cell response to certain infections (eg, TB), immune-mediated diseases, and foreign bodies. Granulomas “wall off” a resistant stimulus without completely eradicating or degrading it → persistent inflammation → fibrosis, organ damage.

HISTOLOGY



Focus of epithelioid cells (activated macrophages with abundant pink cytoplasm) surrounded by lymphocytes and multinucleated giant cells (formed by fusion of several activated macrophages).

Two types:

- Caseating: associated with Central necrosis. Seen with infectious etiologies (eg, TB, fungal).
- Noncaseating **A**: no central necrosis. Seen with autoimmune diseases (eg, sarcoidosis, Crohn disease).


MECHANISM

- 1 APCs present antigens to CD4+ Th cells and secrete IL-12 → CD4+ Th cells differentiate into Th1 cells
- 2 Th1 secretes IFN-γ → macrophage activation
- 3 Macrophages ↑ cytokine secretion (eg, TNF) → formation of epithelioid macrophages and giant cells.

Anti-TNF therapy can cause sequestering granulomas to break down → disseminated disease.

Always test for latent TB before starting anti-TNF therapy.

Associated with hypercalcemia due to ↑ 1α-hydroxylase activity in activated macrophages, resulting in ↑ vitamin D activity.



ETIOLOGIES

INFECTIOUS

Bacterial: *Mycobacteria* (tuberculosis, leprosy), *Bartonella henselae* (cat scratch disease; stellate necrotizing granulomas), *Listeria monocytogenes* (granulomatosis infantiseptica), *Treponema pallidum* (3° syphilis)
 Fungal: endemic mycoses (eg, histoplasmosis)
 Parasitic: schistosomiasis

NONINFECTIOUS

Immune-mediated: sarcoidosis, Crohn disease, 1° biliary cholangitis, subacute (de Quervain/ granulomatous) thyroiditis
 Vasculitis: granulomatosis with polyangiitis (Wegener), eosinophilic granulomatosis with polyangiitis (Churg-Strauss), giant cell (temporal) arteritis, Takayasu arteritis
 Foreign bodies: berylliosis, talcosis, hypersensitivity pneumonitis
 Chronic granulomatous disease

Scar formation

Occurs when repair cannot be accomplished by cell regeneration alone. Nonregenerated cells (2° to severe acute or chronic injury) are replaced by connective tissue. 70–80% of tensile strength regained at 3 months; little tensile strength regained thereafter. Associated with excess TGF-β.


SCAR TYPE	Hypertrophic A	Keloid B
COLLAGEN SYNTHESIS	↑ (type III collagen)	↑↑↑ (types I and III collagen)
COLLAGEN ORGANIZATION	Parallel	Disorganized
EXTENT OF SCAR	Confined to borders of original wound	Extends beyond borders of original wound with “claw-like” projections typically on earlobes, face, upper extremities
RECURRENCE	Infrequent	Frequent
PREDISPOSITION	None	↑ incidence in ethnic groups with darker skin



▶ PATHOLOGY—NEOPLASIA

Neoplasia and neoplastic progression

Uncontrolled, monoclonal proliferation of cells. Can be benign or malignant. Any neoplastic growth has two components: parenchyma (neoplastic cells) and supporting stroma (non-neoplastic; eg, blood vessels, connective tissue).



Normal cells

① Normal cells with basal → apical polarity. See cervical example **A**, which shows normal cells and spectrum of dysplasia, as discussed below.

Dysplasia

② Loss of uniformity in cell size and shape (pleomorphism); loss of tissue orientation; nuclear changes (eg, ↑ nuclear:cytoplasmic ratio) **A**.

Carcinoma in situ/ preinvasive


③ Irreversible severe dysplasia that involves the entire thickness of epithelium but does not penetrate the intact basement membrane **A**.

Invasive carcinoma

④ Cells have invaded basement membrane using collagenases and hydrolases (metalloproteinases). Cell-cell contacts lost by inactivation of E-cadherin.

Metastasis

⑤ Spread to distant organ(s) via lymphatics or blood.



Normal

Mild dysplasia

Moderate dysplasia

Severe dysplasia/
carcinoma in situ

Tumor nomenclature

Carcinoma implies epithelial origin, whereas **sarcoma** denotes mesenchymal origin. Both terms generally imply malignancy.

Benign tumors are usually well-differentiated and well-demarcated, with low mitotic activity, no metastases, and no necrosis.

Malignant tumors (cancers) may show poor differentiation, erratic growth, local invasion, metastasis, and ↓ apoptosis.

Terms for non-neoplastic malformations include hamartoma (disorganized overgrowth of tissues in their native location, eg, Peutz-Jeghers polyps) and choristoma (normal tissue in a foreign location, eg, gastric tissue located in distal ileum in Meckel diverticulum).

CELL TYPE	BENIGN	MALIGNANT
Epithelium	Adenoma, papilloma	Adenocarcinoma, papillary carcinoma
Mesenchyme		
Blood cells		Leukemia, lymphoma
Blood vessels	Hemangioma	Angiosarcoma
Smooth muscle	Leiomyoma	Leiomyosarcoma
Striated muscle	Rhabdomyoma	Rhabdomyosarcoma
Connective tissue	Fibroma	Fibrosarcoma
Bone	Osteoma	Osteosarcoma
Fat	Lipoma	Liposarcoma
Melanocyte	Nevus/mole	Melanoma


Tumor grade vs stage

Differentiation—degree to which a tumor resembles its tissue of origin. Well-differentiated tumors (often less aggressive) closely resemble their tissue of origin, whereas poorly differentiated tumors (often more aggressive) do not.

Anaplasia—complete lack of differentiation of cells in a malignant neoplasm.

Grade


Degree of cellular differentiation and mitotic activity on histology. Ranges from low grade (well-differentiated) to high grade (poorly differentiated, undifferentiated, or anaplastic).

**Stage**

Degree of localization/spread based on site and size of 1° lesion, spread to regional lymph nodes, presence of metastases. Based on clinical (c) or pathologic (p) findings. Stage generally has more prognostic value than grade (eg, a high-stage yet low-grade tumor is usually worse than a low-stage yet high-grade tumor). **Stage** determines **Survival**.

TNM staging system (**Stage = Spread**):

T = **T**umor size/invasiveness, **N** = **N**ode involvement, **M** = **M**etastases, eg, cT3N1M0. Each TNM factor has independent prognostic value; N and M are often most important.




HALLMARK	MECHANISM
Hallmarks of cancer	Cancer is caused by (mostly acquired) DNA mutations that affect fundamental cellular processes (eg, growth, DNA repair, survival).
Growth signal self-sufficiency	<p>Mutations in genes encoding:</p> <ul style="list-style-type: none"> ▪ Proto-oncogenes → ↑ growth factors → autocrine loop (eg, ↑ PDGF in brain tumors) ▪ Growth factor receptors → constitutive signalling (eg, <i>HER2/neu</i> in breast cancer) ▪ Signaling molecules (eg, <i>RAS</i>) ▪ Transcription factors (eg, <i>MYC</i>) ▪ Cell cycle regulators (eg, cyclins, CDKs)
Anti-growth signal insensitivity	<ul style="list-style-type: none"> ▪ Mutations in tumor suppressor genes (eg, <i>Rb</i>) ▪ Loss of E-cadherin function → loss of contact inhibition (eg, <i>NF2</i> mutations)
Evasion of apoptosis	Mutations in genes that regulate apoptosis (eg, <i>TP53</i> , <i>BCL2</i> → follicular B cell lymphoma).
Limitless replicative potential	Reactivation of telomerase → maintenance and lengthening of telomeres → prevention of chromosome shortening and cell aging.
Sustained angiogenesis	↑ pro-angiogenic factors (eg, VEGF) or ↓ inhibitory factors. Factors may be produced by tumor or stromal cells. Vessels can sprout from existing capillaries (neoangiogenesis) or endothelial cells are recruited from bone marrow (vasculogenesis). Vessels may be leaky and/or dilated.
Tissue invasion	Loss of E-cadherin function → loosening of intercellular junctions → metalloproteinases degrade basement membrane and ECM → cells attach to ECM proteins (eg, laminin, fibronectin) → cells migrate through degraded ECM (“locomotion”) → vascular dissemination.
Metastasis	Tumor cells or emboli spread via lymphatics or blood → adhesion to endothelium → extravasation and homing. Site of metastasis can be predicted by site of 1° tumor, as the target organ is often the first-encountered capillary bed. Some cancers show organ tropism (eg, lung cancers commonly metastasize to adrenals).
Warburg effect	Shift of glucose metabolism away from mitochondrial oxidative phosphorylation toward glycolysis.
Immune evasion in cancer	<p>Normally, immune cells can recognize and attack tumor cells. For successful tumorigenesis, tumor cells must evade the immune system. Multiple escape mechanisms exist:</p> <ul style="list-style-type: none"> ▪ ↓ MHC class I expression by tumor cells → cytotoxic T cells are unable to recognize tumor cells. ▪ Tumor cells secrete immunosuppressive factors (eg, TGF-β) and recruit regulatory T cells to down regulate immune response. ▪ Tumor cells up regulate immune checkpoint molecules, which inhibit immune response.

Immune checkpoint interactions

Signals that modulate T cell activation and function → ↓ immune response against tumor cells. Targeted by several cancer immunotherapies. Examples:

- Interaction between PD-1 (on T cells) and PD-L1/2 (on tumor cells or immune cells in tumor microenvironment) → T cell dysfunction (exhaustion). Inhibited by antibodies against PD-1 (eg, pembrolizumab, nivolumab) or PD-L1 (eg, atezolizumab, durvalumab, avelumab).
- CTLA-4 on T cells outcompetes CD28 for B7 on APCs → loss of T cell costimulatory signal. Inhibited by ipilimumab (anti-CTLA-4 antibody).



Cancer epidemiology


Skin cancer (basal > squamous >> melanoma) is the most common cancer (not included below).

	MEN	WOMEN	CHILDREN (AGE 0-14)	NOTES
Cancer incidence	1. Prostate 2. Lung 3. Colon/rectum	1. Breast 2. Lung 3. Colon/rectum	1. Leukemia 2. CNS 3. Neuroblastoma	Lung cancer incidence has ↓ in men, but has not changed significantly in women.
Cancer mortality	1. Lung 2. Prostate 3. Colon/rectum	1. Lung 2. Breast 3. Colon/rectum	1. Leukemia 2. CNS 3. Neuroblastoma	Cancer is the 2nd leading cause of death in the United States (heart disease is 1st).

Common metastases

Most sarcomas spread hematogenously; most carcinomas spread via lymphatics. However, **Four Carcinomas Route Hematogenously: Follicular thyroid carcinoma, Choriocarcinoma, Renal cell carcinoma, and Hepatocellular carcinoma.**

SITE OF METASTASIS	1° TUMOR	NOTES
Brain	Lung > breast > melanoma, colon, kidney	50% of brain tumors are from metastases A B Commonly seen as multiple well-circumscribed tumors at gray/white matter junction
Liver	Colon >> Stomach > Pancreas (Cancer Sometimes Penetrates liver)	Liver C D and lung are the most common sites of metastasis after the regional lymph nodes
Bone	Prostate, Breast > Kidney, Thyroid, Lung (Painful Bones Kill The Lungs)	Bone metastasis E F >> 1° bone tumors (eg, multiple myeloma) Predilection for axial skeleton G Bone metastasis can be: <ul style="list-style-type: none"> ▪ Lytic (eg, thyroid, kidney, non-small cell lung cancer) ▪ Blastic (eg, prostate, small cell lung cancer) ▪ Mixed (eg, breast cancer)



Oncogenes

Gain of function mutation converts proto-oncogene (normal gene) to oncogene → ↑ cancer risk.
Requires damage to only **one** allele of a proto-**oncogene**.

GENE	GENE PRODUCT	ASSOCIATED NEOPLASM
ALK	Receptor tyrosine K inase	Lung Adenocarcinoma (A denocarcinoma of the Lung K inase)
BCR-ABL	Non-receptor tyrosine kinase	CML, ALL
BCL-2	Antiapoptotic molecule (inhibits apoptosis)	Follicular and diffuse large B Cell Lymphomas
BRAF	Serine/threonine kinase	Melanoma, non-Hodgkin lymphoma, papillary thyroid carcinoma, hairy cell leukemia
c-KIT	CytoK ine receptor	Gastrointestinal stromal tumor (GIST)
c-MYC	Transcription factor	Burkitt lymphoma
HER2/neu (c-erbB2)	Receptor tyrosine kinase	Breast and gastric carcinomas
JAK2	Tyrosine kinase	Chronic myeloproliferative disorders
KRAS	GTPase	Colon cancer, lung cancer, pancreatic cancer
MYCL1	Transcription factor	Lung tumor
N-myc (MYCN)	Transcription factor	Neuroblastoma
RET	Receptor tyrosine kinase	MEN 2A and 2B, papillary thyroid carcinoma, pheochromocytoma

Tumor suppressor genes

Loss of function → ↑ cancer risk; both (**two**) alleles of a **tumor** suppressor gene must be lost for expression of disease.

GENE	GENE PRODUCT	ASSOCIATED CONDITION
APC	Negative regulator of β -catenin/WNT pathway	Colorectal cancer (associated with FAP)
BRCA1/BRCA2	BRCA1/BRCA2 proteins	B reast, ovarian, and pancreatic c ancers
CDKN2A	p16, blocks $G_1 \rightarrow S$ phase	Melanoma, pancreatic cancer
DCC	DCC —Deleted in C olon C ancer	Colon cancer
SMAD4 (DPC4)	DPC —Deleted in P ancreatic C ancer	Pancreatic cancer
MEN1	Menin	M ultiple E ndocrine N eoplasia type 1
NF1	Neurofibromin (Ras GTPase activating protein)	N eurofibromatosis type 1
NF2	Merlin (schwannomin) protein	N eurofibromatosis type 2
PTEN	Negative regulator of PI3k/AKT pathway	P rostate, b reas T , and E Ndometrial cancers
Rb	Inhibits E2F; blocks $G_1 \rightarrow S$ phase	R etinoblastoma, osteosarcoma (b one cancer)
TP53	p53, activates p21, blocks $G_1 \rightarrow S$ phase	Most human cancers, Li-Fraumeni syndrome (multiple malignancies at early age, aka, SBLA cancer syndrome: S arcoma, B reast, L eukemia, A drenal gland)
TSC1	Hamartin protein	T uberous s clerosis
TSC2	Tuberin protein	T uberous s clerosis
VHL	Inhibits hypoxia-inducible factor 1 α	v on H ippel- L indau disease
WT1	Urogenital development transcription factor	W ilms t umor (nephroblastoma)

Carcinogens

TOXIN	EXPOSURE	ORGAN	IMPACT
Aflatoxins (<i>Aspergillus</i>)	Stored grains and nuts	Liver	Hepatocellular carcinoma
Alkylating agents	Oncologic chemotherapy	Blood	Leukemia/lymphoma
Aromatic amines (eg, benzidine, 2-naphthylamine)	Textile industry (dyes), cigarette smoke (2-naphthylamine)	Bladder	Transitional cell carcinoma
Arsenic	Herbicides (vineyard workers), metal smelting	Liver Lung Skin	Angiosarcoma Lung cancer Squamous cell carcinoma
Asbestos	Old roofing material, shipyard workers	Lung	Bronchogenic carcinoma > mesothelioma
Cigarette smoke		Bladder Cervix Esophagus Kidney Larynx Lung Oropharynx Pancreas	Transitional cell carcinoma Squamous cell carcinoma Squamous cell carcinoma/ adenocarcinoma Renal cell carcinoma Squamous cell carcinoma Squamous cell and small cell carcinoma Oropharyngeal cancer Pancreatic adenocarcinoma
Ethanol		Esophagus Liver	Squamous cell carcinoma Hepatocellular carcinoma
Ionizing radiation		Thyroid	Papillary thyroid carcinoma, leukemias
Nickel, chromium, beryllium, silica	Occupational exposure	Lung	Lung cancer
Nitrosamines	Smoked foods	Stomach	Gastric cancer
Radon	Byproduct of uranium decay, accumulates in basements	Lung	Lung cancer (2nd leading cause after cigarette smoke)
Vinyl chloride	Used to make PVC pipes (plumbers)	Liver	Angiosarcoma

Oncogenic microbes

Microbe	Associated cancer
EBV	Burkitt lymphoma, Hodgkin lymphoma, nasopharyngeal carcinoma, 1° CNS lymphoma (in immunocompromised patients)
HBV, HCV	Hepatocellular carcinoma
HHV-8	Kaposi sarcoma
HPV	Cervical and penile/anal carcinoma (types 16, 18), head and neck cancer
<i>H pylori</i>	Gastric adenocarcinoma and MALT lymphoma
HTLV-1	Adult T-cell Leukemia/Lymphoma
Liver fluke (<i>Clonorchis sinensis</i>)	Cholangiocarcinoma
<i>Schistosoma haematobium</i>	Squamous cell bladder cancer

Serum tumor markers Tumor markers should not be used as the 1° tool for cancer diagnosis or screening. They may be used to monitor tumor recurrence and response to therapy, but definitive diagnosis is made via biopsy. Some can be associated with non-neoplastic conditions.

MARKER	IMPORTANT ASSOCIATIONS	NOTES
Alkaline phosphatase	Metastases to bone or liver, Paget disease of bone, seminoma (placental ALP).	Exclude hepatic origin by checking LFTs and GGT levels.
α-fetoprotein	Hepatocellular carcinoma, Endodermal sinus (yolk sac) tumor, Mixed germ cell tumor, Ataxia-telangiectasia, Neural tube defects. (HE-MAN is the alpha male!)	Normally made by fetus. Transiently elevated in pregnancy. High levels associated with neural tube and abdominal wall defects, low levels associated with Down syndrome.
hCG	Hydatidiform moles and Choriocarcinomas (Gestational trophoblastic disease), testicular cancer, mixed germ cell tumor.	Produced by syncytiotrophoblasts of the placenta.
CA 15-3/CA 27-29	Breast cancer.	
CA 19-9	Pancreatic adenocarcinoma.	
CA 125	Ovarian cancer.	
Calcitonin	Medullary thyroid carcinoma (alone and in MEN2A, MEN2B).	
CEA	Colorectal and pancreatic cancers. Minor associations: gastric, breast, and medullary thyroid carcinomas.	Carcinoembryonic antigen. Very nonspecific.
Chromogranin	Neuroendocrine tumors.	
LDH	Testicular germ cell tumors, ovarian dysgerminoma, other cancers.	Can be used as an indicator of tumor burden.
Neuron-specific enolase	Neuroendocrine tumors (eg, small cell lung cancer, carcinoid tumor, neuroblastoma)	
PSA	Prostate cancer.	Prostate-specific antigen. Also elevated in BPH and prostatitis. Questionable risk/benefit for screening. Marker for recurrence after treatment.

Important immunohistochemical stains


Determine primary site of origin for metastatic tumors and characterize tumors that are difficult to classify. Can have prognostic and predictive value.

STAIN	TARGET	TUMORS IDENTIFIED
Chromogranin and synaptophysin	Neuroendocrine cells	Small cell carcinoma of the lung, carcinoid tumor
Cytokeratin	Epithelial cells	Epithelial tumors (eg, squamous cell carcinoma)
DesMin	Muscle	Muscle tumors (eg, rhabdomyosarcoma)
GFAP	NeuroGlia (eg, astrocytes, Schwann cells, oligodendrocytes)	Astrocytoma, Glioblastoma
Neurofilament	Neurons	Neuronal tumors (eg, neuroblastoma)
PSA	Prostatic epithelium	Prostate cancer
S-100	Neural crest cells	Melanoma, schwannoma, Langerhans cell histiocytosis
TRAP	Tartrate-resistant acid phosphatase	Hairy cell leukemia
Vimentin	Mesenchymal tissue (eg, fibroblasts, endothelial cells, macrophages)	Mesenchymal tumors (eg, sarcoma), but also many other tumors (eg, endometrial carcinoma, renal cell carcinoma, meningioma)

P-glycoprotein

Also known as multidrug resistance protein 1 (MDR1). Classically seen in adrenocortical carcinoma but also expressed by other cancer cells (eg, colon, liver). Used to pump out toxins, including chemotherapeutic agents (one mechanism of ↓ responsiveness or resistance to chemotherapy over time).

Psammoma bodies



Laminated, concentric spherules with dystrophic calcification **A**, **PSaMMOMa** bodies are seen in:

- Papillary carcinoma of thyroid
- Somatostatinoma
- Meningioma
- Malignant Mesothelioma
- Ovarian serous papillary cystadenocarcinoma
- Prolactinoma (**Milk**)

Cachexia

Weight loss, muscle atrophy, and fatigue that occur in chronic disease (eg, cancer, AIDS, heart failure, COPD). Mediated by TNF-α, IFN-γ, IL-1, and IL-6.

Paraneoplastic syndromes

MANIFESTATION	DESCRIPTION/MECHANISM	MOST COMMONLY ASSOCIATED TUMOR(S)
Musculoskeletal and cutaneous		
Dermatomyositis	Progressive proximal muscle weakness, Gottron papules, heliotrope rash	Adenocarcinomas, especially ovarian
Acanthosis nigricans	Hyperpigmented velvety plaques in axilla and neck	Gastric adenocarcinoma and other visceral malignancies
Sign of Leser-Trélat	Sudden onset of multiple seborrheic keratoses	GI adenocarcinomas and other visceral malignancies
Hypertrophic osteoarthropathy	Abnormal proliferation of skin and bone at distal extremities → clubbing, arthralgia, joint effusions, periostosis of tubular bones	Adenocarcinoma of the lung
Endocrine		
Hypercalcemia	PTHrP ↑ 1,25-(OH) ₂ vitamin D ₃ (calcitriol)	Squamous cell carcinomas of lung, head, and neck; renal, bladder, breast, and ovarian carcinomas Lymphoma
Cushing syndrome	↑ ACTH	Small cell lung cancer
Hyponatremia (SIADH)	↑ ADH	
Hematologic		
Polycythemia	↑ Erythropoietin Paraneoplastic rise to high hematocrit levels	Pheochromocytoma, renal cell carcinoma, HCC, hemangioblastoma, leiomyoma
Pure red cell aplasia	Anemia with low reticulocytes	Thymoma
Good syndrome	Hypogammaglobulinemia	
Trousseau syndrome	Migratory superficial thrombophlebitis	
Nonbacterial thrombotic (marantic) endocarditis	Deposition of sterile platelet thrombi on heart valves	Adenocarcinomas, especially pancreatic
Neuromuscular		
Anti-NMDA receptor encephalitis	Psychiatric disturbance, memory deficits, seizures, dyskinesias, autonomic instability, language dysfunction	Ovarian teratoma
Opsoclonus-myoclonus ataxia syndrome	“Dancing eyes, dancing feet”	Neuroblastoma (children), small cell lung cancer (adults)
Paraneoplastic cerebellar degeneration	Antibodies against antigens in Purkinje cells	Small cell lung cancer (anti-Hu), gynecologic and breast cancers (anti-Yo), and Hodgkin lymphoma (anti-Tr)
Paraneoplastic encephalomyelitis	Antibodies against Hu antigens in neurons	Small cell lung cancer
Lambert-Eaton myasthenic syndrome	Antibodies against presynaptic (P/Q-type) Ca ²⁺ channels at NMJ	
Myasthenia gravis	Antibodies against postsynaptic ACh receptors at NMJ	Thymoma

HIGH-YIELD PRINCIPLES IN

Pharmacology

“One pill makes you larger, and one pill makes you small.”

—Grace Slick

“I was under medication when I made the decision not to burn the tapes.”

—Richard Nixon

“I wondher why ye can always read a doctor’s bill an’ ye niver can read his purscription.”

—Finley Peter Dunne

“One of the first duties of the physician is to educate the masses not to take medicine.”

—William Osler

Preparation for pharmacology questions is straightforward. Know all the mechanisms, clinical use, and important adverse effects of key drugs and their major variants. Obscure derivatives are low-yield. Learn their classic and distinguishing toxicities as well as major drug-drug interactions. Reviewing associated biochemistry, physiology, and microbiology concepts can be useful while studying pharmacology. The exam has a strong emphasis on ANS, CNS, antimicrobial, and cardiovascular agents as well as on NSAIDs, which are covered throughout the text. Specific drug dosages or trade names are generally not testable. The exam may use graphs to test various pharmacology content, so make sure you are comfortable interpreting them.

▶ Pharmacokinetics and Pharmacodynamics	230
▶ Autonomic Drugs	236
▶ Toxicities and Side Effects	248
▶ Miscellaneous	253


▶ PHARMACOLOGY—PHARMACOKINETICS AND PHARMACODYNAMICS

Enzyme kinetics


Michaelis-Menten kinetics

K_m is inversely related to the affinity of the enzyme for its substrate.
 V_{max} is directly proportional to the enzyme concentration.
 Most enzymatic reactions follow a hyperbolic curve (ie, Michaelis-Menten kinetics); however, enzymatic reactions that exhibit a sigmoid curve usually indicate cooperative kinetics (eg, hemoglobin).

[S] = concentration of substrate; V = velocity.



Effects of enzyme inhibition




Lineweaver-Burk plot


The closer to 0 on the Y-axis, the higher the V_{max} .
 The closer to 0 on the X-axis, the higher the K_m .
 The higher the K_m , the lower the affinity.

Competitive inhibitors cross each other, whereas noncompetitive inhibitors do not.

Competitive inhibitors increase K_m .



Effects of enzyme inhibition




	Competitive inhibitors, reversible	Competitive inhibitors, irreversible	Noncompetitive inhibitors
Resemble substrate	Yes	Yes	No
Overcome by ↑ [S]	Yes	No	No
Bind active site	Yes	Yes	No
Effect on V_{max}	Unchanged	↓	↓
Effect on K_m	↑	Unchanged	Unchanged
Pharmacodynamics	↓ potency	↓ efficacy	↓ efficacy

Pharmacokinetics**Bioavailability (F)**

Fraction of administered drug reaching systemic circulation unchanged. For an IV dose, $F = 100\%$.

Orally: F typically $< 100\%$ due to incomplete absorption and first-pass metabolism. Can be calculated from the area under the curve in a plot of plasma concentration over time.

**Volume of distribution (V_d)**

Theoretical volume occupied by the total amount of drug in the body relative to its plasma concentration. Apparent V_d of plasma protein-bound drugs can be altered by liver and kidney disease (\downarrow protein binding, $\uparrow V_d$). Drugs may distribute in more than one compartment.

$$V_d = \frac{\text{amount of drug in the body}}{\text{plasma drug concentration}}$$

V_d	COMPARTMENT	DRUG TYPES
Low	Intravascular	Large/charged molecules; plasma protein bound
Medium	ECF	Small hydrophilic molecules
High	All tissues including fat	Small lipophilic molecules, especially if bound to tissue protein

Clearance (CL)

The volume of plasma cleared of drug per unit time. Clearance may be impaired with defects in cardiac, hepatic, or renal function.

$$CL = \frac{\text{rate of elimination of drug}}{\text{plasma drug concentration}} = V_d \times K_e \text{ (elimination constant)}$$

Half-life ($t_{1/2}$)

The time required to change the amount of drug in the body by $\frac{1}{2}$ during elimination.

In first-order kinetics, a drug infused at a constant rate takes 4–5 half-lives to reach steady state. It takes 3.3 half-lives to reach 90% of the steady-state level.

$$t_{1/2} = \frac{0.7 \times V_d}{CL} \text{ in first-order elimination}$$

# of half-lives	1	2	3	4
% remaining	50%	25%	12.5%	6.25%

Dosage calculations

$$\text{Loading dose} = \frac{C_p \times V_d}{F}$$

$$\text{Maintenance dose} = \frac{C_p \times CL \times \tau}{F}$$


C_p = target plasma concentration at steady state
 τ = dosage interval (time between doses), if not administered continuously

In renal or liver disease, maintenance dose \downarrow and loading dose is usually unchanged.

Time to steady state depends primarily on $t_{1/2}$ and is independent of dose and dosing frequency.

Drug metabolism

Geriatric patients lose phase I first. Patients who are slow acetylators have ↑ side effects from certain drugs because of ↓ rate of metabolism (eg, isoniazid).



Elimination of drugs

Zero-order elimination



Rate of elimination is constant regardless of C_p (ie, constant **amount** of drug eliminated per unit time). C_p ↓ linearly with time. Examples of drugs—**P**henytoin, **E**thanol, and **A**spirin (at high or toxic concentrations).

Capacity-limited elimination. **PEA** (a pea is round, shaped like the “0” in **zero-order**).

First-order elimination

Rate of **F**irst-order elimination is directly proportional to the drug concentration (ie, constant **F**raction of drug eliminated per unit time). C_p ↓ exponentially with time. Applies to most drugs.

Flow-dependent elimination.




Urine pH and drug elimination


Ionized species are trapped in urine and cleared quickly. Neutral forms can be reabsorbed.

Weak acids

Examples: phenobarbital, methotrexate, aspirin (salicylates). Trapped in basic environments. Treat overdose with sodium bicarbonate to alkalinize urine.

**Weak bases**

Examples: TCAs, amphetamines. Trapped in acidic environments. Treat overdose with ammonium chloride to acidify urine.




TCA toxicity is generally treated with sodium bicarbonate to overcome the sodium channel-blocking activity of TCAs, but not for accelerating drug elimination.

pKa


pH at which drugs (weak acid or base) are 50% ionized and 50% nonionized. The pKa represents the strength of the weak acid or base.

Efficacy vs potency**Efficacy**


Maximal effect a drug can produce. Represented by the y-value (V_{\max}). ↑ y-value = ↑ V_{\max} = ↑ efficacy. Unrelated to potency (ie, efficacious drugs can have high or low potency). Partial agonists have less efficacy than full agonists.

**Potency**

Amount of drug needed for a given effect. Represented by the x-value (EC_{50}). Left shifting = ↓ EC_{50} = ↑ potency = ↓ drug needed. Unrelated to efficacy (ie, potent drugs can have high or low efficacy).



Receptor binding



AGONIST WITH	POTENCY	EFFICACY	REMARKS	EXAMPLE
A Competitive antagonist	↓	No change	Can be overcome by ↑ agonist concentration	Diazepam (agonist) + flumazenil (competitive antagonist) on GABA _A receptor.
B Noncompetitive antagonist	No change	↓	Cannot be overcome by ↑ agonist concentration	Norepinephrine (agonist) + phenoxybenzamine (noncompetitive antagonist) on α-receptors.
C Partial agonist (alone)	Independent	↓	Acts at same site as full agonist	Morphine (full agonist) vs buprenorphine (partial agonist) at opioid μ-receptors.

Therapeutic index

Measurement of drug safety.


$$\frac{TD_{50}}{ED_{50}} = \frac{\text{median toxic dose}}{\text{median effective dose}}$$

Therapeutic window—dosage range that can safely and effectively treat disease.

TITE: Therapeutic Index = TD_{50} / ED_{50} .

Safer drugs have higher TI values. Drugs with lower TI values frequently require monitoring (eg, Warfarin, Theophylline, Digoxin, Antiepileptic drugs, Lithium; Warning! These Drugs Are Lethal!).

LD₅₀ (lethal median dose) often replaces TD₅₀ in animal studies.




Types of drug interactions

TERM	DEFINITION	EXAMPLE
Additive	Effect of substances A and B together is equal to the sum of their individual effects	Aspirin and acetaminophen “2 + 2 = 4”
Permissive	Presence of substance A is required for the full effects of substance B	Cortisol on catecholamine responsiveness
Synergistic	Effect of substances A and B together is greater than the sum of their individual effects	Clopidogrel with aspirin “2 + 2 > 4”
Potentiation	Similar to synergism, but drug B with no therapeutic action enhances the therapeutic action of drug A	Carbidopa only blocks enzyme to prevent peripheral conversion of levodopa “2 + 0 > 2”
Antagonistic	Effect of substances A and B together is less than the sum of their individual effects	Ethanol antidote for methanol toxicity “2 + 2 < 4”
Tachyphylactic	Acute decrease in response to a drug after initial/repeated administration	Nitrates, niacin, phenylephrine, LSD, MDMA

▶ PHARMACOLOGY—AUTONOMIC DRUGS

Autonomic receptors



Pelvic splanchnic nerves and CNs III, VII, IX and X are part of the parasympathetic nervous system.

Adrenal medulla is directly innervated by preganglionic sympathetic fibers.

Sweat glands are part of the **sympathetic** pathway but are innervated by **cholinergic** fibers (**sympathetic** nervous system results in a “**chold**” sweat).

Acetylcholine receptors


Nicotinic ACh receptors are ligand-gated Na^+/K^+ channels. Two subtypes: N_N (found in autonomic ganglia, adrenal medulla) and N_M (found in neuromuscular junction of skeletal muscle).

Muscarinic ACh receptors are G-protein-coupled receptors that usually act through 2nd messengers. 5 subtypes: M_{1-5} found in heart, smooth muscle, brain, exocrine glands, and on sweat glands (cholinergic sympathetic).

Micturition control

Micturition center in pons regulates involuntary bladder function via coordination of sympathetic and parasympathetic nervous systems.

- ⊕ sympathetic → ↑ urinary retention
- ⊕ parasympathetic → ↑ urine voiding. Some autonomic drugs act on smooth muscle receptors to treat bladder dysfunction.




DRUGS	MECHANISM	USE
1 Muscarinic antagonists (eg, oxybutynin)	⊖ M ₃ receptor → relaxation of detrusor smooth muscle → ↓ detrusor overactivity	Urgency incontinence
1 Muscarinic agonists (eg, bethanechol)	⊕ M ₃ receptor → contraction of detrusor smooth muscle → ↑ bladder emptying	Urinary retention
2 Sympathomimetics (eg, mirabegron)	⊕ β ₃ receptor → relaxation of detrusor smooth muscle → ↑ bladder capacity	Urgency incontinence
3 α₁-blockers (eg, tamsulosin)	⊖ α ₁ -receptor → relaxation of smooth muscle (bladder neck, prostate) → ↓ urinary obstruction	BPH

G-protein-linked second messengers

RECEPTOR	G-PROTEIN CLASS	MAJOR FUNCTIONS
Adrenergic		
α_1	q	↑ vascular smooth muscle contraction, ↑ pupillary dilator muscle contraction (mydriasis), ↑ intestinal and bladder sphincter muscle contraction
α_2	i	↓ sympathetic (adrenergic) outflow, ↓ insulin release, ↓ lipolysis, ↑ platelet aggregation, ↓ aqueous humor production
β_1	s	↑ heart rate, ↑ contractility (one heart), ↑ renin release, ↑ lipolysis
β_2	s	Vasodilation, bronchodilation (two lungs), ↑ lipolysis, ↑ insulin release, ↑ glycogenolysis, ↓ uterine tone (tocolysis), ↑ aqueous humor production, ↑ cellular K^+ uptake
β_3	s	↑ lipolysis, ↑ thermogenesis in skeletal muscle, ↑ bladder relaxation
Cholinergic		
M_1	q	Mediates higher cognitive functions, stimulates enteric nervous system
M_2	i	↓ heart rate and contractility of atria
M_3	q	↑ exocrine gland secretions (eg, lacrimal, sweat, salivary, gastric acid), ↑ gut peristalsis, ↑ bladder contraction, bronchoconstriction, ↑ pupillary sphincter muscle contraction (miosis), ciliary muscle contraction (accommodation), ↑ insulin release, endothelium-mediated vasodilation
Dopamine		
D_1	s	Relaxes renal vascular smooth muscle, activates direct pathway of striatum
D_2	i	Modulates transmitter release, especially in brain, inhibits indirect pathway of striatum
Histamine		
H_1	q	↑ nasal and bronchial mucus production, ↑ vascular permeability, bronchoconstriction, pruritus, pain
H_2	s	↑ gastric acid secretion
Vasopressin		
V_1	q	↑ vascular smooth muscle contraction
V_2	s	↑ H_2O permeability and reabsorption via upregulating aquaporin-2 in collecting two bules (tubules) of kidney, ↑ release of vWF


“After **q**isses (kisses), you get a **q**iq (kick) out of **s**iq (sick) **s**qs (super kinky sex).”



Autonomic drugs

Release of norepinephrine from a sympathetic nerve ending is modulated by NE itself, acting on presynaptic α_2 -autoreceptors → negative feedback.

Amphetamines use the NE transporter (NET) to enter the presynaptic terminal, where they utilize the vesicular monoamine transporter (VMAT) to enter neurosecretory vesicles. This displaces NE from the vesicles. Once NE reaches a concentration threshold within the presynaptic terminal, the action of NET is reversed, and NE is expelled into the synaptic cleft, contributing to the characteristics and effects of ↑ NE observed in patients taking amphetamines.



Cholinomimetic agents

Watch for exacerbation of COPD, asthma, and peptic ulcers in susceptible patients.

DRUG	ACTION	APPLICATIONS
Direct agonists		
Bethanechol	Activates b ladder smooth muscle; resistant to AChE. No nicotinic activity. “ Bethany, call me to activate your bladder. ”	Urinary retention.
Carbachol	C arbon copy of a cetyl ch oline (but resistant to AChE).	Constricts pupil and relieves intraocular pressure in open-angle glaucoma.
Methacholine	Stimulates m uscarinic receptors in airway when inhaled.	Challenge test for diagnosis of asthma.
Pilocarpine	Contracts ciliary muscle of eye (open-angle glaucoma), pupillary sphincter (closed-angle glaucoma); resistant to AChE, can cross blood-brain barrier (tertiary amine). “You cry, drool, and sweat on your p illow.”	Potent stimulator of sweat, tears, and saliva Open-angle and closed-angle glaucoma, xerostomia (Sjögren syndrome).
Indirect agonists (anticholinesterases)		
Donepezil, rivastigmine, galantamine	↑ ACh.	1st line for Alzheimer disease (Dona Riva dances at the gala).
Edrophonium	↑ ACh.	Historically used to diagnose myasthenia gravis; replaced by anti-AChR Ab (anti-acetylcholine receptor antibody) test.
Neostigmine	↑ ACh. Neo CNS = No CNS penetration (quaternary amine).	Postoperative and neurogenic ileus and urinary retention, myasthenia gravis, reversal of neuromuscular junction blockade (postoperative).
Physostigmine	↑ ACh. Ph reely (freely) crosses blood-brain barrier → CNS (tertiary amine).	Antidote for anticholinergic toxicity; physostigmine “phyxes” atropine overdose.
Pyridostigmine	↑ ACh; ↑ muscle strength. Used with glycopyrrolate, hyoscyamine, or propantheline to control pyridostigmine side effects. Pyridostigmine gets rid of myasthenia gravis.	Myasthenia gravis (long acting); does not penetrate CNS (quaternary amine).
Anticholinesterase poisoning		
Muscarinic effects	D iarrhea, U rination, M iosis, B ronchospasm, B radycardia, E mesis, L acrimation, S weating, S alivation.	DUMBBELSS. Reversed by atropine, a competitive inhibitor. Atropine can cross BBB to relieve CNS symptoms.
Nicotinic effects	Neuromuscular blockade (mechanism similar to succinylcholine).	Reversed by pralidoxime, regenerates AChE via dephosphorylation if given early. Pralidoxime (quaternary amine) does not readily cross BBB.
CNS effects	Respiratory depression, lethargy, seizures, coma.	

Muscarinic antagonists

DRUGS	ORGAN SYSTEMS	APPLICATIONS
Atropine, homatropine, tropicamide	Eye	Produce mydriasis and cycloplegia.
Benztropine, trihexyphenidyl	CNS	P arkinson disease (“ park my Benz ”). Acute dystonia.
Glycopyrrolate	GI, respiratory	Parenteral: preoperative use to reduce airway secretions. Oral: drooling, peptic ulcer.
Hyoscyamine, dicyclomine	GI	Antispasmodics for irritable bowel syndrome.
Ipratropium, tiotropium	Respiratory	COPD, asthma (“ I pray I can breathe soon!”).
Oxybutynin, solifenacin, tolterodine	Genitourinary	Reduce bladder spasms and urge urinary incontinence (overactive bladder).
Scopolamine	CNS	Motion sickness.

Atropine

Muscarinic antagonist. Used to treat bradycardia and for ophthalmic applications.


ORGAN SYSTEM	ACTION	NOTES
Eye	↑ pupil dilation, cycloplegia	Blocks muscarinic effects (DUMBBELSS) of anticholinesterases, but not the nicotinic effects.
Airway	Bronchodilation, ↓ secretions	
Stomach	↓ acid secretion	
Gut	↓ motility	
Bladder	↓ urgency in cystitis	
ADVERSE EFFECTS	<p>↑ body temperature (due to ↓ sweating); ↑ HR; dry mouth; dry, flushed skin; cycloplegia; constipation; disorientation</p> <p>Can cause acute angle-closure glaucoma in elderly (due to mydriasis), urinary retention in men with prostatic hyperplasia, and hyperthermia in infants.</p>	<p>Side effects:</p> <p>Hot as a hare Fast as a fiddle Dry as a bone Red as a beet Blind as a bat Mad as a hatter Full as a flask</p> <p>Jimson weed (<i>Datura</i>) → gardener’s pupil (mydriasis due to plant alkaloids)</p>

Sympathomimetics

DRUG	ACTION	HEMODYNAMIC CHANGES	APPLICATIONS
Direct sympathomimetics			
Albuterol, salmeterol, terbutaline	$\beta_2 > \beta_1$	↑ HR (little effect)	A lbuterol for A cute asthma/COPD. S almeterol for S erial (long-term) asthma/COPD. Terbutaline for acute bronchospasm in asthma and tocolysis.
Dobutamine	$\beta_1 > \beta_2, \alpha$	↔/↓ BP, ↑ HR, ↑ CO	Heart failure (HF), cardiogenic shock (inotropic > chronotropic), cardiac stress testing.
Dopamine	$D_1 = D_2 > \beta > \alpha$	↑ BP (high dose), ↑ HR, ↑ CO	Unstable bradycardia, HF, shock; inotropic and chronotropic effects at lower doses due to β effects; vasoconstriction at high doses due to α effects.
Epinephrine	$\beta > \alpha$	↑ BP (high dose), ↑ HR, ↑ CO	Anaphylaxis, asthma, open-angle glaucoma; α effects predominate at high doses. Significantly stronger effect at β_2 -receptor than norepinephrine.
Fenoldopam	D_1	↓ BP (vasodilation), ↑ HR, ↑ CO	Postoperative hypertension, hypertensive crisis. Vasodilator (coronary, peripheral, renal, and splanchnic). Promotes natriuresis. Can cause hypotension and tachycardia.
Isoproterenol	$\beta_1 = \beta_2$	↓ BP (vasodilation), ↑ HR, ↑ CO	Electrophysiologic evaluation of tachyarrhythmias. Can worsen ischemia. Has negligible α effect.
Midodrine	α_1	↑ BP (vasoconstriction), ↓ HR, ↔/↓ CO	Autonomic insufficiency and postural hypotension. May exacerbate supine hypertension.
Mirabegron	β_3		Urinary urgency or incontinence or overactive bladder. Think “mirab ₃ gron.”
Norepinephrine	$\alpha_1 > \alpha_2 > \beta_1$	↑ BP, ↑ HR, ↔/↑ CO	Hypotension, septic shock.
Phenylephrine	$\alpha_1 > \alpha_2$	↑ BP (vasoconstriction), ↓ HR, ↔/↓ CO	Hypotension (vasoconstrictor), ocular procedures (mydriatic), rhinitis (decongestant), ischemic priapism.
Indirect sympathomimetics			
Amphetamine	Indirect general agonist, reuptake inhibitor, also releases stored catecholamines		Narcolepsy, obesity, ADHD.
Cocaine	Indirect general agonist, reuptake inhibitor		Causes vasoconstriction and local anesthesia. Caution when giving β -blockers if cocaine intoxication is suspected (can lead to unopposed α_1 activation → extreme hypertension, coronary vasospasm).
Ephedrine	Indirect general agonist, releases stored catecholamines		Nasal decongestion (pseudoephedrine), urinary incontinence, hypotension.

Norepinephrine vs isoproterenol

NE \uparrow systolic and diastolic pressures as a result of α_1 -mediated vasoconstriction \rightarrow \uparrow mean arterial pressure \rightarrow reflex bradycardia. However, isoproterenol (rarely used) has little α effect but causes β_2 -mediated vasodilation, resulting in \downarrow mean arterial pressure and \uparrow heart rate through β_1 and reflex activity.




Sympatholytics (α_2 -agonists)


DRUG	APPLICATIONS	ADVERSE EFFECTS
Clonidine, guanfacine	Hypertensive urgency (limited situations), ADHD, Tourette syndrome, symptom control in opioid withdrawal	CNS depression, bradycardia, hypotension, respiratory depression, miosis, rebound hypertension with abrupt cessation
α-methyl dopa	Hypertension in pregnancy	Direct Coombs \oplus hemolysis, drug-induced lupus, hyperprolactinemia
Tizanidine	Relief of spasticity	Hypotension, weakness, xerostomia

α -blockers

DRUG	APPLICATIONS	ADVERSE EFFECTS
Nonselective		
Phenoxybenzamine	Irreversible. Pheochromocytoma (used preoperatively) to prevent catecholamine (hypertensive) crisis	Orthostatic hypotension, reflex tachycardia
Phentolamine	Reversible. Given to patients on MAO inhibitors who eat tyramine-containing foods and for severe cocaine-induced hypertension (2nd line)	
α_1 selective (-osin ending)		
Prazosin, terazosin, doxazosin, tamsulosin	Urinary symptoms of BPH; PTSD (prazosin); hypertension (except tamsulosin)	1st-dose orthostatic hypotension, dizziness, headache
α_2 selective		
Mirtazapine	Depression	Sedation, \uparrow serum cholesterol, \uparrow appetite



Epinephrine response exhibits reversal of mean arterial pressure from a net increase (the α response) to a net decrease (the β_2 response).



Phenylephrine response is suppressed but not reversed because it is a “pure” α -agonist (lacks β -agonist properties).

APPLICATION	ACTIONS	NOTES/EXAMPLES
β-blockers	Acebutolol, atenolol, betaxolol, bisoprolol, carvedilol, esmolol, labetalol, metoprolol, nadolol, nebivolol, pindolol, propranolol, timolol.	
Angina pectoris	↓ heart rate and contractility → ↓ O ₂ consumption	
Glaucoma	↓ production of aqueous humor	Timolol
Heart failure	↓ mortality	Bisoprolol, Carvedilol, Metoprolol (β-blockers Curb Mortality)
Hypertension	↓ cardiac output, ↓ renin secretion (due to β ₁ -receptor blockade on JG cells)	
Hyperthyroidism/ thyroid storm	Symptom control (↓ heart rate, ↓ tremor)	Propranolol
Hypertrophic cardiomyopathy	↓ heart rate → ↑ filling time, relieving obstruction	
Myocardial infarction	↓ O ₂ demand (short-term), ↓ mortality (long-term)	
Supraventricular tachycardia	↓ AV conduction velocity (class II antiarrhythmic)	Metoprolol, esmolol
Variceal bleeding	↓ hepatic venous pressure gradient and portal hypertension (prophylactic use)	Nadolol, propranolol, carvedilol
ADVERSE EFFECTS	Erectile dysfunction, cardiovascular (bradycardia, AV block, HF), CNS (seizures, sleep alterations), dyslipidemia (metoprolol), and asthma/COPD exacerbations	Use of β-blockers for acute cocaine-associated chest pain remains controversial due to unsubstantiated concern for unopposed α-adrenergic stimulation
SELECTIVITY	β ₁ -selective antagonists (β ₁ > β ₂)— a cebutolol (partial agonist), a tenolol, b etaxolol, b isoprolol, e smolol, m etoprolol	Selective antagonists mostly go from A to M (β ₁ with 1st half of alphabet)
	Nonselective antagonists (β ₁ = β ₂)— n adolol, p indolol (partial agonist), p ropranolol, t imolol	NonZ elective antagonists mostly go from N to Z (β ₂ with 2nd half of alphabet)
	Nonselective α- and β-antagonists— c arvedilol, l abetalol	Nonselective α- and β-antagonists have modified suffixes (instead of “-olol”)
	N ebivolol combines cardiac-selective β ₁ -adrenergic blockade with stimulation of β ₃ -receptors (activate n itric o xide synthase in the vasculature and ↓ SVR)	N ebivolol increases NO

Phosphodiesterase inhibitors

Phosphodiesterase (PDE) inhibitors inhibit PDE, which catalyzes the hydrolysis of cAMP and/or cGMP, and thereby increase cAMP and/or cGMP. These inhibitors have varying specificity for PDE isoforms and thus have different clinical uses.

TYPE OF INHIBITOR	MECHANISM OF ACTION	CLINICAL USES	ADVERSE EFFECTS
Nonspecific PDE inhibitor Theophylline	↓ cAMP hydrolysis → ↑ cAMP → bronchial smooth muscle relaxation → bronchodilation	COPD/asthma (rarely used)	Cardiotoxicity (eg, tachycardia, arrhythmia), neurotoxicity (eg, headache), abdominal pain
PDE-5 inhibitors Sildenafil, vardenafil, tadalafil, avanafil	↓ hydrolysis of cGMP → ↑ cGMP → ↑ smooth muscle relaxation by enhancing NO activity → pulmonary vasodilation and ↑ blood flow in corpus cavernosum fills the penis	Erectile dysfunction Pulmonary hypertension BPH (tadalafil only)	Facial flushing, headache, dyspepsia, hypotension in patients taking nitrates; “Hot and sweaty,” then Headache, Heartburn, Hypotension Sildenafil only: cyanopia (blue-tinted vision) via inhibition of PDE-6 in retina
PDE-4 inhibitor Roflumilast	↑ cAMP in neutrophils, granulocytes, and bronchial epithelium	Severe COPD	Abdominal pain, weight loss, mental disorders (eg, depression)
PDE-3 inhibitor Milrinone	In cardiomyocytes: ↑ cAMP → ↑ Ca ²⁺ influx → ↑ inotropy and chronotropy In vascular smooth muscle: ↑ cAMP → MLCK inhibition → vasodilation → ↓ preload and afterload	Acute decompensated HF with cardiogenic shock	Tachycardia, ventricular arrhythmias (thus not for chronic use), hypotension
“Platelet inhibitors” Cilostazol ^a Dipyridamole ^b	In platelets: ↑ cAMP → inhibition of platelet aggregation	Intermittent claudication Stroke or TIA prevention (with aspirin) Cardiac stress testing (dipyridamole only, due to coronary vasodilation) Prevention of coronary stent restenosis	Nausea, headache, facial flushing, hypotension, abdominal pain

^aCilostazol is a PDE-3 inhibitor, but due to its indications is categorized as a platelet inhibitor together with dipyridamole.

^bDipyridamole is a nonspecific PDE inhibitor, leading to inhibition of platelet aggregation. It also prevents adenosine reuptake by platelets → ↑ extracellular adenosine → ↑ vasodilation.

Ingested seafood toxins Toxin actions include **H**istamine release, **T**otal block of Na⁺ channels, or opening of Na⁺ channels to **C**ause depolarization.

TOXIN	SOURCE	ACTION	SYMPTOMS	TREATMENT
Histamine (scombroid poisoning)	Spoiled dark-meat fish such as tuna, mahi-mahi, mackerel, and bonito	Bacterial histidine decarboxylase converts histidine to histamine Frequently misdiagnosed as fish allergy	Mimics anaphylaxis: acute burning sensation of mouth, flushing of face, erythema, urticaria, itching May progress to bronchospasm, angioedema, hypotension	Antihistamines Albuterol and epinephrine if needed
Tetrodotoxin	Pufferfish	Highly potent toxin; binds fast voltage-gated Na ⁺ channels in nerve tissue, preventing depolarization	Nausea, diarrhea, paresthesias, weakness, dizziness, loss of reflexes	Supportive
Ciguatoxin	Reef fish such as barracuda, snapper, and moray eel	Opens Na ⁺ channels, causing depolarization	Nausea, vomiting, diarrhea; perioral numbness; reversal of hot and cold sensations; bradycardia, heart block, hypotension	Supportive

Beers criteria

Widely used criteria developed to reduce potentially inappropriate prescribing and harmful polypharmacy in the geriatric population. Includes > 50 medications that should be avoided in elderly patients due to ↓ efficacy and/or ↑ risk of adverse events. Examples:

- α-blockers (↑ risk of hypotension)
- Anticholinergics, antidepressants, antihistamines, opioids (↑ risk of delirium, sedation, falls, constipation, urinary retention)
- Benzodiazepines (↑ risk of delirium, sedation, falls)
- NSAIDs (↑ risk of GI bleeding, especially with concomitant anticoagulation)
- PPIs (↑ risk of *C difficile* infection)

▶ PHARMACOLOGY—TOXICITIES AND SIDE EFFECTS

Specific toxicity treatments

TOXIN	TREATMENT
Acetaminophen	N-acetylcysteine (replenishes glutathione)
AChE inhibitors, organophosphates	Atropine > pralidoxime
Antimuscarinic, anticholinergic agents	Physostigmine (crosses BBB), control hyperthermia
Arsenic	Dimercaprol, succimer
Benzodiazepines	Flumazenil
β-blockers	Atropine, glucagon, saline
Carbon monoxide	100% O ₂ , hyperbaric O ₂
Copper	“ Penny ”cillamine (penicillamine), trientine (copper penny × 3)
Cyanide	Hydroxocobalamin, nitrites + sodium thiosulfate
Digitalis (digoxin)	Digoxin-specific antibody fragments
Heparin	Protamine sulfate
Iron (Fe)	De fer oxamine, de fer asirox, de fer iprone
Lead	Calcium disodium EDTA, dimercaprol, succimer, penicillamine
Mercury	Di mer caprol, suc cimer
Methanol, ethylene glycol (antifreeze)	Fomepizole > ethanol, dialysis
Methemoglobin	Meth ylene blue, vitamin C (reducing agent)
OpiOids	Nal OxO ne
Salicylates	NaHCO ₃ (alkalinize urine), dialysis
TCA's	NaHCO ₃ (stabilizes cardiac cell membrane)
Warfarin	Vitamin K (delayed effect), PCC (prothrombin complex concentrate)/FFP (immediate effect)

Drug reactions—cardiovascular

DRUG REACTION	CAUSAL AGENTS
Coronary vasospasm	Cocaine , Amphetamines , Sumatriptan , Ergot alkaloids (CASE)
Cutaneous flushing	Vancomycin , Adenosine , Niacin , Ca²⁺ channel blockers , Echinocandins , Nitrates (flushed from VANCEN [dancing]) Red man syndrome —rate-dependent infusion reaction to vancomycin causing widespread pruritic erythema due to histamine release. Manage with diphenhydramine, slower infusion rate.
Dilated cardiomyopathy	Anthracyclines (eg, Doxorubicin , Daunorubicin); prevent with Dexrazoxane
Torsades de pointes	Agents that prolong QT interval: anti Arr rhythmics (class IA, III), anti Bio totics (eg, macrolides), anti“ C ”ychotics (eg, ziprasidone), anti D epressants (eg, TCAs), anti E metics (eg, ondansetron) (ABCDE)

Drug reactions—endocrine/reproductive

DRUG REACTION	CAUSAL AGENTS	NOTES
Adrenocortical insufficiency	HPA suppression 2° to glucocorticoid withdrawal	
Diabetes insipidus	Lithium, demeclocycline	
Hot flashes	SERMs (eg, tamoxifen, clomiphene, raloxifene)	
Hyperglycemia	Tacrolimus, Protease inhibitors, Niacin, HCTZ, Corticosteroids	The People Need Hard Candies
Hyperprolactinemia	Typical antipsychotics (eg, haloperidol), atypical antipsychotics (eg, risperidone), metoclopramide, methyl dopa, reserpine	Presents with hypogonadism (eg, infertility, amenorrhea, erectile dysfunction) and galactorrhea
Hyperthyroidism	Amiodarone, iodine	
Hypothyroidism	Amiodarone, Sulfonamides, Lithium	I AM Suddenly Lethargic
SIADH	Carbamazepine, Cyclophosphamide, SSRIs	Can't Concentrate Serum Sodium

Drug reactions—gastrointestinal

DRUG REACTION	CAUSAL AGENTS	NOTES
Acute cholestatic hepatitis, jaundice	Macrolides (eg, erythromycin)	
Diarrhea	Acamprosate, antidiabetic agents (acarbose, metformin, pramlintide), colchicine, cholinesterase inhibitors, lipid-lowering agents (eg, ezetimibe, orlistat), macrolides (eg, erythromycin), SSRIs, chemotherapy (eg, irinotecan)	
Focal to massive hepatic necrosis	Halothane, <i>Amanita phalloides</i> (death cap mushroom), Valproic acid, Acetaminophen	Liver “HAVAc”
Hepatitis	Rifampin, isoniazid, pyrazinamide, statins, fibrates	
Pancreatitis	Didanosine, Corticosteroids, Alcohol, Valproic acid, Azathioprine, Diuretics (eg, furosemide, HCTZ)	Drugs Causing A Violent Abdominal Distress
Pill-induced esophagitis	Bisphosphonates, ferrous sulfate, NSAIDs, potassium chloride, tetracyclines	Caustic effect minimized with upright posture and adequate water ingestion
Pseudomembranous colitis	Ampicillin, cephalosporins, clindamycin, fluoroquinolones, PPIs	Antibiotics predispose to superinfection by resistant <i>C difficile</i>

Drug reactions—hematologic

DRUG REACTION	CAUSAL AGENTS	NOTES
Agranulocytosis	Dapsone, Clozapine, Carbamazepine, Propylthiouracil, Methimazole, Colchicine, Ganciclovir	Drugs Can Cause Pretty Major Collapse of Granulocytes
Aplastic anemia	Carbamazepine, Methimazole, NSAIDs, Benzene, Chloramphenicol, Propylthiouracil	Can't Make New Blood Cells Properly
Direct Coombs ⊕ hemolytic anemia	Penicillin, methylDopa, Cephalosporins	P Diddy Coombs
Drug reaction with eosinophilia and systemic symptoms (DRESS)	Allopurinol, anticonvulsants, antibiotics, sulfa drugs	Potentially fatal delayed hypersensitivity reaction. Latency period (2- 8 weeks), then fever, morbilliform skin rash, frequent multiorgan involvement. Treatment: withdrawal of offending drug, corticosteroids
Gray baby syndrome	Chloramphenicol	
Hemolysis in G6PD deficiency	Isoniazid, Sulfonamides, Dapsone, Primaquine, Aspirin, Ibuprofen, Nitrofurantoin	Hemolysis IS D PAIN
Megaloblastic anemia	Hydroxyurea, Phenytoin, Methotrexate, Sulfa drugs	You're having a mega blast with PMS
Thrombocytopenia	Heparin, vancomycin, linezolid, quinidine, indinavir, ganciclovir, abciximab	
Thrombotic complications	Combined oral contraceptives, hormone replacement therapy, SERMs (eg, tamoxifen)	Estrogen-mediated side effect

Drug reactions—musculoskeletal/skin/connective tissue

DRUG REACTION	CAUSAL AGENTS	NOTES
Drug-induced lupus	Methyldopa, Minocycline, Hydralazine, Isoniazid, Phenytoin, Sulfa drugs, Etanercept, Procainamide	Lupus Makes My HIPS Extremely Painful
Fat redistribution	Protease inhibitors, Glucocorticoids	Fat PiG
Gingival hyperplasia	Cyclosporine, Ca ²⁺ channel blockers, Phenytoin	Can Cause Puffy gums
Hyperuricemia (gout)	Pyrazinamide, Thiazides, Furosemide, Niacin, Cyclosporine	Painful Tophi and Feet Need Care
Myopathy	Statins, fibrates, niacin, colchicine, daptomycin, hydroxychloroquine, interferon- α , penicillamine, glucocorticoids	
Osteoporosis	Corticosteroids, depot medroxyprogesterone acetate, GnRH agonists, aromatase inhibitors, anticonvulsants, heparin, PPIs	
Photosensitivity	Sulfonamides, Amiodarone, Tetracyclines, 5-FU	SAT For Photo
Rash (Stevens-Johnson syndrome)	Anti-epileptic drugs (especially lamotrigine), allopurinol, sulfa drugs, penicillin	Steven Johnson has epileptic allergy to sulfa drugs and penicillin
Teeth discoloration	Tetracyclines	Teethracyclines
Tendon/cartilage damage	Fluoroquinolones	

Drug reactions—neurologic

DRUG REACTION	CAUSAL AGENTS	NOTES
Cinchonism	Quinidine, quinine	Can present with tinnitus, hearing/vision loss, psychosis, and cognitive impairment
Parkinson-like syndrome	Antipsychotics, Reserpine, Metoclopramide	Cogwheel rigidity of ARM
Peripheral neuropathy	Isoniazid, phenytoin, platinum agents (eg, cisplatin), vincristine	
Idiopathic intracranial hypertension	Growth hormones, tetracyclines, vitamin A	
Seizures	Isoniazid, Bupropion, Imipenem/cilastatin, Tramadol, Enflurane	With seizures, I BITE my tongue
Tardive dyskinesia	Antipsychotics, metoclopramide	
Visual disturbance	Topiramate (blurred vision/diplopia, haloes), Digoxin (yellow-tinged vision), Isoniazid (optic neuritis), Vigabatrin (bilateral visual field defects), PDE-5 inhibitors (blue-tinged vision), Ethambutol (color vision changes)	These Drugs Irritate Very Precious Eyes

Drug reactions—renal/genitourinary

DRUG REACTION	CAUSAL AGENTS	NOTES
Fanconi syndrome	Cisplatin, ifosfamide, expired tetracyclines, tenofovir	
Hemorrhagic cystitis	Cyclophosphamide, ifosfamide	Prevent by coadministering with mesna
Interstitial nephritis	Diuretics (Pee), NSAIDs (Pain-free), Penicillins and cephalosporins, PPIs, rifamPin, and sulfa drugs	Remember the 5 P's

Drug reactions—respiratory

DRUG REACTION	CAUSAL AGENTS	NOTES
Dry cough	ACE inhibitors	
Pulmonary fibrosis	Methotrexate, Nitrofurantoin, Carmustine, Bleomycin, Busulfan, Amiodarone	My Nose Cannot Breathe Bad Air

Drug reactions—multiorgan

DRUG REACTION	CAUSAL AGENTS	NOTES
Antimuscarinic	Atropine, TCAs, H ₁ -blockers, antipsychotics	
Disulfiram-like reaction	1st-generation Sulfonylureas, Procarbazine, certain Cephalosporins, Griseofulvin, Metronidazole	Sorry Pals, Can't Go Mingle
Nephrotoxicity/ototoxicity	Loop diuretics, Aminoglycosides, cisPlatin, Vancomycin, amphotERICin B	Listen And Pee Very TERriBly Cisplatin toxicity may respond to amifostine

Drugs affecting pupil size

↑ pupil size	↓ pupil size
Anticholinergics (eg, atropine, TCAs, tropicamide, scopolamine, antihistamines)	Sympatholytics (eg, α_2 -agonists)
Drugs of abuse (eg, amphetamines, cocaine, LSD), meperidine	Drugs of abuse (eg, heroin/opioids)
Sympathomimetics	Parasympathomimetics (eg, pilocarpine), organophosphates

Cytochrome P-450 interactions (selected)

Inducers (+)	Substrates	Inhibitors (-)
Modafinil	Warfarin	Sodium valproate
Chronic alcohol use	Anti-epileptics	Isoniazid
St. John's wort	Theophylline	Cimetidine
Phenytoin	OCPs	Ketoconazole
Phenobarbital		Fluconazole
Nevirapine		Acute alcohol abuse
Rifampin		Chloramphenicol
Griseofulvin		Erythromycin/clarithromycin
Carbamazepine		Sulfonamides
		Ciprofloxacin
		Omeprazole
		Metronidazole
		Amiodarone
		Ritonavir
		Grapefruit juice
Most chronic alcoholics	War Against The OCPs	SICKFACES.COM (when
Steal Phen-Phen and Never		I Am Really drinking
Refuse Greasy Carbs		Grapefruit juice)

Sulfa drugs

Sulfonamide antibiotics, Sulfasalazine, Probenecid, Furosemide, Acetazolamide, Celecoxib, Thiazides, Sulfonylureas. Patients with sulfa allergies may develop fever, urinary tract infection, Stevens-Johnson syndrome, hemolytic anemia, thrombocytopenia, agranulocytosis, acute interstitial nephritis, and urticaria (hives).

Scary Sulfa Pharm FACTS

▶ PHARMACOLOGY—MISCELLANEOUS

Drug names

ENDING	CATEGORY	EXAMPLE
Antimicrobial		
-bendazole	Antiparasitic/antihelminthic	Mebendazole
-cillin	Transpeptidase inhibitor	Ampicillin
-conazole	Ergosterol synthesis inhibitor	Ketoconazole
-cycline	Protein synthesis inhibitor	Tetracycline
-ivir	Neuraminidase inhibitor	Oseltamivir
-navir	Protease inhibitor	Ritonavir
-ovir	Viral DNA polymerase inhibitor	Acyclovir
-tegravir	Integrase inhibitor	Elvitegravir, raltegravir
-thromycin	Macrolide antibiotic	Azithromycin
CNS		
-apine, -idone	Atypical antipsychotic	Quetiapine, risperidone
-azine	Typical antipsychotic	Thioridazine
-barbital	Barbiturate	Phenobarbital
-ipramine, -triptyline	TCA	Imipramine, amitriptyline
-triptan	5-HT _{1B/1D} agonist	Sumatriptan
-zepam, -zolam	Benzodiazepine	Diazepam, alprazolam
Autonomic		
-chol	Cholinergic agonist	Bethanechol, carbachol
-olol	β-blocker	Propranolol
-stigmine	AChE inhibitor	Neostigmine
-terol	β ₂ -agonist	Albuterol
-zosin	α ₁ -blocker	Prazosin
Cardiovascular		
-afil	PDE-5 inhibitor	Sildenafil
-dipine	Dihydropyridine Ca ²⁺ channel blocker	Amlodipine
-pril	ACE inhibitor	Captopril
-sartan	Angiotensin-II receptor blocker	Losartan
-xaban	Direct factor Xa inhibitor	Apixaban, edoxaban, rivaroxaban
Metabolic		
-glibflozin	SGLT-2 inhibitor	Dapagliflozin, canagliflozin
-glinide	Meglitinide	Repaglinide, nateglinide
-gliptin	DPP-4 inhibitor	Sitagliptin
-glitazone	PPAR-γ activator	Rosiglitazone
-glutide	GLP-1 analog	Liraglutide, albiglutide

Drug names (continued)

ENDING	CATEGORY	EXAMPLE
Other		
-dronate	Bisphosphonate	Alendronate
-prazole	Proton pump inhibitor	Omeprazole
-prost	Prostaglandin analog	Latanoprost
-sentan	Endothelin receptor antagonist	Bosentan
-tidine	H ₂ -antagonist	Cimetidine
-vaptan	ADH antagonist	Tolvaptan

Biologic agents

ENDING	CATEGORY	EXAMPLE
Monoclonal antibodies (-mab)—target overexpressed cell surface receptors		
-ximab	Chimeric human-mouse monoclonal antibody	Rituximab
-zumab	Humanized mouse monoclonal antibody	Bevacizumab
-umab	Human monoclonal antibody	Denosumab
Small molecule inhibitors (-ib)—target intracellular molecules		
-tinib	Tyrosine kinase inhibitor	Imatinib
-zomib	Proteasome inhibitor	Bortezomib
-ciclib	Cyclin-dependent kinase inhibitor	Palbociclib
Receptor fusion proteins (-cept)		
-cept	TNF- α antagonist	Etanercept
Interleukin receptor modulators (-kin)—agonists and antagonists of interleukin receptors		
-leukin	IL-2 agonist/analog	Aldesleukin
-kinra	Interleukin receptor antagonist	Anakinra

HIGH-YIELD PRINCIPLES IN

Public Health Sciences

“Medicine is a science of uncertainty and an art of probability.”

—William Osler

“There are two kinds of statistics: the kind you look up and the kind you make up.”

—Rex Stout

“On a long enough timeline, the survival rate for everyone drops to zero.”

—Chuck Palahniuk

“There are three kinds of lies: lies, damned lies, and statistics.”

—Mark Twain

A heterogenous mix of epidemiology, biostatistics, ethics, law, healthcare delivery, patient safety, quality improvement, and more falls under the heading of public health sciences. Biostatistics and epidemiology are the foundations of evidence-based medicine and are very high yield. Make sure you can quickly apply biostatistical equations such as sensitivity, specificity, and predictive values in a problem-solving format. Also, know how to set up your own 2×2 tables. Quality improvement and patient safety topics were introduced a few years ago on the exam and represent trends in health system science. Medical ethics questions often require application of principles. Typically, you are presented with a patient scenario and then asked how you would respond.

▶ Epidemiology and Biostatistics	256
▶ Ethics	265
▶ The Well Patient	270
▶ Healthcare Delivery	270
▶ Quality and Safety	273

▶ PUBLIC HEALTH SCIENCES—EPIDEMIOLOGY AND BIOSTATISTICS

Observational studies

STUDY TYPE	DESIGN	MEASURES/EXAMPLE
Cross-sectional study	Frequency of disease and frequency of risk-related factors are assessed in the present. Asks, “What is happening?”	Disease prevalence. Can show risk factor association with disease, but does not establish causality.
Case-control study	Compares a group of people with disease to a group without disease. Looks to see if odds of prior exposure or risk factor differ by disease state. Asks, “What happened?”	Odds ratio (OR). Patients with COPD had higher odds of a smoking history than those without COPD.
Cohort study	Compares a group with a given exposure or risk factor to a group without such exposure. Looks to see if exposure or risk factor is associated with later development of disease. Can be prospective or retrospective.	Relative risk (RR). Smokers had a higher risk of developing COPD than nonsmokers. Cohort = relative risk.
Crossover study	Compares the effect of a series of ≥ 2 treatments on a participant. Order in which participants receive treatments is randomized. Washout period occurs between each treatment.	Allows participants to serve as their own controls.
Twin concordance study	Compares the frequency with which both monozygotic twins vs both dizygotic twins develop the same disease.	Measures heritability and influence of environmental factors (“nature vs nurture”).
Adoption study	Compares siblings raised by biological vs adoptive parents.	Measures heritability and influence of environmental factors.

Clinical trial

Experimental study involving humans. Compares therapeutic benefits of ≥ 2 treatments, or of treatment and placebo. Study quality improves when study is randomized, controlled, and double-blinded (ie, neither patient nor doctor knows whether the patient is in the treatment or control group). Triple-blind refers to the additional blinding of the researchers analyzing the data. Four phases (“Does the drug **SWIM?**”).

DRUG TRIALS	TYPICAL STUDY SAMPLE	PURPOSE
Phase I	Small number of either healthy volunteers or patients with disease of interest.	“Is it S afe?” Assesses safety, toxicity, pharmacokinetics, and pharmacodynamics.
Phase II	Moderate number of patients with disease of interest.	“Does it W ork?” Assesses treatment efficacy, optimal dosing, and adverse effects.
Phase III	Large number of patients randomly assigned either to the treatment under investigation or to the standard of care (or placebo).	“Is it as good or better?” Compares the new treatment to the current standard of care (any I mprovement?).
Phase IV	Postmarketing surveillance of patients after treatment is approved.	“Can it stay?” Detects rare or long-term adverse effects (eg, black box warnings). Can result in treatment being withdrawn from M arket.

Evaluation of diagnostic tests

Sensitivity and specificity are fixed properties of a test. PPV and NPV vary depending on disease prevalence in population being tested.

		Disease		
		⊕	⊖	
Test	⊕	TP	FP	PPV = TP / (TP + FP)
	⊖	FN	TN	NPV = TN / (TN + FN)
		Sensitivity = TP / (TP + FN)	Specificity = TN / (TN + FP)	Prevalence = TP + FN / (TP + FN + FP + TN)

Sensitivity (true-positive rate)

Proportion of all people with disease who test positive, or the probability that when the disease is present, the test is positive. Value approaching 100% is desirable for **ruling out** disease and indicates a **low false-negative rate**.

$= TP / (TP + FN)$
 $= 1 - FN \text{ rate}$
SN-N-OUT = highly **SeNsitive** test, when **Negative**, rules **OUT** disease
 High sensitivity test used for screening

Specificity (true-negative rate)

Proportion of all people without disease who test negative, or the probability that when the disease is absent, the test is negative. Value approaching 100% is desirable for **ruling in** disease and indicates a **low false-positive rate**.

$= TN / (TN + FP)$
 $= 1 - FP \text{ rate}$
SP-P-IN = highly **SPecific** test, when **Positive**, rules **IN** disease
 High specificity test used for confirmation after a positive screening test

Positive predictive value


Probability that a person who has a positive test result actually has the disease.

$PPV = TP / (TP + FP)$
 PPV varies directly with pretest probability (baseline risk, such as prevalence of disease):
 high pretest probability → high PPV

Negative predictive value

Probability that a person with a negative test result actually does not have the disease.

$NPV = TN / (TN + FN)$
 NPV varies inversely with prevalence or pretest probability



Possible cutoff values for (+) vs (-) test result
A = 100% sensitivity cutoff value
B = practical compromise between specificity and sensitivity
C = 100% specificity cutoff value

Lowering the cutoff value:	↑ Sensitivity ↑ NPV
B → A (↑ FN ↓ FP)	↓ Specificity ↓ PPV
Raising the cutoff value:	↑ Specificity ↑ PPV
B → C (↑ FN ↓ FP)	↓ Sensitivity ↓ NPV

Likelihood ratio

Likelihood that a given test result would be expected in a patient with the target disorder compared to the likelihood that the same result would be expected in a patient without the target disorder.

$$LR^+ = \frac{\text{sensitivity}}{1 - \text{specificity}} = \frac{TP \text{ rate}}{FP \text{ rate}}$$

$$LR^- = \frac{1 - \text{sensitivity}}{\text{specificity}} = \frac{FN \text{ rate}}{TN \text{ rate}}$$

$LR^+ > 10$ indicates a highly specific test, while $LR^- < 0.1$ indicates a highly sensitive test. LRs can be multiplied with pretest odds of disease to estimate posttest odds.


Quantifying risk

Definitions and formulas are based on the classic 2×2 or contingency table.

		Disease or outcome	
		⊕	⊖
Exposure or intervention	⊕	a	b
	⊖	c	d

TERM	DEFINITION	EXAMPLE	FORMULA								
Odds ratio	Typically used in case-control studies. Represents the odds of exposure among cases (a/c) vs odds of exposure among controls (b/d).	If in a case-control study, 20/30 lung cancer patients and 5/25 healthy individuals report smoking, the OR is 8; so the lung cancer patients are 8 times more likely to have a history of smoking.	$OR = \frac{a/c}{b/d} = \frac{ad}{bc}$ <table border="1"> <tr> <td>a</td> <td>b</td> </tr> <tr> <td>20</td> <td>5</td> </tr> <tr> <td>c</td> <td>d</td> </tr> <tr> <td>10</td> <td>20</td> </tr> </table>	a	b	20	5	c	d	10	20
a	b										
20	5										
c	d										
10	20										
Relative risk	Typically used in cohort studies. Risk of developing disease in the exposed group divided by risk in the unexposed group. RR = 1 → no association between exposure and disease. RR > 1 → exposure associated with ↑ disease occurrence. RR < 1 → exposure associated with ↓ disease occurrence.	If 5/10 people exposed to radiation are diagnosed with cancer, and 1/10 people not exposed to radiation are diagnosed with cancer, the RR is 5; so people exposed to radiation have a 5 times greater risk of developing cancer. For rare diseases (low prevalence), OR approximates RR.	$RR = \frac{a/(a+b)}{c/(c+d)}$ <table border="1"> <tr> <td>a</td> <td>b</td> </tr> <tr> <td>5</td> <td>5</td> </tr> <tr> <td>c</td> <td>d</td> </tr> <tr> <td>1</td> <td>9</td> </tr> </table>	a	b	5	5	c	d	1	9
a	b										
5	5										
c	d										
1	9										
Relative risk reduction	The proportion of risk reduction attributable to the intervention as compared to a control.	If 2% of patients who receive a flu shot develop the flu, while 8% of unvaccinated patients develop the flu, then RR = 2/8 = 0.25, and RRR = 0.75.	$RRR = 1 - RR$								
Attributable risk	The difference in risk between exposed and unexposed groups.	If risk of lung cancer in smokers is 21% and risk in nonsmokers is 1%, then the attributable risk is 20%.	$AR = \frac{a}{a+b} - \frac{c}{c+d}$ $AR\% = \frac{RR - 1}{RR} \times 100$								
Absolute risk reduction	The difference in risk (not the proportion) attributable to the intervention as compared to a control.	If 8% of people who receive a placebo vaccine develop the flu vs 2% of people who receive a flu vaccine, then ARR = 8% - 2% = 6% = 0.06.	$ARR = \frac{c}{c+d} - \frac{a}{a+b}$								
Number needed to treat	Number of patients who need to be treated for 1 patient to benefit. Lower number = better treatment.		$NNT = 1/ARR$								
Number needed to harm	Number of patients who need to be exposed to a risk factor for 1 patient to be harmed. Higher number = safer exposure.		$NNH = 1/AR$								
Case fatality rate	Percentage of deaths occurring among those with disease.	If 4 patients die among 10 cases of meningitis, case fatality rate is 40%.	$CFR\% = \frac{\text{deaths}}{\text{cases}} \times 100$								

Incidence vs prevalence



$$\text{Incidence} = \frac{\# \text{ of new cases}}{\# \text{ of people at risk}} \quad (\text{per unit of time})$$

$$\text{Prevalence} = \frac{\# \text{ of existing cases}}{\text{Total \# of people in a population}} \quad (\text{at a point in time})$$

$$\frac{\text{Prevalence}}{1 - \text{prevalence}} = \text{Incidence rate} \times \text{average duration of disease}$$

Prevalence \approx incidence for short duration disease (eg, common cold).

Prevalence $>$ incidence for chronic diseases, due to large # of existing cases (eg, diabetes).

Incidence looks at new cases (**incidents**).

Prevalence looks at **all** current cases.

Prevalence \sim pretest probability.
 \uparrow prevalence $\rightarrow \uparrow$ PPV and \downarrow NPV.

SITUATION	INCIDENCE	PREVALENCE
\uparrow survival time	—	\uparrow
\uparrow mortality	—	\downarrow
Faster recovery time	—	\downarrow
Extensive vaccine administration	\downarrow	\downarrow
\downarrow risk factors	\downarrow	\downarrow

Precision vs accuracy

Precision (reliability)


The consistency and reproducibility of a test.
The absence of random variation in a test.

Random error \downarrow precision in a test.
 \uparrow precision $\rightarrow \downarrow$ standard deviation.
 \uparrow precision $\rightarrow \uparrow$ statistical power ($1 - \beta$).

Accuracy (validity)

The closeness of test results to the true values.
The absence of systematic error or bias in a test.


Systematic error \downarrow accuracy in a test.



Receiving operating characteristic curve

ROC curve demonstrates how well a diagnostic test can distinguish between 2 groups (eg, disease vs healthy). Plots the true-positive rate (sensitivity) against the false-positive rate (1 – specificity).

The better performing test will have a higher area under the curve (AUC), with the curve closer to the upper left corner.

**Bias and study errors**

TYPE	DEFINITION	EXAMPLES	STRATEGIES TO REDUCE BIAS
Recruiting participants			
Selection bias	Nonrandom sampling or treatment allocation of subjects such that study population is not representative of target population. Most commonly a sampling bias.	Berkson bias —cases and/or controls selected from hospitals are less healthy and have different exposures than general population Attrition bias —participants lost to follow up have a different prognosis than those who complete the study	Randomization Ensure the choice of the right comparison/reference group
Performing study			
Recall bias	Awareness of disorder alters recall by subjects; common in retrospective studies	Patients with disease recall exposure after learning of similar cases	Decrease time from exposure to follow-up
Measurement bias	Information is gathered in a systemically distorted manner	Using a faulty automatic sphygmomanometer to measure BP Hawthorne effect —participants change behavior upon awareness of being observed	Use objective, standardized, and previously tested methods of data collection that are planned ahead of time Use placebo group
Procedure bias	Subjects in different groups are not treated the same	Patients in treatment group spend more time in highly specialized hospital units	Blinding (masking) and use of placebo reduce influence of participants and researchers on procedures and interpretation of outcomes as neither are aware of group assignments
Observer-expectancy bias	Researcher's belief in the efficacy of a treatment changes the outcome of that treatment (aka, Pygmalion effect)	An observer expecting treatment group to show signs of recovery is more likely to document positive outcomes	

Bias and study errors (continued)

TYPE	DEFINITION	EXAMPLES	STRATEGY TO REDUCE BIAS
Interpreting results			
Confounding bias	Factor related to both exposure and outcome (but not on causal path) distorts effect of exposure on outcome (vs effect modification, in which the exposure leads to different outcomes in subgroups stratified by the factor)	An uncontrolled study shows an association between drinking coffee and lung cancer. However, coffee drinkers also smoke more, which can account for the association	Multiple/repeated studies Crossover studies (subjects act as their own controls) Matching (patients with similar characteristics in both treatment and control groups)
Lead-time bias	Early detection is confused with ↑ survival	Early detection makes it seem like survival has increased, but the disease's natural history has not changed	Measure “back-end” survival (adjust survival according to the severity of disease at the time of diagnosis)
Length-time bias	Screening test detects diseases with long latency period, while those with shorter latency period become symptomatic earlier	A slowly progressive cancer is more likely detected by a screening test than a rapidly progressive cancer	A randomized controlled trial assigning subjects to the screening program or to no screening

Statistical distribution**Measures of central tendency**

Mean = (sum of values)/(total number of values).

Most affected by outliers (extreme values).

Median = middle value of a list of data sorted from least to greatest.

If there is an even number of values, the median will be the average of the middle two values.

Mode = most common value.

Least affected by outliers.

Measures of dispersion

Standard deviation = how much variability exists in a set of values, around the mean of these values.

σ = SD; n = sample size.

Variance = (SD)².

Standard error = an estimate of how much variability exists in a (theoretical) set of sample means around the true population mean.


SE = σ/\sqrt{n} .

SE ↓ as n ↑.

Normal distribution

Gaussian, also called bell-shaped.


Mean = median = mode.

**Nonnormal distributions****Bimodal**


Suggests two different populations (eg, metabolic polymorphism such as fast vs slow acetylators; age at onset of Hodgkin lymphoma; suicide rate by age).

**Positive skew**

Typically, mean > median > mode. Asymmetry with longer tail on right.

**Negative skew**

Typically, mean < median < mode. Asymmetry with longer tail on left.

**Statistical hypotheses****Null (H₀)**

Hypothesis of no difference or relationship (eg, there is no association between the disease and the risk factor in the population).

Alternative (H₁)

Hypothesis of some difference or relationship (eg, there is some association between the disease and the risk factor in the population).

Outcomes of statistical hypothesis testing**Correct result**

Stating that there is an effect or difference when one exists (null hypothesis rejected in favor of alternative hypothesis).

Stating that there is no effect or difference when none exists (null hypothesis not rejected).

		Reality	
		H_1	H_0
Study	rejects H_0	Power ($1 - \beta$)	α Type I error
	does not reject H_0	β Type II error	

Blue shading = correct result.

Incorrect result**Type I error (α)**

Stating that there is an effect or difference when none exists (null hypothesis incorrectly rejected in favor of alternative hypothesis).

α is the probability of making a type I error. p is judged against a preset α level of significance (usually 0.05). If $p < 0.05$ for a study outcome, the probability of obtaining that result purely by chance is $< 5\%$.

Statistical significance \neq clinical significance.

Also called false-positive error.

α = you **a**ccused an innocent man.
You can never “prove” the alternate hypothesis, but you can reject the null hypothesis as being very unlikely.

Type II error (β)

Stating that there is not an effect or difference when one exists (null hypothesis is not rejected when it is in fact false).

β is the probability of making a type II error. β is related to statistical power ($1 - \beta$), which is the probability of rejecting the null hypothesis when it is false.

↑ power and ↓ β by:

- ↑ sample size
- ↑ expected effect size
- ↑ precision of measurement

Also called false-negative error.

β = you **b**lindly let the guilty man go free.
If you ↑ sample size, you ↑ power. There is **power in numbers**.

Confidence interval

Range of values within which the true mean of the population is expected to fall, with a specified probability.

CI for sample mean = $\bar{x} \pm Z(SE)$

The 95% CI (corresponding to $\alpha = .05$) is often used. As sample size increases, CI narrows.

For the 95% CI, $Z = 1.96$.

For the 99% CI, $Z = 2.58$.

If the 95% CI for a mean difference between 2 variables includes 0, then there is no significant difference and H_0 is not rejected.

If the 95% CI for odds ratio or relative risk includes 1, H_0 is not rejected.

If the CIs between 2 groups do not overlap → statistically significant difference exists.

If the CIs between 2 groups overlap → usually no significant difference exists.


Meta-analysis

A method of statistical analysis that pools summary data (eg, means, RRs) from multiple studies for a more precise estimate of the size of an effect. Also estimates heterogeneity of effect sizes between studies.

Improves power, strength of evidence, and generalizability of study findings. Limited by quality of individual studies and bias in study selection.

Common statistical tests

t-test	Checks differences between means of 2 groups.	Tea is meant for 2. Example: comparing the mean blood pressure between men and women.
ANOVA	Checks differences between means of 3 or more groups.	3 words: AN alysis Of VA riance. Example: comparing the mean blood pressure between members of 3 different ethnic groups.
Chi-square (χ^2)	Checks differences between 2 or more percentages or proportions of categorical outcomes (not mean values).	Pronounce Chi-tegorical . Example: comparing the percentage of members of 3 different ethnic groups who have essential hypertension.
Fisher's exact test	Checks differences between 2 percentages or proportions of categorical, nominal outcomes. Use instead of chi-square test with small populations.	Example: comparing the percentage of 20 men and 20 women with hypertension.




Pearson correlation coefficient

r is always between -1 and $+1$. The closer the absolute value of r is to 1, the stronger the linear correlation between the 2 variables. Variance is how much the measured values differ from the average value in a data set.

Positive r value → positive correlation (as one variable ↑, the other variable ↑).

Negative r value → negative correlation (as one variable ↑, the other variable ↓).

Coefficient of determination = r^2 (amount of variance in one variable that can be explained by variance in another variable).



► PUBLIC HEALTH SCIENCES—ETHICS

Core ethical principles

Autonomy	Obligation to respect patients as individuals (truth-telling, confidentiality), to create conditions necessary for autonomous choice (informed consent), and to honor their preference in accepting or not accepting medical care.
Beneficence	Physicians have a special ethical (fiduciary) duty to act in the patient's best interest. May conflict with autonomy (an informed patient has the right to decide) or what is best for society (eg, mandatory TB treatment). Traditionally, patient interest supersedes.
Nonmaleficence	"Do no harm." Must be balanced against beneficence; if the benefits outweigh the risks, a patient may make an informed decision to proceed (most surgeries and medications fall into this category).
Justice	To treat persons fairly and equitably. This does not always imply equally (eg, triage).

Informed consent

A process (not just a document/signature) that requires:

- Disclosure: discussion of pertinent information (using medical interpreter, if needed)
- Understanding: ability to comprehend
- Capacity: ability to reason and make one's own decisions (distinct from competence, a legal determination)
- Voluntariness: freedom from coercion and manipulation

Patients must have an intelligent understanding of their diagnosis and the risks/benefits of proposed treatment and alternative options, including no treatment.

Patient must be informed that he or she can revoke written consent at any time, even orally.

Exceptions to informed consent (**WIPE** it away):

- **Waiver**—patient explicitly waives the right of informed consent
- **Legally Incompetent**—patient lacks decision-making capacity (obtain consent from legal surrogate)
- **Therapeutic Privilege**—withholding information when disclosure would severely harm the patient or undermine informed decision-making capacity
- **Emergency situation**—implied consent may apply

Consent for minors

A minor is generally any person < 18 years old. Parental consent laws in relation to healthcare vary by state. In general, parental consent should be obtained, but exceptions exist for emergency treatment (eg, blood transfusions) or if minor is legally emancipated (eg, married, self-supporting, or in the military).

Situations in which parental consent is usually not required:

- **Sex** (contraception, STIs, pregnancy)
- **Drugs** (substance abuse)
- **Rock and roll** (emergency/trauma)

Physicians should always encourage healthy minor-guardian communication.

Physician should seek a minor's assent even if their consent is not required.

Decision-making capacity

Physician must determine whether the patient is psychologically and legally capable of making a particular healthcare decision. Note that decisions made with capacity cannot be revoked simply if the patient later loses capacity. Intellectual disability alone (eg, Down syndrome, autism) is not an exclusion criterion for informed decision-making.

Capacity is determined by a physician for a specific healthcare-related decision (eg, to refuse medical care). Competency is determined by a judge and usually refers to more global categories of decision making (eg, legally unable to make any healthcare-related decision).

Components (think **GIEMSA**):

- Decision is consistent with patient's values and **G**oals
- Patient is **I**nformed (knows and understands)
- Patient **E**xpresses a choice
- Decision is not a result of altered **M**ental status (eg, delirium, psychosis, intoxication), **M**ood disorder
- Decision remains **S**table over time
- Patient is \geq 18 years of **A**ge or otherwise legally emancipated

Advance directives

Instructions given by a patient in anticipation of the need for a medical decision. Details vary per state law.

Oral advance directive

Incapacitated patient's prior oral statements commonly used as guide. Problems arise from variance in interpretation. If patient was informed, directive was specific, patient made a choice, and decision was repeated over time to multiple people, then the oral directive is more valid.

Written advance directive

Specifies specific healthcare interventions that a patient anticipates he or she would accept or reject during treatment for a critical or life-threatening illness. A living will is an example.

Medical power of attorney

Patient designates an agent to make medical decisions in the event that he/she loses decision-making capacity. Patient may also specify decisions in clinical situations. Can be revoked by patient if decision-making capacity is intact. More flexible than a living will.

Do not resuscitate order

DNR order prohibits cardiopulmonary resuscitation (CPR). Other resuscitative measures that may follow (eg, feeding tube) are also typically avoided.

Surrogate decision-maker

If a patient loses decision-making capacity and has not prepared an advance directive, individuals (surrogates) who know the patient must determine what the patient would have done. Priority of surrogates: **spouse** → adult **C**hildren → **P**arents → **S**iblings → other relatives (the **spouse ChiPS** in).

Confidentiality

Confidentiality respects patient privacy and autonomy. If the patient is incapacitated or the situation is emergent, disclosing information to family and friends should be guided by professional judgment of patient's best interest. The patient may voluntarily waive the right to confidentiality (eg, insurance company request).

General principles for exceptions to confidentiality:

- Potential physical harm to others is serious and imminent
- Alternative means to warn or protect those at risk is not possible
- Self-harm is likely
- Steps can be taken to prevent harm

Examples of exceptions to patient confidentiality (many are state specific) include the following (“The physician’s good judgment **SAVED** the day”):

- **S**uicidal/homicidal patients.
 - **A**buse (children, elderly, and/or prisoners).
 - Duty to protect—state-specific laws that sometimes allow physician to inform or somehow protect potential **V**ictim from harm.
 - **E**pileptic patients and other impaired automobile drivers.
 - Reportable **D**iseases (eg, STIs, hepatitis, food poisoning); physicians may have a duty to warn public officials, who will then notify people at risk. Dangerous communicable diseases, such as TB or Ebola, may require involuntary treatment.
-

Ethical situations

SITUATION	APPROPRIATE RESPONSE
Patient is not adherent.	Attempt to identify the reason for nonadherence and determine his/her willingness to change; do not coerce the patient into adhering and do not refer him/her to another physician.
Patient desires an unnecessary procedure.	Attempt to understand why the patient wants the procedure and address underlying concerns. Do not refuse to see the patient and do not refer him/her to another physician. Avoid performing unnecessary procedures.
Patient has difficulty taking medications.	Provide written instructions; attempt to simplify treatment regimens; use teach-back method (ask patient to repeat regimen back to physician) to ensure comprehension.
Family members ask for information about patient's prognosis.	Avoid discussing issues with relatives without the patient's permission.
A patient's family member asks you not to disclose the results of a test if the prognosis is poor because the patient will be "unable to handle it."	Attempt to identify why the family member believes such information would be detrimental to the patient's condition. Explain that as long as the patient has decision-making capacity and does not indicate otherwise, communication of information concerning his/her care will not be withheld. However, if you believe the patient might seriously harm himself/herself or others if informed, then you may invoke therapeutic privilege and withhold the information.
A 17-year-old girl is pregnant and requests an abortion.	Many states require parental notification or consent for minors for an abortion. Unless there are specific medical risks associated with pregnancy, a physician should not sway the patient's decision for, or against, an elective abortion (regardless of maternal age or fetal condition).
A 15-year-old girl is pregnant and wants to keep the child. Her parents want you to tell her to give the child up for adoption.	The patient retains the right to make decisions regarding her child, even if her parents disagree. Provide information to the teenager about the practical issues of caring for a baby. Discuss the options, if requested. Encourage discussion between the teenager and her parents to reach the best decision.
A terminally ill patient requests physician assistance in ending his/her own life.	Overwhelming majority of states refuse involvement in any form of physician-assisted death. Physicians may, however, prescribe medically appropriate analgesics even if they shorten the patient's life.
Patient is suicidal.	Assess the seriousness of the threat. If it is serious, suggest that the patient remain in the hospital voluntarily; patient can be hospitalized involuntarily if he/she refuses.
Patient states that he/she finds you attractive.	Ask direct, closed-ended questions and use a chaperone if necessary. Romantic relationships with patients are never appropriate. It may be necessary to transition care to another physician.
A woman who had a mastectomy says she now feels "ugly."	Find out why the patient feels this way. Do not offer falsely reassuring statements (eg, "You still look good").
Patient is angry about the long time he/she spent in the waiting room.	Acknowledge the patient's anger, but do not take a patient's anger personally. Apologize for any inconvenience. Stay away from efforts to explain the delay.
Patient is upset with the way he/she was treated by another doctor.	Suggest that the patient speak directly to that physician regarding his/her concerns. If the problem is with a member of the office staff, tell the patient you will speak to that person.
An invasive test is performed on the wrong patient.	Regardless of the outcome, a physician is ethically obligated to inform a patient that a mistake has been made.

Ethical situations (continued)

SITUATION	APPROPRIATE RESPONSE
A patient requires a treatment not covered by his/her insurance.	Never limit or deny care because of the expense in time or money. Discuss all treatment options with patients, even if some are not covered by their insurance companies.
A 7-year-old boy loses a sister to cancer and now feels responsible.	At ages 5–7, children begin to understand that death is permanent, that all life functions end completely at death, and that everything that is alive eventually dies. Provide a direct, concrete description of his sister's death. Avoid clichés and euphemisms. Reassure the boy that he is not responsible. Identify and normalize fears and feelings. Encourage play and healthy coping behaviors (eg, remembering her in his own way).
Patient is victim of intimate partner violence.	Ask if patient is safe and has an emergency plan. Do not necessarily pressure patient to leave his or her partner, or disclose the incident to the authorities (unless required by state law).
Patient wants to try alternative or holistic medicine.	Explore any underlying reasons with the patient in a supportive, nonjudgmental manner. Advise the patient of known benefits and risks of treatment, including adverse effects, contraindications, and medication interactions.
Physician colleague presents to work impaired.	If impaired or incompetent, colleague is a threat to patient safety. Report the situation to local supervisory personnel. Should the organization fail to take action, alert the state licensing board.
Patient is officially determined to suffer brain death. Patient's family insists on maintaining life support indefinitely because patient is still moving when touched.	Gently explain to family that there is no chance of recovery, and that brain death is equivalent to death. Movement is due to spinal arc reflex and is not voluntary. Bring case to appropriate ethics board regarding futility of care and withdrawal of life support.
A pharmaceutical company offers you a sponsorship in exchange for advertising its new drug.	Reject this offer. Generally, decline gifts and sponsorships to avoid any appearance of conflict of interest. The AMA Code of Ethics does make exceptions for gifts directly benefitting patients; gifts of minimal value; special funding for medical education of students, residents, fellows; grants whose recipients are chosen by independent institutional criteria; and funds that are distributed without attribution to sponsors.
Patient requests a nonemergent procedure that is against your personal or religious beliefs.	Provide accurate and unbiased information so patients can make an informed decision. Explain to the patient that you do not perform the procedure but offer to refer him/her to another physician.
Mother and 15-year-old daughter are unresponsive following a car accident and are bleeding internally. Father says do not transfuse because they are Jehovah's Witnesses.	Transfuse daughter, but do not transfuse mother. Emergent care can be refused by the healthcare proxy for an adult, particularly when patient preferences are known or reasonably inferred, but not for a minor based solely on faith.
A child presents with injuries inconsistent with parental story.	Contact child protective services and ensure child is in a safe location. Physicians are required by law to report any reasonable suspicion of child abuse or endangerment.

▶ PUBLIC HEALTH SCIENCES—THE WELL PATIENT

Changes in the elderly

Sexual changes:

- Men—slower erection/ejaculation, longer refractory period, but unchanged libido.
- Women—vaginal shortening, thinning, and dryness

Sleep patterns: ↓ REM and slow-wave sleep, ↑ sleep latency, ↑ early awakenings

↑ suicide rate

↓ vision and hearing

↓ immune response

↓ renal, pulmonary, and GI function

↓ muscle mass, ↑ fat

Intelligence does not decrease

▶ PUBLIC HEALTH SCIENCES—HEALTHCARE DELIVERY

Disease prevention

Primary disease prevention

Prevent disease before it occurs (eg, HPV vaccination)

Secondary disease prevention

Screen early for and manage existing but asymptomatic disease (eg, Pap smear for cervical cancer)

Tertiary disease prevention

Treatment to reduce complications from disease that is ongoing or has long-term effects (eg, chemotherapy)

Quaternary disease prevention

Quit (avoid) unnecessary medical interventions to minimize incidental harm (eg, imaging studies, optimizing medications to reduce polypharmacy).

Major medical insurance plans

PLAN	PROVIDERS	PAYMENTS	SPECIALIST CARE
Exclusive provider organization	Restricted to limited panel (except emergencies)		No referral required
Health maintenance organization	Restricted to limited panel (except emergencies)	Denied for any service that does not meet established, evidence-based guidelines	Requires referral from primary care provider
Point of service	Patient can see providers outside network	Higher copays and deductibles for out-of-network services	Requires referral from primary care provider
Preferred provider organization	Patient can see providers outside network	Higher copays and deductibles for all services	No referral required
Accountable care organization	Providers voluntarily enroll	Medicare	Specialists voluntarily enroll

Healthcare payment models

Bundled payment	Healthcare organization receives a set amount per service, regardless of ultimate cost, to be divided among all providers and facilities involved.
Capitation	Physicians receive a set amount per patient assigned to them per period of time, regardless of how much the patient uses the healthcare system. Used by some HMOs.
Discounted fee-for-service	Patient pays for each individual service at a discounted rate predetermined by providers and payers (eg, PPOs).
Fee-for-service	Patient pays for each individual service.
Global payment	Patient pays for all expenses associated with a single incident of care with a single payment. Most commonly used during elective surgeries, as it covers the cost of surgery as well as the necessary pre- and postoperative visits.

Medicare and Medicaid

Medicare and Medicaid—federal social healthcare programs that originated from amendments to the Social Security Act. Medicare is available to patients ≥ 65 years old, < 65 with certain disabilities, and those with end-stage renal disease. Medicaid is joint federal and state health assistance for people with limited income and/or resources.

Medicar**E** is for **E**lderly.
Medicai**D** is for **D**estitute.

The 4 parts of Medicare:

- Part **A**: Hospit**A**l insurance, home hospice care
- Part **B**: **B**asic medical **b**ills (eg, doctor’s fees, diagnostic testing)
- Part **C**: (parts A + B = **C**ombo) delivered by approved private **c**ompanies
- Part **D**: Prescription **D**rugs

Hospice care

Medical care focused on providing comfort and palliation instead of definitive cure. Available to patients on Medicare or Medicaid and in most private insurance plans whose life expectancy is < 6 months. During end-of-life care, priority is given to improving the patient’s comfort and relieving pain (often includes opioid, sedative, or anxiolytic medications). Facilitating comfort is prioritized over potential side effects (eg, respiratory depression). This prioritization of positive effects over negative effects is called the **principle of double effect**.

Common causes of death (US) by age

	< 1 YR	1–14 YR	15–34 YR	35–44 YR	45–64 YR	65+ YR
#1	Congenital malformations	Unintentional injury	Unintentional injury	Unintentional injury	Cancer	Heart disease
#2	Preterm birth	Cancer	Suicide	Cancer	Heart disease	Cancer
#3	Maternal pregnancy complications	Congenital malformations	Homicide	Heart disease	Unintentional injury	Chronic respiratory disease

Conditions with frequent hospital readmissions

Readmissions may be reduced by discharge planning and outpatient follow-up appointments. The table below is based on readmission for any reason within 30 days of discharge.

	MEDICARE	MEDICAID	PRIVATE INSURANCE	UNINSURED
#1	Congestive HF	Mood disorders	Maintenance of chemotherapy or radiotherapy	Mood disorders
#2	Septicemia	Schizophrenia/psychotic disorders	Mood disorders	Alcohol-related disorders
#3	Pneumonia	Diabetes mellitus with complications	Complications of surgical procedures or medical care	Diabetes mellitus with complications

▶ PUBLIC HEALTH SCIENCES—QUALITY AND SAFETY

Safety culture

Organizational environment in which everyone can freely bring up safety concerns without fear of censure. Facilitates error identification.

Event reporting systems collect data on errors for internal and external monitoring.

Human factors design


Forcing functions (those that prevent undesirable actions [eg, connecting feeding syringe to IV tubing]) are the most effective. Standardization improves process reliability (eg, clinical pathways, guidelines, checklists). Simplification reduces wasteful activities (eg, consolidating electronic medical records).

Deficient designs hinder workflow and lead to staff workarounds that bypass safety features (eg, patient ID barcodes affixed to computers due to unreadable wristbands).

PDSA cycle

Process improvement model to test changes in real clinical setting. Impact on patients:

- **P**lan—define problem and solution
- **D**o—test new process
- **S**tudy—measure and analyze data
- **A**ct—integrate new process into workflow




Quality measurements

	MEASURE	EXAMPLE
Structural	Physical equipment, resources, facilities	Number of diabetes educators
Process	Performance of system as planned	Percentage of diabetic patients whose HbA _{1c} was measured in the past 6 months
Outcome	Impact on patients	Average HbA _{1c} of patients with diabetes
Balancing	Impact on other systems/outcomes	Incidence of hypoglycemia among patients who tried an intervention to lower HbA _{1c}

Swiss cheese model

Focuses on systems and conditions rather than an individual's error. The risk of a threat becoming a reality is mitigated by differing layers and types of defenses. Patient harm can occur despite multiple safeguards when "the holes in the cheese line up."



Types of medical errors May involve patient identification, diagnosis, monitoring, nosocomial infection, medications, procedures, devices, documentation, handoffs. Medical errors should be disclosed to patients, independent of immediate outcome (harmful or not).

Active error	Occurs at level of frontline operator (eg, wrong IV pump dose programmed).	Immediate impact.
Latent error	Occurs in processes indirect from operator but impacts patient care (eg, different types of IV pumps used within same hospital).	Accident waiting to happen.
Never event	Adverse event that is identifiable, serious, and usually preventable (eg, scalpel retained in a surgical patient's abdomen).	Major error that should never occur.

Burnout vs fatigue

Burnout	Prolonged, excessive stress → cynicism, detachment, ↓ motivation and interest, sense of failure and helplessness, ↓ immunity. Medical errors due to lack of concern.
Fatigue	Sleep deprivation → ↓ energy and motivation, cognitive impairment. Medical errors due to compromised intellectual function.

Medical error analysis

	DESIGN	METHODS
Root cause analysis	Retrospective approach. Applied after failure event to prevent recurrence.	Uses records and participant interviews to identify all the underlying problems (eg, process, people, environment, equipment, materials, management) that led to an error.
Failure mode and effects analysis	Forward-looking approach. Applied before process implementation to prevent failure occurrence.	Uses inductive reasoning to identify all the ways a process might fail and prioritizes them by their probability of occurrence and impact on patients.

SECTION III

High-Yield Organ Systems

“Symptoms, then, are in reality nothing but the cry from suffering organs.”
—Jean-Martin Charcot

“Man is an intelligence in servitude to his organs.”
—Aldous Huxley

“When every part of the machine is correctly adjusted and in perfect harmony, health will hold dominion over the human organism by laws as natural and immutable as the laws of gravity.”
—Andrew T. Still

▶ Approaching the Organ Systems	276
▶ Cardiovascular	279
▶ Endocrine	325
▶ Gastrointestinal	357
▶ Hematology and Oncology	403
▶ Musculoskeletal, Skin, and Connective Tissue	445
▶ Neurology and Special Senses	489
▶ Psychiatry	553
▶ Renal	577
▶ Reproductive	611
▶ Respiratory	659

▶ APPROACHING THE ORGAN SYSTEMS

In this section, we have divided the High-Yield Facts into the major **Organ Systems**. Within each Organ System are several subsections, including **Embryology**, **Anatomy**, **Physiology**, **Pathology**, and **Pharmacology**. As you progress through each Organ System, refer back to information in the previous subsections to organize these basic science subsections into a “vertically integrated” framework for learning. Below is some general advice for studying the organ systems by these subsections.

Embryology

Relevant embryology is included in each organ system subsection. Embryology tends to correspond well with the relevant anatomy, especially with regard to congenital malformations.

Anatomy

Several topics fall under this heading, including gross anatomy, histology, and neuroanatomy. Do not memorize all the small details; however, do not ignore anatomy altogether. Review what you have already learned and what you wish you had learned. Many questions require two or more steps. The first step is to identify a structure on anatomic cross section, electron micrograph, or photomicrograph. The second step may require an understanding of the clinical significance of the structure.

When studying, stress clinically important material. For example, be familiar with gross anatomy and radiologic anatomy related to specific diseases (eg, Pancoast tumor, Horner syndrome), traumatic injuries (eg, fractures, sensory and motor nerve deficits), procedures (eg, lumbar puncture), and common surgeries (eg, cholecystectomy). There are also many questions on the exam involving x-rays, CT scans, and neuro MRI scans. Many students suggest browsing through a general radiology atlas, pathology atlas, and histology atlas. Focus on learning basic anatomy at key levels in the body (eg, sagittal brain MRI; axial CT of the midthorax, abdomen, and pelvis). Basic neuroanatomy (especially pathways, blood supply, and functional anatomy), associated neuropathology, and neurophysiology have good yield. Please note that many of the photographic images in this book are for illustrative purposes and are not necessarily reflective of Step 1 emphasis.

Physiology

The portion of the examination dealing with physiology is broad and concept oriented and thus does not lend itself as well to fact-based review. Diagrams are often the best study aids, especially given the increasing number of questions requiring the interpretation of diagrams. Learn to apply basic physiologic relationships in a variety of ways (eg, the Fick equation, clearance equations). You are seldom asked to perform complex

calculations. Hormones are the focus of many questions, so learn their sites of production and action as well as their regulatory mechanisms.

A large portion of the physiology tested on the USMLE Step 1 is clinically relevant and involves understanding physiologic changes associated with pathologic processes (eg, changes in pulmonary function with COPD). Thus, it is worthwhile to review the physiologic changes that are found with common pathologies of the major organ systems (eg, heart, lungs, kidneys, GI tract) and endocrine glands.

Pathology

Questions dealing with this discipline are difficult to prepare for because of the sheer volume of material involved. Review the basic principles and hallmark characteristics of the key diseases. Given the clinical orientation of Step 1, it is no longer sufficient to know only the “buzzword” associations of certain diseases (eg, café-au-lait macules and neurofibromatosis); you must also know the clinical descriptions of these findings.

Given the clinical slant of the USMLE Step 1, it is also important to review the classic presenting signs and symptoms of diseases as well as their associated laboratory findings. Delve into the signs, symptoms, and pathophysiology of major diseases that have a high prevalence in the United States (eg, alcoholism, diabetes, hypertension, heart failure, ischemic heart disease, infectious disease). Be prepared to think one step beyond the simple diagnosis to treatment or complications.

The examination includes a number of color photomicrographs and photographs of gross specimens that are presented in the setting of a brief clinical history. However, read the question and the choices carefully before looking at the illustration, because the history will help you identify the pathologic process. Flip through an illustrated pathology textbook, color atlases, and appropriate Web sites in order to look at the pictures in the days before the exam. Pay attention to potential clues such as age, sex, ethnicity, occupation, recent activities and exposures, and specialized lab tests.

Pharmacology

Preparation for questions on pharmacology is straightforward. Learning all the key drugs and their characteristics (eg, mechanisms, clinical use, and important side effects) is high yield. Focus on understanding the prototype drugs in each class. Avoid memorizing obscure derivatives. Learn the “classic” and distinguishing toxicities of the major drugs. Do not bother with drug dosages or trade names. Reviewing associated biochemistry, physiology, and microbiology can be useful while studying pharmacology. There is a strong emphasis on ANS, CNS, antimicrobial, and cardiovascular agents as well as NSAIDs. Much of the material is clinically relevant. Newer drugs on the market are also fair game.

HIGH-YIELD SYSTEMS

Cardiovascular

“As for me, except for an occasional heart attack, I feel as young as I ever did.”

—Robert Benchley

“Hearts will never be practical until they are made unbreakable.”

—The Wizard of Oz

“As the arteries grow hard, the heart grows soft.”

—H. L. Mencken

“Nobody has ever measured, not even poets, how much the heart can hold.”

—Zelda Fitzgerald

“Only from the heart can you touch the sky.”

—Rumi

“It is not the size of the man but the size of his heart that matters.”

—Evander Holyfield

The cardiovascular system is one of the highest yield areas for the boards and, for some students, may be the most challenging. Focusing on understanding the mechanisms instead of memorizing the details can make a big difference, especially for this topic. Pathophysiology of atherosclerosis and heart failure, MOA of drugs (particular physiology interactions) and their adverse effects, ECGs of heart blocks, the cardiac cycle, and the Starling curve are some of the more high-yield topics. Differentiating between systolic and diastolic dysfunction is also very important. Heart murmurs and maneuvers that affect these murmurs have also been high yield and may be asked in a multimedia format.

▶ Embryology	280
▶ Anatomy	283
▶ Physiology	284
▶ Pathology	298
▶ Pharmacology	316


▶ CARDIOVASCULAR—EMBRYOLOGY

Heart morphogenesis First functional organ in vertebrate embryos; beats spontaneously by week 4 of development.

Cardiac looping


Primary heart tube loops to establish left-right polarity; begins in week 4 of development.

Defect in left-right **Dynein** (involved in L/R asymmetry) can lead to **Dextrocardia**, as seen in Kartagener syndrome (1° ciliary **Dyskinesia**).

**Septation of the chambers****Atria**

- Septum primum grows toward endocardial cushions, narrowing foramen primum.
- Foramen secundum forms in septum primum (foramen primum regresses).
- Septum secundum develops on the right side of septum primum, as foramen secundum maintains right-to-left shunt.
- Septum secundum expands and covers most of foramen secundum. The residual foramen is the foramen ovale.
- Remaining portion of septum primum forms the one-way valve of the foramen ovale.
- Septum primum closes against septum secundum, sealing the foramen ovale soon after birth because of \uparrow LA pressure and \downarrow RA pressure.
- Septum secundum and septum primum fuse during infancy/early childhood, forming the atrial septum.


Patent foramen ovale—caused by failure of septum primum and septum secundum to fuse after birth; most are left untreated. Can lead to paradoxical emboli (venous thromboemboli entering the systemic arterial circulation) as can occur in ASD.



Heart morphogenesis (continued)**Ventricles**

- ① Muscular interventricular septum forms. Opening is called interventricular foramen.
- ② Aorticopulmonary septum rotates and fuses with muscular ventricular septum to form membranous interventricular septum, closing interventricular foramen.
- ③ Growth of endocardial cushions separates atria from ventricles and contributes to both atrial septation and membranous portion of the interventricular septum.

Ventricular septal defect—most common congenital cardiac anomaly, usually occurs in membranous septum.

**Outflow tract formation**

Neural crest and endocardial cell migrations
 → truncal and bulbar ridges that spiral and fuse to form aorticopulmonary septum
 → ascending aorta and pulmonary trunk.

Conotruncal abnormalities associated with failure of neural crest cells to migrate:

- Transposition of great vessels.
- Tetralogy of Fallot.
- Persistent truncus arteriosus.

Valve development


Aortic/pulmonary: derived from endocardial cushions of outflow tract.
 Mitral/tricuspid: derived from fused endocardial cushions of the AV canal.

Valvular anomalies may be stenotic, regurgitant, atretic (eg, tricuspid atresia), or displaced (eg, Ebstein anomaly).

Heart embryology

EMBRYONIC STRUCTURE	GIVES RISE TO
Truncus arteriosus	Ascending aorta and pulmonary trunk
Bulbus cordis	Smooth parts (outflow tract) of left and right ventricles
Primitive ventricle	Trabeculated part of left and right ventricles
Primitive atrium	Trabeculated part of left and right atria
Left horn of sinus venosus	Coronary sinus
Right horn of sinus venosus	Smooth part of right atrium (sinus venarum)
Endocardial cushion	Atrial septum, membranous interventricular septum; AV and semilunar valves
Right common cardinal vein and right anterior cardinal vein	Superior vena cava (SVC)
Posterior, subcardinal, and supracardinal veins	Inferior vena cava (IVC)
Primitive pulmonary vein	Smooth part of left atrium

Fetal circulation



Blood in umbilical vein has a PO_2 of ≈ 30 mm Hg and is $\approx 80\%$ saturated with O_2 . Umbilical arteries have low O_2 saturation.

3 important shunts:

- 1 Blood entering fetus through the umbilical vein is conducted via the **ductus venosus** into the IVC, bypassing hepatic circulation.
- 2 Most of the highly **O**xxygenated blood reaching the heart via the IVC is directed through the **foramen O**vale into the left atrium.
- 3 **D**eoxygenated blood from the SVC passes through the RA \rightarrow RV \rightarrow main pulmonary artery \rightarrow **D**uctus arteriosus \rightarrow **D**escending aorta; shunt is due to high fetal pulmonary artery resistance (due partly to low O_2 tension).

At birth, infant takes a breath \rightarrow \downarrow resistance in pulmonary vasculature \rightarrow \uparrow left atrial pressure vs right atrial pressure \rightarrow foramen ovale closes (now called fossa ovalis); \uparrow in O_2 (from respiration) and \downarrow in prostaglandins (from placental separation) \rightarrow closure of ductus arteriosus.

Indomethacin helps **close** the patent **D**uctus arteriosus \rightarrow ligamentum arteriosum (remnant of ductus arteriosus). Come **In** and **close** the **D**oor.


Prostaglandins **E**₁ and **E**₂ **kEEp** PDA open.

Fetal-postnatal derivatives

FETAL STRUCTURE	POSTNATAL DERIVATIVE	NOTES
D uctus arteriosus	Ligamentum arteriosum	Near the left recurrent laryngeal nerve
D uctus venosus	Ligamentum venosum	
F oramen ovale	F ossa ovalis	
A llantois \rightarrow u rachus	Medial umbilical ligament	Urachus is part of allantoic duct between bladder and umbilicus
U mbilical arteries	Medial umbilical ligaments	
U mbilical vein	Ligamentum teres hepatis (round ligament)	Contained in falciform ligament
N otochord	N ucleus pulposus	

▶ CARDIOVASCULAR—ANATOMY

Anatomy of the heart



LA is the most posterior part of the heart **A**; enlargement of the LA (eg, in mitral stenosis) can lead to compression of the esophagus (dysphagia) and/or the left recurrent laryngeal nerve, a branch of the vagus nerve, causing hoarseness (**Ortner syndrome**).

RV is the most anterior part of the heart and most commonly injured in trauma.

Pericardium

Consists of 3 layers (from outer to inner):

- Fibrous pericardium
- Parietal layer of serous pericardium
- Visceral layer of serous pericardium

Pericardial cavity lies between parietal and visceral layers.

Pericardium innervated by phrenic nerve.

Pericarditis can cause referred pain to the neck, arms, or one or both shoulders (often left).

Coronary blood supply

LAD and its branches supply anterior 2/3 of interventricular septum, anterolateral papillary muscle, and anterior surface of LV. Most commonly occluded.

PDA supplies AV node (dependent on dominance), posterior 1/3 of interventricular septum, posterior 2/3 walls of ventricles, and posteromedial papillary muscle.


RCA supplies SA node (blood supply independent of dominance). Infarct may cause nodal dysfunction (bradycardia or heart block). Right (acute) marginal artery supplies RV.

Dominance:

- Right-dominant circulation (85%) = PDA arises from RCA.
- Left-dominant circulation (8%) = PDA arises from LCX.
- Codominant circulation (7%) = PDA arises from both LCX and RCA.

Coronary blood flow peaks in early diastole.

- Key:
- AMA = Acute marginal artery
 - LAD = Left anterior descending artery
 - LCA = Left coronary artery
 - LCX = Left circumflex artery
 - OMA = Obtuse marginal artery
 - PDA = Posterior descending artery
 - PT = Pulmonary trunk
 - PV = Pulmonary vein
 - RCA = Right coronary artery




► CARDIOVASCULAR—PHYSIOLOGY

Cardiac output variables

Stroke volume	Stroke Volume affected by C ontractility, A fterload, and P reload. ↑ SV with: <ul style="list-style-type: none"> ▪ ↑ Contractility (eg, anxiety, exercise) ▪ ↑ Preload (eg, early pregnancy) ▪ ↓ Afterload 	SV CAP. A failing heart has ↓ SV (systolic and/or diastolic dysfunction).
Contractility	Contractility (and SV) ↑ with: <ul style="list-style-type: none"> ▪ Catecholamine stimulation via β_1 receptor: <ul style="list-style-type: none"> ▪ Activated protein kinase A <ul style="list-style-type: none"> → phospholamban phosphorylation → active Ca^{2+} ATPase → ↑ Ca^{2+} storage in sarcoplasmic reticulum ▪ Activated protein kinase A → Ca^{2+} channel phosphorylation → ↑ Ca^{2+} entry → ↑ Ca^{2+}-induced Ca^{2+} release ▪ ↑ intracellular Ca^{2+} ▪ ↓ extracellular Na^+ (↓ activity of $\text{Na}^+/\text{Ca}^{2+}$ exchanger) ▪ Digitalis (blocks Na^+/K^+ pump → ↑ intracellular Na^+ → ↓ $\text{Na}^+/\text{Ca}^{2+}$ exchanger activity → ↑ intracellular Ca^{2+}) 	Contractility (and SV) ↓ with: <ul style="list-style-type: none"> ▪ β_1-blockade (↓ cAMP) ▪ HF with systolic dysfunction ▪ Acidosis ▪ Hypoxia/hypercapnia (↓ PO_2/↑ PCO_2) ▪ Non-dihydropyridine Ca^{2+} channel blockers
Preload	Preload approximated by ventricular EDV; depends on venous tone and circulating blood volume.	Vasodilators (eg, nitroglycerin) ↓ preload.
Afterload	Afterload approximated by MAP. ↑ wall tension per Laplace's law → ↑ pressure → ↑ afterload. LV compensates for ↑ afterload by thickening (hypertrophy) in order to ↓ wall stress.	A rterial vasodilators (eg, hydralazine) ↓ A fterload. ACE inhibitors and ARBs ↓ both preload and afterload. Chronic hypertension (↑ MAP) → LV hypertrophy.
Myocardial oxygen demand	Myocardial O_2 demand is ↑ by: <ul style="list-style-type: none"> ▪ ↑ Contractility ▪ ↑ Afterload (proportional to arterial pressure) ▪ ↑ heart Rate ▪ ↑ Diameter of ventricle (↑ wall tension) 	Wall tension follows Laplace's law: Wall tension = pressure × radius Wall stress = $\frac{\text{pressure} \times \text{radius}}{2 \times \text{wall thickness}}$

Cardiac output equations

	EQUATION	NOTES
Stroke volume	$SV = EDV - ESV$	
Ejection fraction	$EF = \frac{SV}{EDV} = \frac{EDV - ESV}{EDV}$	EF is an index of ventricular contractility (↓ in systolic HF; usually normal in diastolic HF).
Cardiac output	$CO = SV \times HR$ Fick principle: $CO = \frac{\text{rate of O}_2 \text{ consumption}}{(\text{arterial O}_2 \text{ content} - \text{venous O}_2 \text{ content})}$	In early stages of exercise, CO maintained by ↑ HR and ↑ SV. In later stages, CO maintained by ↑ HR only (SV plateaus). Diastole is shortened with ↑↑ HR (eg, ventricular tachycardia) → ↓ diastolic filling time → ↓ SV → ↓ CO.
Pulse pressure	$PP = SBP - DBP$	PP directly proportional to SV and inversely proportional to arterial compliance. ↑ PP in hyperthyroidism, aortic regurgitation, aortic stiffening (isolated systolic hypertension in elderly), obstructive sleep apnea (↑ sympathetic tone), anemia, exercise (transient). ↓ PP in aortic stenosis, cardiogenic shock, cardiac tamponade, advanced HF.
Mean arterial pressure	$MAP = CO \times TPR$	MAP (at resting HR) = $\frac{2}{3} DBP + \frac{1}{3} SBP = DBP + \frac{1}{3} PP$.

Starling curves

Force of contraction is proportional to end-diastolic length of cardiac muscle fiber (preload).

- ↑ contractility with catecholamines, positive inotropes (eg, digoxin).
- ↓ contractility with loss of functional myocardium (eg, MI), β-blockers (acutely), non-dihydropyridine Ca²⁺ channel blockers, dilated cardiomyopathy.

Resistance, pressure, flow

$$\Delta P = Q \times R$$

Similar to Ohm's law: $\Delta V = I \times R$

Volumetric flow rate (Q) = flow velocity (v) \times cross-sectional area (A)

Resistance

$$= \frac{\text{driving pressure } (\Delta P)}{Q} = \frac{8\eta \text{ (viscosity)} \times \text{length}}{\pi r^4}$$

Total resistance of vessels in series:

$$R_T = R_1 + R_2 + R_3 \dots$$

Total resistance of vessels in parallel:

$$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} \dots$$

Capillaries have highest total cross-sectional area and lowest flow velocity.


Pressure gradient drives flow from high pressure to low pressure.

Arterioles account for most of TPR. Veins provide most of blood storage capacity.

Viscosity depends mostly on hematocrit.

Viscosity \uparrow in hyperproteinemic states (eg, multiple myeloma), polycythemia.

Viscosity \downarrow in anemia.


Cardiac and vascular function curves

Intersection of curves = operating point of heart (ie, venous return and CO are equal, as circulatory system is a closed system).

GRAPH	EFFECT	EXAMPLES
A Inotropy	Changes in contractility \rightarrow altered SV \rightarrow altered CO/VR and RA pressure (RAP)	1 Catecholamines, digoxin, exercise \oplus 2 HF with reduced EF, narcotic overdose, sympathetic inhibition \ominus
B Venous return	Changes in circulating volume \rightarrow altered RAP \rightarrow altered SV \rightarrow change in CO	3 Fluid infusion, sympathetic activity \oplus 4 Acute hemorrhage, spinal anesthesia \ominus
C Total peripheral resistance	Changes in TPR \rightarrow altered CO Change in RAP unpredictable.	5 Vasopressors \oplus 6 Exercise, AV shunt \ominus

Changes often occur in tandem, and may be reinforcing (eg, exercise \uparrow inotropy and \downarrow TPR to maximize CO) or compensatory (eg, HF \downarrow inotropy \rightarrow fluid retention to \uparrow preload to maintain CO).


Pressure-volume loops and cardiac cycle



The black loop represents normal cardiac physiology.

Phases—left ventricle:

- 1 Isovolumetric contraction—period between mitral valve closing and aortic valve opening; period of highest O_2 consumption
- 2 Systolic ejection—period between aortic valve opening and closing
- 3 Isovolumetric relaxation—period between aortic valve closing and mitral valve opening
- 4 Rapid filling—period just after mitral valve opening
- 5 Reduced filling—period just before mitral valve closing



Heart sounds:

S1—mitral and tricuspid valve closure. Loudest at mitral area.

S2—aortic and pulmonary valve closure. Loudest at left upper sternal border.

S3—in early diastole during rapid ventricular filling phase. Best heard at apex with patient in left lateral decubitus position. Associated with ↑ filling pressures (eg, MR, AR, HF, thyrotoxicosis) and more common in dilated ventricles (but can be normal in children, young adults, athletes, and pregnancy).

S4—in late diastole (“atrial kick”). Best heard at apex with patient in left lateral decubitus position. High atrial pressure. Associated with ventricular noncompliance (eg, hypertrophy). Left atrium must push against stiff LV wall. Considered abnormal if palpable.

Jugular venous pulse (JVP):

a wave—atrial contraction. Absent in atrial fibrillation (AF).

c wave—RV contraction (closed tricuspid valve bulging into atrium).


x descent—downward displacement of closed tricuspid valve during rapid ventricular ejection phase. Reduced or absent in tricuspid regurgitation and right HF because pressure gradients are reduced.

v wave—↑ right atrial pressure due to filling (“villing”) against closed tricuspid valve.

y descent—RA emptying into RV. Prominent in constrictive pericarditis, absent in cardiac tamponade.


Pressure-volume loops and valvular disease

Aortic stenosis




- ↑ LV pressure
- ↑ ESV
- No change in EDV
- ↓ SV
- Ventricular hypertrophy → ↓ ventricular compliance → ↑ EDP for given EDV

Mitral regurgitation




- No true isovolumetric phase
- ↓ ESV due to ↓ resistance and ↑ regurgitation into LA during systole
- ↑ EDV due to ↑ LA volume/pressure from regurgitation → ↑ ventricular filling
- ↑ SV

Aortic regurgitation



- No true isovolumetric phase
- ↑ EDV
- ↑ SV

Mitral stenosis




- ↑ LA pressure
- ↓ EDV because of impaired ventricular filling
- ↓ ESV
- ↓ SV

Splitting of S2

Physiologic splitting


Inspiration → drop in intrathoracic pressure
→ ↑ venous return → ↑ RV filling → ↑ RV
stroke volume → ↑ RV ejection time
→ delayed closure of pulmonic valve.
↓ pulmonary impedance (↑ capacity of the
pulmonary circulation) also occurs during
inspiration, which contributes to delayed
closure of pulmonic valve.



E = Expiration
I = Inspiration


Wide splitting

Seen in conditions that delay RV emptying (eg,
pulmonic stenosis, right bundle branch block).
Causes delayed pulmonic sound (especially
on inspiration). An exaggeration of normal
splitting.




Fixed splitting

Heard in ASD. ASD → left-to-right shunt
→ ↑ RA and RV volumes → ↑ flow through
pulmonic valve → delayed pulmonic valve
closure (independent of respiration).



Paradoxical splitting

Heard in conditions that delay aortic valve
closure (eg, aortic stenosis, left bundle branch
block). Normal order of semilunar valve
closure is reversed so that P2 sound occurs
before delayed A2 sound. On inspiration,
P2 closes later and moves closer to A2,
“paradoxically” eliminating the split. On
expiration, the split can be heard (opposite to
physiologic splitting).



Auscultation of the heart

Where to listen: **APT M**


Aortic area:

- Systolic murmur**
 Aortic stenosis
 Flow murmur (eg, physiologic murmur)
 Aortic valve sclerosis

Left sternal border:

- Diastolic murmur
 Aortic regurgitation (valvular)
 Pulmonic regurgitation
Systolic murmur
 Hypertrophic cardiomyopathy

- Aortic
- Pulmonic
- Tricuspid
- Mitral



Pulmonic area:

- Systolic ejection murmur**
 Pulmonic stenosis
 Atrial septal defect
 Flow murmur

Tricuspid area:

- Holosystolic murmur**
 Tricuspid regurgitation
 Ventricular septal defect
Diastolic murmur
 Tricuspid stenosis

Mitral area (apex):


- Holosystolic murmur**
 Mitral regurgitation
Systolic murmur
 Mitral valve prolapse
Diastolic murmur
 Mitral stenosis

MANEUVER	CARDIOVASCULAR CHANGES	MURMURS THAT INCREASE WITH MANEUVER	MURMURS THAT DECREASE WITH MANEUVER
Standing Valsalva (strain phase)	↓ preload (↓ LV volume)	MVP (↓ LV volume) HCM (↓ LV volume)	Most murmurs (↓ flow through stenotic or regurgitant valve)
Passive leg raise	↑ preload (↑ LV volume)		
Squatting	↑ preload, ↑ afterload (↑ LV volume)	Most murmurs (↑ flow through stenotic or regurgitant valve)	MVP (↑ LV volume) HCM (↑ LV volume)
Hand grip	↑↑ afterload → ↑ reverse flow across aortic valve (↑ LV volume)	Most other left-sided murmurs (AR, MR, VSD)	AS (↓ transaortic valve pressure gradient) HCM (↑ LV volume)
Inspiration	↑ venous return to right heart, ↓ venous return to left heart	Most right-sided murmurs	Most left-sided murmurs

Heart murmurs


Systolic

Aortic stenosis




Crescendo-decrescendo systolic ejection murmur and soft S2 (ejection click may be present). LV \gg aortic pressure during systole. Loudest at heart base; radiates to carotids. “Pulsus parvus et tardus”—pulses are weak with a delayed peak. Can lead to **S**yncope, **A**ngina, and **D**yspnea on exertion (**SAD**). Most commonly due to age-related calcification in older patients (> 60 years old) or in younger patients with early-onset calcification of bicuspid aortic valve.

Mitral/tricuspid regurgitation




Holosystolic, high-pitched “blowing murmur.”
Mitral—loudest at apex and radiates toward axilla. MR is often due to ischemic heart disease (post-MI), MVP, LV dilatation.
Tricuspid—loudest at tricuspid area. TR commonly caused by RV dilatation. Rheumatic fever and infective endocarditis can cause either MR or TR.

Mitral valve prolapse



Late systolic crescendo murmur with midsystolic click (MC) due to sudden tensing of chordae tendineae as mitral leaflets prolapse into the LA (**C**hordae cause **C**rescendo with **C**lick). Most frequent valvular lesion. Best heard over apex. Loudest just before S2. Usually benign. Can predispose to infective endocarditis. Can be caused by myxomatous degeneration (1° or 2° to connective tissue disease such as Marfan or Ehlers-Danlos syndrome), rheumatic fever (particularly in developing countries), chordae rupture.


Ventricular septal defect



Holosystolic, harsh-sounding murmur. Loudest at tricuspid area. Larger VSDs have a lower intensity murmur than smaller VSDs.


Diastolic

Aortic regurgitation



High-pitched “blowing” early diastolic decrescendo murmur. Best heard at base (aortic root dilation) or left sternal border (valvular disease). Long diastolic murmur, hyperdynamic pulse, and head bobbing when severe and chronic. Wide pulse pressure. Causes include **B**icuspid aortic valve, **E**ndocarditis, **A**ortic root dilation, **R**heumatic fever (**BEAR**). Progresses to left HF.


Mitral stenosis



Follows opening snap (OS; due to abrupt halt in leaflet motion in diastole, after rapid opening due to fusion at leaflet tips). Delayed rumbling mid-to-late diastolic murmur (\downarrow interval between S2 and OS correlates with \uparrow severity). LA \gg LV pressure during diastole.
Often a late (and highly specific) sequela of rheumatic fever. Chronic MS can result in pulmonary congestion/hypertension and LA dilation \rightarrow atrial fibrillation and Ortner syndrome.

Continuous

Patent ductus arteriosus



Continuous machine-like murmur. Best heard at left infraclavicular area. Loudest at S2. Often due to congenital rubella or prematurity.
“**P**DAs (**P**ublic **D**isplays of **A**ffection) are **c**ontinuously annoying”

Myocardial action potential

Phase 0 = rapid upstroke and depolarization—voltage-gated Na^+ channels open.

Phase 1 = initial repolarization—inactivation of voltage-gated Na^+ channels. Voltage-gated K^+ channels begin to open.


Phase 2 = plateau— Ca^{2+} influx through voltage-gated Ca^{2+} channels balances K^+ efflux. Ca^{2+} influx triggers Ca^{2+} release from sarcoplasmic reticulum and myocyte contraction.

Phase 3 = rapid repolarization—massive K^+ efflux due to opening of voltage-gated slow delayed-rectifier K^+ channels and closure of voltage-gated Ca^{2+} channels.

Phase 4 = resting potential—high K^+ permeability through K^+ channels.

In contrast to skeletal muscle:

- Cardiac muscle action potential has a plateau due to Ca^{2+} influx and K^+ efflux.
- Cardiac muscle contraction requires Ca^{2+} influx from ECF to induce Ca^{2+} release from sarcoplasmic reticulum (Ca^{2+} -induced Ca^{2+} release).
- Cardiac myocytes are electrically coupled to each other by gap junctions.



Occurs in all cardiac myocytes except for those in the SA and AV nodes.


Pacemaker action potential

Occurs in the SA and AV nodes. Key differences from the ventricular action potential include:

Phase 0 = upstroke—opening of voltage-gated Ca^{2+} channels. Fast voltage-gated Na^+ channels are permanently inactivated because of the less negative resting potential of these cells. Results in a slow conduction velocity that is used by the AV node to prolong transmission from the atria to ventricles. Phases 1 and 2 are absent.

Phase 3 = repolarization—inactivation of the Ca^{2+} channels and \uparrow activation of K^+ channels $\rightarrow \uparrow \text{K}^+$ efflux.



Phase 4 = slow spontaneous diastolic depolarization due to I_{f} (“funny current”). I_{f} channels responsible for a slow, mixed Na^+/K^+ inward current; different from I_{Na} in phase 0 of ventricular action potential. Accounts for automaticity of SA and AV nodes. The slope of phase 4 in the SA node determines HR. ACh/adenosine \downarrow the rate of diastolic depolarization and \downarrow HR, while catecholamines \uparrow depolarization and \uparrow HR. Sympathetic stimulation \uparrow the chance that I_{f} channels are open and thus \uparrow HR.




Electrocardiogram

Conduction pathway: SA node → atria
 → AV node → bundle of His → right and left bundle branches → Purkinje fibers
 → ventricles; left bundle branch divides into left anterior and posterior fascicles.
 SA node—located at junction of RA and SVC; “pacemaker” inherent dominance with slow phase of upstroke.
 AV node—located in posteroinferior part of interatrial septum. Blood supply usually from RCA. 100-msec delay allows time for ventricular filling.
 Pacemaker rates: SA > AV > bundle of His/Purkinje/ventricles.
 Speed of conduction: **H**is-Purkinje > **A**tria > **V**entricles > **A**V node. **H**e **P**arks **A**t **V**entura **A**venue.

P wave—atrial depolarization.
 PR interval—time from start of atrial depolarization to start of ventricular depolarization (normally 120-200 msec).
 QRS complex—ventricular depolarization (normally < 100 msec).
 QT interval—ventricular depolarization, mechanical contraction of the ventricles, ventricular repolarization.
 T wave—ventricular repolarization. T-wave inversion may indicate ischemia or recent MI.
 J point—junction between end of QRS complex and start of ST segment.
 ST segment—isolectric, ventricles depolarized.
 U wave—prominent in hypokalemia (think hyp“U”kalemia), bradycardia.



Torsades de pointes

Polymorphic ventricular tachycardia, characterized by shifting sinusoidal waveforms on ECG; can progress to ventricular fibrillation (VF). Long QT interval predisposes to torsades de pointes. Caused by drugs, ↓ K⁺, ↓ Mg²⁺, ↓ Ca²⁺, congenital abnormalities. Treatment includes magnesium sulfate.

Drug-induced long QT (**ABCDE**):

- Anti**A**rrhythmics (class IA, III)
- Anti**B**iotics (eg, macrolides)
- Anti**C**hotics (eg, haloperidol)
- Anti**D**epressants (eg, TCAs)
- Anti**E**metics (eg, ondansetron)

Torsades de pointes = twisting of the points

Congenital long QT syndrome

Inherited disorder of myocardial repolarization, typically due to ion channel defects (most commonly loss-of-function mutations affecting K⁺ channels); ↑ risk of sudden cardiac death (SCD) due to torsades de pointes. Includes:


- **Romano-Ward syndrome**—autosomal dominant, pure cardiac phenotype (**no** deafness).
- **Jervell and Lange-Nielsen syndrome**—autosomal recessive, sensorineural deafness.

Brugada syndrome








Autosomal dominant disorder most common in Asian males. ECG pattern of pseudo-right bundle branch block and ST elevations in V₁-V₃. ↑ risk of ventricular tachyarrhythmias and SCD. Prevent SCD with implantable cardioverter-defibrillator (ICD).

Wolff-Parkinson-White syndrome

Most common type of ventricular pre-excitation syndrome. Abnormal fast accessory conduction pathway from atria to ventricle (bundle of Kent) bypasses the rate-slowing AV node → ventricles begin to partially depolarize earlier → characteristic delta wave with widened QRS complex and shortened PR interval on ECG. May result in reentry circuit → supraventricular tachycardia.



ECG tracings


RHYTHM	DESCRIPTION	EXAMPLE
Atrial fibrillation	Chaotic and erratic baseline with no discrete P waves in between irregularly spaced QRS complexes. Irregularly irregular heartbeat. Most common risk factors include hypertension and coronary artery disease (CAD). Occasionally seen after binge drinking (“holiday heart syndrome”). Can lead to thromboembolic events, particularly stroke. Treatment: anticoagulation, rate and rhythm control and/or cardioversion.	
Atrial flutter	A rapid succession of identical, back-to-back atrial depolarization waves. The identical appearance accounts for the “sawtooth” appearance of the flutter waves. Treat like atrial fibrillation +/- catheter ablation.	
Ventricular fibrillation	A completely erratic rhythm with no identifiable waves. Fatal arrhythmia without immediate CPR and defibrillation.	
AV block		
First-degree AV block	The PR interval is prolonged (> 200 msec). Benign and asymptomatic. No treatment required.	
Second-degree AV block		
Mobitz type I (Wenckebach)	Progressive lengthening of PR interval until a beat is “dropped” (a P wave not followed by a QRS complex). Usually asymptomatic. Variable RR interval with a pattern (regularly irregular).	
Mobitz type II	Dropped beats that are not preceded by a change in the length of the PR interval (as in type I). May progress to 3rd-degree block. Often treated with pacemaker.	
Third-degree (complete) AV block	The atria and ventricles beat independently of each other. P waves and QRS complexes not rhythmically associated. Atrial rate > ventricular rate. Usually treated with pacemaker. Can be caused by Lyme disease.	

Atrial natriuretic peptide

Released from **atrial myocytes** in response to \uparrow blood volume and atrial pressure. Acts via cGMP. Causes vasodilation and \downarrow Na^+ reabsorption at the renal collecting tubule. Dilates afferent renal arterioles and constricts efferent arterioles, promoting diuresis and contributing to “aldosterone escape” mechanism.

B-type (brain) natriuretic peptide

Released from **ventricular myocytes** in response to \uparrow tension. Similar physiologic action to ANP, with longer half-life. BNP blood test used for diagnosing HF (very good negative predictive value). Available in recombinant form (nesiritide) for treatment of HF.

Baroreceptors and chemoreceptors**Receptors:**

- Aortic arch transmits via vagus nerve to solitary nucleus of medulla (responds to changes in BP).
- Carotid sinus (dilated region at carotid bifurcation) transmits via glossopharyngeal nerve to solitary nucleus of medulla (responds to changes in BP).

Baroreceptors:


- Hypotension— \downarrow arterial pressure \rightarrow \downarrow stretch \rightarrow \downarrow afferent baroreceptor firing \rightarrow \uparrow efferent sympathetic firing and \downarrow efferent parasympathetic stimulation \rightarrow vasoconstriction, \uparrow HR, \uparrow contractility, \uparrow BP. Important in the response to severe hemorrhage.
- Carotid massage— \uparrow pressure on carotid sinus \rightarrow \uparrow stretch \rightarrow \uparrow afferent baroreceptor firing \rightarrow \uparrow AV node refractory period \rightarrow \downarrow HR.
- Component of Cushing reflex (triad of hypertension, bradycardia, and respiratory depression)— \uparrow intracranial pressure constricts arterioles \rightarrow cerebral ischemia \rightarrow \uparrow pCO_2 and \downarrow pH \rightarrow central reflex sympathetic \uparrow in perfusion pressure (hypertension) \rightarrow \uparrow stretch \rightarrow peripheral reflex baroreceptor-induced bradycardia.

Chemoreceptors:

- Peripheral—carotid and aortic bodies are stimulated by \uparrow PCO_2 , \downarrow pH of blood, and \downarrow PO_2 (< 60 mm Hg).
- Central—are stimulated by changes in pH and PCO_2 of brain interstitial fluid, which in turn are influenced by arterial CO_2 as H^+ cannot cross the blood-brain barrier. Do not directly respond to PO_2 . Central chemoreceptors become less responsive with chronically \uparrow PCO_2 (eg, COPD) \rightarrow \uparrow dependence on peripheral chemoreceptors to detect \downarrow O_2 to drive respiration.

Normal cardiac pressures

Pulmonary capillary wedge pressure (PCWP; in mm Hg) is a good approximation of left atrial pressure. In mitral stenosis, PCWP > LV end diastolic pressure. PCWP is measured with pulmonary artery catheter (Swan-Ganz catheter).



Autoregulation

How blood flow to an organ remains constant over a wide range of perfusion pressures.

ORGAN	FACTORS DETERMINING AUTOREGULATION	
Heart	Local metabolites (vasodilatory): adenosine, NO, CO ₂ , ↓ O ₂	The pulmonary vasculature is unique in that alveolar hypoxia causes vasoconstriction so that only well-ventilated areas are perfused. In other organs, hypoxia causes vasodilation.
Brain	Local metabolites (vasodilatory): CO ₂ (pH)	
Kidneys	Myogenic and tubuloglomerular feedback	
Lungs	Hypoxia causes vasoconstriction	
Skeletal muscle	Local metabolites during exercise (vasodilatory): CHALK CO ₂ , H ⁺ , Adenosine, Lactate, K ⁺ At rest: sympathetic tone in arteries	
Skin	Sympathetic vasoconstriction most important mechanism for temperature control	

Capillary fluid exchange

Starling forces determine fluid movement through capillary membranes:

- P_c = capillary hydrostatic pressure—pushes fluid out of capillary
- P_i = interstitial hydrostatic pressure—pushes fluid into capillary
- π_c = plasma colloid osmotic (oncotic) pressure—pulls fluid into capillary
- π_i = interstitial fluid colloid osmotic pressure—pulls fluid out of capillary


$J_v = \text{net fluid flow} = K_f [(P_c - P_i) - \sigma(\pi_c - \pi_i)]$

K_f = capillary permeability to fluid

σ = reflection coefficient (measure of capillary permeability to protein)

Edema—excess fluid outflow into interstitium commonly caused by:

- ↑ capillary pressure (↑ P_c; eg, HF)
- ↑ capillary permeability (↑ K_f; eg, toxins, infections, burns)
- ↑ interstitial fluid colloid osmotic pressure (↑ π_i; eg, lymphatic blockage)
- ↓ plasma proteins (↓ π_c; eg, nephrotic syndrome, liver failure, protein malnutrition)



► CARDIOVASCULAR—PATHOLOGY

Congenital heart diseases

RIGHT-TO-LEFT SHUNTS

Early cyanosis—“blue babies.” Often diagnosed prenatally or become evident immediately after birth. Usually require urgent surgical treatment and/or maintenance of a PDA.

The **5 T's**:

1. **T**runcus arteriosus (**1** vessel)
2. **T**ransposition (**2** switched vessels)
3. **T**ricuspid atresia (**3** = **Tri**)
4. **T**etralogy of Fallot (**4** = **Tetra**)
5. **TAPVR** (**5** letters in the name)


Persistent truncus arteriosus

Truncus arteriosus fails to divide into pulmonary trunk and aorta due to failure of aorticopulmonary septum formation; most patients have accompanying VSD.


D-transposition of great vessels

Aorta leaves RV (anterior) and pulmonary trunk leaves LV (posterior) → separation of systemic and pulmonary circulations. Not compatible with life unless a shunt is present to allow mixing of blood (eg, VSD, PDA, or patent foramen ovale).

Due to failure of the aorticopulmonary septum to spiral (“egg on a string” appearance on CXR). Without surgical intervention, most infants die within the first few months of life.

**Tricuspid atresia**

Absence of tricuspid valve and hypoplastic RV; requires both ASD and VSD for viability.

Tetralogy of Fallot

Caused by anterosuperior displacement of the infundibular septum. Most common cause of early childhood cyanosis.


- 1 **P**ulmonary infundibular stenosis (most important determinant for prognosis)
- 2 **R**ight ventricular hypertrophy (RVH)—boot-shaped heart on CXR **A**
- 3 **O**verriding aorta
- 4 **V**SD

Pulmonary stenosis forces right-to-left flow across VSD → RVH, “tet spells” (often caused by crying, fever, and exercise due to exacerbation of RV outflow obstruction).

PROVE.

Squatting: ↑ SVR, ↓ right-to-left shunt, improves cyanosis.

Associated with 22q11 syndromes.

**Total anomalous pulmonary venous return**

Pulmonary veins drain into right heart circulation (SVC, coronary sinus, etc); associated with ASD and sometimes PDA to allow for right-to-left shunting to maintain CO.

Ebstein anomaly

Displacement of tricuspid valve leaflets downward into RV, artificially “atrializing” the ventricle. Associated with tricuspid regurgitation, accessory conduction pathways, right-sided HF.

Can be caused by lithium exposure in utero.

Congenital heart diseases (continued)**LEFT-TO-RIGHT SHUNTS**

Acyanotic at presentation; cyanosis may occur years later. Frequency: VSD > ASD > PDA.


Right-to-Left shunts: **eaRLy** cyanosis.

Left-to-Right shunts: “**LateR**” cyanosis.

Ventricular septal defect


Asymptomatic at birth, may manifest weeks later or remain asymptomatic throughout life. Most self resolve; larger lesions **B** may lead to LV overload and HF.

O₂ saturation ↑ in RV and pulmonary artery.

**Atrial septal defect**

Defect in interatrial septum **C**; wide, fixed split S2. Ostium secundum defects most common and usually an isolated finding; ostium primum defects rarer and usually occur with other cardiac anomalies. Symptoms range from none to HF. Distinct from patent foramen ovale in that septa are missing tissue rather than unfused.


O₂ saturation ↑ in RA, RV, and pulmonary artery. May lead to paradoxical emboli (systemic venous emboli use ASD to bypass lungs and become systemic arterial emboli). Associated with Down syndrome.

**Patent ductus arteriosus**


In fetal period, shunt is right to left (normal). In neonatal period, ↓ pulmonary vascular resistance → shunt becomes left to right → progressive RVH and/or LVH and HF.

PDA is normal in utero and normally closes only after birth.


Associated with a continuous, “machine-like” murmur. Patency is maintained by PGE synthesis and low O₂ tension. Uncorrected PDA **D** can eventually result in late cyanosis in the lower extremities (differential cyanosis).

**Eisenmenger syndrome**

Uncorrected left-to-right shunt (VSD, ASD, PDA) → ↑ pulmonary blood flow → pathologic remodeling of vasculature → pulmonary arterial hypertension. RVH occurs to compensate → shunt becomes right to left. Causes late cyanosis, clubbing **E**, and polycythemia. Age of onset varies.

**OTHER ANOMALIES****Coarctation of the aorta**

Aortic narrowing **F** near insertion of ductus arteriosus (“juxtaductal”). Associated with bicuspid aortic valve, other heart defects, and Turner syndrome. Hypertension in upper extremities and weak, delayed pulse in lower extremities (brachial-femoral delay). With age, intercostal arteries enlarge due to collateral circulation; arteries erode ribs → notched appearance on CXR. Complications include HF, ↑ risk of cerebral hemorrhage (berry aneurysms), aortic rupture, and possible endocarditis.



Congenital cardiac defect associations


DISORDER	DEFECT
Alcohol exposure in utero (fetal alcohol syndrome)	VSD, PDA, ASD, tetralogy of Fallot
Congenital rubella	PDA, pulmonary artery stenosis, septal defects
Down syndrome	AV septal defect (endocardial cushion defect), VSD, ASD
Infant of diabetic mother	Transposition of great vessels, VSD
Marfan syndrome	MVP, thoracic aortic aneurysm and dissection, aortic regurgitation
Prenatal lithium exposure	Ebstein anomaly
Turner syndrome	Bicuspid aortic valve, coarctation of aorta
Williams syndrome	Supravalvular aortic stenosis
22q11 syndromes	Truncus arteriosus, tetralogy of Fallot

Hypertension

Persistent systolic BP \geq 130 mm Hg and/or diastolic BP \geq 80 mm Hg.

RISK FACTORS

\uparrow age, obesity, diabetes, physical inactivity, excess salt intake, excess alcohol intake, cigarette smoking, family history; African American > Caucasian > Asian.

FEATURES

90% of hypertension is 1° (essential) and related to \uparrow CO or \uparrow TPR. Remaining 10% mostly 2° to renal/renovascular diseases such as fibromuscular dysplasia (characteristic “string of beads” appearance of renal artery **A**, usually seen in women of child-bearing age) and atherosclerotic renal artery stenosis or to 1° hyperaldosteronism.

Hypertensive urgency—severe (\geq 180/ \geq 120 mm Hg) hypertension without acute end-organ damage.


Hypertensive emergency—severe hypertension with evidence of acute end-organ damage (eg, encephalopathy, stroke, retinal hemorrhages and exudates, papilledema, MI, HF, aortic dissection, kidney injury, microangiopathic hemolytic anemia, eclampsia).

PREDISPOSES TO

CAD, LVH, HF, atrial fibrillation; aortic dissection, aortic aneurysm; stroke; CKD (hypertensive nephropathy); retinopathy.

Hyperlipidemia signs

Xanthomas	Plaques or nodules composed of lipid-laden histiocytes in skin A , especially the eyelids (xanthelasma B).
Tendinous xanthoma	Lipid deposit in tendon C , especially Achilles.
Corneal arcus	Lipid deposit in cornea. Common in elderly (arcus senilis D), but appears earlier in life with hypercholesterolemia.



Arteriosclerosis


Hardening of arteries, with arterial wall thickening and loss of elasticity.

Arteriolosclerosis

Common. Affects small arteries and arterioles. Two types: hyaline (thickening of vessel walls 2° to plasma protein leak into endothelium in essential hypertension or diabetes mellitus **A**) and hyperplastic (“onion skinning” in severe hypertension **B** with proliferation of smooth muscle cells).

Mönckeberg sclerosis (Medial calcific sclerosis)

Uncommon. Affects **M**edium-sized arteries. Calcification of internal elastic lamina and media of arteries → vascular stiffening without obstruction. “Pipestem” appearance on x-ray **C**. Does not obstruct blood flow; intima not involved.



Atherosclerosis

Very common. Disease of elastic arteries and large- and medium-sized muscular arteries; a form of arteriosclerosis caused by buildup of cholesterol plaques in intima.

LOCATION

Abdominal aorta > Coronary artery > Popliteal artery > Carotid artery > circle of Willis.
A CoPy Cat named Willis.

RISK FACTORS


Modifiable: smoking, hypertension, dyslipidemia (↑ LDL, ↓ HDL), diabetes.
Non-modifiable: age, sex (↑ in men and postmenopausal women), family history.

SYMPTOMS


Angina, claudication, but can be asymptomatic.

PROGRESSION

Inflammation important in pathogenesis: endothelial cell dysfunction → macrophage and LDL accumulation → foam cell formation → fatty streaks → smooth muscle cell migration (involves PDGF and FGF), proliferation, and extracellular matrix deposition → fibrous plaque → complex atheromas **A** → calcification (calcium content correlates with risk of complications).

**COMPLICATIONS**


Aneurysms, ischemia, infarcts, peripheral vascular disease, thrombus, emboli.

**Aortic aneurysm**

Localized pathologic dilation of the aorta. May cause abdominal and/or back pain, which is a sign of leaking, dissection, or imminent rupture.

Abdominal aortic aneurysm

Usually associated with atherosclerosis. Risk factors include history of tobacco use, ↑ age, male sex, family history. May present as palpable pulsatile abdominal mass (arrows in **A** point to outer dilated calcified aortic wall, with partial crescent-shaped non-opacification of aorta due to flap/clot). Most often infrarenal (distal to origin of renal arteries).


**Thoracic aortic aneurysm**

Associated with cystic medial degeneration. Risk factors include hypertension, bicuspid aortic valve, connective tissue disease (eg, Marfan syndrome). Also associated with 3° syphilis (obliterative endarteritis of the vasa vasorum). Aortic root dilatation may lead to aortic valve regurgitation.

Traumatic aortic rupture


Due to trauma and/or deceleration injury, most commonly at aortic isthmus (proximal descending aorta just distal to origin of left subclavian artery). X-ray may reveal widened mediastinum.

Aortic dissection



Longitudinal intimal tear forming a false lumen. Associated with hypertension, bicuspid aortic valve, inherited connective tissue disorders (eg, Marfan syndrome). Can present with tearing, sudden-onset chest pain radiating to the back +/- markedly unequal BP in arms. CXR can show mediastinal widening. Can result in organ ischemia, aortic rupture, death. Two types:

- Stanford type **A** (proximal): involves Ascending aorta (red arrow in **A**). May extend to aortic arch or descending aorta (blue arrow in **A**). May result in acute aortic regurgitation or cardiac tamponade. Treatment: surgery.
- Stanford type **B** (distal): involves only descending aorta (Below left subclavian artery). Treatment: β -blockers, then vasodilators.



Ischemic heart disease manifestations

Angina Chest pain due to ischemic myocardium 2° to coronary artery narrowing or spasm; no myocyte necrosis.

- **Stable**—usually 2° to atherosclerosis (≥ 70% occlusion); exertional chest pain in classic distribution (usually with ST depression on ECG), resolving with rest or nitroglycerin.
- **Vasospastic** (also called **Prinzmetal** or **Variant**)—occurs at rest 2° to coronary artery spasm; transient ST elevation on ECG. Smoking is a risk factor; hypertension and hypercholesterolemia are not. Triggers include cocaine, alcohol, and triptans. Treat with Ca²⁺ channel blockers, nitrates, and smoking cessation (if applicable).
- **Unstable**—thrombosis with incomplete coronary artery occlusion; +/- ST depression and/or T-wave inversion on ECG but no cardiac biomarker elevation (unlike NSTEMI); ↑ in frequency or intensity of chest pain or any chest pain at rest.

Coronary steal syndrome Distal to coronary stenosis, vessels are maximally dilated at baseline. Administration of vasodilators (eg, dipyridamole, regadenoson) dilates normal vessels → blood is shunted toward well-perfused areas → ischemia in myocardium perfused by stenosed vessels. Principle behind pharmacologic stress tests with coronary vasodilators.



Sudden cardiac death Death from cardiac causes within 1 hour of onset of symptoms, most commonly due to a lethal arrhythmia (eg, VF). Associated with CAD (up to 70% of cases), cardiomyopathy (hypertrophic, dilated), and hereditary ion channelopathies (eg, long QT syndrome, Brugada syndrome). Prevent with ICD.

Chronic ischemic heart disease Progressive onset of HF over many years due to chronic ischemic myocardial damage.

Myocardial infarction Most often due to rupture of coronary artery atherosclerotic plaque → acute thrombosis. ↑ cardiac biomarkers (CK-MB, troponins) are diagnostic.

Non-ST-segment elevation MI (NSTEMI)
 Subendocardial infarcts
 Subendocardium (inner 1/3) especially vulnerable to ischemia
 ST depression on ECG









ST-segment elevation MI (STEMI)
 Transmural infarcts
 Full thickness of myocardial wall involved
 ST elevation, pathologic Q waves on ECG



Evolution of myocardial infarction

Commonly occluded coronary arteries: LAD > RCA > circumflex.

Symptoms: diaphoresis, nausea, vomiting, severe retrosternal pain, pain in left arm and/or jaw, shortness of breath, fatigue.

TIME	GROSS	LIGHT MICROSCOPE	COMPLICATIONS
0–24 hr	<p>Dark mottling</p>  <p>Occluded artery Infarct Dark mottling; pale with tetrazolium stain</p>	<p>Early coagulative necrosis → cell content released into blood; edema, hemorrhage, wavy fibers</p> <p>Reperfusion injury → free radicals and ↑ Ca²⁺ influx → hypercontraction of myofibrils (dark eosinophilic stripes)</p> 	Ventricular arrhythmia, HF, cardiogenic shock
1–3 days	 <p>Hyperemia</p>	<p>Extensive coagulative necrosis</p> <p>Tissue surrounding infarct shows acute inflammation with neutrophils</p> 	Postinfarction fibrinous pericarditis
3–14 days	 <p>Hyperemic border; central yellow-brown softening</p>	<p>Macrophages, then granulation tissue at margins</p> 	Free wall rupture → tamponade; papillary muscle rupture → mitral regurgitation; interventricular septal rupture due to macrophage-mediated structural degradation → left-to-right shunt LV pseudoaneurysm (risk of rupture)
2 weeks to several months	 <p>Recanalized artery Gray-white scar</p>	<p>Contracted scar complete</p> 	Dressler syndrome, HF, arrhythmias, true ventricular aneurysm (risk of mural thrombus)


Diagnosis of myocardial infarction

In the first 6 hours, ECG is the gold standard. Cardiac troponin I rises after 4 hours (peaks at 24 hr) and is ↑ for 7–10 days; more specific than other protein markers.

CK-MB rises after 6–12 hours (peaks at 16–24 hr) and is predominantly found in myocardium but can also be released from skeletal muscle. Useful in diagnosing reinfarction following acute MI because levels return to normal after 48 hours.


Large MIs lead to greater elevations in troponin I and CK-MB. Exact curves vary with testing procedure.

ECG changes can include ST elevation (STEMI, transmural infarct), ST depression (NSTEMI, subendocardial infarct), hyperacute (peaked) T waves, T-wave inversion, new left bundle branch block, and pathologic Q waves or poor R wave progression (evolving or old transmural infarct).




ECG localization of STEMI

INFARCT LOCATION	LEADS WITH ST-SEGMENT ELEVATIONS OR Q WAVES
Anteroseptal (LAD)	V ₁ –V ₂
Anterolateral (distal LAD)	V ₃ –V ₄
Anterolateral (LAD or LCX)	V ₅ –V ₆
Lateral (LCX)	I, aVL
Inferior (RCA)	II, III, aVF
Posterior (PDA)	V ₇ –V ₉ , ST depression in V ₁ –V ₃ with tall R waves



Myocardial infarction complications

Cardiac arrhythmia	Occurs within the first few days after MI. Important cause of death before reaching the hospital and within the first 24 hours post-MI.
Postinfarction fibrinous pericarditis	1–3 days: friction rub.
Papillary muscle rupture	2–7 days: posteromedial papillary muscle rupture A ↑ risk due to single blood supply from posterior descending artery. Can result in severe mitral regurgitation.
Interventricular septal rupture	3–5 days: macrophage-mediated degradation → VSD → ↑ O ₂ saturation and pressure in RV.
Ventricular pseudoaneurysm formation	3–14 days: free wall rupture contained by adherent pericardium or scar tissue B ; ↓ CO, risk of arrhythmia, embolus from mural thrombus.
Ventricular free wall rupture	5–14 days: free wall rupture C → cardiac tamponade. LV hypertrophy and previous MI protect against free wall rupture. Acute form usually leads to sudden death.
True ventricular aneurysm	2 weeks to several months: outward bulge with contraction (“dyskinesia”), associated with fibrosis.
Dressler syndrome	Several weeks: autoimmune phenomenon resulting in fibrinous pericarditis.
LV failure and pulmonary edema	Can occur 2° to LV infarction, VSD, free wall rupture, papillary muscle rupture with mitral regurgitation.


**Acute coronary syndrome treatments**

Unstable angina/NSTEMI—Anticoagulation (eg, heparin), antiplatelet therapy (eg, aspirin) + ADP receptor inhibitors (eg, clopidogrel), β -blockers, ACE inhibitors, statins. Symptom control with nitroglycerin and morphine.

STEMI—In addition to above, reperfusion therapy most important (percutaneous coronary intervention preferred over fibrinolysis).

Cardiomyopathies

Dilated cardiomyopathy



Most common cardiomyopathy (90% of cases).

Often idiopathic or familial (eg, due to mutation of *TTN* gene encoding the sarcomeric protein titin).

Other etiologies include drugs (eg, alcohol, cocaine, doxorubicin), infection (eg, coxsackie B virus, Chagas disease), ischemia (eg, CAD), systemic conditions (eg, hemochromatosis, sarcoidosis, thyrotoxicosis, wet beriberi), peripartum cardiomyopathy.

Findings: HF, S3, systolic regurgitant murmur, dilated heart on echocardiogram, balloon appearance of heart on CXR.


Treatment: Na⁺ restriction, ACE inhibitors, β-blockers, diuretics, mineralocorticoid receptor blockers (eg, spironolactone), digoxin, ICD, heart transplant.

Leads to systolic dysfunction.

Dilated cardiomyopathy **A** displays eccentric hypertrophy (sarcomeres added in series).

Takotsubo cardiomyopathy: broken heart syndrome—ventricular apical ballooning likely due to increased sympathetic stimulation (eg, stressful situations).

Hypertrophic obstructive cardiomyopathy



60–70% of cases are familial, autosomal dominant (most commonly due to mutations in genes encoding sarcomeric proteins, such as myosin binding protein C and β-myosin heavy chain). Causes syncope during exercise and may lead to sudden death (eg, in young athletes) due to ventricular arrhythmia.

Findings: S4, systolic murmur. May see mitral regurgitation due to impaired mitral valve closure.

Treatment: cessation of high-intensity athletics, use of β-blocker or non-dihydropyridine Ca²⁺ channel blockers (eg, verapamil). ICD if syncope occurs.

Diastolic dysfunction ensues.

Marked ventricular concentric hypertrophy (sarcomeres added in parallel) **B**, often septal predominance. Myofibrillar disarray and fibrosis.

Physiology of HOCM—asymmetric septal hypertrophy and systolic anterior motion of mitral valve → outflow obstruction → dyspnea, possible syncope.

Other causes of concentric LV hypertrophy: chronic HTN, Friedreich ataxia.

Restrictive/infiltrative cardiomyopathy

Postradiation fibrosis, **L**öffler endocarditis, **E**ndocardial fibroelastosis (thick fibroelastic tissue in endocardium of young children), **A**myloidosis, **S**arcoidosis, **H**emochromatosis (although dilated cardiomyopathy is more common) (**P**uppy **LEASH**).

Diastolic dysfunction ensues. Can have low-voltage ECG despite thick myocardium (especially in amyloidosis).

Löffler endocarditis—associated with hypereosinophilic syndrome; histology shows eosinophilic infiltrates in myocardium.

Heart failure



Clinical syndrome of cardiac pump dysfunction → congestion and low perfusion. Symptoms include dyspnea, orthopnea, fatigue; signs include S3 heart sound, rales, jugular venous distention (JVD), pitting edema **A**.

Systolic dysfunction—reduced EF, ↑ EDV; ↓ contractility often 2° to ischemia/MI or dilated cardiomyopathy.

Diastolic dysfunction—preserved EF, normal EDV; ↓ compliance (↑ EDP) often 2° to myocardial hypertrophy.

Right HF most often results from left HF. Cor pulmonale refers to isolated right HF due to pulmonary cause.

ACE inhibitors or angiotensin II receptor blockers, β-blockers (except in acute decompensated HF), and spironolactone ↓ mortality. Loop and thiazide diuretics are used mainly for symptomatic relief. Hydralazine with nitrate therapy improves both symptoms and mortality in select patients.

Left heart failure

Orthopnea Shortness of breath when supine: ↑ venous return from redistribution of blood (immediate gravity effect) exacerbates pulmonary vascular congestion.

Paroxysmal nocturnal dyspnea Breathless awakening from sleep: ↑ venous return from redistribution of blood, reabsorption of peripheral edema, etc.




Pulmonary edema ↑ pulmonary venous pressure → pulmonary venous distention and transudation of fluid. Presence of hemosiderin-laden macrophages (“HF” cells) in lungs.

Right heart failure

Hepatomegaly (nutmeg liver) ↑ central venous pressure → ↑ resistance to portal flow. Rarely, leads to “cardiac cirrhosis.”

Jugular venous distention ↑ venous pressure.



Peripheral edema ↑ venous pressure → fluid transudation.



Shock

Inadequate organ perfusion and delivery of nutrients necessary for normal tissue and cellular function. Initially may be reversible but life threatening if not treated promptly.

	CAUSED BY	SKIN	PCWP (PRELOAD)	CO	SVR (AFTERLOAD)	TREATMENT
Hypovolemic shock	Hemorrhage, dehydration, burns	Cold, clammy	↓↓	↓	↑	IV fluids
Cardiogenic shock	Acute MI, HF, valvular dysfunction, arrhythmia					Inotropes, diuresis
Obstructive shock	Cardiac tamponade, pulmonary embolism, tension pneumothorax	Cold, clammy	↑ or ↓	↓↓	↑	Relieve obstruction
Distributive shock	Sepsis, anaphylaxis CNS injury	Warm Dry	↓ ↓	↑ ↓	↓↓ ↓↓	IV fluids, pressors, epinephrine (anaphylaxis)

Cardiac tamponade

Compression of the heart by fluid (eg, blood, effusions [arrows in **A**] in pericardial space) → ↓ CO. Equilibration of diastolic pressures in all 4 chambers.

Findings: Beck triad (hypotension, distended neck veins, distant heart sounds), ↑ HR, pulsus paradoxus. ECG shows low-voltage QRS and electrical alternans **B** (due to “swinging” movement of heart in large effusion).

Pulsus paradoxus—↓ in amplitude of systolic BP by > 10 mm Hg during inspiration. Seen in constrictive **P**ericarditis, obstructive pulmonary disease (eg, **C**roup, **O**SA, **A**sthma, COPD), cardiac **T**amponade (**Pea COAT**).

Bacterial endocarditis

Acute—*S aureus* (high virulence). Large vegetations on previously normal valves **A**. Rapid onset.

Subacute—viridans streptococci (low virulence). Smaller vegetations on congenitally abnormal or diseased valves. Sequela of dental procedures. Gradual onset.

Symptoms: fever (most common), new murmur, Roth spots (round white spots on retina surrounded by hemorrhage **B**), Osler nodes (Ouchy raised lesions on finger or toe pads **C** due to immune complex deposition), Janeway lesions (small, painless, erythematous lesions on palm or sole) **D**, splinter hemorrhages **E** on nail bed.

Associated with glomerulonephritis, septic arterial or pulmonary emboli.

May be nonbacterial (marantic/thrombotic) 2° to malignancy, hypercoagulable state, or lupus.

FROM JANE with ♥:

Fever

Roth spots

Osler nodes

Murmur

Janeway lesions

Anemia

Nail-bed hemorrhage

Emboli

Requires multiple blood cultures for diagnosis.


If culture ⊖, most likely *Coxiella burnetii*, *Bartonella* spp.


Mitral valve is most frequently involved.

Tricuspid valve endocarditis is associated with IV **drug** abuse (don't "**tri**" **drugs**). Associated with *S aureus*, *Pseudomonas*, and *Candida*.

S bovis (*galloyticus*) is present in colon cancer, *S epidermidis* on prosthetic valves.

Native valve endocarditis may be due to **HACEK** organisms (*Haemophilus*, *Aggregatibacter* [formerly *Actinobacillus*], *Cardiobacterium*, *Eikenella*, *Kingella*).



Rheumatic fever

A consequence of pharyngeal infection with group A β -hemolytic streptococci. Late sequelae include rheumatic heart disease, which affects heart valves—mitral > aortic >> tricuspid (high-pressure valves affected most). Early lesion is mitral valve regurgitation; late lesion is mitral stenosis.

Associated with Aschoff bodies (granuloma with giant cells [blue arrows in **A**]), Anitschkow cells (enlarged macrophages with ovoid, wavy, rod-like nucleus [red arrow in **A**]), \uparrow anti-streptolysin O (ASO) and \uparrow anti-DNase B titers.

Immune mediated (type II hypersensitivity); not a direct effect of bacteria. Antibodies to **M** protein cross-react with self antigens, often myosin (molecular mimicry).

Treatment/prophylaxis: penicillin.

JONES (major criteria):

Joint (migratory polyarthritits)

♥ (carditis)

Nodules in skin (subcutaneous)

Erythema marginatum (evanescent rash with ring margin)


Sydenham chorea

Syphilitic heart disease

3° syphilis disrupts the vasa vasorum of the aorta with consequent atrophy of vessel wall and dilation of aorta and valve ring.

May see calcification of aortic root, ascending aortic arch, and thoracic aorta. Leads to “tree bark” appearance of aorta.

Can result in aneurysm of ascending aorta or aortic arch, aortic insufficiency.

Acute pericarditis

Inflammation of the pericardium [A, red arrows]. Commonly presents with sharp pain, aggravated by inspiration, and relieved by sitting up and leaning forward. Often complicated by pericardial effusion [between yellow arrows in A]. Presents with friction rub. ECG changes include widespread ST-segment elevation and/or PR depression.

Causes include idiopathic (most common; presumed viral), confirmed infection (eg, coxsackievirus B), neoplasia, autoimmune (eg, SLE, rheumatoid arthritis), uremia, cardiovascular (acute STEMI or Dressler syndrome), radiation therapy.

Treatment: NSAIDs, colchicine, glucocorticoids, dialysis (uremia).

Myocarditis

Inflammation of myocardium → global enlargement of heart and dilation of all chambers. Major cause of SCD in adults < 40 years old.

Presentation highly variable, can include dyspnea, chest pain, fever, arrhythmias (persistent tachycardia out of proportion to fever is characteristic).

Multiple causes:

- Viral (eg, adenovirus, coxsackie B, parvovirus B19, HIV, HHV-6); lymphocytic infiltrate with focal necrosis highly indicative of viral myocarditis.
- Parasitic (eg, *Trypanosoma cruzi*, *Toxoplasma gondii*)
- Bacterial (eg, *Borrelia burgdorferi*, *Mycoplasma pneumoniae*, *Corynebacterium diphtheriae*)
- Toxins (eg, carbon monoxide, black widow venom)
- Rheumatic fever
- Drugs (eg, doxorubicin, cocaine)
- Autoimmune (eg, Kawasaki disease, sarcoidosis, SLE, polymyositis/dermatomyositis)


Complications include sudden death, arrhythmias, heart block, dilated cardiomyopathy, HF, mural thrombus with systemic emboli.

Vasculitides

	EPIDEMIOLOGY/PRESENTATION	NOTES
Large-vessel vasculitis		
Giant cell (temporal) arteritis	Usually elderly females. Unilateral headache, possible temporal artery tenderness, jaw claudication. May lead to irreversible blindness due to ophthalmic artery occlusion. Associated with polymyalgia rheumatica.	Most commonly affects branches of carotid artery. Focal granulomatous inflammation A . ↑ ESR. Treat with high-dose corticosteroids prior to temporal artery biopsy to prevent blindness.
Takayasu arteritis	Usually Asian females < 40 years old. “Pulseless disease” (weak upper extremity pulses), fever, night sweats, arthritis, myalgias, skin nodules, ocular disturbances.	Granulomatous thickening and narrowing of aortic arch and proximal great vessels B . ↑ ESR. Treatment: corticosteroids.
Medium-vessel vasculitis		
Buerger disease (thromboangiitis obliterans)	Heavy smokers, males < 40 years old. Intermittent claudication. May lead to gangrene C , autoamputation of digits, superficial nodular phlebitis. Raynaud phenomenon is often present.	Segmental thrombosing vasculitis with vein and nerve involvement. Treatment: smoking cessation.
Kawasaki disease (mucocutaneous lymph node syndrome)	Asian children < 4 years old. C onjunctival injection, R ash (polymorphous → desquamating), A denopathy (cervical), S trawberry tongue (oral mucositis) D , H and-foot changes (edema, erythema), f ever.	CRASH and burn on a Kawasaki . May develop coronary artery aneurysms E ; thrombosis or rupture can cause death. Treatment: IV immunoglobulin and aspirin.
Polyarteritis nodosa	Usually middle-aged men. Hepatitis B seropositivity in 30% of patients. Fever, weight loss, malaise, headache. GI: abdominal pain, melena. Hypertension, neurologic dysfunction, cutaneous eruptions, renal damage.	Typically involves renal and visceral vessels, not pulmonary arteries. Different stages of transmural inflammation with fibrinoid necrosis. Innumerable renal microaneurysms F and spasms on arteriogram (string of pearls appearance). Treatment: corticosteroids, cyclophosphamide.
Small-vessel vasculitis		
Behçet syndrome	High incidence in people of Turkish and eastern Mediterranean descent. Recurrent aphthous ulcers, genital ulcerations, uveitis, erythema nodosum. Can be precipitated by HSV or parvovirus. Flares last 1–4 weeks.	Immune complex vasculitis. Associated with HLA-B51.
Cutaneous small-vessel vasculitis	Occurs 7–10 days after certain medications (penicillin, cephalosporins, phenytoin, allopurinol) or infections (eg, HCV, HIV). Palpable purpura, no visceral involvement.	Immune complex–mediated leukocytoclastic vasculitis; late involvement indicates systemic vasculitis.

Vasculitides (continued)


	EPIDEMIOLOGY/PRESENTATION	NOTES
Small-vessel vasculitis (continued)		
Eosinophilic granulomatosis with polyangiitis (Churg-Strauss)	Asthma, sinusitis, skin nodules or purpura, peripheral neuropathy (eg, wrist/foot drop). Can also involve heart, GI, kidneys (pauci-immune glomerulonephritis).	Granulomatous, necrotizing vasculitis with eosinophilia G . MPO-ANCA/p-ANCA, ↑ IgE level.
Granulomatosis with polyangiitis (Wegener)	Upper respiratory tract: perforation of nasal septum, chronic sinusitis, otitis media, mastoiditis. Lower respiratory tract: hemoptysis, cough, dyspnea. Renal: hematuria, red cell casts.	Triad: <ul style="list-style-type: none"> ▪ Focal necrotizing vasculitis ▪ Necrotizing granulomas in lung and upper airway ▪ Necrotizing glomerulonephritis PR3-ANCA/c-ANCA H (anti-proteinase 3). CXR: large nodular densities. Treatment: cyclophosphamide, corticosteroids.
Immunoglobulin A vasculitis	Also called Henoch-Schönlein purpura. Most common childhood systemic vasculitis. Often follows URI. Classic triad: <ul style="list-style-type: none"> ▪ Skin: palpable purpura on buttocks/legs I ▪ Arthralgias ▪ GI: abdominal pain (associated with intussusception) 	Vasculitis 2° to IgA immune complex deposition. Associated with IgA nephropathy (Berger disease). Treatment: supportive care, possibly corticosteroids.
Microscopic polyangiitis	Necrotizing vasculitis commonly involving lung, kidneys, and skin with pauci-immune glomerulonephritis and palpable purpura. Presentation similar to granulomatosis with polyangiitis but without nasopharyngeal involvement.	No granulomas. MPO-ANCA/p-ANCA J (anti-myeloperoxidase). Treatment: cyclophosphamide, corticosteroids.
Mixed cryoglobulinemia	Often due to viral infections, especially HCV. Triad of palpable purpura, weakness, arthralgias. May also have peripheral neuropathy and renal disease (eg, glomerulonephritis).	C ryoglobulins are immunoglobulins that precipitate in the C old. Vasculitis due to mixed IgG and IgM immune complex deposition.



Cardiac tumors

Most common heart tumor is a metastasis (eg, melanoma).

Myxomas



Most common 1° cardiac tumor in **adults** (arrows in **A**). 90% occur in the atria (mostly left atrium). Myxomas are usually described as a “ball valve” obstruction in the left atrium (associated with multiple syncopal episodes). IL-6 production by tumor → constitutional symptoms (eg, fever, weight loss). May auscultate early diastolic “tumor plop” sound. Histology: gelatinous material, myxoma cells immersed in glycosaminoglycans.

Adults make **myxed** drinks.

Rhabdomyomas

Most frequent 1° cardiac tumor in children (associated with tuberous sclerosis). Histology: hamartomatous growths.

Kussmaul sign

↑ in JVP on inspiration instead of a normal ↓.
 Inspiration → negative intrathoracic pressure not transmitted to heart → impaired filling of right ventricle → blood backs up into vena cava → JVD. May be seen with constrictive pericarditis, restrictive cardiomyopathies, right heart failure, massive pulmonary embolism, right atrial or ventricular tumors.

Hereditary hemorrhagic telangiectasia


Also called Osler-Weber-Rendu syndrome. Autosomal dominant disorder of blood vessels. Findings: blanching lesions (telangiectasias) on skin and mucous membranes, recurrent epistaxis, skin discolorations, arteriovenous malformations (AVMs), GI bleeding, hematuria.

▶ **CARDIOVASCULAR—PHARMACOLOGY**

Hypertension treatment

Primary (essential) hypertension	Thiazide diuretics, ACE inhibitors, angiotensin II receptor blockers (ARBs), dihydropyridine Ca ²⁺ channel blockers.	
Hypertension with heart failure	Diuretics, ACE inhibitors/ARBs, β-blockers (compensated HF), aldosterone antagonists.	β-blockers must be used cautiously in decompensated HF and are contraindicated in cardiogenic shock. In HF, ARBs may be combined with the neprilysin inhibitor sacubitril.
Hypertension with diabetes mellitus	ACE inhibitors/ARBs, Ca ²⁺ channel blockers, thiazide diuretics, β-blockers.	ACE inhibitors/ARBs are protective against diabetic nephropathy. β-blockers can mask hypoglycemia symptoms.
Hypertension in asthma	ARBs, Ca ²⁺ channel blockers, thiazide diuretics, cardioselective β-blockers.	Avoid nonselective β-blockers to prevent β ₂ -receptor–induced bronchoconstriction. Avoid ACE inhibitors to prevent confusion between drug or asthma-related cough.
Hypertension in pregnancy	Hydralazine, labetalol, methyl dopa, nifedipine.	“He likes my neonate.”

Cardiac therapy



FX

Calcium channel blockers	Amlodipine, clevidipine, nifedipine, nimodipine (dihydropyridines, act on vascular smooth muscle); diltiazem, verapamil (non-dihydropyridines, act on heart).
MECHANISM	Block voltage-dependent L-type calcium channels of cardiac and smooth muscle → ↓ muscle contractility. Vascular smooth muscle—amlodipine = nifedipine > diltiazem > verapamil. Heart—verapamil > diltiazem > amlodipine = nifedipine (verapamil = ventricle).
CLINICAL USE	Dihydropyridines (except nimodipine): hypertension, angina (including vasospastic type), Raynaud phenomenon. Nimodipine: subarachnoid hemorrhage (prevents cerebral vasospasm). Nifedipine, clevidipine: hypertensive urgency or emergency. Non-dihydropyridines: hypertension, angina, atrial fibrillation/flutter.
ADVERSE EFFECTS	Gingival hyperplasia. Dihydropyridine: peripheral edema, flushing, dizziness. Non-dihydropyridine: cardiac depression, AV block, hyperprolactinemia (verapamil), constipation.

Hydralazine

MECHANISM	↑ cGMP → smooth muscle relaxation. Vasodilates arterioles > veins; afterload reduction.
CLINICAL USE	Severe hypertension (particularly acute), HF (with organic nitrate). Safe to use during pregnancy. Frequently coadministered with a β-blocker to prevent reflex tachycardia.
ADVERSE EFFECTS	Compensatory tachycardia (contraindicated in angina/CAD), fluid retention, headache, angina, drug-induced lupus.

Hypertensive emergency

Treat with labetalol, clevidipine, fenoldopam, nifedipine, nitroprusside.

Nitroprusside	Short acting vasodilator (arteries = veins); ↑ cGMP via direct release of NO. Can cause cyanide toxicity (releases cyanide).
Fenoldopam	Dopamine D ₁ receptor agonist—coronary, peripheral, renal, and splanchnic vasodilation. ↓ BP, ↑ natriuresis. Also used postoperatively as an antihypertensive. Can cause hypotension and tachycardia.

Nitrates

Nitroglycerin, isosorbide dinitrate, isosorbide mononitrate.

MECHANISM	Vasodilate by ↑ NO in vascular smooth muscle → ↑ in cGMP and smooth muscle relaxation. Dilate veins >> arteries. ↓ preload.
CLINICAL USE	Angina, acute coronary syndrome, pulmonary edema.
ADVERSE EFFECTS	Reflex tachycardia (treat with β-blockers), hypotension, flushing, headache, “Monday disease” in industrial exposure: development of tolerance for the vasodilating action during the work week and loss of tolerance over the weekend → tachycardia, dizziness, headache upon reexposure. Contraindicated in right ventricular infarction, hypertrophic cardiomyopathy, and with concurrent PDE-5 inhibitor use.

Antianginal therapy

Goal is reduction of myocardial O_2 consumption (MVO_2) by ↓ 1 or more of the determinants of MVO_2 : end-diastolic volume, BP, HR, contractility.

COMPONENT	NITRATES	β-BLOCKERS	NITRATES + β-BLOCKERS
End-diastolic volume	↓	No effect or ↑	No effect or ↓
Blood pressure	↓	↓	↓
Contractility	↑ (reflex response)	↓	Little/no effect
Heart rate	↑ (reflex response)	↓	No effect or ↓
Ejection time	↓	↑	Little/no effect
MVO_2	↓	↓	↓↓

Verapamil is similar to β-blockers in effect.

Pindolol and acebutolol are partial β-agonists that should be used with caution in angina.

Ranolazine

MECHANISM	Inhibits the late phase of inward sodium current thereby reducing diastolic wall tension and oxygen consumption. Does not affect heart rate or blood pressure.
CLINICAL USE	Angina refractory to other medical therapies.
ADVERSE EFFECTS	Constipation, dizziness, headache, nausea.


Sacubitril

MECHANISM	A neprilysin inhibitor; prevents degradation of natriuretic peptides, angiotensin II, and substance P → ↑ vasodilation, ↓ ECF volume.
CLINICAL USE	Used in combination with valsartan (an ARB) to treat HFrEF.
ADVERSE EFFECTS	Hypotension, hyperkalemia, cough, dizziness; contraindicated with ACE inhibitors due to angioedema.

Lipid-lowering agents

DRUG	LDL	HDL	TRIGLYCERIDES	MECHANISMS OF ACTION	ADVERSE EFFECTS/PROBLEMS
HMG-CoA reductase inhibitors (eg, atorvastatin, simvastatin)	↓↓↓	↑	↓	Inhibit conversion of HMG-CoA to mevalonate, a cholesterol precursor; ↑ LDL recycling; ↓ mortality in CAD patients	Hepatotoxicity (↑ LFTs), myopathy (esp when used with fibrates or niacin)
Bile acid resins Cholestyramine, colestipol, colesevelam	↓↓	↑ slightly	↑ slightly	Prevent intestinal reabsorption of bile acids; liver must use cholesterol to make more	GI upset, ↓ absorption of other drugs and fat-soluble vitamins
Ezetimibe	↓↓	↑/—	↓/—	Prevents cholesterol absorption at small intestine brush border	Rare ↑ LFTs, diarrhea
Fibrates Gemfibrozil, bezafibrate, fenofibrate	↓	↑	↓↓↓	Upregulate LPL → ↑ TG clearance Activates PPAR-α to induce HDL synthesis	Myopathy (↑ risk with statins), cholesterol gallstones (via inhibition of cholesterol 7α-hydroxylase)
Niacin	↓↓	↑↑	↓	Inhibits lipolysis (hormone-sensitive lipase) in adipose tissue; reduces hepatic VLDL synthesis	Flushed face (↓ by NSAIDs or long-term use) Hyperglycemia Hyperuricemia
PCSK9 inhibitors Alirocumab, evolocumab	↓↓↓	↑	↓	Inactivation of LDL-receptor degradation → ↑ removal of LDL from bloodstream	Myalgias, delirium, dementia, other neurocognitive effects
Fish oil and marine omega-3 fatty acids	↑ slightly	↑ slightly	↓ at high doses	Believed to decrease FFA delivery to liver and decrease activity of TG-synthesizing enzymes	Nausea, fish-like taste

Lipid-lowering agents (continued)




Cardiac glycosides

Digoxin.

MECHANISM

Direct inhibition of Na^+/K^+ ATPase
 → indirect inhibition of $\text{Na}^+/\text{Ca}^{2+}$ exchanger.
 $\uparrow [\text{Ca}^{2+}]_i \rightarrow$ positive inotropy. Stimulates vagus nerve → \downarrow HR.



CLINICAL USE

HF (\uparrow contractility); atrial fibrillation (\downarrow conduction at AV node and depression of SA node).

ADVERSE EFFECTS

Cholinergic effects (nausea, vomiting, diarrhea), blurry **yellow** vision (think van **Glow**), arrhythmias, AV block.
 Can lead to hyperkalemia, which indicates poor prognosis.
 Factors predisposing to toxicity: renal failure (\downarrow excretion), hypokalemia (permissive for digoxin binding at K^+ -binding site on Na^+/K^+ ATPase), drugs that displace digoxin from tissue-binding sites, and \downarrow clearance (eg, verapamil, amiodarone, quinidine).


ANTIDOTE

Slowly normalize K^+ , cardiac pacer, anti-digoxin Fab fragments, Mg^{2+} .


**Antiarrhythmics—
sodium channel
blockers (class I)**

Slow or block (↓) conduction (especially in depolarized cells). ↓ slope of phase 0 depolarization. Are state dependent (selectively depress tissue that is frequently depolarized [eg, tachycardia]).


Class IA	Quinidine, Procainamide, Disopyramide. “The Queen Proclaims Diso’s pyramid. ”
MECHANISM	Moderate Na ⁺ channel blockade. ↑ AP duration, ↑ effective refractory period (ERP) in ventricular action potential, ↑ QT interval, some potassium channel blocking effects.
CLINICAL USE	Both atrial and ventricular arrhythmias, especially re-entrant and ectopic SVT and VT.
ADVERSE EFFECTS	Cinchonism (headache, tinnitus with quinidine), reversible SLE-like syndrome (procainamide), HF (disopyramide), thrombocytopenia, torsades de pointes due to ↑ QT interval.



Class IB	Lidocaine, Mexiletine. “I’d Buy Liddy’s Mexican Tacos. ”
MECHANISM	Weak Na ⁺ channel blockade. ↓ AP duration. Preferentially affect ischemic or depolarized Purkinje and ventricular tissue. Phenytoin can also fall into the IB category.
CLINICAL USE	Acute ventricular arrhythmias (especially post-MI), digitalis-induced arrhythmias. IB is Best post-MI.
ADVERSE EFFECTS	CNS stimulation/depression, cardiovascular depression.



Class IC	Flecainide, Propafenone. “ Can I have Fries, Please. ”
MECHANISM	Strong Na ⁺ channel blockade. Significantly prolongs ERP in AV node and accessory bypass tracts. No effect on ERP in Purkinje and ventricular tissue. Minimal effect on AP duration.
CLINICAL USE	SVTs, including atrial fibrillation. Only as a last resort in refractory VT.
ADVERSE EFFECTS	Proarrhythmic, especially post-MI (contraindicated). IC is C ontraindicated in structural and ischemic heart disease.



Antiarrhythmics— β -blockers (class II)


Metoprolol, propranolol, esmolol, atenolol, timolol, carvedilol.

MECHANISMDecrease SA and AV nodal activity by \downarrow cAMP, \downarrow Ca^{2+} currents. Suppress abnormal pacemakers by \downarrow slope of phase 4.AV node particularly sensitive— \uparrow PR interval. Esmolol very short acting.**CLINICAL USE**

SVT, ventricular rate control for atrial fibrillation and atrial flutter.

ADVERSE EFFECTS

Impotence, exacerbation of COPD and asthma, cardiovascular effects (bradycardia, AV block, HF), CNS effects (sedation, sleep alterations). May mask the signs of hypoglycemia.

Metoprolol can cause dyslipidemia. Propranolol can exacerbate vasospasm in vasospastic angina. β -blockers (except the nonselective α - and β -antagonists carvedilol and labetalol) cause unopposed α_1 -agonism if given alone for pheochromocytoma or for cocaine toxicity (unsubstantiated). Treat β -blocker overdose with saline, atropine, glucagon.**Antiarrhythmics—potassium channel blockers (class III)**

Amiodarone, Ibutilide, Dofetilide, Sotalol.

AIDS.**MECHANISM** \uparrow AP duration, \uparrow ERP, \uparrow QT interval.**CLINICAL USE**

Atrial fibrillation, atrial flutter; ventricular tachycardia (amiodarone, sotalol).


ADVERSE EFFECTSSotalol—torsades de pointes, excessive β blockade.

Ibutilide—torsades de pointes.

Amiodarone—pulmonary fibrosis, hepatotoxicity, hypothyroidism or hyperthyroidism (amiodarone is 40% iodine by weight), acts as hapten (corneal deposits, blue/gray skin deposits resulting in photodermatitis), neurologic effects, constipation, cardiovascular effects (bradycardia, heart block, HF).

Remember to check PFTs, LFTs, and TFTs when using amiodarone.


Amiodarone is lipophilic and has class I, II, III, and IV effects.



**Antiarrhythmics—
calcium channel
blockers (class IV)**

Diltiazem, Verapamil

MECHANISM	Decrease conduction Velocity, ↑ ERP, ↑ PR interval.
CLINICAL USE	Prevention of nodal arrhythmias (eg, SVT), rate control in atrial fibrillation.
ADVERSE EFFECTS	Constipation, flushing, edema, cardiovascular effects (HF, AV block, sinus node depression).

**Other antiarrhythmics**

Adenosine	↑ K ⁺ out of cells → hyperpolarizing the cell and ↓ I _{Ca} , decreasing AV node conduction. Drug of choice in diagnosing/terminating certain forms of SVT. Very short acting (~ 15 sec). Effects blunted by theophylline and caffeine (both are adenosine receptor antagonists). Adverse effects include flushing, hypotension, chest pain, sense of impending doom, bronchospasm.
Magnesium	Effective in torsades de pointes and digoxin toxicity.

Ivabradine

MECHANISM	I vabradine prolongs slow depolarization (phase “ I V”) by selectively inhibiting “funny” sodium channels (I _f).
CLINICAL USE	Chronic stable angina in patients who cannot take β-blockers. Chronic HFrEF.
ADVERSE EFFECTS	Luminous phenomena/visual brightness, hypertension, bradycardia.

HIGH-YIELD SYSTEMS

Endocrine

“If you skew the endocrine system, you lose the pathways to self.”

—Hilary Mantel

“We have learned that there is an endocrinology of elation and despair, a chemistry of mystical insight, and, in relation to the autonomic nervous system, a meteorology and even . . . an astro-physics of changing moods.”

—Aldous Huxley

“Chocolate causes certain endocrine glands to secrete hormones that affect your feelings and behavior by making you happy.”

—Elaine Sherman, *Book of Divine Indulgences*

The endocrine system comprises widely distributed organs that work in a highly integrated manner to orchestrate a state of hormonal equilibrium within the body. Generally speaking, endocrine diseases can be classified either as diseases of underproduction or overproduction, or as conditions involving the development of mass lesions—which themselves may be associated with underproduction or overproduction of hormones. Therefore, study the endocrine system first by learning the glands, their hormones, and their regulation, and then by integrating disease manifestations with diagnosis and management. Take time to learn the multisystem connections.

▶ Embryology	326
▶ Anatomy	327
▶ Physiology	328
▶ Pathology	338
▶ Pharmacology	352

▶ ENDOCRINE—EMBRYOLOGY

Thyroid development




Thyroid diverticulum arises from floor of primitive pharynx and descends into neck. Connected to tongue by thyroglossal duct, which normally disappears but may persist as cysts or the pyramidal lobe of thyroid. Foramen cecum is normal remnant of thyroglossal duct.

Most common ectopic thyroid tissue site is the tongue (lingual thyroid). Removal may result in hypothyroidism if it is the only thyroid tissue present.

Thyroglossal duct cyst **A** presents as an anterior midline neck mass that moves with swallowing or protrusion of the tongue (vs persistent cervical sinus leading to pharyngeal cleft cyst in lateral neck).

Thyroid follicular cells derived from endoderm.



▶ ENDOCRINE—ANATOMY

Pituitary gland

Anterior pituitary (adenohypophysis)

Secretes FSH, LH, ACTH, TSH, prolactin, GH, and β -endorphin. Melanotropin (MSH) secreted from intermediate lobe of pituitary. Derived from oral ectoderm (Rathke pouch).

- α subunit—hormone subunit common to TSH, LH, FSH, and hCG.
- β subunit—determines hormone specificity.


Proopiomelanocortin derivatives— β -endorphin, ACTH, and MSH. Go **pro** with a **BAM!**
FLAT PiG: FSH, LH, ACTH, TSH, PRL, GH.
B-FLAT: Basophils—FSH, LH, ACTH, TSH.
Acid PiG: Acidophils — PRL, GH.

Posterior pituitary (neurohypophysis)

Stores and releases vasopressin (antidiuretic hormone, or ADH) and oxytocin, both made in the hypothalamus (supraoptic and paraventricular nuclei) and transported to posterior pituitary via neurophysins (carrier proteins). Derived from **neuroectoderm**.

Adrenal cortex and medulla

Adrenal cortex (derived from mesoderm) and medulla (derived from neural crest).




GFR corresponds with **S**alt (mineralocorticoids), **S**ugar (glucocorticoids), and **S**ex (androgens).
“The deeper you go, **the sweeter it gets.**”


► ENDOCRINE—PHYSIOLOGY

Hypothalamic-pituitary hormones

HORMONE	FUNCTION	CLINICAL NOTES
ADH	↑ water permeability of distal convoluted tubule and collecting duct cells in kidney to ↑ water reabsorption	Stimulus for secretion is ↑ plasma osmolality, except in SIADH, in which ADH is elevated despite ↓ plasma osmolality
CRH	↑ ACTH, MSH, β-endorphin	↓ in chronic exogenous steroid use
Dopamine	↓ prolactin, TSH	Also called prolactin-inhibiting factor Dopamine antagonists (eg, antipsychotics) can cause galactorrhea due to hyperprolactinemia
GHRH	↑ GH	Analog (tesamorelin) used to treat HIV-associated lipodystrophy
GnRH	↑ FSH, LH	Suppressed by hyperprolactinemia Tonic GnRH analog (eg, leuprolide) suppresses hypothalamic–pituitary–gonadal axis. Pulsatile GnRH leads to puberty, fertility
MSH	↑ melanogenesis by melanocytes	Causes hyperpigmentation in Cushing disease, as MSH and ACTH share the same precursor molecule, proopiomelanocortin
Oxytocin	Causes uterine contractions during labor. Responsible for milk letdown reflex in response to suckling.	Modulates fear, anxiety, social bonding, mood, and depression
Prolactin	↓ GnRH Stimulates lactogenesis.	Pituitary prolactinoma → amenorrhea, osteoporosis, hypogonadism, galactorrhea Breastfeeding → ↑ prolactin → ↓ GnRH → delayed postpartum ovulation (natural contraception)
Somatostatin	↓ GH, TSH	Also called growth hormone inhibiting hormone (GHIH) Analogues used to treat acromegaly
TRH	↑ TSH, prolactin	↑ TRH (eg, in 1°/2° hypothyroidism) may increase prolactin secretion → galactorrhea



Growth hormone



Also called somatotropin. Secreted by anterior pituitary.

Stimulates linear growth and muscle mass through IGF-1 (somatomedin C) secretion by liver. ↑ insulin resistance (diabetogenic).

Released in pulses in response to growth hormone–releasing hormone (GHRH).

Secretion ↑ during exercise, deep sleep, puberty, hypoglycemia, CKD.

Secretion ↓ by glucose, somatostatin, somatomedin (regulatory molecule secreted by liver in response to GH acting on target tissues).

Excess secretion of GH (eg, pituitary adenoma) may cause acromegaly (adults) or gigantism (children). Treatment: somatostatin analogs (eg, octreotide) or surgery.

Antidiuretic hormone

Also called vasopressin.

SOURCE

Synthesized in hypothalamus (supraoptic and paraventricular nuclei), stored and secreted by posterior pituitary.

FUNCTION

Regulates blood pressure (V_1 -receptors) and serum osmolality (V_2 -receptors). Primary function is serum osmolality regulation (ADH ↓ serum osmolality, ↑ urine osmolality) via regulation of aquaporin channel insertion in principal cells of renal collecting duct.

ADH level is ↓ in central diabetes insipidus (DI), normal or ↑ in nephrogenic DI.

Nephrogenic DI can be caused by mutation in V_2 -receptor.


Desmopressin (ADH analog) is a treatment for central DI and nocturnal enuresis.

REGULATION

Plasma osmolality (1°); hypovolemia.

Prolactin

SOURCE	Secreted mainly by anterior pituitary.	Structurally homologous to growth hormone.
FUNCTION	Stimulates milk production in breast; inhibits ovulation in females and spermatogenesis in males by inhibiting GnRH synthesis and release.	Excessive amounts of prolactin associated with ↓ libido.
REGULATION	Prolactin secretion from anterior pituitary is tonically inhibited by dopamine from tuberoinfundibular pathway of hypothalamus. Prolactin in turn inhibits its own secretion by ↑ dopamine synthesis and secretion from hypothalamus. TRH ↑ prolactin secretion (eg, in 1° or 2° hypothyroidism).	Dopamine agonists (eg, bromocriptine) inhibit prolactin secretion and can be used in treatment of prolactinoma. Dopamine antagonists (eg, most antipsychotics, metoclopramide) and estrogens (eg, OCPs, pregnancy) stimulate prolactin secretion.



Thyroid hormones

Thyroid produces triiodothyronine (T_3) and thyroxine (T_4), iodine-containing hormones that control the body's metabolic rate.

SOURCE

Follicles of thyroid. 5'-deiodinase converts T_4 (the major thyroid product) to T_3 in peripheral tissue (5, 4, 3). Peripheral conversion is inhibited by glucocorticoids, β -blockers, and propylthiouracil (PTU). Reverse T_3 (rT_3) is a metabolically inactive byproduct of the peripheral conversion of T_4 and its production is increased by growth hormone and glucocorticoids. Functions of thyroid peroxidase include oxidation, organification of iodine, and coupling of monoiodotyrosine (MIT) and diiodotyrosine (DIT). Inhibited by PTU and methimazole. $DIT + DIT = T_4$. $DIT + MIT = T_3$. Wolff-Chaikoff effect—excess iodine temporarily turns off thyroid peroxidase \rightarrow $\downarrow T_3/T_4$ production (protective autoregulatory effect).

FUNCTION

Only free hormone is active. T_3 binds nuclear receptor with greater affinity than T_4 . T_3 functions

—7 B's:

- **B**rain maturation
- **B**one growth (synergism with GH)
- **β** -adrenergic effects. $\uparrow \beta_1$ receptors in heart \rightarrow \uparrow CO, HR, SV, contractility; β -blockers alleviate adrenergic symptoms in thyrotoxicosis
- **B**asal metabolic rate \uparrow (via Na^+/K^+ -ATPase activity \rightarrow \uparrow O_2 consumption, RR, body temperature)
- **B**lood sugar (\uparrow glycogenolysis, gluconeogenesis)
- **B**reak down lipids (\uparrow lipolysis)
- Stimulates surfactant synthesis in **B**abies

REGULATION


TRH \oplus TSH release \rightarrow \oplus follicular cells. Thyroid-stimulating immunoglobulin (TSI) may \oplus follicular cells in Graves disease.

Negative feedback primarily by free T_3/T_4 :

- Anterior pituitary \rightarrow \downarrow sensitivity to TRH
- Hypothalamus \rightarrow \downarrow TRH secretion

Thyroxine-binding globulin (TBG) binds most T_3/T_4 in blood. Bound T_3/T_4 = inactive.

- \uparrow TBG in pregnancy, OCP use (estrogen \rightarrow \uparrow TBG) \rightarrow \uparrow total T_3/T_4
- \downarrow TBG in steroid use, nephrotic syndrome



Parathyroid hormone

SOURCE

Chief cells of parathyroid

FUNCTION


↑ free Ca^{2+} in the blood (1° function)
 ↑ Ca^{2+} and PO_4^{3-} absorption in GI system
 ↑ Ca^{2+} and PO_4^{3-} from bone resorption
 ↑ Ca^{2+} reabsorption from DCT
 ↓ PO_4^{3-} reabsorption in PCT
 ↑ $1,25\text{-(OH)}_2\text{D}_3$ (calcitriol) production by activating 1α -hydroxylase in **PCT**
Tri to make D_3 in the **PCT**

PTH ↑ serum Ca^{2+} , ↓ serum PO_4^{3-} , ↑ urine PO_4^{3-} , ↑ urine cAMP
 ↑ RANK-L (receptor activator of NF- κ B ligand) secreted by osteoblasts and osteocytes; binds RANK (receptor) on osteoclasts and their precursors to stimulate osteoclasts and ↑ Ca^{2+} → bone resorption (intermittent PTH release can also stimulate bone formation)

PTH = Phosphate-Trashing Hormone
 PTH-related peptide (PTHrP) functions like PTH and is commonly increased in malignancies (eg, squamous cell carcinoma of the lung, renal cell carcinoma)

REGULATION

↓ serum Ca^{2+} → ↑ PTH secretion
 ↑ serum PO_4^{3-} → ↑ PTH secretion
 ↓ serum Mg^{2+} → ↑ PTH secretion
 ↓↓ serum Mg^{2+} → ↓ PTH secretion
 Common causes of ↓ Mg^{2+} include diarrhea, aminoglycosides, diuretics, alcohol abuse



Calcium homeostasis

Plasma Ca^{2+} exists in three forms:

- Ionized/free (~ 45%, active form)
- Bound to albumin (~ 40%)
- Bound to anions (~ 15%)

↑ pH (less H^+) → albumin binds more Ca^{2+} → ↓ ionized Ca^{2+} (eg, cramps, pain, paresthesias, carpopedal spasm) → ↑ PTH

↓ pH (more H^+) → albumin binds less Ca^{2+} → ↑ ionized Ca^{2+} → ↓ PTH


Ionized/free Ca^{2+} is 1° regulator of PTH; changes in pH alter PTH secretion, whereas changes in albumin concentration do not

Calcitonin

SOURCE	Parafollicular cells (C cells) of thyroid.	Calcitonin opposes actions of PTH. Not important in normal Ca^{2+} homeostasis Calcitonin tones down serum Ca^{2+} levels and keeps it in bones
FUNCTION	↓ bone resorption of Ca^{2+} .	
REGULATION	↑ serum Ca^{2+} → ↑ calcitonin secretion.	

Glucagon

SOURCE	Made by α cells of pancreas.
FUNCTION	Promotes glycogenolysis, gluconeogenesis, lipolysis, ketogenesis. Elevates blood sugar levels to maintain homeostasis when bloodstream glucose levels fall too low (ie, fasting state).
REGULATION	Secreted in response to hypoglycemia. Inhibited by insulin, hyperglycemia, somatostatin.

Insulin**SYNTHESIS**

Preproinsulin (synthesized in RER of pancreatic β cells) \rightarrow cleavage of “presignal” \rightarrow proinsulin (stored in secretory granules) \rightarrow cleavage of proinsulin \rightarrow exocytosis of insulin and C-peptide equally. Insulin and C-peptide are \uparrow in insulinoma and sulfonylurea use, whereas exogenous insulin lacks C-peptide.

FUNCTION

Binds **insulin** receptors (tyrosine kinase activity **1**), **inducing** glucose uptake (carrier-mediated transport) **into** insulin-dependent tissue **2** and gene transcription.

Anabolic effects of insulin:

- \uparrow glucose transport in skeletal muscle and adipose tissue
- \uparrow glycogen synthesis and storage
- \uparrow triglyceride synthesis
- \uparrow Na^+ retention (kidneys)
- \uparrow protein synthesis (muscles)
- \uparrow cellular uptake of K^+ and amino acids
- \downarrow glucagon release
- \downarrow lipolysis in adipose tissue

Unlike glucose, insulin does not cross placenta.

Insulin-dependent glucose transporters:

- GLUT4: adipose tissue, striated muscle (exercise can also \uparrow GLUT4 expression)

Insulin-independent transporters:

- GLUT1: RBCs, brain, cornea, placenta
- GLUT2 (**bidirectional**): β islet cells, liver, kidney, GI tract (think **2-way street**)
- GLUT3: brain, placenta
- GLUT5 (**Fructose**): spermatoocytes, GI tract
- SGLT1/SGLT2 (Na^+ -glucose cotransporters): kidney, small intestine



Brain prefers glucose, but may use ketone bodies during starvation. RBCs utilize glucose, as they lack mitochondria for aerobic metabolism.

BRICK LIPS (insulin-independent glucose uptake): **B**rain, **R**BCs, **I**ntestine, **C**ornea, **K**idney, **L**iver, **I**slet (β) cells, **P**lacenta, **S**permatoocytes.


REGULATION

Glucose is the major regulator of insulin release. \uparrow insulin response with oral vs IV glucose due to incretins (eg, glucagon-like peptide 1 [GLP-1], glucose-dependent insulinotropic polypeptide [GIP]), which are released after meals and \uparrow β cell sensitivity to glucose. Release \downarrow by α_2 , \uparrow by β_2 stimulation (**2** = regulates **insulin**)

Glucose enters β cells **3** \rightarrow \uparrow ATP generated from glucose metabolism **4** closes K^+ channels (target of sulfonylureas) **5** and depolarizes β cell membrane **6**. Voltage-gated Ca^{2+} channels open \rightarrow Ca^{2+} influx **7** and stimulation of insulin exocytosis **8**.



Adrenal steroids and congenital adrenal hyperplasias



^aRate-limiting step.

ENZYME DEFICIENCY	MINERALOCORTICIDS	[K ⁺]	BP	CORTISOL	SEX HORMONES	LABS	PRESENTATION
A 17α-hydroxylase^a	↑	↓	↑	↓	↓	↓ androstenedione	XY: ambiguous genitalia, undescended testes XX: lacks 2° sexual development
B 21-hydroxylase^a	↓	↑	↓	↓	↑	↑ renin activity ↑ 17-hydroxyprogesterone	Most common Presents in infancy (salt wasting) or childhood (precocious puberty) XX: virilization
C 11β-hydroxylase^a	↓ aldosterone ↑ 11-deoxycorticosterone (results in ↑ BP)	↓	↑	↓	↑	↓ renin activity	Presents in infancy (severe hypertension) or childhood (precocious puberty) XX: virilization

^aAll congenital adrenal enzyme deficiencies are autosomal recessive disorders and most are characterized by skin hyperpigmentation (due to ↑ MSH production, which is coproduced and secreted with ACTH) and bilateral adrenal gland enlargement (due to ↑ ACTH stimulation).

If deficient enzyme starts with 1, it causes hypertension; if deficient enzyme ends with 1, it causes virilization in females.

Cortisol

SOURCE

Adrenal zona fasciculata.


FUNCTION

- ↑ **A**ppetite
- ↑ **B**lood pressure:
 - Upregulates α_1 -receptors on arterioles → ↑ sensitivity to norepinephrine and epinephrine (permissive action)
 - At high concentrations, can bind to mineralocorticoid (aldosterone) receptors
- ↑ **I**nsulin resistance (diabetogenic)
- ↑ **G**luconeogenesis, lipolysis, and proteolysis (↓ glucose utilization)
- ↓ **F**ibroblast activity (poor wound healing, ↓ collagen synthesis, ↑ striae)
- ↓ **I**nflammatory and **I**mmune responses:
 - Inhibits production of leukotrienes and prostaglandins
 - Inhibits WBC adhesion → neutrophilia
 - Blocks histamine release from mast cells
 - Eosinopenia, lymphopenia
 - Blocks IL-2 production
- ↓ **B**one formation (↓ osteoblast activity)

Bound to corticosteroid-binding globulin.

Cortisol is **A BIG FIB**.

Exogenous corticosteroids can cause reactivation of TB and candidiasis (blocks IL-2 production).



REGULATION

CRH (hypothalamus) stimulates ACTH release (pituitary) → cortisol production in adrenal zona fasciculata. Excess cortisol ↓ CRH, ACTH, and cortisol secretion.

Chronic stress may induce prolonged cortisol secretion, cortisol resistance, impaired immunocompetency, and dysregulation of HPA axis.

Appetite regulation**Ghrelin**

Stimulates hunger (orexigenic effect) and GH release (via GH secretagog receptor). Produced by stomach. Sleep deprivation, fasting, or Prader-Willi syndrome → ↑ ghrelin production. **G**hrelin makes you hun**gh**re and **gh**row. Acts on lateral area of hypothalamus (hunger center) to ↑ appetite.

Leptin


Satiety hormone. Produced by adipose tissue. Mutation of leptin gene → central obesity. (Obese people have ↑ leptin due to ↑ adipose tissue but also appear resistant to leptin's anorexigenic effect.) Sleep deprivation or starvation → ↓ leptin production. **L**eptin keeps you **thin**. Acts on ventromedial area of hypothalamus (satiety center) to ↓ appetite.

Endocannabinoids

Act at cannabinoid receptors in hypothalamus and nucleus accumbens, two key brain areas for the homeostatic and hedonic control of food intake → ↑ appetite. Exogenous cannabinoids cause “the munchies.”

Signaling pathways of endocrine hormones

cAMP	F SH, L H, A CTH, T SH, C RH, h CG, A DH (V ₂ -receptor), M SH, P TH, C alcitonin, H istamine (H ₂ -receptor), G lucagon, G HRH	F LAT C hAMPs C Hu G G
cGMP	B NP, A NP, E DRF (NO)	B AD G ra M Pa Think vasodilation and diuresis
IP₃	G nRH, O xytocin, A DH (V ₁ -receptor), T RH, H istamine (H ₁ -receptor), A ngiotensin II, G astrin	G OAT H AG
Intracellular receptor	P rogesterone, E strogen, T estosterone, C ortisol, A ldosterone, T ₃ / T ₄ , V itamin D	P ET C AT on T V
Receptor tyrosine kinase	I GF-1, F GF, P DGF, E GF, T GF-β, I nsulin	M AP kinase pathway G et F ound I n the M AP
Nonreceptor tyrosine kinase	P rolactin, I mmunomodulators (eg, cytokines IL-2, IL-6, IFN), G H, G -CSF, E rythropoietin, T hrombopoietin	J AK/ S TAT pathway Think acidophils and cytokines P IGGLET

Signaling pathways of steroid hormones

Steroid hormones are lipophilic and therefore must circulate bound to specific binding globulins, which ↑ their solubility.

In men, ↑ sex hormone-binding globulin (SHBG) lowers free testosterone → gynecomastia.

In women, ↓ SHBG raises free testosterone → hirsutism.

↑ estrogen (eg, OCPs, pregnancy) → ↑ SHBG.

► ENDOCRINE—PATHOLOGY

Syndrome of inappropriate antidiuretic hormone secretion

Characterized by:

- Excessive free water retention
- Euvolemic hyponatremia with continued urinary Na^+ excretion
- Urine osmolality > serum osmolality

Body responds to water retention with ↓ aldosterone and ↑ ANP and BNP → ↑ urinary Na^+ secretion → normalization of extracellular fluid volume → euvolemic hyponatremia. Very low serum Na^+ levels can lead to cerebral edema, seizures. Correct slowly to prevent osmotic demyelination syndrome (formerly called central pontine myelinolysis).

SIADH causes include:

- Ectopic ADH (eg, small cell lung cancer)
- CNS disorders/head trauma
- Pulmonary disease
- Drugs (eg, SSRIs, carbamazepine, cyclophosphamide)

Treatment: fluid restriction (first line), salt tablets, IV hypertonic saline, diuretics, ADH antagonists (eg, conivaptan, tolvaptan, demeclocycline).

Diabetes insipidus

Characterized by intense thirst and polyuria with inability to concentrate urine due to lack of ADH (central) or failure of response to circulating ADH (nephrogenic).

	Central DI	Nephrogenic DI
ETIOLOGY	Pituitary tumor, autoimmune, trauma, surgery, ischemic encephalopathy, idiopathic	Hereditary (ADH receptor mutation), 2° to hypercalcemia, hypokalemia, lithium, demeclocycline (ADH antagonist)
FINDINGS	↓ ADH	Normal or ↑ ADH levels
		Urine specific gravity < 1.006 Urine osmolality < 300 mOsm/kg Serum osmolality > 290 mOsm/kg Hyperosmotic volume contraction
WATER DEPRIVATION TEST ^a	> 50% ↑ in urine osmolality only after administration of ADH analog	Minimal change in urine osmolality, even after administration of ADH analog
TREATMENT	Desmopressin Hydration	HCTZ, indomethacin, amiloride Hydration, dietary salt restriction, avoidance of offending agent

^aNo water intake for 2–3 hr followed by hourly measurements of urine volume and osmolality as well as plasma Na^+ concentration and osmolality. ADH analog (desmopressin) is administered if serum osmolality > 295–300 mOsm/kg, plasma Na^+ ≥ 145 mEq/L, or urine osmolality does not rise despite a rising plasma osmolality.

Hypopituitarism

Undersecretion of pituitary hormones due to:

- Nonsecreting pituitary adenoma, craniopharyngioma
- **Sheehan syndrome**—ischemic infarct of pituitary following postpartum bleeding; pregnancy-induced pituitary growth → ↑ susceptibility to hypoperfusion. Usually presents with failure to lactate, absent menstruation, cold intolerance
- **Empty sella syndrome**—atrophy or compression of pituitary (which lies in the sella turcica), often idiopathic, common in obese women; associated with idiopathic intracranial hypertension
- **Pituitary apoplexy**—sudden hemorrhage of pituitary gland, often in the presence of an existing pituitary adenoma. Usually presents with sudden onset severe headache, visual impairment (eg, bitemporal hemianopia, diplopia due to CN III palsy), and features of hypopituitarism
- Brain injury
- Radiation

Treatment: hormone replacement therapy (corticosteroids, thyroxine, sex steroids, human growth hormone)


Acromegaly

Excess GH in adults. Typically caused by pituitary adenoma.

FINDINGS

Large tongue with deep furrows, deep voice, large hands and feet, coarsening of facial features with aging **A**, frontal bossing, diaphoresis (excessive sweating), impaired glucose tolerance (insulin resistance), hypertension. ↑ risk of colorectal polyps and cancer.

↑ GH in children → gigantism (↑ linear bone growth). HF most common cause of death.

**DIAGNOSIS**


↑ serum IGF-1; failure to suppress serum GH following oral glucose tolerance test; pituitary mass seen on brain MRI.

TREATMENT

Pituitary adenoma resection. If not cured, treat with octreotide (somatostatin analog), pegvisomant (GH receptor antagonist), or dopamine agonists (eg, cabergoline).

Hypothyroidism vs hyperthyroidism


FINDINGS	Hypothyroidism	Hyperthyroidism
METABOLIC	Cold intolerance, ↓ sweating, weight gain (↓ basal metabolic rate → ↓ calorogenesis), hyponatremia (↓ free water clearance)	Heat intolerance, ↑ sweating, weight loss (↑ synthesis of Na ⁺ -K ⁺ ATPase → ↑ basal metabolic rate → ↑ calorogenesis)
SKIN/HAIR	Dry, cool skin (due to ↓ blood flow); coarse, brittle hair; diffuse alopecia; brittle nails; puffy facies and generalized nonpitting edema (myxedema) due to ↑ GAGs in interstitial spaces → ↑ osmotic pressure → water retention	Warm, moist skin (due to vasodilation); fine hair; onycholysis (A); pretibial myxedema in Graves disease
OCULAR	Periorbital edema	Ophthalmopathy in Graves disease (including periorbital edema, exophthalmos), lid lag/retraction (↑ sympathetic stimulation of levator palpebrae superioris and superior tarsal muscle)
GASTROINTESTINAL	Constipation (↓ GI motility), ↓ appetite	Hyperdefecation/diarrhea (↑ GI motility), ↑ appetite
MUSCULOSKELETAL	Hypothyroid myopathy (proximal weakness, ↑ CK), carpal tunnel syndrome, myoedema (small lump rising on the surface of a muscle when struck with a hammer)	Thyrotoxic myopathy (proximal weakness, normal CK), osteoporosis/↑ fracture rate (T ₃ directly stimulates bone resorption)
REPRODUCTIVE	Abnormal uterine bleeding, ↓ libido, infertility	Abnormal uterine bleeding, gynecomastia, ↓ libido, infertility
NEUROPSYCHIATRIC	Hypoactivity, lethargy, fatigue, weakness, depressed mood, ↓ reflexes (delayed/slow relaxing)	Hyperactivity, restlessness, anxiety, insomnia, fine tremors (due to ↑ β-adrenergic activity), ↑ reflexes (brisk)
CARDIOVASCULAR	Bradycardia, dyspnea on exertion (↓ cardiac output)	Tachycardia, palpitations, dyspnea, arrhythmias (eg, atrial fibrillation), chest pain and systolic HTN due to ↑ number and sensitivity of β-adrenergic receptors, ↑ expression of cardiac sarcolemmal ATPase and ↓ expression of phospholamban
LABS	↑ TSH (if 1°) ↓ free T ₃ and T ₄ Hypercholesterolemia (due to ↓ LDL receptor expression)	↓ TSH (if 1°) ↑ free T ₃ and T ₄ ↓ LDL, HDL, and total cholesterol



Hypothyroidism

Hashimoto thyroiditis	Most common cause of hypothyroidism in iodine-sufficient regions; an autoimmune disorder with antithyroid peroxidase (antimicrosomal) and antithyroglobulin antibodies. Associated with HLA-DR3, HLA-DR5, ↑ risk of non-Hodgkin lymphoma (typically of B-cell origin). May be hyperthyroid early in course due to thyrotoxicosis during follicular rupture. Histology: Hürthle cells A , lymphoid aggregates with germinal centers. Findings: moderately enlarged, nontender thyroid.
Postpartum thyroiditis	Self-limited thyroiditis arising up to 1 year after delivery. Presents as transient hyperthyroidism, hypothyroidism, or hyperthyroidism followed by hypothyroidism. Majority of women are euthyroid following resolution. Thyroid usually painless and normal in size. Histology: lymphocytic infiltrate with occasional germinal center formation.
Congenital hypothyroidism (cretinism)	Severe fetal hypothyroidism due to antibody-mediated maternal hypothyroidism, thyroid dysgenesis (most common cause in US; eg, agenesis, ectopy, hypoplasia), iodine deficiency, dyshormonogenetic goiter (commonly due to mutations in thyroid peroxidase). Findings (6 P's): P ot-bellied, P ale, P uffy-faced child B with P rotruding umbilicus, P rotuberant tongue C , and P oor brain development.
Subacute granulomatous thyroiditis (de Quervain)	Self-limited disease often following a flu-like illness (eg, viral infection). May be hyperthyroid early in course, followed by hypothyroidism (permanent in ~15% of cases). Histology: granulomatous inflammation. Findings: ↑ ESR, jaw pain, very tender thyroid. (de Quervain is associated with pain .)
Riedel thyroiditis	Thyroid replaced by fibrous tissue and inflammatory infiltrate D . Fibrosis may extend to local structures (eg, trachea, esophagus), mimicking anaplastic carcinoma. 1/3 of patients are hypothyroid. Considered a manifestation of IgG ₄ -related systemic disease (eg, autoimmune pancreatitis, retroperitoneal fibrosis, noninfectious aortitis). Findings: fixed, hard (rock-like), painless goiter.
Other causes	Iodine deficiency (with goiter E), goitrogens (eg, amiodarone, lithium), Wolff-Chaikoff effect (thyroid gland downregulation in response to ↑ iodide).



Hyperthyroidism**Graves disease**

Most common cause of hyperthyroidism. Thyroid-stimulating immunoglobulin (IgG, can cause transient neonatal hyperthyroidism; type II hypersensitivity) stimulates TSH receptors on thyroid (hyperthyroidism, diffuse goiter), dermal fibroblasts (pretibial myxedema), and orbital fibroblasts (Graves orbitopathy). Activation of T-cells → lymphocytic infiltration of retroorbital space → ↑ cytokines (eg, TNF- α , IFN- γ) → ↑ fibroblast secretion of hydrophilic GAGs → ↑ osmotic muscle swelling, muscle inflammation, and adipocyte count → exophthalmos **A**. Often presents during stress (eg, pregnancy). Associated with HLA-DR3 and HLA-B8. Histology: tall, crowded follicular epithelial cells; scalloped colloid.

Toxic multinodular goiter

Focal patches of hyperfunctioning follicular cells distended with colloid working independently of TSH (due to TSH receptor mutations in 60% of cases). ↑ release of T₃ and T₄. Hot nodules are rarely malignant.

Thyroid storm


Uncommon but serious complication that occurs when hyperthyroidism is incompletely treated/untreated and then significantly worsens in the setting of acute stress such as infection, trauma, surgery. Presents with agitation, delirium, fever, diarrhea, coma, and tachyarrhythmia (cause of death). May see ↑ LFTs. Treat with the **4 P**'s: β -blockers (eg, **P**ropranolol), **P**ropylthiouracil, corticosteroids (eg, **P**rednisolone), **P**otassium iodide (Lugol iodine). Iodide load → ↓ T₄ synthesis → Wolff-Chaikoff effect.

Jod-Basedow phenomenon

Iodine-induced hyperthyroidism. Occurs when a patient with iodine deficiency and partially autonomous thyroid tissue (eg, autonomous nodule) is made iodine replete. Can happen after iodine IV contrast or amiodarone use. Opposite to Wolff-Chaikoff effect.

Causes of goiter

Smooth/diffuse: Graves disease, Hashimoto thyroiditis, iodine deficiency, TSH-secreting pituitary adenoma.
Nodular: toxic multinodular goiter, thyroid adenoma, thyroid cancer, thyroid cyst.


Thyroid adenoma

Benign solitary growth of the thyroid. Most are nonfunctional (“cold”), can rarely cause hyperthyroidism via autonomous thyroid hormone production (“hot” or “toxic”). Most common histology is follicular (arrows in **A**); absence of capsular or vascular invasion (unlike follicular carcinoma).

Thyroid cancer

Typically diagnosed with fine needle aspiration; treated with thyroidectomy. Complications of surgery include hypocalcemia (due to removal of parathyroid glands), transection of recurrent laryngeal nerve during ligation of inferior thyroid artery (leads to dysphagia and dysphonia [hoarseness]), and injury to the external branch of the superior laryngeal nerve during ligation of superior thyroid vascular pedicle (may lead to loss of tenor usually noticeable in professional voice users).

Papillary carcinoma




Most common, excellent prognosis. Empty-appearing nuclei with central clearing (“Orphan Annie” eyes) **A**, psammoma bodies, nuclear grooves (**Papi** and **Moma** adopted **Orphan Annie**). ↑ risk with *RET/PTC* rearrangements and *BRAF* mutations, childhood irradiation. Papillary carcinoma: most **P**revalent, **P**alpable lymph nodes. Good prognosis.

Follicular carcinoma

Good prognosis. Invades thyroid capsule and vasculature (unlike follicular adenoma), uniform follicles; hematogenous spread is common. Associated with *RAS* mutation and *PAX8-PPAR-γ* translocations.

Medullary carcinoma




From parafollicular “C cells”; produces calcitonin, sheets of polygonal cells in an amyloid stroma **B** (stains with Congo red). Associated with MEN 2A and 2B (*RET* mutations).

Undifferentiated/anaplastic carcinoma

Older patients; presents with rapidly enlarging neck mass → compressive symptoms (eg, dyspnea, dysphagia, hoarseness); very poor prognosis. Associated with *TP53* mutation.

Diagnosing parathyroid disease



Hypoparathyroidism

Due to injury to parathyroid glands or their blood supply (usually during surgery), autoimmune destruction, or DiGeorge syndrome. Findings: tetany, hypocalcemia, hyperphosphatemia.

Chvostek sign—tapping of facial nerve (tap the **C**heek) → contraction of facial muscles.

Trousseau sign—occlusion of brachial artery with BP cuff (cuff the **T**riceps) → carpal spasm.

Pseudohypoparathyroidism type 1A—autosomal dominant, maternally transmitted mutations (imprinted *GNAS* gene). *GNAS1*-inactivating mutation (coupled to PTH receptor) that encodes the G_s protein α subunit → inactivation of adenylate cyclase when PTH binds to its receptor → end-organ resistance (kidney and bone) to PTH.

Physical findings: Albright hereditary osteodystrophy (shortened 4th/5th digits **A**, short stature, round face, subcutaneous calcifications, developmental delay).

Labs: ↑ PTH, ↓ Ca^{2+} , ↑ PO_4^{3-} .


Pseudopseudohypoparathyroidism—autosomal dominant, paternally transmitted mutations (imprinted *GNAS* gene) but without end-organ resistance to PTH due to normal maternal allele maintaining renal responsiveness to PTH.

Physical findings: same as Albright hereditary osteodystrophy.

Labs: normal PTH, Ca^{2+} , PO_4^{3-} .

Lab values in hypocalcemia

DISORDER	Ca^{2+}	PO_4^{3-}	PTH
Vitamin D deficiency	↓	↓	↑
Hypoparathyroidism	↓	↑	↓
2° hyperparathyroidism (CKD)	↓	↑	↑
Pseudohypoparathyroidism	↓	↑	↑
Hyperphosphatemia	↓	↑	↑

Hyperparathyroidism**Primary hyperparathyroidism**

Usually due to parathyroid adenoma or hyperplasia. **Hypercalcemia**, hypercalciuria (renal **stones**), polyuria (**thrones**), hypophosphatemia, \uparrow PTH, \uparrow ALP, \uparrow urinary cAMP. Most often asymptomatic. May present with **bone** pain, weakness, constipation (“**groans**”), abdominal/flank pain (kidney stones, acute pancreatitis), neuropsychiatric disturbances (“**psychiatric overtones**”).

Osteitis fibrosa cystica—cystic **bone** spaces filled with brown fibrous tissue **A** (“brown tumor” consisting of osteoclasts and deposited hemosiderin from hemorrhages; causes bone pain). Due to \uparrow PTH, classically associated with 1° (but also seen with 2°) hyperparathyroidism.

“**Stones, thrones, bones, groans, and psychiatric overtones.**”

Secondary hyperparathyroidism

2° hyperplasia due to \downarrow Ca^{2+} absorption and/or \uparrow PO_4^{3-} , most often in chronic kidney disease (causes hypovitaminosis D and hyperphosphatemia \rightarrow \downarrow Ca^{2+}).

Hypocalcemia, hyperphosphatemia in chronic kidney disease (vs hypophosphatemia with most other causes), \uparrow ALP, \uparrow PTH.

Renal osteodystrophy—renal disease \rightarrow 2° and 3° hyperparathyroidism \rightarrow bone lesions.

Tertiary hyperparathyroidism

Refractory (autonomous) hyperparathyroidism resulting from chronic kidney disease. $\uparrow\uparrow$ PTH, \uparrow Ca^{2+} .


Familial hypocalciuric hypercalcemia

Defective G-coupled Ca^{2+} -sensing receptors in multiple tissues (eg, parathyroids, kidneys). Higher than normal Ca^{2+} levels required to suppress PTH. Excessive renal Ca^{2+} reabsorption \rightarrow mild hypercalcemia and hypocalciuria with normal to \uparrow PTH levels.

Diabetes mellitus

ACUTE MANIFESTATIONS	Polydipsia, polyuria, polyphagia, weight loss, DKA (type 1), hyperosmolar hyperglycemic state (type 2). Rarely, can be caused by unopposed secretion of GH and epinephrine. Also seen in patients on glucocorticoid therapy (steroid diabetes).		
CHRONIC COMPLICATIONS	Nonenzymatic glycation: <ul style="list-style-type: none"> Small vessel disease (diffuse thickening of basement membrane) → retinopathy (hemorrhage, exudates, microaneurysms, vessel proliferation), glaucoma, nephropathy. Nodular glomerulosclerosis → progressive proteinuria (initially microalbuminuria; ACE inhibitors and ARBs are renoprotective) and arteriosclerosis (causing hypertension) → chronic kidney disease. Large vessel atherosclerosis, CAD, peripheral vascular occlusive disease, gangrene → limb loss, cerebrovascular disease. MI most common cause of death. Osmotic damage (sorbitol accumulation in organs with aldose reductase and ↓ or absent sorbitol dehydrogenase): <ul style="list-style-type: none"> Neuropathy (motor, sensory [glove and stocking distribution], and autonomic degeneration). Cataracts. 		
DIAGNOSIS	TEST	DIAGNOSTIC CUTOFF	NOTES
	HbA _{1c}	≥ 6.5%	Reflects average blood glucose over prior 3 months
	Fasting plasma glucose	≥ 126 mg/dL	Fasting for > 8 hours
	2-hour oral glucose tolerance test	≥ 200 mg/dL	2 hours after consumption of 75 g of glucose in water

Insulin deficiency or severe insulin insensitivity



Type 1 vs type 2 diabetes mellitus

	Type 1	Type 2
1° DEFECT	Autoimmune T-cell–mediated destruction of β cells (eg, due to presence of glutamic acid decarboxylase antibodies)	\uparrow resistance to insulin, progressive pancreatic β -cell failure
INSULIN NECESSARY IN TREATMENT	Always	Sometimes
AGE (EXCEPTIONS COMMON)	< 30 yr	> 40 yr
ASSOCIATION WITH OBESITY	No	Yes
GENETIC PREDISPOSITION	Relatively weak (50% concordance in identical twins), polygenic	Relatively strong (90% concordance in identical twins), polygenic
ASSOCIATION WITH HLA SYSTEM	Yes, HLA-DR4 and -DR3 (4 – 3 = type 1)	No
GLUCOSE INTOLERANCE	Severe	Mild to moderate
INSULIN SENSITIVITY	High	Low
KETOACIDOSIS	Common	Rare
β -CELL NUMBERS IN THE ISLETS	\downarrow	Variable (with amyloid deposits)
SERUM INSULIN LEVEL	\downarrow	\uparrow initially, but \downarrow in advanced disease
CLASSIC SYMPTOMS OF POLYURIA, POLYDIPSIA, POLYPHAGIA, WEIGHT LOSS	Common	Sometimes
HISTOLOGY	Islet leukocytic infiltrate	Islet amyloid polypeptide (IAPP) deposits

Diabetic ketoacidosis	Insulin absent, ketones present (\rightarrow complications). Insulin noncompliance or \uparrow requirements from \uparrow stress (eg, infection) \rightarrow excess fat breakdown and \uparrow ketogenesis from \uparrow free fatty acids \rightarrow ketone bodies (β -hydroxybutyrate > acetoacetate).
SIGNS/SYMPTOMS	DKA is Deadly: Delirium/psychosis, Kussmaul respirations (rapid, deep breathing), Abdominal pain/nausea/vomiting, Dehydration . Fruity breath odor (due to exhaled acetone).
LABS	Hyperglycemia, \uparrow H ⁺ , \downarrow HCO ₃ ⁻ (\uparrow anion gap metabolic acidosis), \uparrow urine and blood ketone levels, leukocytosis. Normal/ \uparrow serum K ⁺ , but depleted intracellular K ⁺ due to transcellular shift from \downarrow insulin and acidosis. Osmotic diuresis \rightarrow \uparrow K ⁺ loss in urine \rightarrow total body K ⁺ depletion.
COMPLICATIONS	Life-threatening mucormycosis, cerebral edema, cardiac arrhythmias, HF.
TREATMENT	IV fluids, IV insulin, K ⁺ (to replete intracellular stores) +/- glucose to prevent hypoglycemia.

Hyperosmolar hyperglycemic state	Insulin present, ketones absent. Profound hyperglycemia \rightarrow excessive osmotic diuresis \rightarrow dehydration and \uparrow serum osmolality \rightarrow HHS. Classically seen in elderly type 2 diabetics with limited ability to drink.
SIGNS/SYMPTOMS	Thirst, polyuria, lethargy, focal neurologic deficits, seizures.
LABS	Hyperglycemia (often >600 mg/dL), \uparrow serum osmolality (> 320 mOsm/kg), normal pH (no acidosis), no ketones. Normal/ \uparrow serum K ⁺ , \downarrow intracellular K ⁺ .
COMPLICATIONS	Can progress to coma and death if untreated.
TREATMENT	IV fluids, IV insulin, and K ⁺ (to replete intracellular stores).

Cushing syndrome

ETIOLOGY

↑ cortisol due to a variety of causes:



- Exogenous corticosteroids → ↓ ACTH → bilateral adrenal atrophy. Most common cause.
- Primary adrenal adenoma, hyperplasia, or carcinoma → ↓ ACTH → atrophy of uninvolved adrenal gland.
- ACTH-secreting pituitary adenoma (Cushing disease); paraneoplastic ACTH secretion (eg, small cell lung cancer, bronchial carcinoids) → bilateral adrenal hyperplasia. Cushing disease is responsible for the majority of endogenous cases of Cushing syndrome.

FINDINGS

CUSHING Syndrome: ↑ **C**holesterol, ↑ **U**rinary free cortisol, **S**kin changes (thinning, striae **A**), **H**ypertension, **I**mmunosuppression, **N**eoplasm (a cause, not a finding), **G**rowth retardation (in children), ↑ **S**ugar (hyperglycemia, insulin resistance). Also, amenorrhea, moon facies **B**, buffalo hump, osteoporosis, ↑ weight (truncal obesity), hirsutism.

DIAGNOSIS

Screening tests include: ↑ free cortisol on 24-hr urinalysis, ↑ late night salivary cortisol, and no suppression with overnight low-dose dexamethasone test.





Nelson syndrome

Enlargement of pre-existing ACTH-secreting pituitary adenoma after bilateral adrenalectomy for refractory Cushing disease → ↑ ACTH (hyperpigmentation), mass effect (headaches, bitemporal hemianopia).

Treatment: transsphenoidal resection, postoperative pituitary irradiation for residual tumor.

Adrenal insufficiency

Inability of adrenal glands to generate enough glucocorticoids +/- mineralocorticoids for the body's needs. Symptoms include weakness, fatigue, orthostatic hypotension, muscle aches, weight loss, GI disturbances, sugar and/or salt cravings. Treatment: glucocorticoid/mineralocorticoid replacement.

**Primary adrenal insufficiency**

↓ gland function → ↓ cortisol, ↓ aldosterone → hypotension (hyponatremic volume contraction), hyperkalemia, metabolic acidosis, skin/mucosal hyperpigmentation
A (↑ melanin synthesis due to ↑ MSH, a byproduct of ACTH production from POMC).
 ▪ **Acute**—sudden onset (eg, due to massive hemorrhage). May present with shock in acute adrenal crisis.
 ▪ **Chronic**—**Addison disease**. Due to adrenal atrophy or destruction by disease (autoimmune destruction most common in the Western world; TB most common in the developing world).

Primary **P**igments the skin/mucosa.

Associated with autoimmune polyglandular syndromes.

Waterhouse-Friderichsen syndrome—acute 1° adrenal insufficiency due to adrenal hemorrhage associated with septicemia (usually *Neisseria meningitidis*), DIC, endotoxic shock.

Secondary adrenal insufficiency

Seen with ↓ pituitary ACTH production. No skin/mucosal hyperpigmentation (ACTH is not elevated), no hyperkalemia (aldosterone synthesis preserved due to functioning adrenal gland, intact RAAS).

Secondary **S**pares the skin/mucosa.

Tertiary adrenal insufficiency

Seen in patients with chronic exogenous steroid use, precipitated by abrupt withdrawal. Aldosterone synthesis unaffected.

Tertiary from **T**reatment.

Hyperaldosteronism

Increased secretion of aldosterone from adrenal gland. Clinical features include hypertension, ↓ or normal K⁺, metabolic alkalosis. 1° hyperaldosteronism does not directly cause edema due to aldosterone escape mechanism. However, certain 2° causes of hyperaldosteronism (eg, heart failure) impair the aldosterone escape mechanism, leading to worsening of edema.

Primary hyperaldosteronism

Seen with adrenal adenoma (Conn syndrome) or bilateral adrenal hyperplasia. ↑ aldosterone, ↓ renin. Leads to treatment-resistant hypertension.

Secondary hyperaldosteronism


Seen in patients with renovascular hypertension, juxtaglomerular cell tumors (renin-producing), and edema (eg, cirrhosis, heart failure, nephrotic syndrome).

Neuroendocrine tumors

Heterogeneous group of neoplasms originating from neuroendocrine cells (which have traits similar to nerve cells and hormone-producing cells).

Most neoplasms occur in the GI system (eg, carcinoid, gastrinoma), pancreas (eg, insulinoma, glucagonoma), and lungs (eg, small cell carcinoma). Also in thyroid (eg, medullary carcinoma) and adrenals (eg, pheochromocytoma).


Neuroendocrine cells (eg, pancreatic β cells, enterochromaffin cells) share a common biologic function through amine precursor uptake decarboxylase (APUD) despite differences in embryologic origin, anatomic site, and secretory products (eg, chromogranin A, neuron-specific enolase [NSE], synaptophysin, serotonin, histamine, calcitonin). Treatment: surgical resection, somatostatin analogs.

Neuroblastoma

Most common tumor of the adrenal medulla in **children**, usually < 4 years old. Originates from **N**eural crest cells. Occurs anywhere along the sympathetic chain.

Most common presentation is abdominal distension and a firm, irregular mass that can cross the midline (vs Wilms tumor, which is smooth and unilateral). Less likely to develop hypertension than with pheochromocytoma (**N**euroblastoma is **N**ormotensive). Can also present with opsoclonus-myoclonus syndrome (“dancing eyes-dancing feet”).

↑ HVA and VMA (catecholamine metabolites) in urine. Homer-Wright rosettes (neuroblasts surrounding a central lumen **A**) characteristic of neuroblastoma and medulloblastoma. Bombesin and **NSE** ⊕. Associated with amplification of **N**-myc oncogene.

Pheochromocytoma**ETIOLOGY**

Most common tumor of the adrenal medulla in **adults** **A**. Derived from chromaffin cells (arise from neural crest).

May be associated with germline mutations (eg, *NF-1*, *VHL*, *RET* [MEN 2A, 2B]).

Rule of 10's:

10% malignant

10% bilateral

10% extra-adrenal (eg, bladder wall, organ of Zuckerkandl)

10% calcify

10% kids

SYMPTOMS

Most tumors secrete epinephrine, norepinephrine, and dopamine, which can cause episodic hypertension. May also secrete EPO → polycythemia.

Symptoms occur in “spells”—relapse and remit.

Episodic hyperadrenergic symptoms (**5 P's**):

Pressure (↑ BP)

Pain (headache)

Perspiration

Palpitations (tachycardia)

Pallor

FINDINGS

↑ catecholamines and metanephrines (eg, homovanillic acid, vanillylmandelic acid) in urine and plasma.

Chromogranin, synaptophysin and **NSE** ⊕.

TREATMENT

Irreversible α -antagonists (eg, phenoxybenzamine) followed by β -blockers prior to tumor resection. α -blockade must be achieved before giving β -blockers to avoid a hypertensive crisis. **A** before **B**.

Phenoxybenzamine for **pheochromocytoma**.

Multiple endocrine neoplasias

All **MEN** syndromes have autosomal **dominant** inheritance.
 “All **MEN** are **dominant**” (or so they think).

SUBTYPE	CHARACTERISTICS	COMMENTS
MEN 1	<p>Pituitary tumors (prolactin or GH)</p> <p>Pancreatic endocrine tumors—Zollinger-Ellison syndrome, insulinomas, VIPomas, glucagonomas (rare)</p> <p>Parathyroid adenomas</p> <p>Associated with mutation of <i>MEN1</i> (menin, a tumor suppressor, chromosome 11), angiofibromas, collagenomas, meningiomas</p>	
MEN 2A	<p>Parathyroid hyperplasia</p> <p>Medullary thyroid carcinoma—neoplasm of parafollicular C cells; secretes calcitonin; prophylactic thyroidectomy required</p> <p>Pheochromocytoma (secretes catecholamines)</p> <p>Associated with mutation in <i>RET</i> (codes for receptor tyrosine kinase)</p>	
MEN 2B	<p>Medullary thyroid carcinoma</p> <p>Pheochromocytoma</p> <p>Mucosal neuromas A (oral/intestinal ganglioneuromatosis)</p> <p>Associated with marfanoid habitus; mutation in <i>RET</i> gene</p>	




MEN 1 = 3 P's: **P**ituitary, **P**arathyroid, and **P**ancreas

MEN 2A = 2 P's: **P**arathyroid and **P**heochromocytoma

MEN 2B = 1 P: **P**heochromocytoma

Pancreatic islet cell tumors

Insulinoma	<p>Tumor of pancreatic β cells \rightarrow overproduction of insulin \rightarrow hypoglycemia.</p> <p>May see Whipple triad: low blood glucose, symptoms of hypoglycemia (eg, lethargy, syncope, diplopia), and resolution of symptoms after normalization of plasma glucose levels. Symptomatic patients have \downarrow blood glucose and \uparrow C-peptide levels (vs exogenous insulin use). \sim 10% of cases associated with MEN 1 syndrome.</p> <p>Treatment: surgical resection.</p>
Glucagonoma	<p>Tumor of pancreatic α cells \rightarrow overproduction of glucagon.</p> <p>Presents with 6 D's: Dermatitis (necrolytic migratory erythema), Diabetes (hyperglycemia), DVT, Declining weight, Depression, Diarrhea.</p> <p>Treatment: octreotide, surgical resection.</p>
Somatostatinoma	<p>Tumor of pancreatic δ cells \rightarrow overproduction of somatostatin \rightarrow \downarrow secretion of secretin, cholecystokinin, glucagon, insulin, gastrin, gastric inhibitory peptide (GIP).</p> <p>May present with diabetes/glucose intolerance, steatorrhea, gallstones, achlorhydria.</p> <p>Treatment: surgical resection; somatostatin analogs (eg, octreotide) for symptom control.</p>

Carcinoid syndrome

Carcinoid tumors arise from neuroendocrine cells most commonly in the intestine or lung. Rare and does not occur if tumor is limited to the GI tract.

Prominent rosettes (arrow in **A**), chromogranin A \oplus and synaptophysin \oplus .

Neuroendocrine cells secrete 5-HT \rightarrow recurrent diarrhea, wheezing, right-sided valvular heart disease (eg, tricuspid regurgitation, pulmonic stenosis), niacin deficiency (pellagra). 5-HT undergoes hepatic first-pass metabolism and enzymatic breakdown by MAO in the lung.

Treatment: surgical resection, somatostatin analog (eg, octreotide, telotristat) for symptom control.

Rule of thirds:

- 1/3 metastasize
- 1/3 present with 2nd malignancy
- 1/3 are multiple

Zollinger-Ellison syndrome

Gastrin-secreting tumor (gastrinoma) of pancreas or duodenum. Acid hypersecretion causes recurrent ulcers in duodenum and jejunum. Presents with abdominal pain (peptic ulcer disease, distal ulcers), diarrhea (malabsorption). Positive secretin stimulation test: gastrin levels remain elevated after administration of secretin, which normally inhibits gastrin release. May be associated with MEN 1.

► ENDOCRINE—PHARMACOLOGY**Diabetes mellitus therapy**

All patients with diabetes mellitus should receive education on diet, exercise, blood glucose monitoring, and complication management. Treatment differs based on the type of diabetes and glycemic control:

- Type 1 DM—insulin replacement
- Type 2 DM—oral agents (metformin is first line), non-insulin injectables, insulin replacement; weight loss particularly helpful in lowering blood glucose
- Gestational DM—insulin replacement if nutrition therapy and exercise alone fail

Regular (short-acting) insulin is preferred for DKA (IV), hyperkalemia (+ glucose), stress hyperglycemia.


To **N**ormalize **P**ancreatic **F**unction (-gliTs, -gliNs, -gliPs, -gliFs).

DRUG CLASS**MECHANISM****ADVERSE EFFECTS****Insulin preparations**

Rapid acting (1-hr peak): **L**ispro, **A**spart, **G**lulisine (no **LAG**)
 Short acting (2–3 hr peak): regular
 Intermediate acting (4–10 hr peak): NPH
 Long acting (no real peak): detemir, glargine

Bind insulin receptor (tyrosine kinase activity)
 Liver: \uparrow glucose storage as glycogen
 Muscle: \uparrow glycogen, protein synthesis
 Fat: \uparrow TG storage
 Cell membrane: \uparrow K⁺ uptake

Hypoglycemia, lipodystrophy, hypersensitivity reactions (rare), weight gain



Diabetes mellitus therapy (continued)

DRUG CLASS	MECHANISM	ADVERSE EFFECTS
Increase insulin sensitivity		
Biguanides Metformin	Inhibit mGPD → inhibition of hepatic gluconeogenesis and the action of glucagon. ↑ glycolysis, peripheral glucose uptake (↑ insulin sensitivity).	GI upset, lactic acidosis (use with caution in renal insufficiency), vitamin B ₁₂ deficiency. Weight loss (often desired).
Glitazones/ thiazolidinediones “-gliTs” Pioglitazone, rosiglitazone	Activate PPAR-γ (a nuclear receptor) → ↑ insulin sensitivity and levels of adiponectin → regulation of glucose metabolism and fatty acid storage.	Weight gain, edema, HF, ↑ risk of fractures. Delayed onset of action (several weeks). Rosiglitazone: ↑ risk of MI, cardiovascular death.
Increase insulin secretion		
Sulfonylureas (1st gen) Chlorpropamide, tolbutamide		Disulfiram-like reaction (FIRst-generation only). Rarely used.
Sulfonylureas (2nd gen) Glipizide, glyburide	Close K ⁺ channels in pancreatic B cell membrane → cell depolarizes → insulin release via ↑ Ca ²⁺ influx.	Hypoglycemia (↑ risk in renal insufficiency), weight gain.
Meglitinides “-gliNs” Nateglinide, repaglinide		
Increase glucose-induced insulin secretion		
GLP-1 analogs Exenatide, liraglutide	↓ glucagon release, ↓ gastric emptying, ↑ glucose-dependent insulin release.	Nausea, vomiting, pancreatitis. Weight loss (often desired). ↑ satiety (often desired).
DPP-4 inhibitors “-gliPs” Linagliptin, saxagliptin, sitagliptin	Inhibit DPP-4 enzyme that deactivates GLP-1 → ↓ glucagon release, ↓ gastric emptying. ↑ glucose-dependent insulin release.	Respiratory and urinary infections, weight neutral. ↑ satiety (often desired).
Decrease glucose absorption		
Sodium-glucose co-transporter 2 (SGLT2) inhibitors “-gliFs” Canagliflozin, dapagliflozin, empagliflozin	Block reabsorption of glucose in proximal convoluted tubule.	Glucosuria (UTIs, vulvovaginal candidiasis), dehydration (orthostatic hypotension), hyperkalemia, weight loss. Use with caution in renal insufficiency (↓ efficacy with ↓ GFR).
α-glucosidase inhibitors Acarbose, miglitol	Inhibit intestinal brush-border α-glucosidases → delayed carbohydrate hydrolysis and glucose absorption → ↓ postprandial hyperglycemia.	GI upset, bloating. Not recommended in renal insufficiency.
Others		
Amylin analogs Pramlintide	↓ glucagon release, ↓ gastric emptying.	Hypoglycemia, nausea. ↑ satiety (often desired).

Thionamides

Propylthiouracil, methimazole.

MECHANISM	Block thyroid peroxidase, inhibiting the oxidation of iodide as well as the organification and coupling of iodine → inhibition of thyroid hormone synthesis. PTU also blocks 5'-deiodinase → ↓ Peripheral conversion of T ₄ to T ₃ .
CLINICAL USE	Hyperthyroidism. PTU used in first trimester of pregnancy (due to methimazole teratogenicity); methimazole used in second and third trimesters of pregnancy (due to risk of PTU-induced hepatotoxicity). Not used to treat Graves ophthalmopathy (treated with corticosteroids).
ADVERSE EFFECTS	Skin rash, agranulocytosis (rare), aplastic anemia, hepatotoxicity. Methimazole is a possible teratogen (can cause aplasia cutis).

Levothyroxine, liothyronine

MECHANISM	Hormone replacement for T ₄ (levothyroxine) or T ₃ (liothyronine).
CLINICAL USE	Hypothyroidism, myxedema. May be abused for weight loss. Distinguish exogenous hyperthyroidism from endogenous hyperthyroidism by using a combination of TSH receptor antibodies, radioactive iodine uptake, and/or measurement of thyroid blood flow on ultrasound.
ADVERSE EFFECTS	Tachycardia, heat intolerance, tremors, arrhythmias.

Hypothalamic/pituitary drugs

DRUG	CLINICAL USE
Conivaptan, tolvaptan	ADH antagonists SIADH (block action of ADH at V ₂ -receptor)
Demeclocycline	ADH antagonist, a tetracycline SIADH
Desmopressin	Central DI, von Willebrand disease, sleep enuresis, hemophilia A
GH	GH deficiency, Turner syndrome
Oxytocin	Induction of labor (stimulates uterine contractions), control uterine hemorrhage
Somatostatin (octreotide)	Acromegaly, carcinoid syndrome, gastrinoma, glucagonoma, esophageal varices

Fludrocortisone

MECHANISM	Synthetic analog of aldosterone with little glucocorticoid effects.
CLINICAL USE	Mineralocorticoid replacement in 1° adrenal insufficiency.
ADVERSE EFFECTS	Similar to glucocorticoids; also edema, exacerbation of heart failure, hyperpigmentation.

Cinacalcet

MECHANISM	Sensitizes Ca ²⁺ -sensing receptor (CaSR) in parathyroid gland to circulating Ca ²⁺ → ↓ PTH.
CLINICAL USE	2° hyperparathyroidism in patients with CKD receiving hemodialysis, hypercalcemia in 1° hyperparathyroidism (if parathyroidectomy fails), or in parathyroid carcinoma.
ADVERSE EFFECTS	Hypocalcemia.

Sevelamer

MECHANISM	Nonabsorbable phosphate binder that prevents phosphate absorption from the GI tract.
CLINICAL USE	Hyperphosphatemia in CKD.
ADVERSE EFFECTS	Hypophosphatemia, GI upset.

HIGH-YIELD SYSTEMS

Gastrointestinal

“A good set of bowels is worth more to a man than any quantity of brains.”

—Josh Billings

“Man should strive to have his intestines relaxed all the days of his life.”

—Moses Maimonides

“All right, let’s not panic. I’ll make the money by selling one of my livers. I can get by with one.”

—Homer Simpson

When studying the gastrointestinal system, be sure to understand the normal embryology, anatomy, and physiology and how it is affected in the various pathologic diseases. Study not only what a disease entails, but also its specific findings, so that you can differentiate between two similar diseases. For example, what specifically makes ulcerative colitis different than Crohn disease? Also, it is important to understand bile metabolism and which lab values increase or decrease depending on the disease process. Be comfortable with basic interpretation of abdominal x-rays, CT scans, and endoscopic images.

▶ Embryology	358
▶ Anatomy	360
▶ Physiology	371
▶ Pathology	376
▶ Pharmacology	398

▶ GASTROINTESTINAL—EMBRYOLOGY

Normal gastrointestinal embryology

Foregut—esophagus to duodenum at level of pancreatic duct and common bile duct insertion (ampulla of Vater).
 Midgut—lower duodenum to proximal 2/3 of transverse colon.
 Hindgut—distal 1/3 of transverse colon to anal canal above pectinate line.
 Midgut development:

- 6th week—physiologic herniation of midgut through umbilical ring
- 10th week—returns to abdominal cavity + rotates around superior mesenteric artery (SMA), total 270° counterclockwise

Ventral wall defects

Developmental defects due to failure of rostral fold closure (eg, sternal defects [ectopia cordis]), lateral fold closure (eg, omphalocele, gastroschisis), or caudal fold closure (eg, bladder exstrophy).

Gastroschisis

Omphalocele

ETIOLOGY

Extrusion of abdominal contents through abdominal folds (typically right of umbilicus)

Failure of lateral walls to migrate at umbilical ring → persistent midline herniation of abdominal contents into umbilical cord

COVERAGE

Not covered by peritoneum or amnion **A**; “the guts come out of the gap (**schism**) in the letter **G**”

Surrounded by peritoneum **B** (light gray shiny sac); “abdominal contents are **sealed** in the letter **O**”

ASSOCIATIONS

Not associated with chromosome abnormalities; favorable prognosis

Associated with congenital anomalies (eg, trisomies 13 and 18, Beckwith-Wiedemann syndrome) and other structural abnormalities (eg, cardiac, GU, neural tube)




Congenital umbilical hernia

Failure of umbilical ring to close after physiologic herniation of the midgut. Small defects usually close spontaneously.


Tracheoesophageal anomalies

Esophageal atresia (EA) with distal tracheoesophageal fistula (TEF) is the most common (85%) and often presents as polyhydramnios in utero (due to inability of fetus to swallow amniotic fluid). Neonates drool, choke, and vomit with first feeding. TEFs allow air to enter stomach (visible on CXR). Cyanosis is 2° to laryngospasm (to avoid reflux-related aspiration). Clinical test: failure to pass nasogastric tube into stomach.

In **H**-type, the fistula resembles the letter **H**. In pure EA, CXR shows gasless abdomen.



Intestinal atresia




Presents with bilious vomiting and abdominal distension within first 1–2 days of life.

Duodenal atresia—failure to recanalize. Abdominal x-ray **A** shows “double bubble” (dilated stomach, proximal duodenum). Associated with **D**own syndrome.

Jejunal and ileal atresia—disruption of mesenteric vessels (typically SMA) → ischemic necrosis of fetal intestine → segmental resorption: bowel becomes discontinuous. X-ray shows dilated loops of small bowel with air-fluid levels.

Hypertrophic pyloric stenosis




Most common cause of gastric outlet obstruction in infants (1:600). Palpable olive-shaped mass in epigastric region, visible peristaltic waves, and nonbilious projectile vomiting at ~ 2–6 weeks old. More common in firstborn males; associated with exposure to macrolides.

Results in hypokalemic hypochloremic metabolic alkalosis (2° to vomiting of gastric acid and subsequent volume contraction).

Ultrasound shows thickened and lengthened pylorus **A**.

Treatment: surgical incision of pyloric muscles (pyloromyotomy).

Pancreas and spleen embryology




Pancreas—derived from foregut. Ventral pancreatic bud contributes to uncinate process and main pancreatic duct. The dorsal pancreatic bud alone becomes the body, tail, isthmus, and accessory pancreatic duct. Both the ventral and dorsal buds contribute to pancreatic head.

Annular pancreas—abnormal rotation of ventral pancreatic bud forms a ring of pancreatic tissue → encircles 2nd part of duodenum; may cause duodenal narrowing (arrows in **A**) and vomiting.

Pancreas divisum—ventral and dorsal parts fail to fuse at 8 weeks. Common anomaly; mostly asymptomatic, but may cause chronic abdominal pain and/or pancreatitis.

Spleen—arises in mesentery of stomach (hence is mesodermal) but has foregut supply (celiac trunk → splenic artery).





► GASTROINTESTINAL—ANATOMY

Retroperitoneal structures


Retroperitoneal structures **A** are posterior to (and outside of) the peritoneal cavity. Injuries to retroperitoneal structures can cause blood or gas accumulation in retroperitoneal space.

SAD PUCKER:

- Suprarenal (adrenal) glands [not shown]
- Aorta and IVC
- Duodenum (2nd through 4th parts)
- Pancreas (except tail)
- Ureters [not shown]
- Colon (descending and ascending)
- Kidneys
- Esophagus (thoracic portion) [not shown]
- Rectum (partially) [not shown]



Important gastrointestinal ligaments



LIGAMENT	CONNECTS	STRUCTURES CONTAINED	NOTES
Falciform ligament	Liver to anterior abdominal wall	Ligamentum teres hepatis (derivative of fetal umbilical vein), patent paraumbilical veins	Derivative of ventral mesentery
Hepatoduodenal ligament	Liver to duodenum	Portal triad: proper hepatic artery, portal vein, common bile duct	Derivative of ventral mesentery Pringle maneuver—ligament is compressed manually or with a vascular clamp in omental foramen to control bleeding from hepatic inflow source Borders the omental foramen, which connects the greater and lesser sacs Part of lesser omentum
Gastrohepatic ligament	Liver to lesser curvature of stomach	Gastric vessels	Derivative of ventral mesentery Separates greater and lesser sacs on the right May be cut during surgery to access lesser sac Part of lesser omentum
Gastrocolic ligament (not shown)	Greater curvature and transverse colon	Gastroepiploic arteries	Derivative of dorsal mesentery Part of greater omentum
Gastrosplenic ligament	Greater curvature and spleen	Short gastrics, left gastroepiploic vessels	Derivative of dorsal mesentery Separates greater and lesser sacs on the left Part of greater omentum
Splenorenal ligament	Spleen to left pararenal space	Splenic artery and vein, tail of pancreas	Derivative of dorsal mesentery

Digestive tract anatomy


Layers of gut wall **A** (inside to outside—**MSMS**):

- **M**ucosa—epithelium, lamina propria, muscularis mucosa
- **S**ubmucosa—includes **S**ubmucosal nerve plexus (Meissner), **S**ecretes fluid
- **M**uscularis externa—includes **M**yenteric nerve plexus (Auerbach), **M**otility
- **S**erosa (when intraperitoneal), adventitia (when retroperitoneal)

Ulcers can extend into submucosa, inner or outer muscular layer. Erosions are in mucosa only.

Frequency of basal electric rhythm (slow waves), which originate in the interstitial cells of Cajal:

- Stomach—3 waves/min
- Duodenum—12 waves/min
- Ileum—8–9 waves/min

**Digestive tract histology****Esophagus**

Nonkeratinized stratified squamous epithelium. Upper 1/3, striated muscle; middle and lower 2/3 smooth muscle, with some overlap at the transition.

Stomach

Gastric glands **A**.

Duodenum

Villi **B** and microvilli ↑ absorptive surface. Brunner glands (HCO_3^- -secreting cells of submucosa) and crypts of Lieberkühn (contain stem cells that replace enterocytes/goblet cells and Paneth cells that secrete defensins, lysozyme, and TNF).

Jejunum


Villi, crypts of Lieberkühn, and plicae circulares (also present in distal duodenum) **C**.

Ileum


Peyer patches (arrow in **D**; lymphoid aggregates in lamina propria, submucosa), plicae circulares (proximal ileum), and crypts of Lieberkühn. Largest number of goblet cells in the small intestine.

Colon

Crypts of Lieberkühn with abundant goblet cells, but no villi **E**.



Abdominal aorta and branches



Arteries supplying GI structures are single and branch anteriorly.


Arteries supplying non-GI structures are paired and branch laterally and posteriorly.

Two areas of the colon have dual blood supply from distal arterial branches (“watershed regions”) → susceptible in colonic ischemia:

- Splenic flexure—SMA and IMA
- Rectosigmoid junction—the last sigmoid arterial branch from the IMA and superior rectal artery

Nutcracker syndrome—compression of left renal vein between superior mesenteric artery and aorta. Characterized by abdominal (flank) pain and gross hematuria (from rupture of thin-walled renal varicosities).

Superior mesenteric artery syndrome—characterized by intermittent intestinal obstruction symptoms (primarily postprandial pain) when SMA and aorta compress transverse (third) portion of duodenum. Typically occurs in conditions associated with diminished mesenteric fat (eg, low body weight/malnutrition).



Gastrointestinal blood supply and innervation


EMBRYONIC GUT REGION	ARTERY	PARASYMPATHETIC INNERVATION	VERTEBRAL LEVEL	STRUCTURES SUPPLIED
Foregut	Celiac	Vagus	T12/L1	Pharynx (vagus nerve only) and lower esophagus (celiac artery only) to proximal duodenum; liver, gallbladder, pancreas, spleen (mesoderm)
Midgut	SMA	Vagus	L1	Distal duodenum to proximal 2/3 of transverse colon
Hindgut	IMA	Pelvic	L3	Distal 1/3 of transverse colon to upper portion of anal canal

Celiac trunk


Branches of celiac trunk: common hepatic, splenic, and left gastric. These constitute the main blood supply of the foregut.

Strong anastomoses exist between:

- Left and right gastroepiploics
- Left and right gastrics



Portosystemic anastomoses




SITE OF ANASTOMOSIS	CLINICAL SIGN	PORTAL ↔ SYSTEMIC
① Esophagus	Esophageal varices	Left gastric ↔ esophageal (drains into azygos)
② Umbilicus	Caput medusae	Paraumbilical ↔ small epigastric veins of the anterior abdominal wall.
③ Rectum	Anorectal varices	Superior rectal ↔ middle and inferior rectal

Varices of **gut**, **butt**, and **caput** (medusae) are commonly seen with portal hypertension.

④ Treatment with a **transjugular intrahepatic portosystemic shunt (TIPS)** between the portal vein and hepatic vein relieves portal hypertension by shunting blood to the systemic circulation, bypassing the liver. TIPS can precipitate hepatic encephalopathy due to ↓ clearance of ammonia from shunting.

Pectinate line

Also called dentate line. Formed where endoderm (hindgut) meets ectoderm.



Above pectinate line: internal hemorrhoids, adenocarcinoma.



Internal hemorrhoids receive visceral innervation and are therefore **not painful**.

Below pectinate line: external hemorrhoids, anal fissures, squamous cell carcinoma.

External hemorrhoids receive somatic innervation (inferior rectal branch of pudendal nerve) and are therefore **painful** if thrombosed.

Anal fissure—tear in anal mucosa below Pectinate line. **P**ain while **P**ooping; blood on toilet **P**aper. Located **P**osteriorly because this area is **P**oorly **P**erfused. Innervated by **P**udendal nerve. Associated with low-fiber diets and constipation.

Liver tissue architecture



The functional unit of the liver is made up of hexagonally arranged lobules surrounding the central vein with portal triads on the edges (consisting of a portal vein, hepatic artery, bile ducts, as well as lymphatics) **A**.

Apical surface of hepatocytes faces bile canaliculi. Basolateral surface faces sinusoids. Kupffer cells (specialized macrophages) located in sinusoids (black arrows in **B**; yellow arrows show hepatic venule) clear bacteria and damaged or senescent RBCs.

Hepatic stellate (Ito) cells in space of Disse store vitamin A (when quiescent) and produce extracellular matrix (when activated). Responsible for hepatic fibrosis.

Zone I—periportal zone:


- Affected 1st by viral hepatitis
- Best oxygenated, most resistant to circulatory compromise
- Ingested toxins (eg, cocaine)

Zone II—intermediate zone:


- Yellow fever

Zone III—pericentral vein (centrilobular) zone:

- Affected 1st by ischemia (least oxygenated)
- High concentration of cytochrome P-450
- Most sensitive to metabolic toxins (eg, ethanol, CCl₄, halothane, rifampin, acetaminophen)
- Site of alcoholic hepatitis




Biliary structures



Gallstones that reach the confluence of the common bile and pancreatic ducts at the ampulla of Vater can block both the common bile and pancreatic ducts (double duct sign), causing both cholangitis and pancreatitis, respectively.

Tumors that arise in head of pancreas (usually ductal adenocarcinoma) can cause obstruction of common bile duct → enlarged gallbladder with painless jaundice (Courvoisier sign).

Cholangiography shows filling defects in gallbladder (blue arrow) and cystic duct (red arrow) **A**.



Femoral region

ORGANIZATION

Lateral to medial: Nerve-Artery-Vein-Lymphatics.

You go from **lateral to medial** to find your **NAVeL**.


Femoral triangle

Contains femoral nerve, artery, vein.


Venous near the **penis**.

Femoral sheath


Fascial tube 3–4 cm below inguinal ligament. Contains femoral vein, artery, and canal (deep inguinal lymph nodes) but not femoral nerve.



Inguinal canal




Abdominal wall



Hernias


Protrusion of peritoneum through an opening, usually at a site of weakness. Contents may be at risk for incarceration (not reducible back into abdomen/pelvis) and strangulation (ischemia and necrosis). Complicated hernias can present with tenderness, erythema, fever.

Diaphragmatic hernia

Abdominal structures enter the thorax **A**; may occur due to congenital defect of pleuroperitoneal membrane or from trauma. Commonly occurs on left side due to relative protection of right hemidiaphragm by liver. Most commonly a **hiatal hernia**, in which stomach herniates upward through the esophageal hiatus of the diaphragm.


Sliding hiatal hernia—gastroesophageal junction is displaced upward as gastric cardia slides into hiatus; “hourglass stomach.” Most common type. Associated with GERD.

Paraesophageal hiatal hernia—gastroesophageal junction is usually normal but gastric fundus protrudes into the thorax.

**Indirect inguinal hernia**

Goes through the internal (deep) inguinal ring, external (superficial) inguinal ring, and into the groin. Enters internal inguinal ring lateral to inferior epigastric vessels. Caused by failure of processus vaginalis to close (can form hydrocele). May be noticed in **infants** or discovered in adulthood. Much more common in males **B**.

Follows the pathway of testicular descent. Covered by all 3 layers of spermatic fascia.


**Direct inguinal hernia**

Protrudes through inguinal (Hesselbach) triangle. Bulges directly through parietal peritoneum medial to the inferior epigastric vessels but lateral to the rectus abdominis. Goes through external (superficial) inguinal ring only. Covered by external spermatic fascia. Usually occurs in older men due to acquired weakness of transversalis fascia.


MDs don't LIe:

Medial to inferior epigastric vessels = **D**irect hernia.

Lateral to inferior epigastric vessels = **I**ndirect hernia.

**Femoral hernia**

Protrudes below inguinal ligament through femoral canal below and lateral to pubic tubercle. More common in **females**, but overall inguinal hernias are the most common. More likely to present with incarceration or strangulation (vs inguinal hernia).




► GASTROINTESTINAL—PHYSIOLOGY

Gastrointestinal regulatory substances


REGULATORY SUBSTANCE	SOURCE	ACTION	REGULATION	NOTES
Gastrin	G cells (antrum of stomach, duodenum)	<ul style="list-style-type: none"> ↑ gastric H⁺ secretion ↑ growth of gastric mucosa ↑ gastric motility 	<ul style="list-style-type: none"> ↑ by stomach distention/alkalinization, amino acids, peptides, vagal stimulation via gastrin-releasing peptide (GRP) ↓ by pH < 1.5 	<ul style="list-style-type: none"> ↑ by chronic PPI use ↑ in chronic atrophic gastritis (eg, <i>H pylori</i>) ↑↑ in Zollinger-Ellison syndrome (gastrinoma)
Somatostatin	D cells (pancreatic islets, GI mucosa)	<ul style="list-style-type: none"> ↓ gastric acid and pepsinogen secretion ↓ pancreatic and small intestine fluid secretion ↓ gallbladder contraction ↓ insulin and glucagon release 	<ul style="list-style-type: none"> ↑ by acid ↓ by vagal stimulation 	<ul style="list-style-type: none"> Inhibits secretion of various hormones (encourages somato-stasis) Octreotide is an analog used to treat acromegaly, carcinoid syndrome, and variceal bleeding
Cholecystokinin	I cells (duodenum, jejunum)	<ul style="list-style-type: none"> ↑ pancreatic secretion ↑ gallbladder contraction ↓ gastric emptying ↑ sphincter of Oddi relaxation 	<ul style="list-style-type: none"> ↑ by fatty acids, amino acids 	Acts on neural muscarinic pathways to cause pancreatic secretion
Secretin	S cells (duodenum)	<ul style="list-style-type: none"> ↑ pancreatic HCO₃⁻ secretion ↓ gastric acid secretion ↑ bile secretion 	<ul style="list-style-type: none"> ↑ by acid, fatty acids in lumen of duodenum 	↑ HCO ₃ ⁻ neutralizes gastric acid in duodenum, allowing pancreatic enzymes to function
Glucose-dependent insulinotropic peptide	K cells (duodenum, jejunum)	<ul style="list-style-type: none"> Exocrine: ↓ gastric H⁺ secretion Endocrine: ↑ insulin release 	<ul style="list-style-type: none"> ↑ by fatty acids, amino acids, oral glucose 	<ul style="list-style-type: none"> Also called gastric inhibitory peptide (GIP) Oral glucose load ↑ insulin compared to IV equivalent due to GIP secretion
Motilin	Small intestine	Produces migrating motor complexes (MMCs)	<ul style="list-style-type: none"> ↑ in fasting state 	Motilin receptor agonists (eg, erythromycin) are used to stimulate intestinal peristalsis.
Vasoactive intestinal polypeptide	Parasympathetic ganglia in sphincters, gallbladder, small intestine	<ul style="list-style-type: none"> ↑ intestinal water and electrolyte secretion ↑ relaxation of intestinal smooth muscle and sphincters 	<ul style="list-style-type: none"> ↑ by distention and vagal stimulation ↓ by adrenergic input 	VIPoma —non- α , non- β islet cell pancreatic tumor that secretes VIP; associated with Watery Diarrhea, Hypokalemia, Achlorhydria (WDHA syndrome)
Nitric oxide		<ul style="list-style-type: none"> ↑ smooth muscle relaxation, including lower esophageal sphincter (LES) 		Loss of NO secretion is implicated in ↑ LES tone of achalasia
Ghrelin	Stomach	<ul style="list-style-type: none"> ↑ appetite (“ghrowlin’ stomach”) 	<ul style="list-style-type: none"> ↑ in fasting state ↓ by food 	<ul style="list-style-type: none"> ↑ in Prader-Willi syndrome ↓ after gastric bypass surgery

Gastrointestinal secretory products

PRODUCT	SOURCE	ACTION	REGULATION	NOTES
Intrinsic factor	Parietal cells (stomach A)	Vitamin B ₁₂ -binding protein (required for B ₁₂ uptake in terminal ileum)		Autoimmune destruction of parietal cells → chronic gastritis and pernicious anemia.
Gastric acid	Parietal cells (stomach)	↓ stomach pH	↑ by histamine, vagal stimulation (ACh), gastrin ↓ by somatostatin, GIP, prostaglandin, secretin	
Pepsin	Chief cells (stomach)	Protein digestion	↑ by vagal stimulation (ACh), local acid	Pepsinogen (inactive) is converted to pepsin (active) in the presence of H ⁺ .
Bicarbonate	Mucosal cells (stomach, duodenum, salivary glands, pancreas) and Brunner glands (duodenum)	Neutralizes acid	↑ by pancreatic and biliary secretion with secretin	Trapped in mucus that covers the gastric epithelium.



Locations of gastrointestinal secretory cells




Gastrin ↑ acid secretion primarily through its effects on enterochromaffin-like (ECL) cells (leading to histamine release) rather than through its direct effect on parietal cells.

Pancreatic secretions

Isotonic fluid; low flow → high Cl⁻, high flow → high HCO₃⁻.

ENZYME	ROLE	NOTES
α-amylase	Starch digestion	Secreted in active form
Lipases	Fat digestion	
Proteases	Protein digestion	Includes trypsin, chymotrypsin, elastase, carboxypeptidases Secreted as proenzymes also called zymogens
Trypsinogen	Converted to active enzyme trypsin → activation of other proenzymes and cleaving of additional trypsinogen molecules into active trypsin (positive feedback loop)	Converted to trypsin by enterokinase/ enteropeptidase, a brush-border enzyme on duodenal and jejunal mucosa

Carbohydrate absorption




Only monosaccharides (glucose, galactose, fructose) are absorbed by enterocytes. Glucose and galactose are taken up by SGLT1 (Na⁺ dependent). Fructose is taken up via Facilitated diffusion by GLUT5. All are transported to blood by GLUT2.


D-xylose absorption test: simple sugar that requires intact mucosa for absorption, but does not require digestive enzymes. Helps distinguish GI mucosal damage from other causes of malabsorption.

Vitamin and mineral absorption

Iron	Absorbed as Fe ²⁺ in duodenum	Iron F, B, Bro
Folate	Absorbed in small bowel	Clinically relevant in patients with small bowel disease or after resection (eg, vitamin B ₁₂ deficiency following terminal ileum resection)
Vitamin B₁₂	Absorbed in terminal ileum along with bile salts, requires intrinsic factor	



Peyer patches



Unencapsulated lymphoid tissue **A** found in lamina propria and submucosa of ileum. Contain specialized **M** cells that sample and present antigens to **immune** cells. B cells stimulated in germinal centers of Peyer patches differentiate into IgA-secreting plasma cells, which ultimately reside in lamina propria. IgA receives protective secretory component and is then transported across the epithelium to the gut to deal with intraluminal antigen.

Think of **IgA**, the **Intra-gut Antibody**

Bile

Composed of bile salts (bile acids conjugated to glycine or taurine, making them water soluble), phospholipids, cholesterol, bilirubin, water, and ions. Cholesterol 7 α -hydroxylase catalyzes rate-limiting step of bile acid synthesis.

Functions:

- Digestion and absorption of lipids and fat-soluble vitamins
- Cholesterol excretion (body's 1^o means of eliminating cholesterol)
- Antimicrobial activity (via membrane disruption)


↓ absorption of enteric bile salts at distal ileum (as in short bowel syndrome, Crohn disease) prevents normal fat absorption
 Calcium, which normally binds oxalate, binds fat instead, so free oxalate is absorbed by gut → ↑ frequency of calcium oxalate kidney stones

Bilirubin


Heme is metabolized by heme oxygenase to biliverdin, which is subsequently reduced to bilirubin. Unconjugated bilirubin is removed from blood by liver, conjugated with glucuronate, and excreted in bile.

Direct bilirubin: conjugated with glucuronic acid; water soluble (**d**issolves in water).

Indirect bilirubin: unconjugated; water **i**nsoluble.



► GASTROINTESTINAL—PATHOLOGY

Sialolithiasis


Stone(s) in salivary gland duct **A**. Can occur in 3 major salivary glands (parotid, submandibular, sublingual). Single stone more common in submandibular gland (Wharton duct).

Presents as recurrent pre-/periprandial pain and swelling in affected gland.

Caused by dehydration or trauma.


Treat conservatively with NSAIDs, gland massage, warm compresses, sour candies (to promote salivary flow).

Sialadenitis—inflammation of salivary gland due to obstruction, infection, or immune-mediated mechanisms.

Salivary gland tumors

Most are benign and commonly affect parotid gland (80-85%). Nearly half of all submandibular gland neoplasms and most sublingual and minor salivary gland tumors are malignant. Typically present as painless mass/swelling. Facial paralysis or pain suggests malignant involvement.

- **Pleomorphic adenoma** (benign mixed tumor)—most common salivary gland tumor **A**. Composed of chondromyxoid stroma and epithelium and recurs if incompletely excised or ruptured intraoperatively. May undergo malignant transformation.
- **Mucoepidermoid carcinoma**—most common malignant tumor, has mucinous and squamous components.
- **Warthin tumor** (papillary cystadenoma lymphomatosum)—benign cystic tumor with **germinal** centers. Typically found in **smokers**. Bilateral in 10%; multifocal in 10%. “**Warriors** from **Germany** love **smoking**.”

Achalasia

Failure of LES to relax due to degeneration of inhibitory neurons (containing NO and VIP) in the myenteric (Auerbach) plexus of the esophageal wall.

Manometry findings include uncoordinated or absent peristalsis with high LES resting pressure → progressive dysphagia to solids and liquids (vs obstruction—solids only). Barium swallow shows dilated esophagus with an area of distal stenosis (“bird’s beak” **A**).

Associated with ↑ risk of esophageal cancer.


A-achalasia = absence of relaxation.

2° achalasia (pseudoachalasia) may arise from Chagas disease (*T cruzi* infection) or extraesophageal malignancies (mass effect or paraneoplastic).



Chagas disease can cause **achalasia**.

Esophageal pathologies


Diffuse esophageal spasm	Spontaneous, nonperistaltic (uncoordinated) contractions of the esophagus with normal LES pressure. Presents with dysphagia and angina-like chest pain. Barium swallow reveals “corkscrew” esophagus. Manometry is diagnostic. Treatment includes nitrates and CCBs.
Eosinophilic esophagitis	Infiltration of eosinophils in the esophagus often in atopic patients. Food allergens → dysphagia, food impaction. Esophageal rings and linear furrows often seen on endoscopy. Typically unresponsive to GERD therapy.
Esophageal perforation	Most commonly iatrogenic following esophageal instrumentation. Noniatrogenic causes include spontaneous rupture, foreign body ingestion, trauma, malignancy. May present with pneumomediastinum (arrows in A). Subcutaneous emphysema may be due to dissecting air (signs include crepitus in the neck region or chest wall). Boerhaave syndrome —transmural, usually distal esophageal rupture due to violent retching.
Esophageal strictures	Associated with caustic ingestion, acid reflux, and esophagitis.
Esophageal varices	Dilated submucosal veins (red arrows in B C) in lower 1/3 of esophagus 2° to portal hypertension. Common in cirrhotics, may be source of life-threatening hematemesis.
Esophagitis	Associated with reflux, infection in immunocompromised (<i>Candida</i> : white pseudomembrane D ; HSV-1: punched-out ulcers; CMV: linear ulcers), caustic ingestion, or pill-induced esophagitis (eg, bisphosphonates, tetracycline, NSAIDs, iron, and potassium chloride).
Gastroesophageal reflux disease	Commonly presents as heartburn, regurgitation, dysphagia. May also present as chronic cough, hoarseness (laryngopharyngeal reflux). Associated with asthma. Transient decreases in LES tone.
Mallory-Weiss syndrome	Partial thickness, longitudinal lacerations of gastroesophageal junction, confined to mucosa/submucosa, due to severe vomiting. Often presents with hematemesis. Usually found in alcoholics and bulimics.
Plummer-Vinson syndrome	Triad of D ysphagia, I ron deficiency anemia, E sophageal webs. ↑ risk of esophageal S quamous cell carcinoma (“ Plumber DIES ”). May be associated with glossitis.
Schatzki rings	Rings formed at gastroesophageal junction, typically due to chronic acid reflux. Can present with dysphagia.
Sclerodermal esophageal dysmotility	Esophageal smooth muscle atrophy → ↓ LES pressure and dysmotility → acid reflux and dysphagia → stricture, Barrett esophagus, and aspiration. Part of CREST syndrome.



Barrett esophagus



Specialized intestinal metaplasia **A**—replacement of nonkeratinized stratified squamous epithelium with intestinal epithelium (nonciliated columnar with goblet cells [stained blue in **B**]) in distal esophagus. Due to chronic gastroesophageal reflux disease (GERD). Associated with ↑ risk of esophageal adenocarcinoma.




Esophageal cancer

Typically presents with progressive dysphagia (first solids, then liquids) and weight loss. Aggressive course due to lack of serosa in esophageal wall, allowing rapid extension. Poor prognosis due to advanced disease at presentation.


CANCER	PART OF ESOPHAGUS AFFECTED	RISK FACTORS	PREVALENCE
Squamous cell carcinoma	Upper 2/3	Alcohol, hot liquids, caustic strictures, smoking, achalasia	More common worldwide
Adenocarcinoma	Lower 1/3	Chronic GERD, Barrett esophagus, obesity, smoking, achalasia	More common in America

Gastritis

Acute gastritis	Erosions can be caused by: <ul style="list-style-type: none"> ▪ NSAIDs—↓ PGE₂ → ↓ gastric mucosa protection ▪ Burns (Curling ulcer)—hypovolemia → mucosal ischemia ▪ Brain injury (Cushing ulcer)—↑ vagal stimulation → ↑ ACh → ↑ H⁺ production 	Especially common among alcoholics and patients taking daily NSAIDs (eg, patients with rheumatoid arthritis) Burned by the Curling iron Always Cushion the brain
Chronic gastritis	Mucosal inflammation, often leading to atrophy (hypochlorhydria → hypergastrinemia) and intestinal metaplasia (↑ risk of gastric cancers)	
<i>H pylori</i>	Most common. ↑ risk of peptic ulcer disease, MALT lymphoma	Affects antrum first and spreads to body of stomach
Autoimmune	Autoantibodies to the H ⁺ /K ⁺ ATPase on parietal cells and to intrinsic factor. ↑ risk of pernicious anemia	Affects body/fundus of stomach

Ménétrier disease

Hyperplasia of gastric mucosa → hypertrophied rugae (look like brain gyri **A**). Causes excess mucus production with resultant protein loss and parietal cell atrophy with ↓ acid production. Precancerous. Presents with **W**eight loss, **A**norexia, **V**omiting, **E**pigastric pain, **E**dema (due to protein loss) (**WAVEE**).

Gastric cancer

Most commonly gastric adenocarcinoma; lymphoma, GI stromal tumor, carcinoid (rare). Early aggressive local spread with node/liver metastases. Often presents late, with weight loss, abdominal pain, early satiety, and in some cases acanthosis nigricans or Leser-Trélat sign. Associated with blood type A.

- Intestinal—associated with *H pylori*, dietary nitrosamines (smoked foods), tobacco smoking, achlorhydria, chronic gastritis. Commonly on lesser curvature; looks like ulcer with raised margins.
- Diffuse—not associated with *H pylori*; most cases due to E-cadherin mutation; signet ring cells (mucin-filled cells with peripheral nuclei) **A**; stomach wall grossly thickened and leathery (linitis plastica).

Virchow node—involvement of left supraclavicular node by metastasis from stomach.

Krukenberg tumor—bilateral metastases to ovaries. Abundant mucin-secreting, signet ring cells.

Sister Mary Joseph nodule—subcutaneous periumbilical metastasis.

Blumer shelf—palpable mass on digital rectal exam suggesting metastasis to rectouterine pouch (pouch of Douglas).

Peptic ulcer disease

	Gastric ulcer	Duodenal ulcer
PAIN	Can be G reater with meals—weight loss	D ecreases with meals—weight gain
<i>H. PYLORI</i> INFECTION	~ 70%	~ 90%
MECHANISM	↓ mucosal protection against gastric acid	↓ mucosal protection or ↑ gastric acid secretion
OTHER CAUSES	NSAIDs	Zollinger-Ellison syndrome
RISK OF CARCINOMA	↑	Generally benign
OTHER	Biopsy margins to rule out malignancy	

Ulcer complications**Hemorrhage**


Gastric, duodenal (posterior > anterior). Most common complication.
Ruptured gastric ulcer on the lesser curvature of stomach → bleeding from left gastric artery.
An ulcer on the posterior wall of duodenum → bleeding from gastroduodenal artery.

Obstruction

Pyloric channel, duodenal.


Perforation

Duodenal (anterior > posterior).
Anterior duodenal ulcers can perforate into the anterior abdominal cavity, potentially leading to pneumoperitoneum.
May see free air under diaphragm (pneumoperitoneum) **A** with referred pain to the shoulder via irritation of phrenic nerve.



Malabsorption syndromes

Can cause diarrhea, steatorrhea, weight loss, weakness, vitamin and mineral deficiencies. Screen for fecal fat (eg, Sudan stain).

Celiac disease

Gluten-sensitive enteropathy, celiac sprue. Autoimmune-mediated intolerance of gliadin (gluten protein found in wheat) → malabsorption and steatorrhea. Associated with HLA-DQ2, HLA-DQ8, northern European descent, dermatitis herpetiformis, ↓ bone density.

Findings: IgA anti-tissue transglutaminase (IgA tTG), anti-endomysial, anti-deamidated gliadin peptide antibodies; villous atrophy, crypt hyperplasia **A**, and intraepithelial lymphocytosis. Moderately ↑ risk of malignancy (eg, T-cell lymphoma).

↓ mucosal absorption primarily affects distal duodenum and/or proximal jejunum. D-xylose test: passively absorbed in proximal small intestine; blood and urine levels ↓ with mucosa defects or bacterial overgrowth, normal in pancreatic insufficiency. Treatment: gluten-free diet.

Lactose intolerance

Lactase deficiency. Normal-appearing villi, except when 2° to injury at tips of villi (eg, viral enteritis). Osmotic diarrhea with ↓ stool pH (colonic bacteria ferment lactose).

Lactose hydrogen breath test: ⊕ for lactose malabsorption if post-lactose breath hydrogen value rises > 20 ppm compared with baseline.

Pancreatic insufficiency


Due to chronic pancreatitis, cystic fibrosis, obstructing cancer. Causes malabsorption of fat and fat-soluble vitamins (A, D, E, K) as well as vitamin B₁₂.

↓ duodenal bicarbonate (and pH) and fecal elastase.

Tropical sprue

Similar findings as celiac sprue (affects small bowel), but responds to antibiotics. Cause is unknown, but seen in residents of or recent visitors to tropics.

↓ mucosal absorption affecting duodenum and jejunum but can involve ileum with time. Associated with megaloblastic anemia due to folate deficiency and, later, B₁₂ deficiency.


Whipple disease

Infection with *Tropheryma whipplei* (intracellular gram ⊕); PAS ⊕ foamy macrophages in intestinal lamina propria **B**, mesenteric nodes. Cardiac symptoms, Arthralgias, and Neurologic symptoms are common. Diarrhea/steatorrhea occur later in disease course. Most common in older men.

PAS the foamy Whipped cream in a CAN.

Inflammatory bowel diseases

	Crohn disease	Ulcerative colitis
LOCATION	Any portion of the GI tract, usually the terminal ileum and colon. Skip lesions, rectal sparing .	Colitis = colon inflammation. Continuous colonic lesions, always with rectal involvement.
GROSS MORPHOLOGY	Transmural inflammation → fistulas. Cobblestone mucosa, creeping fat , bowel wall thickening (“string sign” on barium swallow x-ray A), linear ulcers, fissures.	Mucosal and submucosal inflammation only. Friable mucosa with superficial and/or deep ulcerations (compare normal B with diseased C). Loss of haustra → “lead pipe” appearance on imaging.
MICROSCOPIC MORPHOLOGY	Noncaseating granulomas and lymphoid aggregates. Th1 mediated.	Crypt abscesses and ulcers, bleeding, no granulomas. Th2 mediated.
COMPLICATIONS	Malabsorption/malnutrition, colorectal cancer (↑ risk with pancolitis). Fistulas (eg, enterovesical fistulae, which can cause recurrent UTI and pneumaturia), phlegmon/abscess, strictures (causing obstruction), perianal disease.	Fulminant colitis, toxic megacolon, perforation.
INTESTINAL MANIFESTATION	Diarrhea that may or may not be bloody.	Bloody diarrhea.
EXTRAIESTINAL MANIFESTATIONS	Rash (pyoderma gangrenosum, erythema nodosum), eye inflammation (episcleritis, uveitis), oral ulcerations (aphthous stomatitis), arthritis (peripheral, spondylitis). Kidney stones (usually calcium oxalate), gallstones. May be ⊕ for anti- <i>Saccharomyces cerevisiae</i> antibodies (ASCA).	1° sclerosing cholangitis. Associated with p-ANCA.
TREATMENT	Corticosteroids, azathioprine, antibiotics (eg, ciprofloxacin, metronidazole), biologics (eg, infliximab, adalimumab).	5-aminosalicylic preparations (eg, mesalamine), 6-mercaptopurine, infliximab, colectomy.
	For Crohn , think of a fat granny and an old crone skipping down a cobblestone road away from the wreck (rectal sparing). Stones are more common in Crohns .	Ulcerative colitis causes ULCCERS : U lcers L arge intestine C ontinuous, C olorectal carcinoma, C rypt abscesses E xtends proximally R ed diarrhea S clerosing cholangitis




Irritable bowel syndrome

Recurrent abdominal pain associated with ≥ 2 of the following:

- Related to defecation
- Change in stool frequency
- Change in form (consistency) of stool

No structural abnormalities. Most common in middle-aged women. Chronic symptoms may be diarrhea-predominant, constipation-predominant, or mixed. Pathophysiology is multifaceted. First-line treatment is lifestyle modification and dietary changes.

Appendicitis



Acute inflammation of the appendix (yellow arrows in **A**), can be due to obstruction by fecalith (red arrow in **A**) (in adults) or lymphoid hyperplasia (in children).

Proximal obstruction of appendiceal lumen produces closed-loop obstruction \rightarrow \uparrow intraluminal pressure \rightarrow stimulation of visceral afferent nerve fibers at T8-T10 \rightarrow initial diffuse periumbilical pain \rightarrow inflammation extends to serosa and irritates parietal peritoneum. Pain localized to RLQ/McBurney point (1/3 the distance from right anterior superior iliac spine to umbilicus). Nausea, fever; may perforate \rightarrow peritonitis; may elicit psoas, obturator, and Rovsing signs, guarding and rebound tenderness on exam.

Differential: diverticulitis (elderly), ectopic pregnancy (use hCG to rule out), pseudoappendicitis. Treatment: appendectomy.

Diverticula of the GI tract

Diverticulum

Blind pouch **A** protruding from the alimentary tract that communicates with the lumen of the gut. Most diverticula (esophagus, stomach, duodenum, colon) are acquired and are termed “false diverticula.”

“True” diverticulum—all gut wall layers outpouch (eg, Meckel).

“False” diverticulum or pseudodiverticulum—only mucosa and submucosa outpouch. Occur especially where vasa recta perforate muscularis externa.

Diverticulosis

Many false diverticula of the colon **B**, commonly sigmoid. Common (in $\sim 50\%$ of people > 60 years). Caused by \uparrow intraluminal pressure and focal weakness in colonic wall. Associated with obesity and diets low in fiber, high in total fat/red meat.

Often asymptomatic or associated with vague discomfort.

Complications include diverticular bleeding (painless hematochezia), diverticulitis.

Diverticulitis

Inflammation of diverticula with wall thickening (red arrows in **C**) classically causing LLQ pain, fever, leukocytosis. Treat with antibiotics.

Complications: abscess, fistula (colovesical fistula \rightarrow pneumaturia), obstruction (inflammatory stenosis), perforation (white arrows in **C**) (\rightarrow peritonitis).




Zenker diverticulum




Pharyngoesophageal false diverticulum **A**. Esophageal dysmotility causes herniation of mucosal tissue at Killian triangle between the thyropharyngeal and cricopharyngeal parts of the inferior pharyngeal constrictor. Presenting symptoms: dysphagia, obstruction, gurgling, aspiration, foul breath, neck mass. Most common in elderly males.

Elder MIKE has bad breath:

- Elderly**
- Males**
- Inferior pharyngeal constrictor**
- Killian triangle**
- Esophageal dysmotility**
- Halitosis**



Meckel diverticulum




True diverticulum. Persistence of the vitelline (omphalomesenteric) duct. May contain ectopic acid-secreting gastric mucosa and/or pancreatic tissue. Most common congenital anomaly of GI tract. Can cause hematochezia/melena (less common), RLQ pain, intussusception, volvulus, or obstruction near terminal ileum.

Contrast with omphalomesenteric cyst = cystic dilation of vitelline duct.
Diagnosis: ^{99m}Tc-pertechnetate scan (aka Meckel scan) for uptake by heterotopic gastric mucosa.

The rule of **2**'s:
2 times as likely in males.
2 inches long.
2 feet from the ileocecal valve.
2% of population.
 Commonly presents in first **2** years of life.
 May have **2** types of epithelia (gastric/pancreatic).


Hirschsprung disease






Congenital megacolon characterized by lack of ganglion cells/enteric nervous plexuses (Auerbach and Meissner plexuses) in distal segment of colon. Due to failure of neural crest cell migration. Associated with loss of function mutations in *RET*.

Presents with bilious emesis, abdominal distention, and failure to pass meconium within 48 hours → chronic constipation. Normal portion of the colon proximal to the aganglionic segment is dilated, resulting in a “transition zone.”

Risk ↑ with Down syndrome.
 Explosive expulsion of feces (squirt sign) → empty rectum on digital exam.
 Diagnosed by absence of ganglionic cells on rectal suction biopsy.
 Treatment: resection.
RET mutation in the **REcTum**.

Malrotation

Anomaly of midgut rotation during fetal development → improper positioning of bowel (small bowel clumped on the right side) **A**, formation of fibrous bands (Ladd bands). Can lead to volvulus, duodenal obstruction.

**Intussusception**

Telescoping **A** of proximal bowel segment into a distal segment, commonly at the ileocecal junction. Most commonly idiopathic, but may be due to lead point.

Compromised blood supply → intermittent, severe, abdominal pain often with “currant jelly” dark red stools.

Majority of cases in infants, unusual in adults.


Most common pathologic lead point:

- Children—Meckel diverticulum
- Adults—intraluminal mass/tumor

On physical exam, patient may draw their legs to chest to ease pain, sausage shaped mass on palpation.


Imaging—Ultrasound/CT may show “target sign.” **B**

May be associated with IgA vasculitis (HSP), recent viral infection (eg, adenovirus; Peyer patch hypertrophy creates lead point).

Volvulus

Twisting of portion of bowel around its mesentery; can lead to obstruction and infarction. Can occur throughout the GI tract.

- Midgut volvulus more common in infants and children (**minors**)
- Sigmoid volvulus (coffee bean sign on x-ray **A**) more common in **seniors** (elderly)

**Other intestinal disorders****Acute mesenteric ischemia**

Critical blockage of intestinal blood flow (often embolic occlusion of SMA) → small bowel necrosis **A** → abdominal pain out of proportion to physical findings. May see red “currant jelly” stools.

Adhesion

Fibrous band of scar tissue; commonly forms after surgery. Most common cause of small bowel obstruction, demonstrated by multiple dilated small bowel loops on x-ray (arrows in **B**).

Angiodysplasia

Tortuous dilation of vessels **C** → hematochezia. Most often found in the right-sided colon. More common in older patients. Confirmed by angiography. Associated with end-stage renal disease, von Willebrand disease, aortic stenosis.

Chronic mesenteric ischemia

“Intestinal angina”: atherosclerosis of celiac artery, SMA, or IMA → intestinal hypoperfusion → postprandial epigastric pain → food aversion and weight loss.

Colonic ischemia

Reduction in intestinal blood flow causes ischemia. Crampy abdominal pain followed by hematochezia. Commonly occurs at watershed areas (splenic flexure, rectosigmoid junction). Typically affects elderly. Thumbprint sign on imaging due to mucosal edema/hemorrhage.

Ileus


Intestinal hypomotility without obstruction → constipation and ↓ flatus; distended/tympanic abdomen with ↓ bowel sounds. Associated with abdominal surgeries, opiates, hypokalemia, sepsis. Treatment: bowel rest, electrolyte correction, cholinergic drugs (stimulate intestinal motility).

Meconium ileus

Meconium plug obstructs intestine, prevents stool passage at birth. Associated with cystic fibrosis.


Necrotizing enterocolitis

Seen in premature, formula-fed infants with immature immune system. Necrosis of intestinal mucosa (most commonly terminal ileum and proximal colon) with possible perforation, which can lead to pneumatosis intestinalis (arrows in **D**), pneumoperitoneum, portal venous gas.



Colonic polyps Growths of tissue within the colon **A**. Grossly characterized as flat, sessile, or pedunculated on the basis of protrusion into colonic lumen. Generally classified by histologic type.

HISTOLOGIC TYPE	CHARACTERISTICS
Generally non-neoplastic	
Hamartomatous polyps	Solitary lesions do not have significant risk of transformation. Growths of normal colonic tissue with distorted architecture. Associated with Peutz-Jeghers syndrome and juvenile polyposis.
Hyperplastic polyps	Most common; generally smaller and predominantly located in rectosigmoid region. Occasionally evolves into serrated polyps and more advanced lesions.
Inflammatory pseudopolyps	Due to mucosal erosion in inflammatory bowel disease.
Mucosal polyps	Small, usually < 5 mm. Look similar to normal mucosa. Clinically insignificant.
Submucosal polyps	May include lipomas, leiomyomas, fibromas, and other lesions.
Malignant potential	
Adenomatous polyps	Neoplastic, via chromosomal instability pathway with mutations in <i>APC</i> and <i>KRAS</i> . Tubular B histology has less malignant potential than villous C (“villous histology is villainous”); tubulovillous has intermediate malignant potential. Usually asymptomatic; may present with occult bleeding.
Serrated polyps	Neoplastic. Characterized by CpG island methylator phenotype (CIMP; cytosine base followed by guanine, linked by a phosphodiester bond). Defect may silence <i>MMR</i> gene (DNA mismatch repair) expression. Mutations lead to microsatellite instability and mutations in <i>BRAF</i> . “Saw-tooth” pattern of crypts on biopsy. Up to 20% of cases of sporadic CRC.




Polyposis syndromes

Familial adenomatous polyposis	Autosomal dominant mutation of <i>APC</i> tumor suppressor gene on chromosome 5q22. 2-hit hypothesis. Thousands of polyps arise starting after puberty; pancolonic; always involves rectum. Prophylactic colectomy or else 100% progress to CRC.
Gardner syndrome	FAP + osseous and soft tissue tumors (eg, osteomas of skull or mandible), congenital hypertrophy of retinal pigment epithelium, impacted/supernumerary teeth.
Turcot syndrome	FAP or Lynch syndrome + malignant CNS tumor (eg, medulloblastoma, glioma). Turcot = Turban .
Peutz-Jeghers syndrome	Autosomal dominant syndrome featuring numerous hamartomas throughout GI tract, along with hyperpigmented macules on mouth, lips, hands, genitalia. Associated with ↑ risk of breast and GI cancers (eg, colorectal, stomach, small bowel, pancreatic).
Juvenile polyposis syndrome	Autosomal dominant syndrome in children (typically < 5 years old) featuring numerous hamartomatous polyps in the colon, stomach, small bowel. Associated with ↑ risk of CRC.

Lynch syndrome

Previously called hereditary nonpolyposis colorectal cancer (HNPCC). Autosomal dominant mutation of mismatch repair genes (eg, *MLH1*, *MSH2*) with subsequent microsatellite instability. ~ 80% progress to CRC. Proximal colon is always involved. Associated with endometrial, ovarian, and skin cancers.

Colorectal cancer**DIAGNOSIS**

Iron deficiency anemia in males (especially > 50 years old) and postmenopausal females raises suspicion.

Screening:

- Low risk: screen at age 50 with colonoscopy (polyp seen in **A**); alternatives include flexible sigmoidoscopy, fecal occult blood testing (FOBT), fecal immunochemical testing (FIT), FIT-fecal DNA, CT colonography
- Patients with a first-degree relative who has colon cancer: screen at age 40 with colonoscopy, or 10 years prior to the relative's presentation
- Patients with IBD: distinct screening protocol

“Apple core” lesion seen on barium enema x-ray **B**.

CEA tumor marker: good for monitoring recurrence, should not be used for screening.

EPIDEMIOLOGY

Most patients are > 50 years old. ~ 25% have a family history.

PRESENTATION

Rectosigmoid > ascending > descending.

Right side (cecal, ascending) associated with occult bleeding; left side (rectosigmoid) associated with hematochezia and obstruction (narrower lumen).

Ascending—exophytic mass, iron deficiency anemia, weight loss.

Descending—infiltrating mass, partial obstruction, colicky pain, hematochezia.


Can present with *S bovis* (*gallolyticus*) bacteremia/endocarditis or as an episode of diverticulitis.

RISK FACTORS


Adenomatous and serrated polyps, familial cancer syndromes, IBD, tobacco use, diet of processed meat with low fiber.

Molecular pathogenesis of colorectal cancer

Chromosomal instability pathway: mutations in *APC* cause FAP and most sporadic cases of CRC via adenoma-carcinoma sequence; (firing order of events is “AK-53”).
 Microsatellite instability pathway: mutations or methylation of mismatch repair genes (eg, *MLH1*) cause Lynch syndrome and some sporadic CRC (via serrated polyp pathway).
 Overexpression of COX-2 has been linked to colorectal cancer, NSAIDs may be chemopreventive.




Cirrhosis and portal hypertension



Cirrhosis—diffuse bridging fibrosis (via stellate cells) and regenerative nodules (red arrows in **A**; white arrows show splenomegaly) disrupt normal architecture of liver; ↑ risk for hepatocellular carcinoma (white arrow in **B**). Etiologies include alcohol, nonalcoholic steatohepatitis, chronic viral hepatitis, autoimmune hepatitis, biliary disease, genetic/metabolic disorders.

Portal hypertension—↑ pressure in portal venous system. Etiologies include cirrhosis (most common cause in Western countries), vascular obstruction (eg, portal vein thrombosis, Budd-Chiari syndrome), schistosomiasis.



Spontaneous bacterial peritonitis

Also called 1° bacterial peritonitis. Common and potentially fatal bacterial infection in patients with cirrhosis and ascites. Often asymptomatic, but can cause fevers, chills, abdominal pain, ileus, or worsening encephalopathy. Commonly caused by gram ⊖ organisms (eg, *E coli*, *Klebsiella*) or less commonly gram ⊕ *Streptococcus*.

Diagnosis: paracentesis with ascitic fluid absolute neutrophil count (ANC) > 250 cells/mm³.

Empiric first-line treatment is 3rd generation cephalosporin (eg, cefotaxime).

Serum markers of liver pathology

ENZYMES RELEASED IN LIVER DAMAGE

Aspartate aminotransferase and alanine aminotransferase	↑ in most liver disease: ALT > AST ↑ in alcoholic liver disease: AST > ALT (AST usually will not exceed 500 U/L in alcoholic hepatitis) AST > ALT in nonalcoholic liver disease suggests progression to advanced fibrosis or cirrhosis ↑↑↑ aminotransferases (>1000 U/L): differential includes drug-induced liver injury (eg, acetaminophen toxicity), ischemic hepatitis, acute viral hepatitis, autoimmune hepatitis
Alkaline phosphatase	↑ in cholestasis (eg, biliary obstruction), infiltrative disorders, bone disease
γ-glutamyl transpeptidase	↑ in various liver and biliary diseases (just as ALP can), but not in bone disease; associated with alcohol use

FUNCTIONAL LIVER MARKERS

Bilirubin	↑ in various liver diseases (eg, biliary obstruction, alcoholic or viral hepatitis, cirrhosis), hemolysis
Albumin	↓ in advanced liver disease (marker of liver's biosynthetic function)
Prothrombin time	↑ in advanced liver disease (↓ production of clotting factors, thereby measuring the liver's biosynthetic function)
Platelets	↓ in advanced liver disease (↓ thrombopoietin, liver sequestration) and portal hypertension (splenomegaly/splenic sequestration)

Reye syndrome

Rare, often fatal childhood hepatic encephalopathy.

Associated with viral infection (especially VZV and influenza) that has been treated with aspirin. Aspirin metabolites ↓ β-oxidation by reversible inhibition of mitochondrial enzymes.

Findings: mitochondrial abnormalities, fatty liver (microvesicular fatty changes), hypoglycemia, vomiting, hepatomegaly, coma.

Avoid aspirin in children, except in those with Kawasaki disease.

Salicylates aren't a ray (**Reye**) of sun**SHINE** for kids:

Steatosis of liver/hepatocytes
Hypoglycemia/**H**epatomegaly
Infection (VZV, influenza)
Not awake (coma)
Encephalopathy

Alcoholic liver disease**Hepatic steatosis**

Macrovesicular fatty change **A** that may be reversible with alcohol cessation.



Alcoholic hepatitis

Requires sustained, long-term consumption. Swollen and necrotic hepatocytes with neutrophilic infiltration. Mallory bodies **B** (intracytoplasmic eosinophilic inclusions of damaged keratin filaments).

Make a toAST with alcohol:
AST > ALT (ratio usually > 2:1).

Alcoholic cirrhosis

Final and usually irreversible form. Sclerosis around central vein (arrows in **C**) may be seen in early disease. Regenerative nodules surrounded by fibrous bands in response to chronic liver injury → portal hypertension and end-stage liver disease.

**Nonalcoholic fatty liver disease**

Metabolic syndrome (insulin resistance); obesity → fatty infiltration of hepatocytes **A** → cellular “ballooning” and eventual necrosis. May cause cirrhosis and HCC. Independent of alcohol use.

ALT > AST (Lipids)

Hepatic encephalopathy

Cirrhosis → portosystemic shunts → ↓ NH₃ metabolism → neuropsychiatric dysfunction. Reversible neuropsychiatric dysfunction ranging from disorientation/asterixis (mild) to difficult arousal or coma (severe).

Triggers:

- ↑ NH₃ production and absorption (due to GI bleed, constipation, infection).
- ↓ NH₃ removal (due to renal failure, diuretics, bypassed hepatic blood flow post-TIPS).


Treatment: lactulose (↑ NH₄⁺ generation) and rifaximin (↓ NH₃-producing gut bacteria).

Hepatocellular carcinoma/hepatoma

Most common 1° malignant tumor of liver in adults **A**. Associated with HBV (+/- cirrhosis) and all other causes of cirrhosis (including HCV, alcoholic and nonalcoholic fatty liver disease, autoimmune disease, hemochromatosis, Wilson disease, α_1 -antitrypsin deficiency) and specific carcinogens (eg, aflatoxin from *Aspergillus*). May lead to Budd-Chiari syndrome.

Findings: jaundice, tender hepatomegaly, ascites, polycythemia, anorexia. Spreads hematogenously.


Diagnosis: \uparrow α -fetoprotein; ultrasound or contrast CT/MRI **B**, biopsy.

**Other liver tumors****Angiosarcoma**

Malignant tumor of endothelial origin; associated with exposure to arsenic, vinyl chloride.

Cavernous hemangioma

Most common benign liver tumor (venous malformation) **A**; typically occurs at age 30–50 years. Biopsy contraindicated because of risk of hemorrhage.

**Hepatic adenoma**


Rare, benign liver tumor, often related to oral contraceptive or anabolic steroid use; may regress spontaneously or rupture (abdominal pain and shock).

Metastases

GI malignancies, breast and lung cancer. Most common overall; metastases are rarely solitary.

Budd-Chiari syndrome

Thrombosis or compression of hepatic veins with centrilobular congestion and necrosis \rightarrow congestive liver disease (hepatomegaly, ascites, varices, abdominal pain, liver failure). Absence of JVD. Associated with hypercoagulable states, polycythemia vera, postpartum state, HCC. May cause nutmeg liver (mottled appearance).

 α_1 -antitrypsin deficiency

Misfolded gene product protein aggregates in hepatocellular ER \rightarrow cirrhosis with PAS \oplus globules **A** in liver. Codominant trait. Often presents in young patients with liver damage and dyspnea without a history of smoking.

In lungs, \downarrow α_1 -antitrypsin \rightarrow uninhibited elastase in alveoli \rightarrow \downarrow elastic tissue \rightarrow panacinar emphysema.

Jaundice

Abnormal yellowing of the skin and/or sclera **A** due to bilirubin deposition. Hyperbilirubinemia 2° to ↑ production or ↓ clearance (impaired hepatic uptake, conjugation, excretion).

HOT Liver—common causes of ↑ bilirubin level:
Hemolysis
Obstuction
Tumor
Liver disease

Conjugated (direct) hyperbilirubinemia

Biliary tract obstruction: gallstones, cholangiocarcinoma, pancreatic or liver cancer, liver fluke.
 Biliary tract disease:
 ▪ 1° sclerosing cholangitis
 ▪ 1° biliary cholangitis
 Excretion defect: Dubin-Johnson syndrome, Rotor syndrome.

Unconjugated (indirect) hyperbilirubinemia

Hemolytic, physiologic (newborns), Crigler-Najjar, Gilbert syndrome.

Mixed (direct and indirect) hyperbilirubinemia

Hepatitis, cirrhosis.

Physiologic neonatal jaundice

At birth, immature UDP-glucuronosyltransferase → unconjugated hyperbilirubinemia → jaundice/kernicterus (deposition of unconjugated, lipid-soluble bilirubin in the brain, particularly basal ganglia).
 Occurs after first 24 hours of life and usually resolves without treatment in 1–2 weeks.
 Treatment: phototherapy (non-UV) isomerizes unconjugated bilirubin to water-soluble form.

Biliary atresia

Most common reason for pediatric liver transplantation.
 Fibro-obliterative destruction of extrahepatic bile ducts → cholestasis.
 Often presents as a newborn with persistent jaundice after 2 weeks of life, darkening urine, acholic stools, hepatomegaly.
 Labs: ↑ direct bilirubin and GGT.

Hereditary hyperbilirubinemias

All autosomal recessive.

1 Gilbert syndrome

Mildly ↓ UDP-glucuronosyltransferase conjugation and impaired bilirubin uptake. Asymptomatic or mild jaundice usually with stress, illness, or fasting. ↑ unconjugated bilirubin without overt hemolysis.

Relatively common, benign condition.

2 Crigler-Najjar syndrome, type I

Absent UDP-glucuronosyltransferase. Presents early in life, but some patients may not have neurologic signs until later in life.

Findings: jaundice, kernicterus (bilirubin deposition in brain), ↑ unconjugated bilirubin.

Treatment: plasmapheresis and phototherapy (does not conjugate UCB; but does ↑ polarity and ↑ water solubility to allow excretion). Liver transplant is curative.


Type II is less severe and responds to phenobarbital, which ↑ liver enzyme synthesis.


3 Dubin-Johnson syndrome

Conjugated hyperbilirubinemia due to defective liver excretion. Grossly black (**D**ark) liver due to impaired excretion of epinephrine metabolites. Benign.

4 Rotor syndrome

Similar to Dubin-Johnson syndrome, but milder in presentation without black (**R**egular) liver. Due to impaired hepatic uptake and excretion.




Wilson disease

Also called hepatolenticular degeneration. Autosomal recessive mutations in hepatocyte copper-transporting ATPase (*ATP7B* gene; chromosome 13) → ↓ copper incorporation into apoceruloplasmin and excretion into bile → ↓ serum ceruloplasmin. Copper accumulates, especially in liver, brain, cornea, kidneys; ↑ urine copper.

Presents before age 40 with liver disease (eg, hepatitis, acute liver failure, cirrhosis), neurologic disease (eg, dysarthria, dystonia, tremor, parkinsonism), psychiatric disease, Kayser-Fleischer rings (deposits in Descemet membrane of cornea) **A**, hemolytic anemia, renal disease (eg, Fanconi syndrome).

Treatment: chelation with penicillamine or trientine, oral zinc. Liver transplant in acute liver failure related to Wilson disease.

Hemochromatosis

Autosomal recessive. On *HFE* gene, located on chromosome 6; associated with HLA-A3. Leads to abnormal **iron** sensing and ↑ intestinal absorption (↑ ferritin, ↑ iron, ↓ TIBC → ↑ transferrin saturation). Iron overload can also be 2° to chronic transfusion therapy (eg, β-thalassemia major). Iron accumulates, especially in liver, pancreas, skin, heart, pituitary, joints. Hemosiderin (iron) can be identified on liver MRI or biopsy with Prussian blue stain **A**.

Presents after age 40 when total body iron > 20 g; iron loss through menstruation slows progression in women. Classic triad of cirrhosis, diabetes mellitus, skin pigmentation (“bronze diabetes”). Also causes restrictive cardiomyopathy (classic) or dilated cardiomyopathy (reversible), hypogonadism, arthropathy (calcium pyrophosphate deposition; especially metacarpophalangeal joints). HCC is common cause of death.


Treatment: repeated phlebotomy, iron (**Fe**) chelation with deferasirox, deferoxamine, deferi-prone.

Biliary tract disease

May present with pruritus, jaundice, dark urine, light-colored stool, hepatosplenomegaly. Typically with cholestatic pattern of LFTs (↑ conjugated bilirubin, ↑ cholesterol, ↑ ALP, ↑ GGT).

	PATHOLOGY	EPIDEMIOLOGY	ADDITIONAL FEATURES
Primary sclerosing cholangitis	Unknown cause of concentric “onion skin” bile duct fibrosis → alternating strictures and dilation with “beading” of intra- and extrahepatic bile ducts on ERCP, magnetic resonance cholangiopancreatography (MRCP).	Classically in middle-aged men with ulcerative colitis.	Associated with ulcerative colitis. p-ANCA ⊕. ↑ IgM. Can lead to 2° biliary cholangitis. ↑ risk of cholangiocarcinoma and gallbladder cancer.
Primary biliary cholangitis	Autoimmune reaction → lymphocytic infiltrate +/- granulomas → destruction of lobular bile ducts.	Classically in middle-aged women.	Anti-mitochondrial antibody ⊕, ↑ IgM. Associated with other autoimmune conditions (eg, Hashimoto thyroiditis, rheumatoid arthritis, celiac disease). Treatment: ursodiol.
Secondary biliary cirrhosis	Extrahepatic biliary obstruction → ↑ pressure in intrahepatic ducts → injury/ fibrosis and bile stasis.	Patients with known obstructive lesions (gallstones, biliary strictures, pancreatic carcinoma).	May be complicated by ascending cholangitis.

Cholelithiasis and related pathologies



↑ cholesterol and/or bilirubin, ↓ bile salts, and gallbladder stasis all cause stones.

2 types of stones:


- Cholesterol stones (radiolucent with 10–20% opaque due to calcifications)—80% of stones. Associated with obesity, Crohn disease, advanced age, estrogen therapy, multiparity, rapid weight loss, Native American origin.
- Pigment stones **A** (black = radiopaque, Ca^{2+} bilirubinate, hemolysis; brown = radiolucent, infection). Associated with Crohn disease, chronic hemolysis, alcoholic cirrhosis, advanced age, biliary infections, total parenteral nutrition (TPN).

Risk factors (**4 F's**):

1. **F**emale
2. **F**at
3. **F**ertile (multiparity)
4. **F**orty

Most common complication is cholecystitis; can also cause acute pancreatitis, ascending cholangitis.

Diagnose with ultrasound. Treat with elective cholecystectomy if symptomatic.



Rx

RELATED PATHOLOGIES


Biliary colic

Associated with nausea/vomiting and dull RUQ pain. Neurohormonal activation (eg, by CCK after a fatty meal) triggers contraction of gallbladder, forcing stone into cystic duct. Labs are normal, ultrasound shows cholelithiasis.

Choledocholithiasis

Presence of gallstone(s) in common bile duct, often leading to elevated ALP, GGT, direct bilirubin, and/or AST/ALT.

Cholecystitis



Acute or chronic inflammation of gallbladder.


Calculous cholecystitis—most common type; due to gallstone impaction in the cystic duct resulting in inflammation and gallbladder wall thickening (arrows in **B**); can produce 2° infection.

Acalculous cholecystitis—due to gallbladder stasis, hypoperfusion, or infection (CMV); seen in critically ill patients.

Murphy sign: inspiratory arrest on RUQ palpation due to pain. Pain may radiate to right shoulder (due to irritation of phrenic nerve). ↑ ALP if bile duct becomes involved (eg, ascending cholangitis).

Diagnose with ultrasound or cholescintigraphy (HIDA scan). Failure to visualize gallbladder on HIDA scan suggests obstruction.

Gallstone ileus—fistula between gallbladder and GI tract → stone enters GI lumen → obstructs at ileocecal valve (narrowest point); can see air in biliary tree (pneumobilia). Rigler triad: radiographic findings of pneumobilia, small bowel obstruction, gallstone (usually in iliac fossa).

Cholelithiasis and related pathologies (continued)**Porcelain gallbladder**

Calcified gallbladder due to chronic cholecystitis; usually found incidentally on imaging **C**.


Treatment: prophylactic cholecystectomy generally recommended due to ↑ risk of gallbladder cancer (mostly adenocarcinoma).

Ascending cholangitis

Infection of biliary tree usually due to obstruction that leads to stasis/bacterial overgrowth.

Charcot triad of cholangitis includes jaundice, fever, RUQ pain.

Reynolds pentad is Charcot triad plus altered mental status and shock (hypotension).



Acute pancreatitis

Autodigestion of pancreas by pancreatic enzymes (**A** shows pancreas [yellow arrows] surrounded by edema [red arrows]).

Causes: **I**diopathic, **G**allstones, **E**thanol, **T**rauma, **S**teroids, **M**umps, **A**utoimmune disease, **S**corpion sting, **H**ypercalcemia/**H**ypertriglyceridemia (> 1000 mg/dL), **E**RPC, **D**rugs (eg, sulfa drugs, NRTIs, protease inhibitors). **I GET SMASHED**.

Diagnosis by 2 of 3 criteria: acute epigastric pain often radiating to the back, ↑ serum amylase or lipase (more specific) to 3× upper limit of normal, or characteristic imaging findings.

Complications: pseudocyst **B** (lined by granulation tissue, not epithelium), abscess, necrosis, hemorrhage, infection, organ failure (ALI/ARDS, shock, renal failure), hypocalcemia (precipitation of Ca²⁺ soaps).



**Chronic pancreatitis**

Chronic inflammation, atrophy, calcification of the pancreas **A**. Major causes include alcohol abuse and genetic predisposition (ie, cystic fibrosis); can be idiopathic. Complications include pancreatic insufficiency and pseudocysts.

Pancreatic insufficiency (typically when <10% pancreatic function) may manifest with steatorrhea, fat-soluble vitamin deficiency, diabetes mellitus.

Amylase and lipase may or may not be elevated (almost always elevated in acute pancreatitis).

Pancreatic adenocarcinoma



Very aggressive tumor arising from pancreatic ducts (disorganized glandular structure with cellular infiltration **A**); often metastatic at presentation, with average survival ~ 1 year after diagnosis. Tumors more common in pancreatic head **B** (lead to obstructive jaundice). Associated with CA 19-9 tumor marker (also CEA, less specific).

Risk factors:

- Tobacco use
- Chronic pancreatitis (especially > 20 years)
- Diabetes
- Age > 50 years
- Jewish and African-American males


Often presents with:

- Abdominal pain radiating to back
- Weight loss (due to malabsorption and anorexia)
- Migratory thrombophlebitis—redness and tenderness on palpation of extremities (Trousseau syndrome)
- Obstructive jaundice with palpable, nontender gallbladder (Courvoisier sign)

Treatment: Whipple procedure (pancreaticoduodenectomy), chemotherapy, radiation therapy.

▶ GASTROINTESTINAL—PHARMACOLOGY

Acid suppression therapy



Histamine-2 blockers Cimetidine, ranitidine, famotidine, nizatidine. Take H₂ blockers before you **dine**. Think “**table for 2**” to remember H₂.

MECHANISM	Reversible block of histamine H ₂ -receptors → ↓ H ⁺ secretion by parietal cells.
CLINICAL USE	Peptic ulcer, gastritis, mild esophageal reflux.
ADVERSE EFFECTS	Cimetidine is a potent inhibitor of cytochrome P-450 (multiple drug interactions); it also has antiandrogenic effects (prolactin release, gynecomastia, impotence, ↓ libido in males); can cross blood-brain barrier (confusion, dizziness, headaches) and placenta. Both cimetidine and ranitidine ↓ renal excretion of creatinine. Other H ₂ blockers are relatively free of these effects.

Proton pump inhibitors Omeprazole, lansoprazole, esomeprazole, pantoprazole, dexlansoprazole.

MECHANISM	Irreversibly inhibit H ⁺ /K ⁺ ATPase in stomach parietal cells.
CLINICAL USE	Peptic ulcer, gastritis, esophageal reflux, Zollinger-Ellison syndrome, component of therapy for <i>H pylori</i> , stress ulcer prophylaxis.
ADVERSE EFFECTS	↑ risk of <i>C difficile</i> infection, pneumonia, acute interstitial nephritis. Vitamin B ₁₂ malabsorption; ↓ serum Mg ²⁺ and ↓ Ca ²⁺ absorption (potentially leading to increased fracture risk in elderly).

Antacids Can affect absorption, bioavailability, or urinary excretion of other drugs by altering gastric and urinary pH or by delaying gastric emptying. All can cause hypokalemia. Overuse can also cause the following problems:

Aluminum hydroxide	Constipation, Hypophosphatemia, Osteodystrophy, Proximal muscle weakness, Seizures	Aluminum amount of feces CHOPS
Calcium carbonate	Hypercalcemia (milk-alkali syndrome), rebound acid ↑	Can chelate and ↓ effectiveness of other drugs (eg, tetracycline)
Magnesium hydroxide	Diarrhea, hyporeflexia, hypotension, cardiac arrest	Mg²⁺ = Must go 2 the bathroom

Bismuth, sucralfate

MECHANISM	Bind to ulcer base, providing physical protection and allowing HCO ₃ ⁻ secretion to reestablish pH gradient in the mucous layer. Sucralfate requires acidic environment, not given with PPIs/H ₂ blockers.
CLINICAL USE	↑ ulcer healing, travelers' diarrhea (bismuth). Bismuth also used in quadruple therapy for <i>H pylori</i> gastritis.

Misoprostol

MECHANISM	PGE ₁ analog. ↑ production and secretion of gastric mucous barrier, ↓ acid production.
CLINICAL USE	Prevention of NSAID-induced peptic ulcers (NSAIDs block PGE ₁ production). Also used off-label for induction of labor (ripens cervix).
ADVERSE EFFECTS	Diarrhea. Contraindicated in women of childbearing potential (abortifacient).

Octreotide

MECHANISM	Long-acting somatostatin analog; inhibits secretion of various splanchnic vasodilatory hormones.
CLINICAL USE	Acute variceal bleeds, acromegaly, VIPoma, carcinoid tumors.
ADVERSE EFFECTS	Nausea, cramps, steatorrhea. ↑ risk of cholelithiasis due to CCK inhibition.

Sulfasalazine

MECHANISM	A combination of sulfapyridine (antibacterial) and 5-aminosalicylic acid (anti-inflammatory). Activated by colonic bacteria.
CLINICAL USE	Ulcerative colitis, Crohn disease (colitis component).
ADVERSE EFFECTS	Malaise, nausea, sulfonamide toxicity, reversible oligospermia.

Loperamide

MECHANISM	Agonist at μ -opioid receptors; slows gut motility. Poor CNS penetration (low addictive potential).
CLINICAL USE	Diarrhea.
ADVERSE EFFECTS	Constipation, nausea.

Ondansetron

MECHANISM	5-HT ₃ antagonist; ↓ vagal stimulation. Powerful central-acting antiemetic.
CLINICAL USE	Control vomiting postoperatively and in patients undergoing cancer chemotherapy.
ADVERSE EFFECTS	Headache, constipation, QT interval prolongation, serotonin syndrome.

Metoclopramide

MECHANISM	D ₂ receptor antagonist. ↑ resting tone, contractility, LES tone, motility, promotes gastric emptying. Does not influence colon transport time.
CLINICAL USE	Diabetic and postoperative gastroparesis, antiemetic, persistent GERD.
ADVERSE EFFECTS	↑ parkinsonian effects, tardive dyskinesia. Restlessness, drowsiness, fatigue, depression, diarrhea. Drug interaction with digoxin and diabetic agents. Contraindicated in patients with small bowel obstruction, Parkinson disease (due to D ₂ -receptor blockade), ↓ seizure threshold.

Orlistat

MECHANISM	Inhibits gastric and pancreatic lipase → ↓ breakdown and absorption of dietary fats. Taken with fat-containing meals.
CLINICAL USE	Weight loss.
ADVERSE EFFECTS	Abdominal pain, flatulence, bowel urgency/frequent bowel movements, steatorrhea; ↓ absorption of fat-soluble vitamins.

Laxatives		Indicated for constipation or patients on opiates requiring a bowel regimen.	
	EXAMPLES	MECHANISM	ADVERSE EFFECTS
Bulk-forming laxatives	Psyllium, methylcellulose	Soluble fibers draw water into gut lumen, forming a viscous liquid that promotes peristalsis	Bloating
Osmotic laxatives	Magnesium hydroxide, magnesium citrate, polyethylene glycol, lactulose	Provides osmotic load to draw water into GI lumen Lactulose also treats hepatic encephalopathy: gut flora degrade lactulose into metabolites (lactic acid, acetic acid) that promote nitrogen excretion as NH_4^+	Diarrhea, dehydration; may be abused by bulimics
Stimulants	Senna	Enteric nerve stimulation → colonic contraction	Diarrhea, melanosis coli
Emollients	Docusate	Promotes incorporation of water and fat into stool	Diarrhea

Aprepitant

MECHANISM	Substance P antagonist. Blocks NK_1 (neurokinin-1) receptors in brain.
CLINICAL USE	Antiemetic for chemotherapy-induced nausea and vomiting.

HIGH-YIELD SYSTEMS

Hematology and Oncology

“You’re always somebody’s type! (blood type, that is)”

—BloodLink

“All the soarings of my mind begin in my blood.”

—Rainer Maria Rilke

“The best blood will at some time get into a fool or a mosquito.”

—Austin O’Malley

When studying hematology, pay close attention to the many cross connections to immunology. Make sure you master the different types of anemias. Be comfortable interpreting blood smears. When reviewing oncologic drugs, focus on mechanisms and adverse effects rather than details of clinical uses, which may be lower yield.

Please note that solid tumors are covered in their respective organ system chapters.

▶ Embryology	404
▶ Anatomy	406
▶ Physiology	410
▶ Pathology	414
▶ Pharmacology	435

▶ HEMATOLOGY AND ONCOLOGY—EMBRYOLOGY

Fetal erythropoiesis


- Fetal erythropoiesis occurs in:
- **Y**olk sac (3–8 weeks)
 - **L**iver (6 weeks–birth)
 - **S**pleen (10–28 weeks)
 - **B**one marrow (18 weeks to adult)

Young **L**iver **S**ynthesizes **B**lood.
















Hemoglobin development

Embryonic globins: ζ and ϵ .
 Fetal hemoglobin (HbF) = $\alpha_2\gamma_2$.
 Adult hemoglobin (HbA₁) = $\alpha_2\beta_2$.
 HbF has higher affinity for O₂ due to less avid binding of 2,3-BPG, allowing HbF to extract O₂ from maternal hemoglobin (HbA₁ and HbA₂) across the placenta. HbA₂ ($\alpha_2\delta_2$) is a form of adult hemoglobin present in small amounts.

From fetal to adult hemoglobin:
Alpha **A**lways; **G**amma **G**oes, **B**ecomes **B**eta.



Blood groups

	ABO classification				Rh classification	
	A	B	AB	O	Rh ⁺	Rh ⁻
RBC type						
Group antigens on RBC surface	A 	B 	A & B 	NONE	Rh (D) 	NONE
Antibodies in plasma	Anti-B  IgM	Anti-A  IgM	NONE	Anti-A  Anti-B  IgM, IgG	NONE	Anti-D  IgG
Clinical relevance	Receive B or AB → hemolytic reaction	Receive A or AB → hemolytic reaction	Universal recipient of RBCs; universal donor of plasma	Receive any non-O → hemolytic reaction Universal donor of RBCs; universal recipient of plasma	Can receive either Rh ⁺ or Rh ⁻ blood	Treat mother with anti-D IgG during and after each pregnancy to prevent anti-D IgG formation


Hemolytic disease of the newborn

Also known as erythroblastosis fetalis.


	Rh hemolytic disease of the newborn	ABO hemolytic disease of the newborn
INTERACTION	Rh ⁻ mother; Rh ⁺ fetus.	Type O mother; type A or B fetus.
MECHANISM	First pregnancy: mother exposed to fetal blood (often during delivery) → formation of maternal anti-D IgG. Subsequent pregnancies: anti-D IgG crosses the placenta → attacks fetal RBCs → hemolysis in the fetus.	Pre-existing maternal anti-A and/or anti-B IgG antibodies cross placenta → hemolysis in the fetus.
PRESENTATION	Hydrops fetalis, jaundice shortly after birth, kernicterus.	Mild jaundice in the neonate within 24 hours of birth. Unlike Rh HDN, can occur in firstborn babies and is usually less severe.
TREATMENT/PREVENTION	Prevent by administration of anti-D IgG to Rh ⁻ pregnant women during third trimester and early postpartum period (if fetus Rh ⁺). Prevents maternal anti-D IgG production.	Treatment: phototherapy or exchange transfusion.

▶ HEMATOLOGY AND ONCOLOGY—ANATOMY

Hematopoiesis




Neutrophils



Acute inflammatory response cells. Numbers ↑ in bacterial infections. Phagocytic. Multilobed nucleus **A**. Specific granules contain leukocyte alkaline phosphatase (LAP), collagenase, lysozyme, and lactoferrin. Azurophilic granules (lysosomes) contain proteinases, acid phosphatase, myeloperoxidase, and β-glucuronidase.

Hypersegmented neutrophils (nucleus has 6+ lobes) are seen in vitamin B₁₂/ folate deficiency. A left shift with ↑ band cells (immature neutrophils) reflects states of ↑ myeloid proliferation (eg, bacterial infections, CML). Important neutrophil chemotactic agents: C5a, IL-8, LTB₄, kallikrein, platelet-activating factor.

Erythrocytes

Carry O_2 to tissues and CO_2 to lungs. Anucleate and lack organelles; biconcave **A**, with large surface area-to-volume ratio for rapid gas exchange. Life span of 120 days. Source of energy is glucose (90% used in glycolysis, 10% used in HMP shunt). Membranes contain Cl^-/HCO_3^- antiporter, which allow RBCs to export HCO_3^- and transport CO_2 from the periphery to the lungs for elimination.

Eryth = red; *cyte* = cell.


Erythrocytosis = polycythemia = \uparrow Hct.

Anisocytosis = varying sizes.

Poikilocytosis = varying shapes.

Reticulocyte = immature RBC; reflects erythroid proliferation.

Bluish color (polychromasia) on Wright-Giemsa stain of reticulocytes represents residual ribosomal RNA.

Thrombocytes (platelets)

Involved in 1^o hemostasis. Small cytoplasmic fragments **A** derived from megakaryocytes. Life span of 8–10 days. When activated by endothelial injury, aggregate with other platelets and interact with fibrinogen to form platelet plug. Contain dense granules (Ca^{2+} , ADP, Serotonin, Histamine; **CASH**) and α granules (vWF, fibrinogen, fibronectin, platelet factor 4). Approximately $\frac{1}{3}$ of platelet pool is stored in the spleen.


Thrombocytopenia or \downarrow platelet function results in petechiae.

vWF receptor: GpIb.

Fibrinogen receptor: GpIIb/IIIa.

Thrombopoietin stimulates megakaryocyte proliferation.


Alfa granules contain vWF, fibrinogen, fibronectin, platelet factor four.

Monocytes

Found in blood, differentiate into macrophages in tissues.

Large, kidney-shaped nucleus **A**. Extensive “frosted glass” cytoplasm.

Mono = one (nucleus); *cyte* = cell.


Macrophages

Phagocytose bacteria, cellular debris, and senescent RBCs. Long life in tissues. Differentiate from circulating blood monocytes **A**. Activated by γ -interferon. Can function as antigen-presenting cell via MHC II. Important cellular component of granulomas (eg, TB, sarcoidosis).

Macro = large; *phage* = eater.

Macrophage naming varies by specific tissue type (eg, Kupffer cells in liver, histiocytes in connective tissue, Langerhans cells in skin, osteoclasts in bone, microglial cells in brain).

Lipid A from bacterial LPS binds CD14 on macrophages to initiate septic shock.

Eosinophils

Defend against helminthic infections (major basic protein). Bilobate nucleus. Packed with large eosinophilic granules of uniform size **A**. Highly phagocytic for antigen-antibody complexes.

Produce histaminase, major basic protein (MBP, a helminthotoxin), eosinophil peroxidase, eosinophil cationic protein, and eosinophil-derived neurotoxin.

Eosin = pink dye; *philic* = loving.

Causes of eosinophilia = **PACCMAN**:

Parasites

Asthma


Eosinophilic granulomatosis with polyangiitis
(Churg-Strauss syndrome)

Chronic adrenal insufficiency

Myeloproliferative disorders

Allergic processes


Neoplasia (eg, Hodgkin lymphoma)

Basophils

Mediate allergic reaction. Densely basophilic granules **A** contain heparin (anticoagulant) and histamine (vasodilator). Leukotrienes synthesized and released on demand.

Basophilic—stains readily with **basic** stains.

Basophilia is uncommon, but can be a sign of myeloproliferative disorders, particularly CML.


Mast cells

Mediate local tissue allergic reactions. Contain basophilic granules **A**. Originate from same precursor as basophils but are not the same cell type. Can bind the Fc portion of IgE to membrane. Activated by tissue trauma, C3a and C5a, surface IgE cross-linking by antigen (IgE receptor aggregation) → degranulation → release of histamine, heparin, tryptase, and eosinophil chemotactic factors.


Involved in type I hypersensitivity reactions.

Cromolyn sodium prevents mast cell degranulation (used for asthma prophylaxis).


Vancomycin, opioids, and radiocontrast dye can elicit IgE-independent mast cell degranulation.

Dendritic cells


Highly phagocytic antigen-presenting cells (APCs) **A**. Function as link between innate and adaptive immune systems. Express MHC class II and Fc receptors on surface.

Lymphocytes

Refer to B cells, T cells, and NK cells. B cells and T cells mediate adaptive immunity. NK cells are part of the innate immune response. Round, densely staining nucleus with small amount of pale cytoplasm **A**.


Natural killer cells

Important in innate immunity, especially against intracellular pathogens. Larger than B and T cells, with distinctive cytoplasmic lytic granules (containing perforin and granzymes) that, when released, act on target cells to induce apoptosis. Distinguish between healthy and infected cells by identifying cell surface proteins (induced by stress, malignant transformation, or microbial infections).

B cells

Mediate humoral immune response. Originate from stem cells in bone marrow and matures in marrow. Migrate to peripheral lymphoid tissue (follicles of lymph nodes, white pulp of spleen, unencapsulated lymphoid tissue). When antigen is encountered, B cells differentiate into plasma cells (which produce antibodies) and memory cells. Can function as an APC.

B = Bone marrow.


T cells

Mediate cellular immune response. Originate from stem cells in the bone marrow, but mature in the thymus. Differentiate into cytotoxic T cells (express CD8, recognize MHC I), helper T cells (express CD4, recognize MHC II), and regulatory T cells. CD28 (costimulatory signal) necessary for T-cell activation. Most circulating lymphocytes are T cells (80%).

T = Thymus.

CD4+ helper T cells are the primary target of HIV.

Rule of 8: MHC II × CD4 = 8;
MHC I × CD8 = 8.


Plasma cells

Produce large amounts of antibody specific to a particular antigen. “Clock-face” chromatin distribution and eccentric nucleus, abundant RER, and well-developed Golgi apparatus (arrows in **A**). Found in bone marrow and normally do not circulate in peripheral blood.

Multiple myeloma is a plasma cell dyscrasia.

▶ HEMATOLOGY AND ONCOLOGY—PHYSIOLOGY

Hemoglobin electrophoresis




On a gel, hemoglobin migrates from the negatively charged cathode to the positively charged anode. HbA migrates the farthest, followed by HbF, HbS, and HbC. This is because the missense mutations in HbS and HbC replace glutamic acid \ominus with valine (neutral) and lysine \oplus , respectively, making HbC and HbS more positively charged than HbA.

A Fat Santa Claus can't (cathode → anode) go far.



Platelet plug formation (primary hemostasis)



Thrombogenesis

Formation of insoluble fibrin mesh.


Aspirin irreversibly inhibits cyclooxygenase, thereby inhibiting TXA₂ synthesis.

Clopidogrel, prasugrel, and ticlopidine inhibit ADP-induced expression of GpIIb/IIIa by irreversibly blocking P2Y₁₂ receptor.


Abciximab, eptifibatide, and tirofiban inhibit GpIIb/IIIa directly.

Ristocetin activates vWF to bind GpIb. Failure of aggregation with ristocetin assay occurs in von Willebrand disease and Bernard-Soulier syndrome.

vWF carries/protects factor VIII; volksWagen Factories make gr8 cars.




Coagulation and kinin pathways



Vitamin K-dependent coagulation

Procoagulation



Anticoagulation

Vitamin K deficiency: ↓ synthesis of factors II, VII, IX, X, protein C, protein S.

Warfarin inhibits vitamin K epoxide reductase. Vitamin K administration can potentially reverse inhibitory effect of warfarin on clotting factor synthesis (delayed). FFP or PCC administration reverses action of warfarin immediately and can be given with vitamin K in cases of severe bleeding.

Neonates lack enteric bacteria, which produce vitamin K. Early administration of vitamin K overcomes neonatal deficiency/coagulopathy.

Factor VII (seven)—shortest half-life.

Factor II (two)—longest (tallest) half-life.

Antithrombin inhibits thrombin (factor IIa) and factors VIIa, IXa, Xa, XIa, XIIa.







Heparin enhances the activity of antithrombin. Principal targets of antithrombin: thrombin and factor Xa.

Factor V Leiden mutation produces a factor V resistant to inhibition by activated protein C.





tPA is used clinically as a thrombolytic.

▶ HEMATOLOGY AND ONCOLOGY—PATHOLOGY






RBC morphology

TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Acanthocytes ("spur cells")		Liver disease, abetalipoproteinemia	Projections of varying size at irregular intervals.
Echinocytes ("burr cells")		Liver disease, ESRD, pyruvate kinase deficiency	Smaller and more uniform projections than acanthocytes
Dacryocytes ("teardrop cells")		Bone marrow infiltration (eg, myelofibrosis)	RBC "sheds a tear " because it's mechanically squeezed out of its home in the bone marrow
Schistocytes (eg, "helmet" cells)		MAHAs (eg, DIC, TTP/HUS, HELLP syndrome), mechanical hemolysis (eg, heart valve prosthesis)	Fragmented RBCs
Degmacytes ("bite cells")		G6PD deficiency	Due to removal of Heinz bodies by splenic macrophages
Elliptocytes		Hereditary elliptocytosis	Caused by mutation in genes encoding RBC membrane proteins (eg, spectrin)


RBC morphology (continued)

TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Spherocytes		Hereditary spherocytosis, autoimmune hemolytic anemia	Small, spherical cells without central pallor
Macro-ovalocytes		Megaloblastic anemia (also hypersegmented PMNs)	
Target cells		HbC disease, Asplenia, Liver disease, Thalassemia	“ HALT ,” said the hunter to his target
Sickle cells		Sickle cell anemia	Sickling occurs with low O ₂ conditions (eg, high altitude, acidosis)

RBC inclusions

Bone marrow			
TYPE	EXAMPLE	ASSOCIATED PATHOLOGY	NOTES
Iron granules (eg, in ringed sideroblasts)		Sideroblastic anemias (eg, lead poisoning, myelodysplastic syndromes, alcoholism)	Perinuclear mitochondria with excess iron (forming ring in ringed sideroblasts) Require Prussian blue stain to be visualized
Peripheral smear			
Howell-Jolly bodies		Functional hyposplenia (eg, sickle cell disease), asplenia	Basophilic nuclear remnants (do not contain iron) Usually removed by splenic macrophages
Basophilic stippling		Sideroblastic anemias, thalassemias	Basophilic ribosomal precipitates (do not contain iron)
Pappenheimer bodies		Sideroblastic anemia	Basophilic granules (contain iron)
Heinz bodies		G6PD deficiency	Denatured and precipitated hemoglobin (contain iron) Phagocytic removal of Heinz bodies → bite cells Requires supravital stain (eg, crystal violet) to be visualized

Anemias



Reticulocyte index

Also called corrected reticulocyte count. Used to correct falsely elevated reticulocyte count in anemia. Measures appropriate bone marrow response to anemic conditions (effective erythropoiesis). High reticulocyte index (RI) indicates compensatory RBC production; low RI indicates inadequate response to correct anemia. Calculated as:

$$RI = \text{reticulocyte \%} \times \frac{\text{actual Hct}}{\text{normal Hct}}$$

[normal Hct ≈ 45%]

Microcytic,**hypochromic anemias**

MCV < 80 fL.

Iron deficiency

↓ iron due to chronic bleeding (eg, GI loss, menorrhagia), malnutrition, absorption disorders, GI surgery (eg, gastrectomy), or ↑ demand (eg, pregnancy) → ↓ final step in heme synthesis.
 Labs: ↓ iron, ↑ TIBC, ↓ ferritin, ↑ free erythrocyte protoporphyrin, ↑ RDW, ↓ RI. Microcytosis and hypochromasia (↑ central pallor) **A**.
 Symptoms: fatigue, conjunctival pallor **B**, pica (persistent craving and compulsive eating of nonfood substances), spoon nails (koilonychia).
 May manifest as glossitis, cheilosis, **Plummer-Vinson syndrome** (triad of iron deficiency anemia, esophageal webs, and dysphagia).

α-thalassemia

α-globin gene deletions on chromosome 16 → ↓ α-globin synthesis. *cis* deletion (deletions occur on same chromosome) prevalent in Asian populations; *trans* deletion (deletions occur on separate chromosomes) prevalent in African populations. Normal is αα/αα.

NUMBER OF α-GLOBIN GENES DELETED	DISEASE	CLINICAL OUTCOME
1 (α α/α −)	α-thalassemia minima	No anemia (silent carrier)
2 (α −/α −; <i>trans</i>) or (α α/− −; <i>cis</i>)	α-thalassemia minor	Mild microcytic, hypochromic anemia; <i>cis</i> deletion may worsen outcome for the carrier's offspring
3 (− −/− α)	Hemoglobin H disease (HbH); excess β-globin forms β ₄	Moderate to severe microcytic hypochromic anemia
4 (− −/− −)	Hemoglobin Barts disease; no α-globin, excess γ-globin forms γ ₄	Hydrops fetalis; incompatible with life

β-thalassemia

Point mutations in splice sites and promoter sequences on chromosome 11 → ↓ β-globin synthesis. Prevalent in Mediterranean populations.

β-thalassemia minor (heterozygote): β chain is underproduced. Usually asymptomatic. Diagnosis confirmed by ↑ HbA₂ (> 3.5%) on electrophoresis.

β-thalassemia major (homozygote): β chain is absent → severe microcytic, hypochromic anemia with target cells and increased anisopoikilocytosis **C** requiring blood transfusion (2° hemochromatosis). Marrow expansion (“crew cut” on skull x-ray) → skeletal deformities (eg, “chipmunk” facies). Extramedullary hematopoiesis → hepatosplenomegaly. ↑ risk of parvovirus B19–induced aplastic crisis. ↑ HbF (α₂γ₂), HbA₂ (α₂δ₂). HbF is protective in the infant and disease becomes symptomatic only after 6 months, when fetal hemoglobin declines.

HbS/β-thalassemia heterozygote: mild to moderate sickle cell disease depending on amount of β-globin production.

Microcytic, hypochromic anemias (continued)**Lead poisoning**

Lead inhibits ferrochelatase and ALA dehydratase → ↓ heme synthesis and ↑ RBC protoporphyrin. Also inhibits rRNA degradation → RBCs retain aggregates of rRNA (basophilic stippling).

Symptoms of **LEAD** poisoning:

- **L**ead **L**ines on gingivae (Burton lines) and on metaphyses of long bones **D** on x-ray.
- **E**ncephalopathy and **E**rythrocyte basophilic stippling.
- **A**bdominal colic and sideroblastic **A**nemia.
- **D**rops—wrist and foot drop. **D**imercaprol and **E**DTA are 1st line of treatment.

Succimer used for chelation for kids (It “**sucks**” to be a kid who eats lead).


Exposure risk ↑ in old houses with chipped paint.

Sideroblastic anemia

Causes: genetic (eg, X-linked defect in ALA synthase gene), acquired (myelodysplastic syndromes), and reversible (alcohol is most common; also lead poisoning, vitamin B₆ deficiency, copper deficiency, drugs [eg, isoniazid, linezolid]).

Lab findings: ↑ iron, normal/↓ TIBC, ↑ ferritin. Ringed sideroblasts (with iron-laden, Prussian blue–stained mitochondria) seen in bone marrow **E**. Peripheral blood smear: basophilic stippling of RBCs. Some acquired variants may be normocytic or macrocytic.

Treatment: pyridoxine (B₆, cofactor for ALA synthase).

**Interpretation of iron studies**

	Iron deficiency	Chronic disease	Hemochromatosis	Pregnancy/OCP use
Serum iron	↓	↓	↑	—
Transferrin or TIBC	↑	↓ ^a	↓	↑
Ferritin	↓	↑	↑	—
% transferrin saturation (serum iron/TIBC)	↓↓	—/↓	↑↑	↓

↑↓ = 1° disturbance.

Transferrin—**t**ransports iron in blood.


TIBC—indirectly measures transferrin.

Ferritin—1° iron storage protein of body.

^aEvolutionary reasoning—pathogens use circulating iron to thrive. The body has adapted a system in which iron is stored within the cells of the body and prevents pathogens from acquiring circulating iron.

Macrocytic anemias

MCV > 100 fL.

	DESCRIPTION	FINDINGS
Megaloblastic anemia 	Impaired DNA synthesis → maturation of nucleus of precursor cells in bone marrow delayed relative to maturation of cytoplasm. Causes: vitamin B ₁₂ deficiency, folate deficiency, medications (eg, hydroxyurea, phenytoin, methotrexate, sulfa drugs).	RBC macrocytosis, hypersegmented neutrophils (arrow in A), glossitis.
Folate deficiency	Causes: malnutrition (eg, alcoholics), malabsorption, drugs (eg, methotrexate, trimethoprim, phenytoin), ↑ requirement (eg, hemolytic anemia, pregnancy).	↑ homocysteine, normal methylmalonic acid. No neurologic symptoms (vs B ₁₂ deficiency).
Vitamin B₁₂ (cobalamin) deficiency	Causes: pernicious anemia, malabsorption (eg, Crohn disease), pancreatic insufficiency, gastrectomy, insufficient intake (eg, veganism), <i>Diphyllobothrium latum</i> (fish tapeworm).	↑ homocysteine, ↑ methylmalonic acid. Neurologic symptoms: reversible dementia, subacute combined degeneration (due to involvement of B ₁₂ in fatty acid pathways and myelin synthesis): spinocerebellar tract, lateral corticospinal tract, dorsal column dysfunction. Folate supplementation in vitamin B ₁₂ deficiency can correct the anemia, but worsens neurologic symptoms. Historically diagnosed with the Schilling test, a test that determines if the cause is dietary insufficiency vs malabsorption. Anemia 2° to insufficient intake may take several years to develop due to liver's ability to store B ₁₂ (as opposed to folate deficiency).
Orotic aciduria	Inability to convert orotic acid to UMP (de novo pyrimidine synthesis pathway) because of defect in UMP synthase. Autosomal recessive. Presents in children as failure to thrive, developmental delay, and megaloblastic anemia refractory to folate and B ₁₂ . No hyperammonemia (vs ornithine transcarbamylase deficiency—↑ orotic acid with hyperammonemia).	Orotic acid in urine. Treatment: uridine monophosphate or uridine triacetate to bypass mutated enzyme.
Nonmegaloblastic anemia	Macrocytic anemia in which DNA synthesis is normal. Causes: alcoholism, liver disease.	RBC macrocytosis without hypersegmented neutrophils.
Diamond-Blackfan anemia	A congenital form of pure red cell aplasia. Rapid-onset anemia within 1st year of life due to intrinsic defect in erythroid progenitor cells.	↑ % HbF (but ↓ total Hb). Short stature, craniofacial abnormalities, and upper extremity malformations (triphalangeal thumbs) in up to 50% of cases.

Normocytic, normochromic anemias

Normocytic, normochromic anemias are classified as nonhemolytic or hemolytic. The hemolytic anemias are further classified according to the cause of the hemolysis (intrinsic vs extrinsic to the RBC) and by the location of the hemolysis (intravascular vs extravascular). Hemolysis can lead to increases in LDH, reticulocytes, unconjugated bilirubin, pigmented gallstones, and urobilinogen in urine.

Intravascular hemolysis


Findings: ↓ haptoglobin, ↑ schistocytes on blood smear. Characteristic hemoglobinuria, hemosiderinuria, and urobilinogen in urine. Notable causes are mechanical hemolysis (eg, prosthetic valve), paroxysmal nocturnal hemoglobinuria, microangiopathic hemolytic anemias.

Extravascular hemolysis

Mechanism: macrophages in spleen clear RBCs. Findings: spherocytes in peripheral smear (most commonly due to hereditary spherocytosis and autoimmune hemolytic anemia), no hemoglobinuria/hemosiderinuria. Can present with urobilinogen in urine.


Nonhemolytic, normocytic anemias

	DESCRIPTION	FINDINGS
Anemia of chronic disease	Inflammation (eg, ↑ IL-6) → ↑ hepcidin (released by liver, binds ferroportin on intestinal mucosal cells and macrophages, thus inhibiting iron transport) → ↓ release of iron from macrophages and ↓ iron absorption from gut. Associated with conditions such as chronic infections, neoplastic disorders, chronic kidney disease, and autoimmune diseases (eg, SLE, rheumatoid arthritis).	↓ iron, ↓ TIBC, ↑ ferritin. Normocytic, but can become microcytic. Treatment: address underlying cause of inflammation, judicious use of blood transfusion, consider erythropoiesis-stimulating agents such as EPO (eg, in chronic kidney disease).
Aplastic anemia	Caused by failure or destruction of hematopoietic stem cells due to: <ul style="list-style-type: none"> ▪ Radiation and drugs (eg, benzene, chloramphenicol, alkylating agents, antimetabolites) ▪ Viral agents (eg, EBV, HIV, hepatitis viruses) ▪ Fanconi anemia (autosomal recessive DNA repair defect → bone marrow failure); normocytosis or macrocytosis on CBC ▪ Idiopathic (immune mediated, 1° stem cell defect); may follow acute hepatitis 	↓ reticulocyte count, ↑ EPO. Pancytopenia characterized by anemia, leukopenia, and thrombocytopenia (not to be confused with aplastic crisis, which causes anemia only). Normal cell morphology, but hypocellular bone marrow with fatty infiltration A (dry bone marrow tap). Symptoms: fatigue, malaise, pallor, purpura, mucosal bleeding, petechiae, infection. Treatment: withdrawal of offending agent, immunosuppressive regimens (eg, antithymocyte globulin, cyclosporine), bone marrow allograft, RBC/platelet transfusion, bone marrow stimulation (eg, GM-CSF).





Intrinsic hemolytic anemias

	DESCRIPTION	FINDINGS
Hereditary spherocytosis	Primarily autosomal dominant. Due to defect in proteins interacting with RBC membrane skeleton and plasma membrane (eg, ankyrin, band 3, protein 4.2, spectrin). Small, round RBCs with less surface area and no central pallor (↑ MCHC) → premature removal by spleen (extravascular hemolysis).	Splenomegaly, aplastic crisis (parvovirus B19 infection). Labs: ↓ mean fluorescence of RBCs in eosin 5-maleimide (EMA) binding test, ↑ fragility in osmotic fragility test. Normal to ↓ MCV with abundance of RBCs. Treatment: splenectomy.
G6PD deficiency	X-linked recessive. G6PD defect → ↓ NADPH → ↓ reduced glutathione → ↑ RBC susceptibility to oxidative stress (eg, sulfa drugs, antimalarials, fava beans) → hemolysis. Causes extravascular and intravascular hemolysis.	Back pain, hemoglobinuria a few days after oxidant stress . Labs: blood smear shows RBCs with Heinz bodies and bite cells. “ Stress makes me eat bites of fava beans with Heinz ketchup.”
Pyruvate kinase deficiency	Autosomal recessive. Pyruvate kinase defect → ↓ ATP → rigid RBCs → extravascular hemolysis. Increases levels of 2,3-BPG → ↓ hemoglobin affinity for O ₂ .	Hemolytic anemia in a newborn.
Paroxysmal nocturnal hemoglobinuria	Hematopoietic stem cell mutation → ↑ complement-mediated intravascular hemolysis, especially at night. Acquired <i>PIGA</i> mutation → impaired GPI anchor synthesis for decay-accelerating factor (DAF/CD55) and membrane inhibitor of reactive lysis (MIRL/CD59), which protect RBC membrane from complement.	Triad: Coombs ⊖ hemolytic anemia, pancytopenia, venous thrombosis (eg, Budd-Chiari syndrome). Pink/red urine in morning. Associated with aplastic anemia, acute leukemias. Labs: CD55/59 ⊖ RBCs on flow cytometry. Treatment: eculizumab (targets terminal complement protein C5).
Sickle cell anemia	Point mutation in β-globin gene → single amino acid substitution (glutamic acid → valine). Mutant HbA is termed HbS. Causes extravascular and intravascular hemolysis. Pathogenesis: low O ₂ , high altitude, or acidosis precipitates sickling (deoxygenated HbS polymerizes) → anemia, vaso-occlusive disease. Newborns are initially asymptomatic because of ↑ HbF and ↓ HbS. Heterozygotes (sickle cell trait) have resistance to malaria. 8% of African Americans carry an HbS allele. Sickle cells are crescent-shaped RBCs A . “Crew cut” on skull x-ray due to marrow expansion from ↑ erythropoiesis (also seen in thalassemias).	Complications in sickle cell disease: <ul style="list-style-type: none"> ▪ Aplastic crisis (transient arrest of erythropoiesis due to parvovirus B19). ▪ Autosplenectomy (Howell-Jolly bodies) → ↑ risk of infection by encapsulated organisms (eg, <i>S pneumoniae</i>). ▪ Splenic infarct/sequestration crisis. ▪ <i>Salmonella</i> osteomyelitis. ▪ Painful vaso-occlusive crises: dactylitis (painful swelling of hands/feet), priapism, acute chest syndrome (respiratory distress, new pulmonary infiltrates on CXR, common cause of death), avascular necrosis, stroke. ▪ Sickling in renal medulla (↓ Po₂) → renal papillary necrosis → hematuria. Hb electrophoresis: ↓↓ HbA, ↑ HbF, ↑↑ HbS. Treatment: hydroxyurea (↑ HbF), hydration.
HbC disease	Glutamic acid-to-lyCine (lysine) mutation in β-globin. Causes extravascular hemolysis.	Patients with HbSC (1 of each mutant gene) have milder disease than HbSS patients. Blood smear in homozygotes: hemoglobin C crystals inside RBCs, target cells.



Extrinsic hemolytic anemias

	DESCRIPTION	FINDINGS
<p>Autoimmune hemolytic anemia</p> 	<p>A normocytic anemia that is usually idiopathic and Coombs ⊕. Two types:</p> <ul style="list-style-type: none"> ▪ Warm AIHA—chronic anemia in which IgG causes RBC agglutination. Seen in SLE and CLL and with certain drugs (eg, α-methyl dopa). “Warm weather is Good.” ▪ Cold AIHA—acute anemia in which IgM + complement causes RBC agglutination upon exposure to cold → painful, blue fingers and toes. Seen in CLL, <i>Mycoplasma pneumoniae</i> infections, infectious Mononucleosis. 	<p>Spherocytes and agglutinated RBCs A on peripheral blood smear.</p> <p>Warm AIHA treatment: steroids, rituximab, splenectomy (if refractory).</p> <p>Cold AIHA treatment: cold avoidance, rituximab.</p> <p>Direct Coombs test—anti-Ig antibody (Coombs reagent) added to patient’s RBCs. RBCs agglutinate if RBCs are coated with Ig.</p> <p>For comparison, Indirect Coombs test—normal RBCs added to patient’s serum. If serum has anti-RBC surface Ig, RBCs agglutinate when Coombs reagent added.</p>



<p>Microangiopathic hemolytic anemia</p>	<p>RBCs are damaged when passing through obstructed or narrowed vessels. Causes intravascular hemolysis. Seen in DIC, TTP/HUS, SLE, HELLP syndrome, hypertensive emergency.</p>	<p>Schistocytes (eg, “helmet cells”) are seen on peripheral blood smear due to mechanical destruction (<i>schisto</i> = to split) of RBCs.</p>
<p>Macroangiopathic hemolytic anemia</p>	<p>Prosthetic heart valves and aortic stenosis may also cause hemolytic anemia 2° to mechanical destruction of RBCs.</p>	<p>Schistocytes on peripheral blood smear.</p>
<p>Hemolytic anemia due to infection</p>	<p>↑ destruction of RBCs (eg, malaria, <i>Babesia</i>).</p>	

Leukopenias

CELL TYPE	CELL COUNT	CAUSES
Neutropenia	Absolute neutrophil count < 1500 cells/mm ³ Severe infections typical when < 500 cells/mm ³	Sepsis/postinfection, drugs (including chemotherapy), aplastic anemia, SLE, radiation
Lymphopenia	Absolute lymphocyte count < 1500 cells/mm ³ (< 3000 cells/mm ³ in children)	HIV, DiGeorge syndrome, SCID, SLE, corticosteroids ^a , radiation, sepsis, postoperative
Eosinopenia	Absolute eosinophil count < 30 cells/mm ³	Cushing syndrome, corticosteroids ^a

^aCorticosteroids cause neutrophilia, despite causing eosinopenia and lymphopenia. Corticosteroids ↓ activation of neutrophil adhesion molecules, impairing migration out of the vasculature to sites of inflammation. In contrast, corticosteroids sequester eosinophils in lymph nodes and cause apoptosis of lymphocytes.



Neutrophil left shift


↑ neutrophil precursors, such as band cells and metamyelocytes, in peripheral blood. Usually seen with neutrophilia in the acute response to infection or inflammation. Called **leukoerythroblastic reaction** when left shift is seen with immature RBCs. Occurs with severe anemia (physiologic response) or marrow response (eg, fibrosis, tumor taking up space in marrow).

A left shift is a shift to a more immature cell in the maturation process.

Heme synthesis, porphyrias, and lead poisoning

The porphyrias are hereditary or acquired conditions of defective heme synthesis that lead to the accumulation of heme precursors. Lead inhibits specific enzymes needed in heme synthesis, leading to a similar condition.

CONDITION	AFFECTED ENZYME	ACCUMULATED SUBSTRATE	PRESENTING SYMPTOMS
<p>Lead poisoning</p> 	Ferrochelatase and ALA dehydratase	Protoporphyrin, ALA (blood)	<p>Microcytic anemia (basophilic stippling in peripheral smear A, ringed sideroblasts in bone marrow), GI and kidney disease. Children—exposure to lead paint → mental deterioration. Adults—environmental exposure (eg, batteries, ammunition) → headache, memory loss, demyelination (peripheral neuropathy).</p>
<p>Acute intermittent porphyria</p>	Porphobilinogen deaminase, previously called uroporphyrinogen I synthase (autosomal dominant mutation)	Porphobilinogen, ALA	<p>Symptoms (5 P's):</p> <ul style="list-style-type: none"> ▪ Painful abdomen ▪ Port wine–colored Pee ▪ Polyneuropathy ▪ Psychological disturbances ▪ Precipitated by drugs (eg, cytochrome P-450 inducers), alcohol, starvation <p>Treatment: hemin and glucose.</p>
<p>Porphyria cutanea tarda</p> 	Uroporphyrinogen decarboxylase	Uroporphyrin (tea-colored urine)	<p>Blistering cutaneous photosensitivity and hyperpigmentation B. Most common porphyria. Exacerbated with alcohol consumption. Causes: familial, hepatitis C. Treatment: phlebotomy, sun avoidance, antimalarials (eg, hydroxychloroquine).</p>



↓ heme → ↑ ALA synthase activity
 ↑ heme → ↓ ALA synthase activity




Iron poisoning

	Acute	Chronic
FINDINGS	High mortality rate associated with accidental ingestion by children (adult iron tablets may look like candy).	Seen in patients with 1° (hereditary) or 2° (eg, chronic blood transfusions for thalassemia or sickle cell disease) hemochromatosis.
MECHANISM	Cell death due to formation of free radicals and peroxidation of membrane lipids.	
SYMPTOMS/SIGNS	Abdominal pain, vomiting, GI bleeding. Radiopaque pill seen on x-ray. May progress to anion gap metabolic acidosis and multiorgan failure. Leads to scarring with GI obstruction.	Arthropathy, cirrhosis, cardiomyopathy, diabetes mellitus and skin pigmentation (“bronze diabetes”), hypogonadism.
TREATMENT	Chelation (eg, deferoxamine, deferasirox), gastric lavage.	Phlebotomy (patients without anemia) or chelation.

Coagulation disorders

PT—tests function of common and extrinsic pathway (factors I, II, V, VII, and X). Defect → ↑ **PT** (Play **T**ennis outside [extrinsic pathway]).
 INR (international normalized ratio) = patient PT/control PT. 1 = normal, > 1 = prolonged. Most common test used to follow patients on warfarin, which prolongs INR.
 PTT—tests function of common and **in**trinsic pathway (all factors except VII and XIII). Defect → ↑ **PTT** (Play **T**able **T**ennis **i**nside).
 Coagulation disorders can be due to clotting factor deficiencies or acquired factor inhibitors. Diagnosed with a mixing study, in which normal plasma is added to patient’s plasma. Clotting factor deficiencies should correct (the PT or PTT returns to within the appropriate normal range), whereas factor inhibitors will not correct.

DISORDER	PT	PTT	MECHANISM AND COMMENTS
<p>Hemophilia A, B, or C</p> 	—	↑	<p>Intrinsic pathway coagulation defect (↑ PTT).</p> <ul style="list-style-type: none"> ▪ A: deficiency of factor VIII; X-linked recessive. ▪ B: deficiency of factor IX; X-linked recessive. ▪ C: deficiency of factor XI; autosomal recessive. <p>Hemorrhage in hemophilia—hemarthroses (bleeding into joints, eg, knee A), easy bruising, bleeding after trauma or surgery (eg, dental procedures). Treatment: desmopressin + factor VIII concentrate (A); factor IX concentrate (B); factor XI concentrate (C).</p>
Vitamin K deficiency	↑	↑	<p>General coagulation defect. Bleeding time normal. ↓ activity of factors II, VII, IX, X, protein C, protein S.</p>

Platelet disorders

All platelet disorders have ↑ bleeding time (BT), mucous membrane bleeding, and microhemorrhages (eg, petechiae, epistaxis). Platelet count (PC) is usually low, but may be normal in qualitative disorders.

DISORDER	PC	BT	NOTES
Bernard-Soulier syndrome	–/↓	↑	Defect in adhesion. ↓ GpIb → ↓ platelet-to-vWF adhesion. Labs: abnormal ristocetin test, large platelets.
Glanzmann thrombasthenia	–	↑	Defect in aggregation. ↓ GpIIb/IIIa (↓ integrin $\alpha_{IIb}\beta_3$) → ↓ platelet-to-platelet aggregation and defective platelet plug formation. Labs: blood smear shows no platelet clumping.
Immune thrombocytopenia	↓	↑	Destruction of platelets in spleen. Anti-GpIIb/IIIa antibodies → splenic macrophages phagocytose platelets. May be idiopathic or 2° to autoimmune disorders (eg, SLE), viral illness (eg, HIV, HCV), malignancy (eg, CLL), or drug reactions. Labs: ↑ megakaryocytes on bone marrow biopsy, ↓ platelet count. Treatment: steroids, IVIG, rituximab, TPO receptor agonists (eg, eltrombopag, romiplostim), or splenectomy for refractory ITP.

Thrombotic microangiopathies

Disorders overlap significantly in symptomatology.

	Thrombotic thrombocytopenic purpura	Hemolytic-uremic syndrome
EPIDEMIOLOGY	Typically females	Typically children
PATHOPHYSIOLOGY	Inhibition or deficiency of ADAMTS13 (a vWF metalloprotease) → ↓ degradation of vWF multimers → ↑ large vWF multimers → ↑ platelet adhesion and aggregation (microthrombi formation)	Commonly caused by Shiga-like toxin from EHEC (serotype O157:H7) infection
PRESENTATION	Triad of thrombocytopenia (↓ platelets), microangiopathic hemolytic anemia (↓ Hb, schistocytes, ↑ LDH), acute kidney injury (↑ Cr)	
DIFFERENTIATING SYMPTOMS	Triad + fever + neurologic symptoms	Triad + bloody diarrhea
LABS	Normal PT and PTT helps distinguish TTP and HUS (coagulation pathway is not activated) from DIC (coagulation pathway is activated)	
TREATMENT	Plasmapheresis, steroids, rituximab	Supportive care

Mixed platelet and coagulation disorders

DISORDER	PC	BT	PT	PTT	NOTES
von Willebrand disease	—	↑	—	—/↑	Intrinsic pathway coagulation defect: ↓ vWF → ↑ PTT (vWF carries/protects factor VIII). Defect in platelet plug formation: ↓ vWF → defect in platelet-to-vWF adhesion. Autosomal dominant. Mild but most common inherited bleeding disorder. No platelet aggregation with ristocetin cofactor assay. Treatment: desmopressin, which releases vWF stored in endothelium.
Disseminated intravascular coagulation	↓	↑	↑	↑	Widespread clotting factor activation → deficiency in clotting factors → bleeding state. Causes: S nake bites, S epsis (gram ⊖), T rauma, O bstetric complications, acute P ancreatitis, M alignancy, N ephrotic syndrome, T ransfusion (SSTOP Making New Thrombi). Labs: schistocytes, ↑ fibrin degradation products (D-dimers), ↓ fibrinogen, ↓ factors V and VIII.

Hereditary thrombosis syndromes leading to hypercoagulability

DISEASE	DESCRIPTION
Antithrombin deficiency	Autosomal dominant inherited deficiency of antithrombin: has no direct effect on the PT, PTT, or thrombin time but diminishes the increase in PTT following heparin administration. Can also be acquired: renal failure/nephrotic syndrome → antithrombin loss in urine → ↓ inhibition of factors IIa and Xa.
Factor V Leiden	Autosomal dominant, most common cause of inherited hypercoagulability in Caucasians. Production of mutant factor V (guanine → adenine DNA point mutation → Arg506Gln mutation near the cleavage site) that is resistant to degradation by activated protein C. Complications include DVT, cerebral vein thrombosis, recurrent pregnancy loss.
Protein C or S deficiency	↓ ability to inactivate factors Va and VIIIa. ↑ risk of thrombotic skin necrosis with hemorrhage after administration of warfarin. If this occurs, think protein C deficiency. Together, protein C Cancels, and protein S Stops, coagulation.
Prothrombin gene mutation	Mutation in 3' untranslated region → ↑ production of prothrombin → ↑ plasma levels and venous clots.

Blood transfusion therapy

COMPONENT	DOSAGE EFFECT	CLINICAL USE
Packed RBCs	↑ Hb and O ₂ carrying capacity	Acute blood loss, severe anemia
Platelets	↑ platelet count (↑ ~ 5000/mm ³ /unit)	Stop significant bleeding (thrombocytopenia, qualitative platelet defects)
Fresh frozen plasma/prothrombin complex concentrate	↑ coagulation factor levels; FFP contains all coagulation factors and plasma proteins; PCC generally contains factors II, VII, IX, and X, as well as protein C and S	Cirrhosis, immediate anticoagulation reversal
Cryoprecipitate	Contains fibrinogen, factor VIII, factor XIII, vWF, and fibronectin	Coagulation factor deficiencies involving fibrinogen and factor VIII


Blood transfusion risks include infection transmission (low), transfusion reactions, iron overload (may lead to 2° hemochromatosis), hypocalcemia (citrate is a Ca²⁺ chelator), and hyperkalemia (RBCs may lyse in old blood units).

Leukemia vs lymphoma

Leukemia	Lymphoid or myeloid neoplasm with widespread involvement of bone marrow. Tumor cells are usually found in peripheral blood.
Lymphoma	Discrete tumor mass arising from lymph nodes. Presentations often blur definitions.

Hodgkin vs non-Hodgkin lymphoma

Hodgkin	Non-Hodgkin
Both may present with constitutional (“B”) signs/symptoms: low-grade fever, night sweats, weight loss.	
Localized, single group of nodes with contiguous spread (stage is strongest predictor of prognosis). Better prognosis.	Multiple lymph nodes involved; extranodal involvement common; noncontiguous spread. Worse prognosis.
Characterized by Reed-Sternberg cells.	Majority involve B cells; a few are of T-cell lineage.
Bimodal distribution: young adulthood and > 55 years; more common in men except for nodular sclerosing type.	Can occur in children and adults.
Associated with EBV.	May be associated with autoimmune diseases and viral infections (eg, HIV, EBV, HTLV).

Hodgkin lymphoma


Contains Reed-Sternberg cells: distinctive tumor giant cells; binucleate or bilobed with the 2 halves as mirror images (“owl eyes” **A**). RS cells are CD15+ and CD30+ B-cell origin. 2 owl eyes × 15 = 30.

SUBTYPE	NOTES
Nodular sclerosis	Most common
Lymphocyte rich	Best prognosis
Mixed cellularity	Eosinophilia, seen in immunocompromised patients
Lymphocyte depleted	Seen in immunocompromised patients

Non-Hodgkin lymphoma

TYPE	OCCURS IN	GENETICS	COMMENTS
Neoplasms of mature B cells			
Burkitt lymphoma	Adolescents or young adults	t(8;14)—translocation of <i>c-myc</i> (8) and heavy-chain Ig (14)	“Starry sky” appearance, sheets of lymphocytes with interspersed “tingible body” macrophages (arrows in A). Associated with EBV. Jaw lesion B in endemic form in Africa; pelvis or abdomen in sporadic form.
Diffuse large B-cell lymphoma	Usually older adults, but 20% in children	Mutations in <i>BCL-2</i> , <i>BCL-6</i>	Most common type of non-Hodgkin lymphoma in adults.
Follicular lymphoma	Adults	t(14;18)—translocation of heavy-chain Ig (14) and <i>BCL-2</i> (18)	Indolent course with painless “waxing and waning” lymphadenopathy. Bcl-2 normally inhibits apoptosis.
Mantle cell lymphoma	Adult males >> adult females	t(11;14)—translocation of cyclin D1 (11) and heavy-chain Ig (14), CD5+	Very aggressive, patients typically present with late-stage disease.
Marginal zone lymphoma	Adults	t(11;18)	Associated with chronic inflammation (eg, Sjögren syndrome, chronic gastritis [MALT lymphoma]).
Primary central nervous system lymphoma	Adults	EBV related; associated with HIV/AIDS	Considered an AIDS-defining illness. Variable presentation: confusion, memory loss, seizures. CNS mass (often single, ring-enhancing lesion on MRI) in immunocompromised patients C , needs to be distinguished from toxoplasmosis via CSF analysis or other lab tests.
Neoplasms of mature T cells			
Adult T-cell lymphoma	Adults	Caused by HTLV (associated with IV drug abuse)	Adults present with cutaneous lesions; common in Japan (T -cell in T okyo), West Africa, and the Caribbean. Lytic bone lesions, hypercalcemia.
Mycosis fungoides/Sézary syndrome	Adults		Mycosis fungoides: skin patches and plaques D (cutaneous T-cell lymphoma), characterized by atypical CD4+ cells with “cerebriform” nuclei and intraepidermal neoplastic cell aggregates (Pautrier microabscess). May progress to Sézary syndrome (T-cell leukemia).



Plasma cell dyscrasias

Characterized by monoclonal immunoglobulin (Ig) overproduction due to plasma cell disorder. Labs: serum protein electrophoresis (SPEP) or free light chain (FLC) assay for initial tests (M spike on SPEP represents overproduction of a monoclonal Ig fragment). For urinalysis, use 24-hr urine protein electrophoresis (UPEP) to detect light chain, as routine urine dipstick detects only albumin.

Confirm with bone marrow biopsy.

Multiple myeloma

Overproduction of IgG (55% of cases) > IgA.

Clinical features: **CRAB**

- Hyper**C**alcemia
- **R**enal involvement
- **A**nemia
- Bone lytic lesions (“punched out” on X-ray **A**) → **B**ack pain.

Peripheral blood smear shows Rouleaux formation **B** (RBCs stacked like poker chips).

Urinalysis shows Ig light chains (Bence Jones proteinuria) with \ominus urine dipstick.

Bone marrow analysis shows > 10% monoclonal plasma cells with clock-face chromatin **C** and intracytoplasmic inclusions containing IgG.

Complications: \uparrow infection risk, 1° amyloidosis (AL).

Waldenstrom macroglobulinemia

Overproduction of IgM (macroglobulinemia because IgM is the **largest** Ig).

Clinical features:

- Peripheral neuropathy
- No CRAB findings
- Hyperviscosity syndrome:
 - Headache
 - Blurry vision
 - Raynaud phenomenon
 - Retinal hemorrhages

Bone marrow analysis shows >10% small lymphocytes with IgM-containing vacuoles (lymphoplasmacytic lymphoma).

Complication: thrombosis.


Monoclonal gammopathy of undetermined significance


Overproduction of any Ig type.

Usually asymptomatic. No CRAB findings.

Bone marrow analysis shows < 10% monoclonal plasma cells.

Complication: 1-2% risk per year of transitioning to multiple myeloma.



Myelodysplastic syndromes

Stem cell disorders involving ineffective hematopoiesis → defects in cell maturation of nonlymphoid lineages. Caused by de novo mutations or environmental exposure (eg, radiation, benzene, chemotherapy). Risk of transformation to AML.

Pseudo-Pelger-Huet anomaly—neutrophils with bilobed (“duet”) nuclei **A**. Typically seen after chemotherapy.

Leukemias

Unregulated growth and differentiation of WBCs in bone marrow → marrow failure → anemia (↓ RBCs), infections (↓ mature WBCs), and hemorrhage (↓ platelets). Usually presents with ↑ circulating WBCs (malignant leukocytes in blood); rare cases present with normal/↓ WBCs. Leukemic cell infiltration of liver, spleen, lymph nodes, and skin (leukemia cutis) possible.

TYPE

NOTES

Lymphoid neoplasms**Acute lymphoblastic leukemia/lymphoma**

Most frequently occurs in children; less common in adults (worse prognosis). T-cell ALL can present as mediastinal mass (presenting as SVC-like syndrome). Associated with Down syndrome. Peripheral blood and bone marrow have ↑↑↑ lymphoblasts **A**. TdT+ (marker of pre-T and pre-B cells), CD10+ (marker of pre-B cells). Most responsive to therapy. May spread to CNS and testes. t(12;21) → better prognosis.

Chronic lymphocytic leukemia/small lymphocytic lymphoma

Age > 60 years. Most common adult leukemia. CD20+, CD23+, CD5+ B-cell neoplasm. Often asymptomatic, progresses slowly; smudge cells **B** in peripheral blood smear; autoimmune hemolytic anemia. **CLL = Crushed Little Lymphocytes** (smudge cells). Richter transformation—CLL/SLL transformation into an aggressive lymphoma, most commonly diffuse large B-cell lymphoma (DLBCL).

Hairy cell leukemia

Adult males. Mature B-cell tumor. Cells have filamentous, hair-like projections (fuzzy appearing on LM **C**). Peripheral lymphadenopathy is uncommon. Causes marrow fibrosis → dry tap on aspiration. Patients usually present with massive splenomegaly and pancytopenia. Stains **TRAP** (tartrate-resistant acid phosphatase) ⊕ (**trapped in a hairy situation**). TRAP stain largely replaced with flow cytometry. Associated with *BRAF* mutations. Treatment: cladribine, pentostatin.


Myeloid neoplasms**Acute myelogenous leukemia**

Median onset 65 years. Auer rods **D**; myeloperoxidase ⊕ cytoplasmic inclusions seen mostly in APL (formerly M3 AML); ↑↑↑ circulating myeloblasts on peripheral smear. Risk factors: prior exposure to alkylating chemotherapy, radiation, myeloproliferative disorders, Down syndrome. APL: t(15;17), responds to all-*trans* retinoic acid (vitamin A) and arsenic, which induce differentiation of promyelocytes; DIC is a common presentation.

Leukemias (continued)

Chronic myelogenous leukemia

Peak incidence: 45–85 years; median age: 64 years. Defined by the Philadelphia chromosome (t[9;22], *BCR-ABL*) and myeloid stem cell proliferation. Presents with dysregulated production of mature and maturing granulocytes (eg, neutrophils, metamyelocytes, myelocytes, basophils **E**) and splenomegaly. May accelerate and transform to AML or ALL (“blast crisis”). Very low leukocyte alkaline phosphatase (LAP) as a result of low activity in malignant neutrophils, vs benign neutrophilia (leukemoid reaction) in which LAP is ↑ due to ↑ leukocyte count with neutrophilia in response to stressors (eg, infections, medications, severe hemorrhage). Responds to bcr-abl tyrosine kinase inhibitors (eg, imatinib).



Chronic myeloproliferative disorders

Malignant hematopoietic neoplasms with varying impacts on WBCs and myeloid cell lines.

Polycythemia vera

Primary polycythemia. Disorder of ↑ RBCs, usually due to acquired *JAK2* mutation. May present as intense itching after shower (aquagenic pruritus). Rare but classic symptom is erythromelalgia (severe, burning pain and red-blue coloration) due to episodic blood clots in vessels of the extremities **A**. ↓ EPO (vs 2° polycythemia, which presents with endogenous or artificially ↑ EPO). Treatment: phlebotomy, hydroxyurea, ruxolitinib (*JAK1/2* inhibitor).


Essential thrombocythemia

Characterized by massive proliferation of megakaryocytes and platelets. Symptoms include bleeding and thrombosis. Blood smear shows markedly increased number of platelets, which may be large or otherwise abnormally formed **B**. Erythromelalgia may occur.

Myelofibrosis

Obliteration of bone marrow with fibrosis **C** due to ↑ fibroblast activity. Associated with massive splenomegaly and “teardrop” RBCs **D**. “Bone marrow **cries** because it’s fibrosed and is a dry tap.”

	RBCs	WBCs	PLATELETS	PHILADELPHIA CHROMOSOME	<i>JAK2</i> MUTATIONS
Polycythemia vera	↑	↑	↑	⊖	⊕
Essential thrombocythemia	–	–	↑	⊖	⊕ (30–50%)
Myelofibrosis	↓	Variable	Variable	⊖	⊕ (30–50%)
CML	↓	↑	↑	⊕	⊖



Polycythemia

	PLASMA VOLUME	RBC MASS	O ₂ SATURATION	EPO LEVELS	ASSOCIATIONS
Relative	↓	–	–	–	Dehydration, burns.
Appropriate absolute	–	↑	↓	↑	Lung disease, congenital heart disease, high altitude.
Inappropriate absolute	–	↑	–	↑	Exogenous EPO: athlete abuse (“blood doping”). Inappropriate EPO secretion: malignancy (eg, renal cell carcinoma, hepatocellular carcinoma).
Polycythemia vera	↑	↑↑	–	↓	EPO ↓ in PCV due to negative feedback suppressing renal EPO production.


↑↓ = 1° disturbance

Chromosomal translocations


TRANSLOCATION	ASSOCIATED DISORDER	NOTES
t(8;14)	Burkitt (Burk-8) lymphoma (<i>c-myc</i> activation)	The Ig heavy chain genes on chromosome 14 are constitutively expressed. When other genes (eg, <i>c-myc</i> and <i>BCL-2</i>) are translocated next to this heavy chain gene region, they are overexpressed.
t(11;14)	Mantle cell lymphoma (cyclin D1 activation)	
t(11;18)	Marginal zone lymphoma	
t(14;18)	Follicular lymphoma (<i>BCL-2</i> activation)	
t(15;17)	APL (M3 type of AML; responds to all-trans retinoic acid)	
t(9;22) (Philadelphia chromosome)	CML (<i>BCR-ABL</i> hybrid), ALL (less common, poor prognostic factor); Philadelphia CreaML cheese	

Langerhans cell histiocytosis

Collective group of proliferative disorders of Langerhans cells. Presents in a child as lytic bone lesions **A** and skin rash or as recurrent otitis media with a mass involving the mastoid bone. Cells are functionally immature and do not effectively stimulate primary T cells via antigen presentation. Cells express S-100 (mesodermal origin) and CD1a. Birbeck granules (“tennis rackets” or rod shaped on EM) are characteristic **B**.




Tumor lysis syndrome



Oncologic emergency triggered by massive tumor cell lysis, most often in lymphomas/leukemias. Release of K^+ → hyperkalemia, release of PO_4^{3-} → hyperphosphatemia, hypocalcemia due to Ca^{2+} sequestration by PO_4^{3-} . \uparrow nucleic acid breakdown → hyperuricemia → acute kidney injury. Prevention and treatment include aggressive hydration, allopurinol, rasburicase.

Hemophagocytic lymphohistiocytosis



Systemic overactivation of macrophages and cytotoxic T cells → fever, pancytopenia, hepatosplenomegaly, $\uparrow\uparrow\uparrow$ serum ferritin levels. Can be inherited or 2° to strong immunologic activation (eg, after EBV infection, malignancy). Bone marrow biopsy shows macrophages phagocytosing marrow elements **A**.

▶ HEMATOLOGY AND ONCOLOGY—PHARMACOLOGY

Direct thrombin inhibitors

Bivalirudin, Argatroban, Dabigatran (only oral agent in class).

MECHANISM

Directly inhibits activity of free and clot-associated thrombin.

CLINICAL USE

Venous thromboembolism, atrial fibrillation. Can be used in HIT, when heparin is **BAD** for the patient. Does not require lab monitoring.

ADVERSE EFFECTS

Bleeding; can reverse dabigatran with idarucizumab. Consider PCC and/or antifibrinolytics (eg, tranexamic acid) if no reversal agent available.

Heparin

MECHANISM	Activates antithrombin, which ↓ action of IIa (thrombin) and factor Xa. Short half-life.
CLINICAL USE	Immediate anticoagulation for pulmonary embolism (PE), acute coronary syndrome, MI, deep venous thrombosis (DVT). Used during pregnancy (does not cross placenta). Follow PTT.
ADVERSE EFFECTS	Bleeding, thrombocytopenia (HIT), osteoporosis, drug-drug interactions. For rapid reversal (antidote), use protamine sulfate (positively charged molecule that binds negatively charged heparin).
NOTES	Low-molecular-weight heparins (eg, enoxaparin, dalteparin)—act predominantly on factor Xa. Fondaparinux acts only on factor Xa. Have better bioavailability and 2–4× longer half life than unfractionated heparin; can be administered subcutaneously and without laboratory monitoring. LMWHs undergo renal clearance (vs hepatic clearance of unfractionated heparin) and are contraindicated in renal insufficiency. Not easily reversible.

Heparin-induced thrombocytopenia (HIT) type 2—development of IgG antibodies against heparin-bound platelet factor 4 (PF4). Antibody-heparin-PF4 complex activates platelets → thrombosis and thrombocytopenia. Highest risk with unfractionated heparin. **HIT type 1** characterized by nonimmunologic milder drop in platelet count, usually asymptomatic.

Warfarin

MECHANISM	Inhibits epoxide reductase, which interferes with γ -carboxylation of vitamin K–dependent clotting factors II, VII, IX, X, and proteins C, S. Metabolism affected by polymorphisms in the gene for vitamin K epoxide reductase complex (<i>VKORC1</i>). In laboratory assay, has effect on EX trinsic pathway and ↑ PT . Long half-life.	The EX -Presiden T went to war (farin).
CLINICAL USE	Chronic anticoagulation (eg, venous thromboembolism prophylaxis, and prevention of stroke in atrial fibrillation). Not used in pregnant women (because warfarin, unlike heparin, crosses placenta). Follow PT/INR.	
ADVERSE EFFECTS	Bleeding, teratogenic, skin/tissue necrosis A , drug-drug interactions. Initial risk of hypercoagulation: protein C has a shorter half-life than factors II and X. Existing protein C depletes before existing factors II and X deplete, and before warfarin can reduce factors II and X production → hypercoagulation. Skin/tissue necrosis within first few days of large doses believed to be due to small vessel microthrombosis.	For reversal of warfarin, give vitamin K. For rapid reversal, give fresh frozen plasma (FFP) or PCC. Heparin “bridging”: heparin frequently used when starting warfarin. Heparin’s activation of antithrombin enables anticoagulation during initial, transient hypercoagulable state caused by warfarin. Initial heparin therapy reduces risk of recurrent venous thromboembolism and skin/tissue necrosis. Metabolized by cytochrome P-450.



Heparin vs warfarin

	Heparin	Warfarin
ROUTE OF ADMINISTRATION	Parenteral (IV, SC)	Oral
SITE OF ACTION	Blood	Liver
ONSET OF ACTION	Rapid (seconds)	Slow, limited by half-lives of normal clotting factors
MECHANISM OF ACTION	Activates antithrombin, which ↓ the action of IIa (thrombin) and factor Xa	Impairs synthesis of vitamin K–dependent clotting factors II, VII, IX, and X, and anti-clotting proteins C and S
DURATION OF ACTION	Hours	Days
AGENTS FOR REVERSAL	Protamine sulfate	Vitamin K, FFP, PCC
MONITORING	PTT (intrinsic pathway)	PT/INR (extrinsic pathway)
CROSSES PLACENTA	No	Yes (teratogenic)

Direct factor Xa inhibitors

	Api X aban, rivaro X aban.
MECHANISM	Bind to and directly inhibit factor Xa .
CLINICAL USE	Treatment and prophylaxis for DVT and PE; stroke prophylaxis in patients with atrial fibrillation. Oral agents do not usually require coagulation monitoring.
ADVERSE EFFECTS	Bleeding. Reverse with ande X anet alfa.

Thrombolytics

	Alteplase (tPA), reteplase (rPA), streptokinase, tenecteplase (TNK-tPA).
MECHANISM	Directly or indirectly aid conversion of plasminogen to plasmin, which cleaves thrombin and fibrin clots. ↑ PT, ↑ PTT, no change in platelet count.
CLINICAL USE	Early MI, early ischemic stroke, direct thrombolysis of severe PE.
ADVERSE EFFECTS	Bleeding. Contraindicated in patients with active bleeding, history of intracranial bleeding, recent surgery, known bleeding diatheses, or severe hypertension. Nonspecific reversal with antifibrinolytics (eg, aminocaproic acid, tranexamic acid), platelet transfusions, and factor corrections (eg, cryoprecipitate, FFP, PCC).

ADP receptor inhibitors

	Clopidogrel, prasugrel, ticagrelor (reversible), ticlopidine.
MECHANISM	Irreversibly block ADP (P2Y ₁₂) receptor, which prevents subsequent platelet aggregation. Prevent expression of glycoproteins IIb/IIIa on platelet surface.
CLINICAL USE	Acute coronary syndrome; coronary stenting. ↓ incidence or recurrence of thrombotic stroke.
ADVERSE EFFECTS	Neutropenia (ticlopidine). TTP may be seen.

Glycoprotein IIb/IIIa inhibitors

Abciximab, eptifibatide, tirofiban.

MECHANISM

Bind to the glycoprotein receptor IIb/IIIa (fibrinogen receptor) on activated platelets, preventing aggregation. Abciximab is made from monoclonal antibody Fab fragments.


CLINICAL USE

Unstable angina, percutaneous coronary intervention.


ADVERSE EFFECTS

Bleeding, thrombocytopenia.

Cancer therapy—cell cycle



Cancer therapy—targets



Antitumor antibiotics

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Bleomycin	Induces free radical formation → breaks in DNA strands.	Testicular cancer, Hodgkin lymphoma.	Pulmonary fibrosis, skin hyperpigmentation. Minimal myelosuppression.
Dactinomycin (actinomycin D)	Intercalates into DNA, preventing RNA synthesis.	Wilms tumor, Ewing sarcoma, rhabdomyosarcoma. Used for childhood tumors.	Myelosuppression.
Anthracyclines (eg, doxorubicin, daunorubicin)	Generate free radicals. Intercalate in DNA → breaks in DNA → ↓ replication. Interferes with topoisomerase II enzyme.	Solid tumors, leukemias, lymphomas.	Cardiotoxicity (dilated cardiomyopathy), myelosuppression, alopecia. Dexrazoxane (iron chelating agent) used to prevent cardiotoxicity.

Antimetabolites

DRUG	MECHANISM ^a	CLINICAL USE	ADVERSE EFFECTS
Azathioprine, 6-mercaptopurine	Purine (thiol) analogs → ↓ de novo purine synthesis. Activated by HGPRT. Azathioprine is metabolized into 6-MP.	Preventing organ rejection, rheumatoid arthritis, IBD, SLE; used to wean patients off steroids in chronic disease and to treat steroid-refractory chronic disease.	Myelosuppression; GI, liver toxicity. Azathioprine and 6-MP are metabolized by xanthine oxidase; thus both have ↑ risk of toxicity with allopurinol or febuxostat.
Cladribine	Purine analog → multiple mechanisms (eg, inhibition of DNA polymerase, DNA strand breaks).	Hairy cell leukemia.	Myelosuppression, nephrotoxicity, and neurotoxicity.
Cytarabine (arabinofuranosyl cytidine)	Pyrimidine analog → DNA chain termination. At higher concentrations, inhibits DNA polymerase.	Leukemias (AML), lymphomas.	Myelosuppression with megaloblastic anemia. CYT arabine causes pan CYT openia.
5-fluorouracil	Pyrimidine analog bioactivated to 5-FdUMP, which covalently complexes with thymidylate synthase and folic acid. Capecitabine is a prodrug. This complex inhibits thymidylate synthase → ↓ dTMP → ↓ DNA synthesis.	Colon cancer, pancreatic cancer, actinic keratosis, basal cell carcinoma (topical). Effects enhanced with the addition of leucovorin.	Myelosuppression, palmar-plantar erythrodysesthesia (hand-foot syndrome).
Methotrexate	Folic acid analog that competitively inhibits dihydrofolate reductase → ↓ dTMP → ↓ DNA synthesis.	Cancers: leukemias (ALL), lymphomas, choriocarcinoma, sarcomas. Non-neoplastic: ectopic pregnancy, medical abortion (with misoprostol), rheumatoid arthritis, psoriasis, IBD, vasculitis.	Myelosuppression, which is reversible with leucovorin (folinic acid) “rescue.” Hepatotoxicity. Mucositis (eg, mouth ulcers). Pulmonary fibrosis. Folate deficiency, which may be teratogenic (neural tube defects) without supplementation. Nephrotoxicity.

^aAll are S-phase specific except cladribine, which is cell cycle nonspecific.

Alkylating agents

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Busulfan	Cross-links DNA.	Used to ablate patient's bone marrow before bone marrow transplantation.	Severe myelosuppression (in almost all cases), pulmonary fibrosis, hyperpigmentation.
Cyclophosphamide, ifosfamide	Cross-link DNA at guanine. Require bioactivation by liver. A nitrogen mustard.	Solid tumors, leukemia, lymphomas, rheumatic disease (eg, SLE, granulomatosis with polyangiitis).	Myelosuppression; SIADH; Fanconi syndrome (ifosfamide); hemorrhagic cystitis and bladder cancer, prevented with mesna (sulfhydryl group of mesna binds toxic metabolites) and adequate hydration.
Nitrosoureas (eg, carmustine, lomustine)	Require bioactivation. Cross blood-brain barrier → CNS. Cross-link DNA.	Brain tumors (including glioblastoma multiforme).	CNS toxicity (convulsions, dizziness, ataxia).
Procarbazine	Cell cycle phase–nonspecific alkylating agent, mechanism unknown. Also a weak MAO inhibitor.	Hodgkin lymphoma, brain tumors.	Bone marrow suppression, pulmonary toxicity, leukemia, disulfiram-like reaction, tyramine-induced hypertensive crisis with consumption of tyramine-rich foods (eg, aged cheese, wine, fava beans).

Microtubule inhibitors

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Paclitaxel, other taxanes	Hyper stabilize polymerized microtubules in M phase so that mitotic spindle cannot break down (anaphase cannot occur).	Ovarian and breast carcinomas.	Myelosuppression, neuropathy, hypersensitivity. Taxes stabilize society.
Vincristine, vinblastine	Vinca alkaloids that bind β -tubulin and inhibit its polymerization into microtubules → prevent mitotic spindle formation (M-phase arrest).	Solid tumors, leukemias, Hodgkin and non-Hodgkin lymphomas.	Vincristine : neurotoxicity (areflexia, peripheral neuritis), constipation (including paralytic ileus). Crisps the nerves. Vinblastine : bone marrow suppression. Blasts the bone marrow.

Cisplatin, carboplatin, oxaliplatin

MECHANISM	Cross-link DNA.
CLINICAL USE	Testicular, bladder, ovary, GI, and lung carcinomas.
ADVERSE EFFECTS	Nephrotoxicity (including Fanconi syndrome), peripheral neuropathy, ototoxicity. Prevent nephrotoxicity with amifostine (free radical scavenger) and chloride (saline) diuresis.

Etoposide, teniposide

MECHANISM	Inhibit topoisomerase II → ↑ DNA degradation (cell cycle arrest in G ₂ and S phases).
CLINICAL USE	Solid tumors (particularly testicular and small cell lung cancer), leukemias, lymphomas.
ADVERSE EFFECTS	Myelosuppression, alopecia.

Irinotecan, topotecan

MECHANISM	Inhibit topoisomerase I and prevent DNA unwinding and replication.
CLINICAL USE	Colon cancer (irinotecan); ovarian and small cell lung cancers (topotecan).
ADVERSE EFFECTS	Severe myelosuppression, diarrhea.

Hydroxyurea

MECHANISM	Inhibits ribonucleotide reductase → ↓ DNA Synthesis (S-phase specific).
CLINICAL USE	Myeloproliferative disorders (eg, CML, polycythemia vera), sickle cell disease (↑ HbF).
ADVERSE EFFECTS	Severe myelosuppression, megaloblastic anemia.

Bevacizumab

MECHANISM	Monoclonal antibody against VEGF. Inhibits angiogenesis (BeVacizumab inhibits Blood Vessel formation).
CLINICAL USE	Solid tumors (eg, colorectal cancer, renal cell carcinoma), wet age-related macular degeneration.
ADVERSE EFFECTS	Hemorrhage, blood clots, and impaired wound healing.

Erlotinib

MECHANISM	EGFR tyrosine kinase inhibitor.
CLINICAL USE	Non-small cell lung cancer.
ADVERSE EFFECTS	Rash, diarrhea.

Cetuximab, panitumumab

MECHANISM	Monoclonal antibodies against EGFR.
CLINICAL USE	Stage IV colorectal cancer (wild-type KRAS), head and neck cancer.
ADVERSE EFFECTS	Rash, elevated LFTs, diarrhea.

Imatinib, dasatinib, nilotinib

MECHANISM	Tyrosine kinase inhibitors of bcr-abl (encoded by Philadelphia chromosome fusion gene in CML) and <i>c-kit</i> (common in GI stromal tumors).
CLINICAL USE	CML, GI stromal tumors (GISTs).
ADVERSE EFFECTS	Fluid retention.

Rituximab

MECHANISM	Monoclonal antibody against CD20, which is found on most B-cell neoplasms.
CLINICAL USE	Non-Hodgkin lymphoma, CLL, ITP, rheumatoid arthritis, TTP, AIHA.
ADVERSE EFFECTS	↑ risk of progressive multifocal leukoencephalopathy.

Bortezomib, carfilzomib

MECHANISM	Proteasome inhibitors, induce arrest at G2-M phase and apoptosis.
CLINICAL USE	Multiple myeloma, mantle cell lymphoma.
ADVERSE EFFECTS	Peripheral neuropathy, herpes zoster reactivation.

Tamoxifen, raloxifene

MECHANISM	Selective estrogen receptor modulators (SERMs)—receptor antagonists in breast and agonists in bone. Block the binding of estrogen to ER ⊕ cells.
CLINICAL USE	Breast cancer treatment (tamoxifen only) and prevention. Raloxifene also useful to prevent osteoporosis.
ADVERSE EFFECTS	Tamoxifen—partial agonist in endometrium, which ↑ the risk of endometrial cancer. Raloxifene —no ↑ in endometrial carcinoma (so you can relax!), because it is an estrogen receptor antagonist in endometrial tissue. Both ↑ risk of thromboembolic events (eg, DVT, PE) and “hot flashes.”

Trastuzumab


MECHANISM	Monoclonal antibody against HER-2 (<i>c-erbB2</i>), a tyrosine kinase receptor. Helps kill cancer cells that overexpress HER-2 through inhibition of HER-2 initiated cellular signaling and antibody-dependent cytotoxicity.
CLINICAL USE	HER-2 ⊕ breast cancer and gastric cancer (tras2zumab).
ADVERSE EFFECTS	Dilated cardiomyopathy. “ Heartceptin ” damages the heart .

Dabrafenib, vemurafenib

MECHANISM	Small molecule inhibitors of <i>BRAF</i> oncogene ⊕ melanoma. VEmuRAF-enib is for V600E -mutated <i>BRAF</i> inhibition . Often co-administered with MEK inhibitors (eg, trametinib).
CLINICAL USE	Metastatic melanoma.

Rasburicase

MECHANISM	Recombinant uricase that catalyzes metabolism of uric acid to allantoin.
CLINICAL USE	Prevention and treatment of tumor lysis syndrome.

Key chemotoxicities

Cisplatin/Carboplatin → ototoxicity

Vincristine → peripheral neuropathy

Bleomycin, Busulfan → pulmonary fibrosis

Doxorubicin → cardiotoxicity

Trastuzumab → cardiotoxicity

Cisplatin/Carboplatin → nephrotoxicity

CYclophosphamide → hemorrhagic cystitis

Nonspecific common toxicities of nearly all cytotoxic chemotherapies include myelosuppression (neutropenia, anemia, thrombocytopenia), GI toxicity (nausea, vomiting, mucositis), alopecia.

HIGH-YIELD SYSTEMS

Musculoskeletal, Skin, and Connective Tissue

“Rigid, the skeleton of habit alone upholds the human frame.”

—Virginia Woolf

“Beauty may be skin deep, but ugly goes clear to the bone.”

—Redd Foxx

“The function of muscle is to pull and not to push, except in the case of the genitals and the tongue.”

—Leonardo da Vinci


“To thrive in life you need three bones. A wishbone. A backbone. And a funny bone.”

—Reba McEntire

This chapter provides information you will need to understand certain anatomical dysfunctions, rheumatic diseases, and dermatologic conditions. Be able to interpret 3D anatomy in the context of radiologic imaging. For the rheumatic diseases, create instructional cases or personas that include the most likely presentation and symptoms: risk factors, gender, important markers (eg, autoantibodies), and other epidemiologic factors. Doing so will allow you to answer the higher order questions that are likely to be asked on the exam.

▶ Anatomy and Physiology	446
▶ Pathology	459
▶ Dermatology	473
▶ Pharmacology	485

► MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—ANATOMY AND PHYSIOLOGY


Rotator cuff muscles

Shoulder muscles that form the rotator cuff:

- **S**upraspinatus (suprascapular nerve)—abducts arm initially (before the action of the deltoid); most common rotator cuff injury (trauma or degeneration and impingement → tendinopathy or tear [arrow in **A**]), assessed by “empty/full can” test
- **I**nfraspinatus (suprascapular nerve)—externally rotates arm; pitching injury
- **t**eres minor (axillary nerve)—adducts and externally rotates arm
- **S**ubscapularis (upper and lower subscapular nerves)—internally rotates and adducts arm

Innervated primarily by C5-C6.

SI tS (small t is for teres **minor**).


**Arm abduction**

DEGREE	MUSCLE	NERVE
0°–15°	Supraspinatus	Suprascapular
15°–100°	Deltoid	Axillary
> 90°	Trapezius	Accessory
> 100°	S erratus A nterior	L ong T horacic (SALT)

Upper extremity nerves

NERVE	CAUSES OF INJURY	PRESENTATION
Axillary (C5-C6)	Fractured surgical neck of humerus Anterior dislocation of humerus	Flattened deltoid Loss of arm abduction at shoulder ($> 15^\circ$) Loss of sensation over deltoid and lateral arm
Musculocutaneous (C5-C7)	Upper trunk compression	↓ biceps (C5-6) reflex Weakness of forearm flexion and supination Loss of sensation over lateral forearm
Radial (C5-T1)	Compression of axilla, eg, due to crutches or sleeping with arm over chair (“Saturday night palsy”) Midshaft fracture of humerus Repetitive pronation/supination of forearm, eg, due to screwdriver use (“finger drop”)	Wrist drop: loss of elbow, wrist, and finger extension ↓ grip strength (wrist extension necessary for maximal action of flexors) Loss of sensation over posterior arm/forearm and dorsal hand
Median (C5-T1)	Supracondylar fracture of humerus → proximal lesion of the nerve Carpal tunnel syndrome and wrist laceration → distal lesion of the nerve	“Ape hand” and “Pope’s blessing” Loss of wrist flexion, flexion of lateral fingers, thumb opposition, lumbricals of index and middle fingers Loss of sensation over thenar eminence and dorsal and palmar aspects of lateral $3\frac{1}{2}$ fingers with proximal lesion
Ulnar (C8-T1)	Fracture of medial epicondyle of humerus “funny bone” (proximal lesion) Fractured hook of hamate (distal lesion) from fall on outstretched hand	“Ulnar claw” on digit extension Radial deviation of wrist upon flexion (proximal lesion) Loss of wrist flexion, flexion of medial fingers, abduction and adduction of fingers (interossei), actions of medial 2 lumbrical muscles Loss of sensation over medial $1\frac{1}{2}$ fingers including hypothenar eminence
Recurrent branch of median nerve (C5-T1)	Superficial laceration of palm	“Ape hand” Loss of thenar muscle group: opposition, abduction, and flexion of thumb No loss of sensation


Humerus fractures, proximally to distally, follow the **ARM** (Axillary → Radial → Median)



R

Brachial plexus lesions

- 1 Erb palsy (“waiter’s tip”)
- 2 Klumpke palsy (claw hand)
- 3 Wrist drop
- 4 Winged scapula
- 5 Deltoid paralysis
- 6 “Saturday night palsy” (wrist drop)
- 7 Difficulty flexing elbow, variable sensory loss
- 8 Decreased thumb function, “Pope’s blessing”
- 9 Intrinsic muscles of hand, claw hand



Randy
Travis
Drinks
Cold
Beer

CONDITION	INJURY	CAUSES	MUSCLE DEFICIT	FUNCTIONAL DEFICIT	PRESENTATION
Erb palsy (“waiter’s tip”)	Traction or tear of upper trunk : C5-C6 roots	Infants—lateral traction on neck during delivery Adults—trauma	D eltoid, supraspinatus I nfraspinatus B iceps brachii H erb gets DIB s on t ips	Abduction (arm hangs by side) Lateral rotation (arm medially rotated) Flexion, supination (arm extended and pronated)	
Klumpke palsy	Traction or tear of lower trunk : C8-T1 roots	Infants—upward force on arm during delivery Adults—trauma (eg, grabbing a tree branch to break a fall)	Intrinsic hand muscles: lumbricals, interossei, thenar, hypothenar	Total claw hand: lumbricals normally flex MCP joints and extend DIP and PIP joints	
Thoracic outlet syndrome	Compression of lower trunk and subclavian vessels, most commonly within the scalene triangle	Cervical rib (arrows in A , Pancoast tumor)	Same as Klumpke palsy	Atrophy of intrinsic hand muscles; ischemia, pain, and edema due to vascular compression	A
Winged scapula	Lesion of long thoracic nerve, roots C5-C7 (“ w ings of h eaven”)	Axillary node dissection after mastectomy, stab wounds	Serratus anterior	Inability to anchor scapula to thoracic cage → cannot abduct arm above horizontal position B	B



Wrist region




Scaphoid, Lunate, Triquetrum, Pisiform, Hamate, Capitate, Trapezoid, Trapezium **A**. (So Long To Pinky, Here Comes The Thumb)

Scaphoid (palpable in anatomic snuff box **B**) is the most commonly fractured carpal bone, typically due to a fall on an outstretched hand. Complications of proximal scaphoid fractures include avascular necrosis and nonunion due to retrograde blood supply from a branch of the radial artery. Fracture not always seen on initial x-ray.

Dislocation of lunate may cause acute carpal tunnel syndrome.



Hand muscles

Thenar (median)—**O**pponens pollicis, **A**bductor pollicis brevis, **F**lexor pollicis brevis, superficial head (deep head by ulnar nerve).

Hypothenar (ulnar)—**O**pponens digiti minimi, **A**bductor digiti minimi, **F**lexor digiti minimi brevis.



Dorsal interossei (ulnar)—abduct the fingers.

Palmar interossei (ulnar)—adduct the fingers.

Lumbricals (1st/2nd, median; 3rd/4th, ulnar)—flex at the MCP joint, extend PIP and DIP joints.





Both groups perform the same functions:

Oppose, **A**bduct, and **F**lex (**OAF**).

DAB = **D**orsals **AB**duct.

PAD = **P**almars **AD**duct.

Distortions of the hand At rest, a balance exists between the extrinsic flexors and extensors of the hand, as well as the intrinsic muscles of the hand—particularly the lumbrical muscles (flexion of MCP, extension of DIP and PIP joints).
 “Clawing”—seen best with **distal** lesions of median or ulnar nerves. Remaining extrinsic flexors of the digits exaggerate the loss of the lumbricals → fingers extend at MCP, flex at DIP and PIP joints.
 Deficits less pronounced in **proximal** lesions; deficits present during voluntary flexion of the digits.



SIGN	“Ulnar claw”	“Pope’s blessing”	“Median claw”	“OK gesture”
PRESENTATION				
CONTEXT	Extending fingers/at rest	Making a fist	Extending fingers/at rest	Making a fist
LOCATION OF LESION	Distal ulnar nerve	Proximal median nerve	Distal median nerve	Proximal ulnar nerve

Note: Atrophy of the thenar eminence (unopposable thumb → “ape hand”) can be seen in median nerve lesions, while atrophy of the hypothenar eminence can be seen in ulnar nerve lesions.




Actions of hip muscles

ACTION	MUSCLES
Abductors	Gluteus medius, gluteus minimus
Adductors	Adductor magnus, adductor longus, adductor brevis
Extensors	Gluteus maximus, semitendinosus, semimembranosus
Flexors	Iliopsoas, rectus femoris, tensor fascia lata, pectineus, sartorius
Internal rotation	Gluteus medius, gluteus minimus, tensor fascia latae
External rotation	Iliopsoas, gluteus maximus, piriformis, obturator

Lower extremity nerves


NERVE	INNERVATION	CAUSE OF INJURY	PRESENTATION/COMMENTS
Iliohypogastric (T12-L1)	Sensory—suprapubic region Motor—transversus abdominis and internal oblique	Abdominal surgery	Burning or tingling pain in surgical incision site radiating to inguinal and suprapubic region
Genitofemoral nerve (L1-L2)	Sensory—scrotum/labia majora, medial thigh Motor—cremaster	Laparoscopic surgery	↓ upper medial thigh and anterior thigh sensation beneath the inguinal ligament (lateral part of the femoral triangle); absent cremasteric reflex
Lateral femoral cutaneous (L2-L3)	Sensory—anterior and lateral thigh	Tight clothing, obesity, pregnancy, pelvic procedures	↓ thigh sensation (anterior and lateral)
Obturator (L2-L4) 	Sensory—medial thigh Motor—obturator externus, adductor longus, adductor brevis, gracilis, pectineus, adductor magnus	Pelvic surgery	↓ thigh sensation (medial) and adduction
Femoral (L2-L4) 	Sensory—anterior thigh, medial leg Motor—quadriceps, iliacus, pectineus, sartorius	Pelvic fracture	↓ leg extension (↓ patellar reflex)
Sciatic (L4-S3)	Motor—semitendinosus, semimembranosus, biceps femoris, adductor magnus	Herniated disc, posterior hip dislocation	Splits into common peroneal and tibial nerves







Lower extremity nerves (*continued*)

NERVE	INNERVATION	CAUSE OF INJURY	PRESENTATION/COMMENTS
Common (fibular) peroneal (L4-S2) 	Superficial peroneal nerve: <ul style="list-style-type: none"> ▪ Sensory—dorsum of foot (except webspace between hallux and 2nd digit) ▪ Motor—peroneus longus and brevis Deep peroneal nerve: <ul style="list-style-type: none"> ▪ Sensory—webspace between hallux and 2nd digit ▪ Motor—tibialis anterior 	Trauma or compression of lateral aspect of leg, fibular neck fracture	PED = P eroneal E verts and D orsiflexes; if injured, foot drop PED Loss of sensation on dorsum of foot Foot drop —inverted and plantarflexed at rest, loss of eversion and dorsiflexion; “steppage gait”
Tibial (L4-S3) 	Sensory—sole of foot Motor—biceps femoris (long head), triceps surae, plantaris, popliteus, flexor muscles of foot	Knee trauma, Baker cyst (proximal lesion); tarsal tunnel syndrome (distal lesion)	TIP = T ibial I nverts and P lantarflexes; if injured, can't stand on TIP toes Inability to curl toes and loss of sensation on sole; in proximal lesions, foot everted at rest with loss of inversion and plantar flexion
Superior gluteal (L4-S1) 	Motor—gluteus medius, gluteus minimus, tensor fascia latae	Iatrogenic injury during intramuscular injection to superomedial gluteal region (prevent by choosing superolateral quadrant, preferably anterolateral region)	Trendelenburg sign/gait—pelvis tilts because weight-bearing leg cannot maintain alignment of pelvis through hip abduction Lesion is contralateral to the side of the hip that drops, ipsilateral to extremity on which the patient stands
Inferior gluteal (L5-S2)	Motor—gluteus maximus	Posterior hip dislocation	Difficulty climbing stairs, rising from seated position; loss of hip extension
Pudendal (S2-S4)	Sensory—perineum Motor—external urethral and anal sphincters	Stretch injury during childbirth, prolonged cycling, horseback riding	↓ sensation in perineum and genital area; can cause fecal and/or urinary incontinence Can be blocked with local anesthetic during childbirth using ischial spine as a landmark for injection

Knee exam


Lateral femoral condyle to anterior tibia: **ACL**.
 Medial femoral condyle to posterior tibia: **PCL**.
LAMP.



TEST	PROCEDURE	
Anterior drawer sign	Bending knee at 90° angle, ↑ anterior gliding of tibia (relative to femur) due to ACL injury Lachman test also tests ACL, but is more sensitive (↑ anterior gliding of tibia [relative to femur] with knee bent at 30° angle)	 ACL tear x
Posterior drawer sign	Bending knee at 90° angle, ↑ posterior gliding of tibia due to PCL injury	 PCL tear x
Abnormal passive abduction	Knee either extended or at ~ 30° angle, lateral (valgus) force → medial space widening of tibia → MCL injury	 Abduction (valgus) force MCL tear x
Abnormal passive adduction	Knee either extended or at ~ 30° angle, medial (varus) force → lateral space widening of tibia → LCL injury	 Adduction (varus) force LCL tear x
McMurray test	During flexion and extension of knee with rotation of tibia/foot (LIME): <ul style="list-style-type: none"> ▪ Pain, “popping” on internal rotation and varus force → Lateral meniscal tear (Internal rotation stresses lateral meniscus) ▪ Pain, “popping” on external rotation and valgus force → Medial meniscal tear (External rotation stresses medial meniscus) 	 Internal rotation and varus force Lateral meniscal tear x  External rotation and valgus force Medial meniscal tear x

Ankle sprains


Anterior **T**alo**F**ibular ligament—most common ankle sprain overall, classified as a low ankle sprain. Due to overinversion/supination of foot.
Anterior inferior tibiofibular ligament—most common high ankle sprain.
Always **T**ears **F**irst.



Signs of lumbosacral radiculopathy

Paresthesia and weakness related to specific lumbosacral spinal nerves. Intervertebral disc (nucleus pulposus) herniates posterolaterally through annulus fibrosus (outer ring) into central canal due to thin posterior longitudinal ligament and thicker anterior longitudinal ligament along midline of vertebral bodies. Nerve affected is usually below the level of herniation.

Disc level herniation	L3-L4	L4-L5	L5-S1
Nerve root affected	L4	L5	S1
Dermatome affected			
Clinical findings	Weakness of knee extension ↓ patellar reflex	Weakness of dorsiflexion Difficulty in heel walking	Weakness of plantar flexion Difficulty in toe walking ↓ Achilles reflex




Neurovascular pairing

Nerves and arteries are frequently named together by the bones/regions with which they are associated. The following are exceptions to this naming convention.

LOCATION	NERVE	ARTERY
Axilla/lateral thorax	Long thoracic	Lateral thoracic
Surgical neck of humerus	Axillary	Posterior circumflex
Midshaft of humerus	Radial	Deep brachial
Distal humerus/cubital fossa	Median	Brachial
Popliteal fossa	Tibial	Popliteal
Posterior to medial malleolus	Tibial	Posterior tibial

Motoneuron action potential to muscle contraction

T-tubules are extensions of plasma membrane in contact with the sarcoplasmic reticulum, allowing for coordinated contraction of striated muscles.




- 1** Action potential opens presynaptic voltage-gated Ca^{2+} channels, inducing acetylcholine (ACh) release.
- 2** Postsynaptic ACh binding leads to muscle cell depolarization at the motor end plate.
- 3** Depolarization travels over the entire muscle cell and deep into the muscle via the T-tubules.
- 4** Membrane depolarization induces conformational changes in the voltage-sensitive dihydropyridine receptor (DHPR) and its mechanically coupled ryanodine receptor (RR) \rightarrow Ca^{2+} release from the sarcoplasmic reticulum into the cytoplasm.
- 5** Tropomyosin is blocking myosin-binding sites on the actin filament. Released Ca^{2+} binds to troponin C (TnC), shifting tropomyosin to expose the myosin-binding sites.
- 6** The myosin head binds strongly to actin, forming a crossbridge. P_i is then released, initiating the power stroke.
- 7** During the power stroke, force is produced as myosin pulls on the thin filament **A**. Muscle shortening occurs, with shortening of **H** and **I** bands and between **Z** lines (**HIZ** shrinkage). The **A** band remains the same length (**A** band is **A**lways the same length). ADP is released at the end of the power stroke.
- 8** Binding of a new ATP molecule causes detachment of myosin head from actin filament. Ca^{2+} is re-sequestered.
- 9** ATP hydrolysis into ADP and P_i results in myosin head returning to high-energy position (cocked). The myosin head can bind to a new site on actin to form a crossbridge if Ca^{2+} remains available.

Types of muscle fibers


	Type I	Type II
CONTRACTION VELOCITY	Slow	Fast
FIBER COLOR	Red	White
PREDOMINANT METABOLISM	Oxidative phosphorylation → sustained contraction	Anaerobic glycolysis
MITOCHONDRIA, MYOGLOBIN	↑	↓
TYPE OF TRAINING	Endurance training	Weight/resistance training, sprinting
NOTES	Think “I slow red ox”	

Vascular smooth muscle contraction and relaxation



Muscle proprioceptors Specialized sensory receptors that relay information about muscle dynamics.

	Muscle spindle	Golgi tendon organ
PATHWAY	<p>① ↑ length and speed of stretch → ② via dorsal root ganglion (DRG) → ③ activation of inhibitory interneuron and α motor neuron → ④ simultaneous inhibition of antagonist muscle (prevents overstretching) and activation of agonist muscle (contraction).</p>	<p>① ↑ tension → ② via DRG → ③ activation of inhibitory interneuron → ④ inhibition of agonist muscle (reduced tension within muscle and tendon)</p>
LOCATION	Body of muscle/type Ia and II sensory axons	Tendons/type Ib sensory axons
ACTIVATION BY	↑ muscle stretch	↑ muscle force



Bone formation

Endochondral ossification

Bones of axial skeleton, appendicular skeleton, and base of skull. Cartilaginous model of bone is first made by chondrocytes. Osteoclasts and osteoblasts later replace with woven bone and then remodel to lamellar bone. In adults, woven bone occurs after fractures and in Paget disease. Defective in achondroplasia.

Membranous ossification

Bones of calvarium, facial bones, and clavicle. Woven bone formed directly without cartilage. Later remodeled to lamellar bone.

Cell biology of bone

Osteoblast	B uilds b one by secreting collagen and catalyzing mineralization in alkaline environment via ALP. Differentiates from mesenchymal stem cells in periosteum. Osteoblastic activity measured by bone ALP, osteocalcin, propeptides of type I procollagen.
Osteoclast	Dissolves (“ c rushes”) bone by secreting H ⁺ and collagenases. Differentiates from a fusion of monocyte/macrophage lineage precursors. RANK receptors on osteoclasts are stimulated by RANKL (RANK ligand, expressed on osteoblasts). OPG (osteoprotegerin, a RANKL decoy receptor) binds RANKL to prevent RANK-RANKL interaction → ↓ osteoclast activity.
Parathyroid hormone	At low, intermittent levels, exerts anabolic effects (building bone) on osteoblasts and osteoclasts (indirect). Chronically ↑ PTH levels (1° hyperparathyroidism) cause catabolic effects (osteitis fibrosa cystica).
Estrogen	Inhibits apoptosis in bone-forming osteoblasts and induces apoptosis in bone-resorbing osteoclasts. Causes closure of epiphyseal plate during puberty. Estrogen deficiency (surgical or postmenopausal) → ↑ cycles of remodeling and bone resorption → ↑ risk of osteoporosis.

▶ MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—PATHOLOGY**Overuse injuries of the elbow**

Medial epicondylitis (golfer’s elbow)	Repetitive flexion (forehand shots) or idiopathic → pain near medial epicondyle.
Lateral epicondylitis (tennis elbow)	Repetitive e xtension (backhand shots) or idiopathic → pain near lateral epicondyle.

Wrist and hand injuries

Metacarpal neck fracture	Also called boxer’s fracture. Common fracture caused by direct blow with a closed fist (eg, from punching a wall). Most commonly seen in 4th and 5th metacarpals A .
---------------------------------	---

**Carpal tunnel syndrome**

Entrapment of median nerve in carpal tunnel (between transverse carpal ligament and carpal bones) → nerve compression → paresthesia, pain, and numbness in distribution of median nerve. Thenar eminence atrophies **B** but sensation spared, because palmar cutaneous branch enters hand external to carpal tunnel.

Suggested by ⊕ Tinel sign (percussion of wrist causes tingling) and Phalen maneuver (90° flexion of wrist causes tingling). Associated with pregnancy (due to edema), rheumatoid arthritis, hypothyroidism, diabetes, acromegaly, dialysis-related amyloidosis; may be associated with repetitive use.

Guyon canal syndrome

Compression of ulnar nerve at wrist. Classically seen in cyclists due to pressure from handlebars.


Clavicle fractures

Common in children and as birth trauma. Usually caused by a fall on outstretched hand or by direct trauma to shoulder. Weakest point at the junction of middle and lateral thirds; fractures at the middle third segment are most common. Presents as shoulder drop, shortened clavicle (lateral fragment is depressed due to arm weight and medially rotated by arm adductors [eg, pectoralis major]).

Common hip and knee conditions

“Unhappy triad”

Common injury in contact sports due to lateral force applied to a planted foot. Consists of damage to the ACL, MCL, and medial meniscus (attached to MCL). However, lateral meniscus involvement is more common than medial meniscus involvement in conjunction with ACL and MCL injury. Presents with acute pain and signs of joint instability.




Prepatellar bursitis

Inflammation of the prepatellar bursa in front of the kneecap (red arrow in B). Can be caused by repeated trauma or pressure from excessive kneeling (also called “housemaid’s knee”).

Baker cyst


Popliteal fluid collection (red arrow in C) in gastrocnemius-semimembranosus bursa commonly communicating with synovial space and related to chronic joint disease (eg, osteoarthritis, rheumatoid arthritis).



Common musculoskeletal conditions

De Quervain tenosynovitis	Noninflammatory thickening of abductor pollicis longus and extensor pollicis brevis tendons → pain or tenderness at radial styloid. ⊕ Finkelstein test (pain at radial styloid with active or passive stretch of thumb tendons). ↑ risk in new mothers, golfers, racquet sport players, “thumb” texters.
Ganglion cyst	Fluid-filled swelling overlying joint or tendon sheath, most commonly at dorsal side of wrist. Arises from herniation of dense connective tissue.
Iliotibial band syndrome	Overuse injury of lateral knee that occurs primarily in runners. Pain develops 2° to friction of iliotibial band against lateral femoral epicondyle.
Limb compartment syndrome	↑ pressure within fascial compartment of a limb → venous outflow obstruction and arteriolar collapse → anoxia and necrosis. Causes include significant long bone fractures, reperfusion injury, animal venoms. Presents with severe pain and tense, swollen compartments with passive stretch of muscles in the affected compartment. Motor deficits are late sign of irreversible muscle and nerve damage.
Medial tibial stress syndrome	Also called shin splints. Common cause of shin pain and diffuse tenderness in runners and military recruits. Caused by bone resorption that outpaces bone formation in tibial cortex.
Plantar fasciitis	Inflammation of plantar aponeurosis characterized by heel pain (worse with first steps in the morning or after period of inactivity) and tenderness.



Childhood musculoskeletal conditions

Developmental dysplasia of the hip	Abnormal acetabulum development in newborns. Major risk factor includes breech presentation. Results in hip instability/dislocation. Commonly tested with Ortolani and Barlow maneuvers (manipulation of newborn hip reveals a “clunk”). Confirmed via ultrasound (x-ray not used until ~4–6 months because cartilage is not ossified).
Legg-Calvé-Perthes disease	Idiopathic avascular necrosis of femoral head. Commonly presents between 5–7 years with insidious onset of hip pain that may cause child to limp. More common in males (4:1 ratio). Initial x-ray often normal.
Osgood-Schlatter disease	Also called traction apophysitis. Overuse injury caused by repetitive strain and chronic avulsion of the secondary ossification center of proximal tibial tubercle. Occurs in adolescents after growth spurt. Common in running and jumping athletes. Presents with progressive anterior knee pain.
	
Patellofemoral syndrome	Overuse injury that commonly presents in young, female athletes as anterior knee pain. Exacerbated by prolonged sitting or weight-bearing on a flexed knee. Treatment: NSAIDs, thigh muscle strengthening.
Radial head subluxation	Also called nursemaid’s elbow. Common elbow injury in children < 5 years. Caused by a sudden pull on the arm → immature annular ligament slips over head of radius. Injured arm held in extended/slightly flexed and pronated position.
Slipped capital femoral epiphysis	Classically presents in an obese young adolescent with hip/knee pain and altered gait. Increased axial force on femoral head → epiphysis displaces relative to femoral neck (like a scoop of ice cream slipping off a cone). Diagnosed via x-ray.

Common pediatric fractures

Greenstick fracture Incomplete fracture extending partway through width of bone **A** following bending stress; bone fails on tension side; compression side intact (compare to torus fracture). Bone is bent like a **green twig**.

Torus (buckle) fracture Axial force applied to immature bone → cortex buckles on compression (concave) side and fractures **B**. Tension (convex) side remains solid (intact).



Achondroplasia

Failure of longitudinal bone growth (endochondral ossification) → short limbs. Membranous ossification is not affected → large head relative to limbs. Constitutive activation of fibroblast growth factor receptor (FGFR3) actually inhibits chondrocyte proliferation. > 85% of mutations occur sporadically; autosomal dominant with full penetrance (homozygosity is lethal). Associated with ↑ paternal age. Most common cause of short-limbed dwarfism.

Osteoporosis



Trabecular (spongy) and cortical bone lose mass despite normal bone mineralization and lab values (serum Ca^{2+} and PO_4^{3-}).



Most commonly due to ↑ bone resorption related to ↓ estrogen levels and old age. Can be 2° to drugs (eg, steroids, alcohol, anticonvulsants, anticoagulants, thyroid replacement therapy) or other conditions (eg, hyperparathyroidism, hyperthyroidism, multiple myeloma, malabsorption syndromes, anorexia).


Diagnosed by bone mineral density measurement by DEXA (dual-energy X-ray absorptiometry) at the lumbar spine, total hip, and femoral neck, with a T-score of ≤ -2.5 or by a fragility fracture (eg, fall from standing height, minimal trauma) at hip or vertebra. One time screening recommended in women ≥ 65 years old.

Prophylaxis: regular weight-bearing exercise and adequate Ca^{2+} and vitamin D intake throughout adulthood.

Treatment: bisphosphonates, teriparatide, SERMs, rarely calcitonin; denosumab (monoclonal antibody against RANKL).

Can lead to **vertebral compression fractures** **A**—acute back pain, loss of height, kyphosis. Also can present with fractures of femoral neck, distal radius (Colles fracture).



Osteopetrosis

Failure of normal bone resorption due to defective osteoclasts → thickened, dense bones that are prone to fracture. Mutations (eg, carbonic anhydrase II) impair ability of osteoclast to generate acidic environment necessary for bone resorption. Overgrowth of cortical bone fills marrow space → pancytopenia, extramedullary hematopoiesis. Can result in cranial nerve impingement and palsies due to narrowed foramina.

X-rays show diffuse symmetric sclerosis (bone-in-bone, “stone bone” **A**). Bone marrow transplant is potentially curative as osteoclasts are derived from monocytes.



Osteomalacia/rickets

Defective mineralization of osteoid (osteomalacia) or cartilaginous growth plates (rickets, only in children). Most commonly due to vitamin D deficiency.

X-rays show osteopenia and “Looser zones” (pseudofractures) in osteomalacia, epiphyseal widening and metaphyseal cupping/fraying in rickets. Children with rickets have pathologic bow legs (genu varum **A**), bead-like costochondral junctions (rachitic rosary **B**), craniotabes (soft skull).

↓ vitamin D → ↓ serum Ca^{2+} → ↑ PTH secretion → ↓ serum PO_4^{3-} .

Hyperactivity of osteoblasts → ↑ ALP.

**Osteitis deformans**


Also called Paget disease of bone. Common, localized disorder of bone remodeling caused by ↑ osteoclastic activity followed by ↑ osteoblastic activity that forms poor-quality bone. Serum Ca^{2+} , phosphorus, and PTH levels are normal. ↑ ALP. Mosaic pattern of woven and lamellar bone (osteocytes within lacunae in chaotic juxtapositions); long bone chalk-stick fractures. ↑ blood flow from ↑ arteriovenous shunts may cause high-output heart failure. ↑ risk of osteosarcoma.

Hat size can be increased due to skull thickening **A**; hearing loss is common due to auditory foramen narrowing.


Stages of Paget disease:

- Lytic—osteoclasts
- Mixed—osteoclasts + osteoblasts
- Sclerotic—osteoblasts
- Quiescent—minimal osteoclast/osteoblast activity

Treatment: bisphosphonates.

Avascular necrosis of bone

Infarction of bone and marrow, usually very painful. Most common site is femoral head (watershed zone) **A** (due to insufficiency of medial circumflex femoral artery). Causes include **C**orticosteroids, **A**lcoholism, **S**ickle cell disease, **T**rauma, **S**LE, “the **B**ends” (caisson/decompression disease), **L**Egg-**C**alvé-**P**erthes disease (idiopathic), **G**aucher disease, **S**lipped capital femoral epiphysis—**C**ASTS **B**end **L**EGS.



Lab values in bone disorders

DISORDER	SERUM Ca ²⁺	PO ₄ ³⁻	ALP	PTH	COMMENTS
Osteoporosis	—	—	—	—	↓ bone mass
Osteopetrosis	—/↓	—	—	—	Dense, brittle bones. Ca ²⁺ ↓ in severe, malignant disease
Paget disease of bone	—	—	↑	—	Abnormal “mosaic” bone architecture
Osteitis fibrosa cystica Primary hyperparathyroidism	↑	↓	↑	↑	“Brown tumors” due to fibrous replacement of bone, subperiosteal thinning Idiopathic or parathyroid hyperplasia, adenoma, carcinoma
Secondary hyperparathyroidism	↓	↑	↑	↑	Often as compensation for CKD (↓ PO ₄ ³⁻ excretion and production of activated vitamin D)
Osteomalacia/rickets	↓	↓	↑	↑	Soft bones; vitamin D deficiency also causes 2° hyperparathyroidism
Hypervitaminosis D	↑	↑	—	↓	Caused by oversupplementation or granulomatous disease (eg, sarcoidosis)


↑ ↓ = 1° change.

Primary bone tumors Metastatic disease is more common than 1° bone tumors. Benign **bone** tumors that start with **O** are more common in **boys**.

TUMOR TYPE	EPIDEMIOLOGY	LOCATION	CHARACTERISTICS
Benign tumors			
Osteochondroma	Most common benign bone tumor Males < 25 years old	Metaphysis of long bones	Lateral bony projection of growth plate (continuous with marrow space) covered by cartilaginous cap A Rarely transforms to chondrosarcoma
Osteoma	Middle age	Surface of facial bones	Associated with Gardner syndrome
Osteoid osteoma	Adults < 25 years old Males > females	Cortex of long bones	Presents as bone pain (worse at night) that is relieved by NSAIDs Bony mass (< 2 cm) with radiolucent osteoid core B
Osteoblastoma	Males > females	Vertebrae	Similar histology to osteoid osteoma Larger size (> 2 cm), pain unresponsive to NSAIDs
Chondroma		Medulla of small bones of hand and feet	Benign tumor of cartilage
Giant cell tumor	20–40 years old	Epiphysis of long bones (often in knee region)	Locally aggressive benign tumor Neoplastic mononuclear cells that express RANKL and reactive multinucleated giant (osteoclast-like) cells. “Osteoclastoma” “Soap bubble” appearance on x-ray C

Primary bone tumors (continued)


TUMOR TYPE	EPIDEMIOLOGY	LOCATION	CHARACTERISTICS
Malignant tumors			
Osteosarcoma (osteogenic sarcoma)	Accounts for 20% of 1° bone cancers. Peak incidence of 1° tumor in males < 20 years. Less common in elderly; usually 2° to predisposing factors, such as Paget disease of bone, bone infarcts, radiation, familial retinoblastoma, Li-Fraumeni syndrome.	Metaphysis of long bones (often in knee region).	Pleomorphic osteoid-producing cells (malignant osteoblasts). Presents as painful enlarging mass or pathologic fractures. Codman triangle D (from elevation of periosteum) or sunburst pattern on x-ray E (think of an osteocod (bone fish) swimming in the sun). Aggressive. 1° usually responsive to treatment (surgery, chemotherapy), poor prognosis for 2°.
Chondrosarcoma		Medulla of pelvis, proximal femur and humerus.	Tumor of malignant chondrocytes.
Ewing sarcoma	Most common in Caucasians. Generally boys < 15 years old.	Diaphysis of long bones (especially femur), pelvic flat bones.	Anaplastic small blue cells of neuroectodermal origin (resemble lymphocytes) F . Differentiate from conditions with similar morphology (eg, lymphoma, chronic osteomyelitis) by testing for t(11;22) (fusion protein EWS-FLI1). “Onion skin” periosteal reaction in bone. Aggressive with early metastases, but responsive to chemotherapy. 11 + 22 = 33 (Patrick Ewing ’s jersey number).



Osteoarthritis vs rheumatoid arthritis

	Osteoarthritis	Rheumatoid arthritis
PATHOGENESIS	Mechanical—wear and tear destroys articular cartilage (degenerative joint disorder) → inflammation with inadequate repair. Chondrocytes mediate degradation and inadequate repair.	Autoimmune—inflammation A induces formation of pannus (proliferative granulation tissue), which erodes articular cartilage and bone.
PREDISPOSING FACTORS	Age, female, obesity, joint trauma.	Female, HLA-DR4 (4-walled “rheum”), smoking. ⊕ rheumatoid factor (IgM antibody that targets IgG Fc region; in 80%), anti-cyclic citrullinated peptide antibody (more specific).
PRESENTATION	Pain in weight-bearing joints after use (eg, at the end of the day), improving with rest. Asymmetric joint involvement. Knee cartilage loss begins medially (“bowlegged”). No systemic symptoms.	Pain, swelling, and morning stiffness lasting > 1 hour, improving with use. Symmetric joint involvement. Systemic symptoms (fever, fatigue, weight loss). Extraarticular manifestations common.*
JOINT FINDINGS	Osteophytes (bone spurs), joint space narrowing, subchondral sclerosis and cysts. Synovial fluid noninflammatory (WBC < 2000/mm ³). Development of Heberden nodes B (at DIP) and Bouchard nodes C (at PIP), and 1st CMC; not MCP.	Erosions, juxta-articular osteopenia, soft tissue swelling, subchondral cysts, joint space narrowing. Deformities: cervical subluxation, ulnar finger deviation, swan neck D , boutonniere E . Involves MCP, PIP, wrist; not DIP or 1st CMC.
TREATMENT	Activity modification, acetaminophen, NSAIDs, intra-articular glucocorticoids.	NSAIDs, glucocorticoids, disease-modifying agents (eg, methotrexate, sulfasalazine), biologic agents (eg, TNF- α inhibitors).

*Extraarticular manifestations include rheumatoid nodules (fibrinoid necrosis with palisading histiocytes) in subcutaneous tissue and lung (+ pneumoconiosis → Caplan syndrome), interstitial lung disease, pleuritis, pericarditis, anemia of chronic disease, neutropenia + splenomegaly (Felty syndrome), AA amyloidosis, Sjögren syndrome, scleritis, carpal tunnel syndrome.



Gout**FINDINGS**

Acute inflammatory monoarthritis caused by precipitation of monosodium urate crystals in joints **A**. Risk factors: male sex, hypertension, obesity, diabetes, dyslipidemia, alcohol use. Strongest risk factor is hyperuricemia, which can be caused by:

- Underexcretion of uric acid (90% of patients)—largely idiopathic, potentiated by renal failure; can be exacerbated by certain medications (eg, thiazide diuretics).
- Overproduction of uric acid (10% of patients)—Lesch-Nyhan syndrome, PRPP excess, ↑ cell turnover (eg, tumor lysis syndrome), von Gierke disease.

Crystals are needle shaped and ⊖ birefringent under polarized light (yellow under parallel light, blue under perpendicular light **B**). Serum uric acid levels may be normal during an acute attack.



SYMPTOMS

Asymmetric joint distribution. Joint is swollen, red, and painful. Classic manifestation is painful MTP joint of big toe (podagra). Tophus formation **C** (often on external ear, olecranon bursa, or Achilles tendon). Acute attack tends to occur after a large meal with foods rich in purines (eg, red meat, seafood), trauma, surgery, dehydration, diuresis, or alcohol consumption (alcohol metabolites compete for same excretion sites in kidney as uric acid → ↓ uric acid secretion and subsequent buildup in blood).

TREATMENT

Acute: NSAIDs (eg, indomethacin), glucocorticoids, colchicine.

Chronic (preventive): xanthine oxidase inhibitors (eg, allopurinol, febuxostat).

**Calcium pyrophosphate deposition disease**

Previously called pseudogout. Deposition of calcium pyrophosphate crystals within the joint space. Occurs in patients > 50 years old; both sexes affected equally. Usually idiopathic, sometimes associated with hemochromatosis, hyperparathyroidism, joint trauma.

Pain and swelling with acute inflammation (pseudogout) and/or chronic degeneration (pseudo-osteoarthritis). Most commonly affected joint is the knee.

Chondrocalcinosis (cartilage calcification) on x-ray.

Crystals are rhomboid and weakly ⊕ birefringent under polarized light (blue when parallel to light) **A**.


Acute treatment: NSAIDs, colchicine, glucocorticoids.

Prophylaxis: colchicine.

The **blue P's**—**blue** (when **Parallel**), **Positive** birefringence, calcium **P**rophosphate, **P**seudogout

Systemic juvenile idiopathic arthritis

Systemic arthritis seen in < 16 year olds. Usually presents with daily spiking fevers, salmon-pink macular rash, arthritis (commonly 2+ joints). Associated with anterior uveitis. Frequently presents with leukocytosis, thrombocytosis, anemia, ↑ ESR, ↑ CRP. Treatment: NSAIDs, steroids, methotrexate, TNF inhibitors.

Sjögren syndrome

Autoimmune disorder characterized by destruction of exocrine glands (especially lacrimal and salivary) by lymphocytic infiltrates **A**. Predominantly affects women 40–60 years old.

Findings:

- Inflammatory joint pain
- Keratoconjunctivitis sicca (↓ tear production and subsequent corneal damage)
- Xerostomia (↓ saliva production) → mucosal atrophy, fissuring of the tongue **B**
- Presence of antinuclear antibodies, rheumatoid factor (can be positive in the absence of rheumatoid arthritis), antiribonucleoprotein antibodies: SS-A (anti-Ro) and/or SS-B (anti-La)
- Bilateral parotid enlargement

Anti-SSA and anti-SSB may also be seen in SLE.

A common 1° disorder or a 2° syndrome associated with other autoimmune disorders (eg, rheumatoid arthritis, SLE, systemic sclerosis).

Complications: dental caries; mucosa-associated lymphoid tissue (MALT) lymphoma (may present as parotid enlargement).

Focal lymphocytic sialadenitis on labial salivary gland biopsy can confirm diagnosis.

Septic arthritis

S aureus, *Streptococcus*, and *Neisseria gonorrhoeae* are common causes. Affected joint is swollen **A**, red, and painful. Synovial fluid purulent (WBC > 50,000/mm³).

Gonococcal arthritis—STI that presents as either purulent arthritis (eg, knee) or triad of polyarthralgia, tenosynovitis (eg, hand), dermatitis (eg, pustules).

Seronegative spondyloarthritis

Arthritis without rheumatoid factor (no anti-IgG antibody). Strong association with HLA-B27 (MHC class I serotype). Subtypes (**PAIR**) share variable occurrence of inflammatory back pain (associated with morning stiffness, improves with exercise), peripheral arthritis, enthesitis (inflamed insertion sites of tendons, eg, Achilles), dactylitis (“sausage fingers”), uveitis.

Psoriatic arthritis

Associated with skin psoriasis and nail lesions. Asymmetric and patchy involvement **A**. Dactylitis and “pencil-in-cup” deformity of DIP on x-ray **B**.

Seen in fewer than 1/3 of patients with psoriasis.

Ankylosing spondylitis

Symmetric involvement of spine and sacroiliac joints → ankylosis (joint fusion), uveitis, aortic regurgitation.

Bamboo spine (vertebral fusion) **C**.

Costovertebral and costosternal ankylosis may cause restrictive lung disease. Monitor degree of reduced chest wall expansion to assess disease severity.

More common in males.

Inflammatory bowel disease

Crohn disease and ulcerative colitis are often associated with spondyloarthritis.

Reactive arthritis

Formerly called Reiter syndrome. Classic triad:


- **Conjunctivitis**
- **Urethritis**
- **Arthritis**

“Can’t see, can’t pee, can’t bend my knee.”
Shigella, *Yersinia*, *Chlamydia*, *Campylobacter*,
Salmonella (**ShY ChiCS**).



Systemic lupus erythematosus

Systemic, remitting, and relapsing autoimmune disease. Organ damage primarily due to a type III hypersensitivity reaction and, to a lesser degree, a type II hypersensitivity reaction. Associated with deficiency of early complement proteins (eg, C1q, C4, C2) → ↓ clearance of immune complexes. Classic presentation: rash, joint pain, and fever in a female of reproductive age (especially of African-American or Hispanic descent).



Libman-Sacks Endocarditis—nonbacterial, verrucous thrombi usually on mitral or aortic valve and can be present on either surface of the valve (but usually on undersurface). **LSE** in **SLE**.

Lupus nephritis (glomerular deposition of DNA-anti-DNA immune complexes) can be nephritic or nephrotic (causing hematuria or proteinuria). Most common and severe type is diffuse proliferative.

Common causes of death in SLE: **Renal disease** (most common), **Infections**, **Cardiovascular disease** (accelerated CAD).

In an anti-SSA ⊕ pregnant woman, ↑ risk of newborn developing **neonatal lupus** → congenital heart block, periorbital/diffuse rash, transaminitis, and cytopenias at birth.

RASH OR PAIN:

- Rash** (malar **A** or discoid **B**)
- Arthritis** (nonerosive)
- Serositis** (eg, pleuritis, pericarditis)
- Hematologic disorders** (eg, cytopenias)
- Oral/nasopharyngeal ulcers** (usually painless)
- Renal disease**
- Photosensitivity**
- Antinuclear antibodies**
- Immunologic disorder** (anti-dsDNA, anti-Sm, antiphospholipid)
- Neurologic disorders** (eg, seizures, psychosis)

Lupus patients die with **Redness In** their **Cheeks**.

Mixed connective tissue disease

Features of SLE, systemic sclerosis, and/or polymyositis. Associated with anti-U1 RNP antibodies (speckled ANA).

Antiphospholipid syndrome

1° or 2° autoimmune disorder (most commonly in SLE).

Diagnosed based on clinical criteria including history of thrombosis (arterial or venous) or spontaneous abortion along with laboratory findings of lupus anticoagulant, anticardiolipin, anti-β₂ glycoprotein I antibodies.

Treatment: systemic anticoagulation.

Anticardiolipin antibodies can cause false-positive VDRL/RPR.

Lupus anticoagulant can cause prolonged PTT that is not corrected by the addition of normal platelet-free plasma.

Polymyalgia rheumatica**SYMPTOMS**

Pain and stiffness in proximal muscles (eg, shoulders, hips), often with fever, malaise, weight loss. Does not cause muscular weakness. More common in women > 50 years old; associated with giant cell (temporal) arteritis.

FINDINGS

↑ ESR, ↑ CRP, normal CK.

TREATMENT

Rapid response to low-dose corticosteroids.

Fibromyalgia

Most common in women 20–50 years old. Chronic, widespread musculoskeletal pain associated with “tender points,” stiffness, paresthesias, poor sleep, fatigue, cognitive disturbance (“fibro fog”). Treatment: regular exercise, antidepressants (TCAs, SNRIs), neuropathic pain agents (eg, gabapentin).

**Polymyositis/
dermatomyositis**

Nonspecific: ⊕ ANA, ↑ CK. Specific: ⊕ anti-Jo-1 (histidyl-tRNA synthetase), ⊕ anti-SRP (signal recognition particle), ⊕ anti-Mi-2 (helicase).

Polymyositis

Progressive symmetric proximal muscle weakness, characterized by endomysial inflammation with CD8+ T cells. Most often involves shoulders.


Dermatomyositis

Clinically similar to polymyositis, but also involves Gottron papules **A**, photodistributed facial erythema (eg, heliotrope [violaceous] edema of the eyelids **B**), “shawl and face” rash **C**, darkening and thickening of fingertips and sides resulting in irregular, “dirty”-appearing marks. ↑ risk of occult malignancy. Perimysial inflammation and atrophy with CD4+ T cells.



Neuromuscular junction diseases

	Myasthenia gravis	Lambert-Eaton myasthenic syndrome
FREQUENCY	Most common NMJ disorder	Uncommon
PATHOPHYSIOLOGY	Autoantibodies to postsynaptic ACh receptor	Autoantibodies to presynaptic Ca^{2+} channel → ↓ ACh release
CLINICAL	Fatigable muscle weakness—ptosis; diplopia; proximal weakness; respiratory muscle involvement → dyspnea; bulbar muscle involvement → dysphagia, difficulty chewing Spared reflexes Worsens with muscle use	Proximal muscle weakness, autonomic symptoms (dry mouth, constipation, impotence) Hyporeflexia Improves with muscle use
ASSOCIATED WITH	Thymoma, thymic hyperplasia	Small cell lung cancer
ACHE INHIBITOR ADMINISTRATION	Reverses symptoms (pyridostigmine for treatment)	Minimal effect



Raynaud phenomenon


↓ blood flow to skin due to arteriolar (small vessel) vasospasm in response to cold or stress: color change from white (ischemia) to blue (hypoxia) to red (reperfusion). Most often in the fingers **A** and toes. Called **Raynaud disease** when 1° (idiopathic), **Raynaud syndrome** when 2° to a disease process such as mixed connective tissue disease, SLE, or CREST syndrome (limited form of systemic sclerosis). Digital ulceration (critical ischemia) seen in 2° Raynaud syndrome. Treat with calcium²⁺ channel blockers.



Scleroderma

Systemic sclerosis. Triad of autoimmunity, noninflammatory vasculopathy, and collagen deposition with fibrosis. Commonly sclerosis of skin, manifesting as puffy, taut skin **A** without wrinkles, fingertip pitting **B**. Can involve other systems, eg, renal (scleroderma renal crisis; treat with ACE inhibitors), pulmonary (interstitial fibrosis, pulmonary HTN), GI (esophageal dysmotility and reflux), cardiovascular. 75% female. 2 major types:


- **Diffuse scleroderma**—widespread skin involvement, rapid progression, early visceral involvement. Associated with anti-Scl-70 antibody (anti-DNA topoisomerase-I antibody) and anti-RNA polymerase III.
- **Limited scleroderma**—limited skin involvement confined to fingers and face. Also with **CREST** syndrome: **C**alcinosis cutis **C**, anti-**C**entromere antibody, **R**aynaud phenomenon, **E**sophageal dysmotility, **S**clerodactyly, and **T**elangiectasia. More benign clinical course.




▶ MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—DERMATOLOGY

Skin layers

Skin has 3 layers: epidermis, dermis, subcutaneous fat (hypodermis, subcutis).
Epidermal layers: **C**ome, **L**et's **G**et **S**un **B**urned.



Epithelial cell junctions



Tight junctions (zonula occludens) **A**—prevents paracellular movement of solutes; composed of claudins and occludins.

Adherens junction (belt desmosome, zonula adherens) **B**—forms “belt” connecting actin cytoskeletons of adjacent cells with **CAD**herins (Ca^{2+} -dependent **ad**hesion proteins). Loss of E-cadherin promotes metastasis.

Desmosome (spot desmosome, macula adherens) **C**—structural support via intermediate filament interactions. Autoantibodies to desmoglein 1 and/or 3 → pemphigus vulgaris.

Gap junction **D**—channel proteins called connexons permit electrical and chemical communication between cells.

Hemidesmosome **E**—connects keratin in basal cells to underlying basement membrane. Autoantibodies → **bullo**us pemphigoid. (Hemidesmosomes are down “**bullo**w.”)

Integrins—membrane proteins that maintain **integrity** of basolateral membrane by binding to collagen, laminin, and fibronectin in basement membrane.

Dermatologic macroscopic terms

LESION	CHARACTERISTICS	EXAMPLES
Macule	Flat lesion with well-circumscribed change in skin color < 1 cm	Freckle (ephelide), labial macule A
Patch	Macule > 1 cm	Large birthmark (congenital nevus) B
Papule	Elevated solid skin lesion < 1 cm	Mole (nevus) C , acne
Plaque	Papule > 1 cm	Psoriasis D
Vesicle	Small fluid-containing blister < 1 cm	Chickenpox (varicella), shingles (zoster) E
Bulla	Large fluid-containing blister > 1 cm	Bullous pemphigoid F
Pustule	Vesicle containing pus	Pustular psoriasis G
Wheal	Transient smooth papule or plaque	Hives (urticaria) H
Scale	Flaking off of stratum corneum	Eczema, psoriasis, SCC I
Crust	Dry exudate	Impetigo J

**Dermatologic microscopic terms**

LESION	CHARACTERISTICS	EXAMPLES
Hyperkeratosis	↑ thickness of stratum corneum	Psoriasis, calluses
Parakeratosis	Retention of nuclei in stratum corneum	Psoriasis, actinic keratosis
Hypergranulosis	↑ thickness of stratum granulosum	Lichen planus
Spongiosis	Epidermal accumulation of edematous fluid in intercellular spaces	Eczematous dermatitis
Acantholysis	Separation of epidermal cells	Pemphigus vulgaris
Acanthosis	Epidermal hyperplasia (↑ spinosum)	Acanthosis nigricans, psoriasis

Pigmented skin disorders

Albinism	Normal melanocyte number with ↓ melanin production A due to ↓ tyrosinase activity or defective tyrosine transport. ↑ risk of skin cancer.
Melasma (chloasma)	Acquired hyperpigmentation associated with pregnancy (“mask of pregnancy” B) or OCP use. More common in women with darker complexions.
Vitiligo	Irregular patches of complete depigmentation C . Caused by destruction of melanocytes (believed to be autoimmune). Associated with other autoimmune disorders.

**Seborrheic dermatitis**

Erythematous, well-demarcated plaques **A** with greasy yellow scales in areas rich in sebaceous glands, such as scalp, face, and periorcular region. Common in both infants (cradle cap) and adults, associated with Parkinson disease. Sebaceous glands are not inflamed, but play a role in disease development. Possibly associated with *Malassezia* spp. Treatment: topical antifungals and corticosteroids.




Common skin disorders

Acne	Multifactorial etiology—↑ sebum/androgen production, abnormal keratinocyte desquamation, <i>Cutibacterium acnes</i> colonization of the pilosebaceous unit (comedones), and inflammation (papules/pustules A , nodules, cysts). Treatment: retinoids, benzoyl peroxide, and antibiotics.
Atopic dermatitis (eczema)	Type I hypersensitivity reaction. Pruritic eruption, commonly on skin flexures. Associated with other atopic diseases (asthma, allergic rhinitis, food allergies); ↑ serum IgE. Mutations in filaggrin gene predispose (via skin barrier dysfunction). Often appears on face in infancy B and then in antecubital fossa C in children and adults.
Allergic contact dermatitis	Type IV hypersensitivity reaction secondary to contact allergen (eg, nickel D , poison ivy, neomycin E).
Melanocytic nevus	Common mole. Benign, but melanoma can arise in congenital or atypical moles. Intradermal nevi are papular F . Junctional nevi are flat macules G .
Pseudofolliculitis barbae	Foreign body inflammatory facial skin disorder characterized by firm, hyperpigmented papules and pustules that are painful and pruritic. Located on cheeks, jawline, and neck. Commonly occurs as a result of shaving (“razor bumps”), primarily affects African-American males.
Psoriasis	Papules and plaques with silvery scaling H , especially on knees and elbows. Acanthosis with parakeratotic scaling (nuclei still in stratum corneum), Munro microabscesses. ↑ stratum spinosum, ↓ stratum granulosum. Auspitz sign (I)—pinpoint bleeding spots from exposure of dermal papillae when scales are scraped off. Associated with nail pitting and psoriatic arthritis.
Rosacea	Inflammatory facial skin disorder characterized by erythematous papules and pustules J , but no comedones. May be associated with facial flushing in response to external stimuli (eg, alcohol, heat). Phymatous rosacea can cause rhinophyma (bulbous deformation of nose).
Seborrheic keratosis	Flat, greasy, pigmented squamous epithelial proliferation of immature keratinocytes with keratin-filled cysts (horn cysts) K . Looks “stuck on.” Lesions occur on head, trunk, and extremities. Common benign neoplasm of older persons. Leser-Trélat sign L —rapid onset of multiple seborrheic keratoses, indicates possible malignancy (eg, GI adenocarcinoma).
Verrucae	Warts; caused by low-risk HPV strains. Soft, tan-colored, cauliflower-like papules M . Epidermal hyperplasia, hyperkeratosis, koilocytosis. Condyloma acuminatum on anus or genitals N .
Urticaria	Hives. Pruritic wheals that form after mast cell degranulation O . Characterized by superficial dermal edema and lymphatic channel dilation.



Vascular tumors of skin

Angiosarcoma	Rare blood vessel malignancy typically occurring in the head, neck, and breast areas. Usually in elderly, on sun-exposed areas. Associated with radiation therapy and chronic postmastectomy lymphedema. Hepatic angiosarcoma associated with vinyl chloride and arsenic exposures. Very aggressive and difficult to resect due to delay in diagnosis.
Bacillary angiomatosis	Benign capillary skin papules A found in AIDS patients. Caused by <i>Bartonella</i> infections. Frequently mistaken for Kaposi sarcoma, but has neutrophilic infiltrate.
Cherry hemangioma	Benign capillary hemangioma B commonly appearing in middle-aged adults. Does not regress. Frequency ↑ with age.
Glomus tumor	Benign, painful, red-blue tumor, commonly under fingernails C . Arises from modified smooth muscle cells of the thermoregulatory glomus body.
Kaposi sarcoma	Endothelial malignancy most commonly affecting the skin, mouth, GI tract, respiratory tract. Classically seen in older Eastern European males, patients with AIDS, and organ transplant patients. Associated with HHV-8 and HIV. Rarely mistaken for bacillary angiomatosis, but has lymphocytic infiltrate.
Pyogenic granuloma	Polypoid lobulated capillary hemangioma D that can ulcerate and bleed. Associated with trauma and pregnancy.
Strawberry hemangioma	Benign capillary hemangioma of infancy E . Appears in first few weeks of life (1/200 births); grows rapidly and regresses spontaneously by 5–8 years old.



Skin infections**Bacterial infections**

Impetigo	Very superficial skin infection. Usually from <i>S aureus</i> or <i>S pyogenes</i> . Highly contagious. Honey-colored crusting A . Bullous impetigo B has bullae and is usually caused by <i>S aureus</i> .
Erysipelas	Infection involving upper dermis and superficial lymphatics, usually from <i>S pyogenes</i> . Presents with well-defined, raised demarcation between infected and normal skin C .
Cellulitis	Acute, painful, spreading infection of deeper dermis and subcutaneous tissues. Usually from <i>S pyogenes</i> or <i>S aureus</i> . Often starts with a break in skin from trauma or another infection D .
Abscess	Collection of pus from a walled-off infection within deeper layers of skin E . Offending organism is almost always <i>S aureus</i> .
Necrotizing fasciitis	Deeper tissue injury, usually from anaerobic bacteria or <i>S pyogenes</i> . Pain may be out of proportion to exam findings. Results in crepitus from methane and CO ₂ production. “Flesh-eating bacteria.” Causes bullae and skin necrosis → violaceous color of bullae, surrounding skin F . Surgical emergency.
Staphylococcal scalded skin syndrome	Exotoxin destroys keratinocyte attachments in stratum granulosum only (vs toxic epidermal necrolysis, which destroys epidermal-dermal junction). Characterized by fever and generalized erythematous rash with sloughing of the upper layers of the epidermis G that heals completely. ⊕ Nikolsky sign (separation of epidermis upon manual stroking of skin). Commonly seen in newborns and children/adults with renal insufficiency.


Viral infections

Herpes	Herpes virus infections (HSV1 and HSV2) of skin can occur anywhere from mucosal surfaces to normal skin. These include herpes labialis, herpes genitalis, herpetic whitlow H (finger).
Molluscum contagiosum	Umbilicated papules I caused by a poxvirus. While frequently seen in children, it may be sexually transmitted in adults.
Varicella zoster virus	Causes varicella (chickenpox) and zoster (shingles). Varicella presents with multiple crops of lesions in various stages from vesicles to crusts. Zoster is a reactivation of the virus in dermatomal distribution (unless it is disseminated).
Hairy leukoplakia	Irregular, white, painless plaques on lateral tongue that cannot be scraped off J . EBV mediated. Occurs in HIV-positive patients, organ transplant recipients. Contrast with thrush (scrapable) and leukoplakia (precancerous).



Autoimmune blistering skin disorders

	Pemphigus vulgaris	Bullous pemphigoid
PATHOPHYSIOLOGY	Potentially fatal. Most commonly seen in older adults. Type II hypersensitivity reaction. IgG antibodies against desmoglein-1 and/or desmoglein-3 (component of desmosomes, which connect keratinocytes in the stratum spinosum).	Less severe than pemphigus vulgaris. Most commonly seen in older adults. Type II hypersensitivity reaction. IgG antibodies against hemidesmosomes (epidermal basement membrane; antibodies are “ bullo ” the epidermis).
GROSS MORPHOLOGY	Flaccid intraepidermal bullae A caused by acantholysis (separation of keratinocytes, “row of tombstones” on H&E stain); oral mucosa is involved. Nikolsky sign ⊕.	Tense blisters C containing eosinophils; oral mucosa spared. Nikolsky sign ⊖.
IMMUNOFLUORESCENCE	Reticular pattern around epidermal cells B .	Linear pattern at epidermal-dermal junction D .



Other blistering skin disorders

Dermatitis herpetiformis	Pruritic papules, vesicles, and bullae (often found on elbows, knees, buttocks) A . Deposits of IgA at tips of dermal papillae. Associated with celiac disease. Treatment: dapsone, gluten-free diet.
Erythema multiforme	Associated with infections (eg, <i>Mycoplasma pneumoniae</i> , HSV), drugs (eg, sulfa drugs, β -lactams, phenytoin). Presents with multiple types of lesions—macules, papules, vesicles, target lesions (look like targets with multiple rings and dusky center showing epithelial disruption) B .
Stevens-Johnson syndrome	Characterized by fever, bullae formation and necrosis, sloughing of skin at dermal-epidermal junction (\oplus Nikolsky), high mortality rate. Typically mucous membranes are involved C D . Targetoid skin lesions may appear, as seen in erythema multiforme. Usually associated with adverse drug reaction. Toxic epidermal necrolysis (TEN) E F is more severe form of SJS involving > 30% body surface area. 10–30% involvement denotes SJS-TEN.




Miscellaneous skin disorders

Acanthosis nigricans	Epidermal hyperplasia causing symmetric, hyperpigmented thickening of skin, especially in axilla or on neck A B . Associated with insulin resistance (eg, diabetes, obesity, Cushing syndrome, PCOS), visceral malignancy (eg, gastric adenocarcinoma).
Actinic keratosis	Premalignant lesions caused by sun exposure. Small, rough, erythematous or brownish papules or plaques C D . Risk of squamous cell carcinoma is proportional to degree of epithelial dysplasia.
Erythema nodosum	Painful, raised inflammatory lesions of subcutaneous fat (panniculitis), usually on anterior shins. Often idiopathic, but can be associated with sarcoidosis, coccidioidomycosis, histoplasmosis, TB, streptococcal infections E , leprosy F , inflammatory bowel disease.
Lichen Planus	P ruritic, P urple, P olygonal P lanar P apules and P laques are the 6 P 's of lichen P lanus G H . Mucosal involvement manifests as Wickham striae (reticular white lines) and hypergranulosis. Sawtooth infiltrate of lymphocytes at dermal-epidermal junction. Associated with hepatitis C.
Pityriasis rosea	“Herald patch” I followed days later by other scaly erythematous plaques, often in a “Christmas tree” distribution on trunk J . Multiple pink plaques with collarette scale. Self-resolving in 6–8 weeks.
Sunburn	Acute cutaneous inflammatory reaction due to excessive UV irradiation. Causes DNA mutations, inducing apoptosis of keratinocytes. UVB is dominant in sun B urn, UVA in t A nning and photo A ging. Exposure to UVA and UVB ↑ risk of skin cancer.



Rule of 9's

The extent of a burn injury can be estimated as a percentage of the body surface area.

**Burn classification**

DEPTH	INVOLVEMENT	APPEARANCE	SENSATION
Superficial burn	Epidermis only	Similar to sunburn; localized, painful, dry, blanching redness with no blisters	Painful
Superficial partial-thickness	All of epidermis and some dermis	Blisters, blanches with pressure, swollen, warm	Painful to temperature and air
Deep partial-thickness burn	All of epidermis and some dermis	Blisters (easily unroofed), does not blanch with pressure	Painless; perception of pressure only
Full-thickness burn	All of skin (epidermis and dermis)	White, waxy, dry, inelastic, leathery, does not blanch with pressure	Painless; perception of deep pressure only
Deeper injury burn	All of skin and at least partial involvement of muscle and/or fascia	White, dry, inelastic, does not blanch with pressure	Painless; some perception of deep pressure

Skin cancer

Basal cell carcinoma more common above **upper lip**

Squamous cell carcinoma more common below **lower lip**

Sun exposure strongly predisposes to skin cancer.



Basal cell carcinoma

Most common skin cancer. Found in sun-exposed areas of body (eg, face). Locally invasive, but rarely metastasizes. Waxy, pink, pearly nodules, commonly with telangiectasias, rolled borders **A**, central crusting or ulceration. BCCs also appear as nonhealing ulcers with infiltrating growth **B** or as a scaling plaque (superficial BCC) **C**. Basal cell tumors have “palisading” (aligned) nuclei **D**.

Keratoacanthoma

Seen in middle-aged and elderly individuals. Rapidly growing, resembles squamous cell carcinoma. Presents as dome-shaped nodule with keratin-filled center. Grows rapidly (4-6 weeks) and may spontaneously regress **E**.

Melanoma

Common tumor with significant risk of metastasis. S-100 tumor marker. Associated with dysplastic nevi; fair-skinned persons are at ↑ risk. Depth of tumor (Breslow thickness) correlates with risk of metastasis. Look for the **ABCDEs**: **A**symmetry, **B**order irregularity, **C**olor variation, **D**iameter > 6 mm, and **E**volution over time. At least 4 different types of melanoma, including superficial spreading **F**, nodular **G**, lentigo maligna **H**, and acral lentiginous (highest prevalence in African-Americans and Asians) **I**. Often driven by activating mutation in BRAF kinase. Primary treatment is excision with appropriately wide margins. Metastatic or unresectable melanoma in patients with *BRAF V600E* mutation may benefit from vemurafenib, a BRAF kinase inhibitor.

Squamous cell carcinoma


Second most common skin cancer. Associated with immunosuppression, chronic nonhealing wounds, and occasionally arsenic exposure. Commonly appears on face **J**, lower lip **K**, ears, hands. Locally invasive, may spread to lymph nodes, and will rarely metastasize. Ulcerative red lesions. Histopathology: keratin “pearls” **L**.

Actinic keratosis, a scaly plaque, is a precursor to squamous cell carcinoma.



▶ MUSCULOSKELETAL, SKIN, AND CONNECTIVE TISSUE—PHARMACOLOGY

Arachidonic acid pathways



LTB_4 is a **neutrophil** chemotactic agent. **Neutrophils** arrive “**B4**” others.
 PGI_2 inhibits platelet aggregation and promotes vasodilation. **Platelet-Gathering Inhibitor.**

Acetaminophen

MECHANISM	Reversibly inhibits cyclooxygenase, mostly in CNS. Inactivated peripherally.
CLINICAL USE	Antipyretic, analgesic, but not anti-inflammatory. Used instead of aspirin to avoid Reye syndrome in children with viral infection.
ADVERSE EFFECTS	Overdose produces hepatic necrosis; acetaminophen metabolite (NAPQI) depletes glutathione and forms toxic tissue byproducts in liver. N-acetylcysteine is antidote—regenerates glutathione.

Aspirin

MECHANISM	NSAID that irreversibly inhibits cyclooxygenase (both COX-1 and COX-2) by covalent acetylation → ↓ synthesis of TXA ₂ and prostaglandins. ↑ bleeding time. No effect on PT, PTT. Effect lasts until new platelets are produced.
CLINICAL USE	Low dose (< 300 mg/day): ↓ platelet aggregation. Intermediate dose (300–2400 mg/day): antipyretic and analgesic. High dose (2400–4000 mg/day): anti-inflammatory.
ADVERSE EFFECTS	Gastric ulceration, tinnitus (CN VIII), allergic reactions (especially in patients with asthma or nasal polyps). Chronic use can lead to acute kidney injury, interstitial nephritis, GI bleeding. Risk of Reye syndrome in children treated with aspirin for viral infection. Toxic doses cause respiratory alkalosis early, but transitions to mixed metabolic acidosis-respiratory alkalosis. Treatment of overdose: NaHCO ₃ .

Celecoxib

MECHANISM	Reversibly and selectively inhibits the cyclooxygenase (COX) isoform 2 (“ Selecoxib ”), which is found in inflammatory cells and vascular endothelium and mediates inflammation and pain; spares COX-1, which helps maintain gastric mucosa. Thus, does not have the corrosive effects of other NSAIDs on the GI lining. Spares platelet function as TXA ₂ production is dependent on COX-1.
CLINICAL USE	Rheumatoid arthritis, osteoarthritis.
ADVERSE EFFECTS	↑ risk of thrombosis, sulfa allergy.

Nonsteroidal anti-inflammatory drugs

Ibuprofen, naproxen, indomethacin, ketorolac, diclofenac, meloxicam, piroxicam.

MECHANISM	Reversibly inhibit cyclooxygenase (both COX-1 and COX-2). Block prostaglandin synthesis.
CLINICAL USE	Antipyretic, analgesic, anti-inflammatory. Indomethacin is used to close a PDA.
ADVERSE EFFECTS	Interstitial nephritis, gastric ulcer (prostaglandins protect gastric mucosa), renal ischemia (prostaglandins vasodilate afferent arteriole), aplastic anemia.

Leflunomide

MECHANISM	Reversibly inhibits dihydroorotate dehydrogenase, preventing pyrimidine synthesis. Suppresses T-cell proliferation.
CLINICAL USE	Rheumatoid arthritis, psoriatic arthritis.
ADVERSE EFFECTS	Diarrhea, hypertension, hepatotoxicity, teratogenicity.

Bisphosphonates

Alendronate, ibandronate, risedronate, zoledronate.


MECHANISM	Pyrophosphate analogs; bind hydroxyapatite in bone, inhibiting osteoclast activity.
CLINICAL USE	Osteoporosis, hypercalcemia, Paget disease of bone, metastatic bone disease, osteogenesis imperfecta.
ADVERSE EFFECTS	Esophagitis (if taken orally, patients are advised to take with water and remain upright for 30 minutes), osteonecrosis of jaw, atypical femoral stress fractures.

Teriparatide

MECHANISM	Recombinant PTH analog, ↑ osteoblastic activity when administered in pulsatile fashion.
CLINICAL USE	Osteoporosis. Causes ↑ bone growth compared to antiresorptive therapies (eg, bisphosphonates).
ADVERSE EFFECTS	↑ risk of osteosarcoma (avoid use in patients with Paget disease of the bone or unexplained elevation of alkaline phosphatase). Avoid in patients who have had prior cancers or radiation therapy. Transient hypercalcemia.

Gout drugs**Chronic gout drugs (preventive)**

Probenecid	Inhibits reabsorption of uric acid in proximal convoluted tubule (also inhibits secretion of penicillin). Can precipitate uric acid calculi.
Allopurinol	Competitive inhibitor of xanthine oxidase → ↓ conversion of hypoxanthine and xanthine to urate. Also used in lymphoma and leukemia to prevent tumor lysis–associated urate nephropathy. ↑ concentrations of xanthine oxidase active metabolites, azathioprine, and 6-MP.
Pegloticase	Recombinant uricase catalyzing uric acid to allantoin (a more water-soluble product).
Febuxostat	Inhibits xanthine oxidase.
Acute gout drugs	
NSAIDs	Any NSAID. Use salicylates with caution (may decrease uric acid excretion, particularly at low doses).
Glucocorticoids	Oral, intra-articular, or parenteral.
Colchicine	Binds and stabilizes tubulin to inhibit microtubule polymerization, impairing neutrophil chemotaxis and degranulation. Acute and prophylactic value. GI, neuromyopathic side effects.

Prevent A Painful Flare.**TNF- α inhibitors**

DRUG	MECHANISM	CLINICAL USE	ADVERSE EFFECTS
Etanercept	Fusion protein (decoy receptor for TNF- α + IgG ₁ Fc), produced by recombinant DNA. Etanercept intercepts TNF.	Rheumatoid arthritis, psoriasis, ankylosing spondylitis	Predisposition to infection, including reactivation of latent TB, since TNF is important in granuloma formation and stabilization. Can also lead to drug-induced lupus.
Infliximab, adalimumab, certolizumab, golimumab	Anti-TNF- α monoclonal antibody.	Inflammatory bowel disease, rheumatoid arthritis, ankylosing spondylitis, psoriasis	

HIGH-YIELD SYSTEMS

Neurology and Special Senses

“We are all now connected by the Internet, like neurons in a giant brain.”
—Stephen Hawking

“Anything’s possible if you’ve got enough nerve.”
—J.K. Rowling, *Harry Potter and the Order of the Phoenix*

“I like nonsense; it wakes up the brain cells.”
—Dr. Seuss

“I believe in an open mind, but not so open that your brains fall out.”
—Arthur Hays Sulzberger

“The chief function of the body is to carry the brain around.”
—Thomas Edison


“Exactly how [the brain] operates remains one of the biggest unsolved mysteries, and it seems the more we probe its secrets, the more surprises we find.”
—Neil deGrasse Tyson

Understand the difference between upper motor neuron (UMN) and lower motor neuron (LMN) findings and the underlying anatomy. Know the major motor, sensory, cerebellar and visual pathways and their respective locations in the CNS. Connect key neurological associations with certain pathologies (eg, cerebellar lesions, stroke manifestations, Brown-Séquard syndrome). Recognize common findings on MRI/CT (eg, ischemic and hemorrhagic stroke) and on neuropathology (eg, neurofibrillary tangles and Lewy bodies). High-yield medications include those used to treat epilepsy, Parkinson disease, migraine, and pain (eg, opioids).

▶ Embryology	490
▶ Anatomy and Physiology	493
▶ Pathology	511
▶ Otology	533
▶ Ophthalmology	534
▶ Pharmacology	544

▶ NEUROLOGY—EMBRYOLOGY

Neural development



Notochord induces overlying ectoderm to differentiate into neuroectoderm and form neural plate. Neural plate gives rise to neural tube and neural crest cells.


Notochord becomes nucleus pulposus of intervertebral disc in adults.

Alar plate (dorsal): sensory; regulated by TGF- β (including bone morphogenetic protein [BMP])
 Basal plate (ventral): motor; regulated by sonic hedgehog gene (*SHH*)

Same orientation as spinal cord

Regional specification of developing brain

Telencephalon is the 1st part. **D**iencephalon is the 2nd part. The rest are arranged alphabetically: **m**esencephalon, **m**etencephalon, **m**yelencephalon.




Central and peripheral nervous systems origins

Neuroepithelia in neural tube—CNS neurons, ependymal cells (inner lining of ventricles, make CSF), oligodendrocytes, astrocytes.


Neural crest—PNS neurons, Schwann cells, glia, melanocytes, adrenal medulla.

Mesoderm—Microglia (like Macrophages).

Neural tube defects	Neuropores fail to fuse (4th week) → persistent connection between amniotic cavity and spinal canal. Associated with maternal diabetes and folate deficiency. ↑ α -fetoprotein (AFP) in amniotic fluid and maternal serum (except spina bifida occulta = normal AFP). ↑ acetylcholinesterase (AChE) in amniotic fluid is a helpful confirmatory test.
Spina bifida occulta	Failure of caudal neuropore to close, but no herniation. Usually seen at lower vertebral levels. Dura is intact. Associated with tuft of hair or skin dimple at level of bony defect.
Meningocele	Meninges (but no neural tissue) herniate through bony defect.
Myelomeningocele	Meninges and neural tissue (eg, cauda equina) herniate through bony defect.
Myeloschisis	Also called rachischisis. Exposed, unfused neural tissue without skin/meningeal covering.
Anencephaly	Failure of rostral neuropore to close → no forebrain, open calvarium. Clinical findings: polyhydramnios (no swallowing center in brain).



Holoprosencephaly



Failure of the embryonic forebrain (prosencephalon) to separate into 2 cerebral hemispheres; usually occurs during weeks 5–6. May be related to mutations in sonic hedgehog signaling pathway. Associated with other midline defects including cleft lip/palate (moderate form) and cyclopia (severe form). ↑ risk for pituitary dysfunction (eg, diabetes insipidus). Can be seen with Patau syndrome (trisomy 13). MRI reveals monoventricle **A** and fusion of basal ganglia (star in **A**).

Lissencephaly


Failure of neuronal migration resulting in a “smooth brain” that lacks sulci and gyri. May be associated with microcephaly, ventriculomegaly.

Posterior fossa malformations

Chiari I malformation Ectopia of cerebellar **tonsils** inferior to foramen magnum (**1** structure) **A**. Congenital, usually asymptomatic in childhood, manifests in adulthood with headaches and cerebellar symptoms. Associated with spinal cavitations (eg, syringomyelia).

Chiari II malformation Herniation of cerebellar **vermis** and **tonsils** (**2** structures) through foramen magnum with aqueductal stenosis → noncommunicating hydrocephalus. Usually associated with lumbosacral myelomeningocele (may present as paralysis/sensory loss at and below the level of the lesion). More severe than Chiari I, usually presents early in life.

Dandy-Walker malformation Agenesis of cerebellar vermis → cystic enlargement of 4th ventricle (arrow in **B**) that fills the enlarged posterior fossa. Associated with noncommunicating hydrocephalus, spina bifida.





Syringomyelia



Cystic cavity (syrinx) within central canal of spinal cord (yellow arrows in **A**). Fibers crossing in anterior white commissure (spinothalamic tract) are typically damaged first. Results in a “cape-like,” bilateral, symmetrical loss of pain and temperature sensation in upper extremities (fine touch sensation is preserved).

Associated with Chiari I malformation (red arrow in **A** shows low-lying cerebellar tonsils), scoliosis and other congenital malformations; acquired causes include trauma and tumors. Most common location cervical > thoracic >> lumbar. **Syrinx** = tube, as in “syringe.”



Tongue development

1st and 2nd pharyngeal arches form anterior 2/3 (thus sensation via CN V₃, taste via CN VII).
3rd and 4th pharyngeal arches form posterior 1/3 (thus sensation and taste mainly via CN IX, extreme posterior via CN X).

Motor innervation is via CN XII to hyoglossus (retracts and depresses tongue), **genioglossus** (**protrudes** tongue), and **styloglossus** (draws sides of tongue upward to create a trough for swallowing).

Motor innervation is via CN X to palatoglossus (elevates posterior tongue during swallowing).

Taste—CN VII, IX, X (solitary nucleus).

Pain—CN V₃, IX, X.


Motor—CN X, XII.

The **Genie** comes **out** of the lamp in **style**.

▶ NEUROLOGY—ANATOMY AND PHYSIOLOGY**Neurons**

Signal-transmitting cells of the nervous system. Permanent cells—do not divide in adulthood.


Signal-relaying cells with dendrites (receive input), cell bodies, and axons (send output). Cell bodies and dendrites can be seen on Nissl staining (stains RER). RER is not present in the axon. Neuron markers: neurofilament protein, synaptophysin.

Astrocytes

Most common glial cell type in CNS. Physical support, repair, extracellular K⁺ buffer, removal of excess neurotransmitter, component of blood-brain barrier, glycogen fuel reserve buffer. Reactive gliosis in response to neural injury.

Derived from neuroectoderm.

Astrocyte marker: GFAP.


Microglia

Phagocytic scavenger cells of CNS (mesodermal, mononuclear origin). Activation in response to tissue damage → release of inflammatory mediators (eg, nitric oxide, glutamate). Not readily discernible by Nissl stain.

HIV-infected microglia fuse to form multinucleated giant cells in CNS seen in HIV-associated dementia.

Ependymal cells

Ciliated simple columnar glial cells line the ventricles and central canal of spinal cord. Apical surfaces are covered in cilia (which circulate CSF) and microvilli (which help with CSF absorption). Specialized ependymal cells (choroid plexus) produce CSF.


Myelin

↑ conduction velocity of signals transmitted down axons → saltatory conduction of action potential at the nodes of Ranvier, where there are high concentrations of Na^+ channels.

In CNS (including CN II), myelin is synthesized by oligodendrocytes; in PNS (including CN III-XII), myelin is synthesized by Schwann cells.

Wraps and insulates axons (arrow in **A**): ↑ space constant and ↑ conduction velocity.


COPS: CNS = Oligodendrocytes, PNS = Schwann cells.

Schwann cells

Promote axonal regeneration. Derived from neural crest.

Each “Schwone” cell myelinates only **1** PNS axon.

Injured in Guillain-Barré syndrome.

Oligodendrocytes

Myelinate axons of neurons in CNS. Each oligodendrocyte can myelinate many axons (~ 30). Predominant type of glial cell in white matter.

Derived from neuroectoderm.


“Fried egg” appearance histologically.

Injured in multiple sclerosis, progressive multifocal leukoencephalopathy (PML), leukodystrophies.

Sensory receptors

RECEPTOR TYPE	SENSORY NEURON FIBER TYPE	LOCATION	SENSES
Free nerve endings	Aδ —fast, myelinated fibers C —slow, unmyelinated A Delta plane is fast, but a tax C is slow	All skin, epidermis, some viscera	Pain, temperature
Meissner corpuscles	Large, myelinated fibers; adapt quickly	Glabrous (hairless) skin	Dynamic, fine/light touch, position sense, low-frequency vibration
Pacinian corpuscles	Large, myelinated fibers; adapt quickly	Deep skin layers, ligaments, joints	High-frequency vibration, pressure
Merkel discs	Large, myelinated fibers; adapt slowly	Finger tips, superficial skin	Pressure, deep static touch (eg, shapes, edges), position sense
Ruffini corpuscles	Dendritic endings with capsule; adapt slowly	Finger tips, joints	Pressure, slippage of objects along surface of skin, joint angle change

Peripheral nerve




Endoneurium—thin, supportive connective tissue that ensheathes and supports individual myelinated nerve fibers.

Endo = inner
Peri = around
Epi = outer

Perineurium (blood-nerve **P**ermeability barrier)—surrounds a fascicle of nerve fibers.

Epineurium—dense connective tissue that surrounds entire nerve (fascicles and blood vessels).

Chromatolysis




Reaction of neuronal cell body to axonal injury. Changes reflect ↑ protein synthesis in effort to repair the damaged axon. Characterized by:

- Round cellular swelling **A**
- Displacement of the nucleus to the periphery
- Dispersion of Nissl substance throughout cytoplasm

Wallerian degeneration—disintegration of the axon and myelin sheath distal to site of axonal injury with macrophages removing debris.


Proximal to the injury, the axon retracts, and the cell body sprouts new protrusions that grow toward other neurons for potential reinnervation. Serves as a preparation for axonal regeneration and functional recovery.



Neurotransmitter changes with disease

	LOCATION OF SYNTHESIS	ANXIETY	DEPRESSION	SCHIZOPHRENIA	ALZHEIMER DISEASE	HUNTINGTON DISEASE	PARKINSON DISEASE
Acetylcholine	Basal nucleus of Meynert				↓	↓	↑
Dopamine	Ventral tegmentum, SNc		↓	↑		↑	↓
GABA	Nucleus accumbens	↓				↓	
Norepinephrine	Locus ceruleus (pons)	↑	↓				
Serotonin	Raphe nuclei (medulla, pons)	↓	↓				↓

Meninges




Three membranes that surround and protect the brain and spinal cord:

- Dura mater—thick outer layer closest to skull. Derived from mesoderm.
- Arachnoid mater—middle layer, contains web-like connections. Derived from neural crest.
- Pia mater—thin, fibrous inner layer that firmly adheres to brain and spinal cord. Derived from neural crest.

CSF flows in the subarachnoid space, located between arachnoid and pia mater.

Epidural space—potential space between the dura mater and skull/vertebral column containing fat and blood vessels. Site of blood collection associated with middle meningeal artery injury.

Blood-brain barrier



Prevents circulating blood substances (eg, bacteria, drugs) from reaching the CSF/CNS. Formed by 3 structures:

- Tight junctions between nonfenestrated capillary endothelial cells
- Basement membrane
- Astrocyte foot processes

Glucose and amino acids cross slowly by carrier-mediated transport mechanisms.

Nonpolar/lipid-soluble substances cross rapidly via diffusion.

Circumventricular organs with fenestrated capillaries and no blood-brain barrier allow molecules in blood to affect brain function (eg, area postrema—vomiting after chemotherapy; OVLT [organum vasculosum lamina terminalis]—osmoreceptors) or neurosecretory products to enter circulation (eg, neurohypophysis—ADH release).

Infarction and/or neoplasm destroys endothelial cell tight junctions → vasogenic edema.

Hyperosmolar agents (eg, mannitol) can disrupt the BBB → ↑ permeability of medications.

Vomiting center

Coordinated by nucleus tractus solitarius (NTS) in the medulla, which receives information from the chemoreceptor trigger zone (CTZ, located within area postrema in 4th ventricle), GI tract (via vagus nerve), vestibular system, and CNS.

CTZ and adjacent vomiting center nuclei receive input from 5 major receptors: muscarinic (M_1), dopamine (D_2), histamine (H_1), serotonin ($5-HT_3$), and neurokinin (NK-1) receptors.

- $5-HT_3$, D_2 , and NK-1 antagonists used to treat chemotherapy-induced vomiting.
- H_1 and M_1 antagonists treat motion sickness; H_1 antagonists treat hyperemesis gravidarum.

Sleep physiology

Sleep cycle is regulated by the circadian rhythm, which is driven by suprachiasmatic nucleus (SCN) of the hypothalamus. Circadian rhythm controls nocturnal release of ACTH, prolactin, melatonin, norepinephrine: SCN → norepinephrine release → pineal gland → ↑ melatonin. SCN is regulated by environment (eg, light).

Two stages: rapid-eye movement (REM) and non-REM.

Alcohol, benzodiazepines, and barbiturates are associated with ↓ REM sleep and N3 sleep; norepinephrine also ↓ REM sleep.


Benzodiazepines are useful for night terrors and sleepwalking by ↓ N3 and REM sleep.

SLEEP STAGE (% OF TOTAL SLEEP TIME IN YOUNG ADULTS)	DESCRIPTION	EEG WAVEFORM AND NOTES
Awake (eyes open)	Alert, active mental concentration.	Beta (highest frequency, lowest amplitude)
Awake (eyes closed)		Alpha
Non-REM sleep		
Stage N1 (5%)	Light sleep.	Theta
Stage N2 (45%)	Deeper sleep; when bruxism (“ twoth ” [tooth grinding] occurs).	Sleep spindles and K complexes
Stage N3 (25%)	Deepest non-REM sleep (slow-wave sleep); sleepwalking , night terrors, and bedwetting occur (wee and flee in N3).	Delta (lowest frequency, highest amplitude)
REM sleep (25%)	Loss of motor tone, ↑ brain O ₂ use, variable pulse/BP, ↑ ACh. REM is when dreaming, nightmares, and penile/clitoral tumescence occur; may serve memory processing function. Extraocular movements due to activity of PPRF (paramedian pontine reticular formation/conjugate gaze center). Occurs every 90 minutes, and duration ↑ through the night.	Beta Changes in elderly: ↓ REM sleep time, ↓ N3. Changes in depression: ↑ REM sleep time, ↓ REM latency, ↓ N3, repeated nighttime awakenings, early morning awakening (terminal insomnia). Changes in narcolepsy: ↓ REM latency. At night, BATS Drink Blood

Hypothalamus	Maintains homeostasis by regulating T hirst and water balance, controlling A denohypophysis (anterior pituitary) and N eurohypophysis (posterior pituitary) release of hormones produced in the hypothalamus, and regulating H unger, A utonomic nervous system, T emperature, and S exual urges (TAN HATS). Inputs (areas not protected by blood-brain barrier): OVLT (senses change in osmolarity), area postrema (found in dorsal medulla, responds to emetics).	
Lateral nucleus	Hunger. Destruction → anorexia, failure to thrive (infants). Stimulated by ghrelin, inhibited by leptin.	L ateral injury makes you L ean.
Ventromedial nucleus	Satiety. Destruction (eg, craniopharyngioma) → hyperphagia. Stimulated by leptin.	V entro M edial injury makes you V ery M assive.
Anterior nucleus	Cooling, parasympathetic.	A/C = A nterior C ooling.
Posterior nucleus	Heating, sympathetic.	H eating controlled by P osterior nucleus (“ H ot P ot”).
Suprachiasmatic nucleus	Circadian rhythm.	SCN is a S un- C ensing N ucleus.
Supraoptic and paraventricular nuclei	Synthesize ADH and oxytocin.	SAD POX : S upraoptic = A DH, P araventricular = O xytocin ADH and oxytocin are carried by neurophysins down axons to posterior pituitary, where these hormones are stored and released.
Preoptic nucleus	Thermoregulation, sexual behavior. Releases GnRH.	Failure of GnRH-producing neurons to migrate from olfactory pit → Kallmann syndrome.

Thalamus Major relay for all ascending sensory information except olfaction.

NUCLEI	INPUT	SENSES	DESTINATION	MNEMONIC
Ventral Postero-Lateral nucleus	Spinothalamic and dorsal columns/medial lemniscus	V ibration, P ain, P ressure, P roprioception, L ight touch, temperature	1° somatosensory cortex	
Ventral postero-Medial nucleus	Trigeminal and gustatory pathway	F ace sensation, taste	1° somatosensory cortex	M akeup goes on the f ace
Lateral geniculate nucleus	CN II, optic chiasm, optic tract	Vision	1° visual cortex (calcarine sulcus)	L ateral = L ight
Medial geniculate nucleus	Superior olive and inferior colliculus of tectum	Hearing	Auditory cortex of temporal lobe	M edial = M usic
Ventral lateral nucleus	Cerebellum, basal ganglia	Motor	Motor cortex	

Limbic system

Collection of neural structures involved in emotion, long-term memory, olfaction, behavior modulation, ANS function.


Consists of hippocampus (red arrows in **A**), amygdalae, mammillary bodies, anterior thalamic nuclei, cingulate gyrus (yellow arrows in **A**), entorhinal cortex. Responsible for **F**eeding, **F**leeing, **F**ighting, **F**eeling, and **S**ex.

The famous **5 F**'s.

Dopaminergic pathways

Commonly altered by drugs (eg, antipsychotics) and movement disorders (eg, Parkinson disease).

PATHWAY	SYMPTOMS OF ALTERED ACTIVITY	NOTES
Mesocortical	↓ activity → “negative” symptoms (eg, anergia, apathy, lack of spontaneity)	Antipsychotic drugs have limited effect
Mesolimbic	↑ activity → “positive” symptoms (eg, delusions, hallucinations)	1° therapeutic target of antipsychotic drugs → ↓ positive symptoms (eg, in schizophrenia)
Nigrostriatal	↓ activity → extrapyramidal symptoms (eg, dystonia, akathisia, parkinsonism, tardive dyskinesia)	Major dopaminergic pathway in brain Significantly affected by movement disorders and antipsychotic drugs
Tuberoinfundibular	↓ activity → ↑ prolactin → ↓ libido, sexual dysfunction, galactorrhea, gynecomastia (in men)	

Cerebellum

Modulates movement; aids in coordination and balance **A**.

Input:

- Contralateral cortex via middle cerebellar peduncle
- Ipsilateral proprioceptive information via inferior cerebellar peduncle from spinal cord

Output:


- The only output of cerebellar cortex = Purkinje cells (always **i**nhibitory) → deep nuclei of cerebellum → contralateral cortex via superior cerebellar peduncle
- Deep nuclei (lateral → medial)—**D**entate, **E**mboliform, **G**lobose, **F**astigial

Lateral lesions—affect voluntary movement of extremities (**lateral** structures); when injured, propensity to fall toward injured (ipsilateral) side.

Medial lesions (eg, vermis, fastigial nuclei, flocculonodular lobe)—truncal ataxia (wide-based cerebellar gait), nystagmus, head tilting. Generally result in bilateral motor deficits affecting axial and proximal limb musculature (**medial** structures).

Don't **E**at **G**reasy **F**oods

Basal ganglia





Important in voluntary movements and adjusting posture **A**.
Receives cortical input, provides negative feedback to cortex to modulate movement.
Striatum = putamen (motor) + **C**audate (cognitive).
Lentiform = putamen + globus pallidus.

D₁ Receptor = **D**IRect pathway.
Indirect (**D**₂) = **I**nhibitory.


Direct (excitatory) pathway—SNc input to the striatum via the nigrostriatal dopaminergic pathway releases GABA, which inhibits GABA release from the GPi, disinhibiting the **T**halamus via the GPi (↑ motion).

Indirect (inhibitory) pathway—SNc input to the striatum via the nigrostriatal dopaminergic pathway releases GABA that disinhibits STN via GPe inhibition, and STN stimulates GPi to inhibit the thalamus (↓ motion).

Dopamine binds to **D**₁, stimulating the excitatory pathway, and to **D**₂, inhibiting the inhibitory pathway → ↑ motion.



Cerebral cortex regions



Cerebral perfusion

Relies on tight autoregulation. Primarily driven by PCO_2 (PO_2 also modulates perfusion in severe hypoxia).


Also relies on a pressure gradient between mean arterial pressure (MAP) and intracranial pressure (ICP). \downarrow blood pressure or \uparrow ICP \rightarrow \downarrow cerebral perfusion pressure (CPP).

Therapeutic hyperventilation \rightarrow $\downarrow PCO_2$ \rightarrow vasoconstriction \rightarrow \downarrow cerebral blood flow \rightarrow \downarrow ICP. May be used to treat acute cerebral edema (eg, 2° to stroke) unresponsive to other interventions.

CPP = MAP – ICP. If CPP = 0, there is no cerebral perfusion \rightarrow brain death.


Hypoxemia increases CPP only if $PO_2 < 50$ mm Hg.

CPP is directly proportional to PCO_2 until $PCO_2 > 90$ mm Hg.





Homunculus


Topographic representation of motor and sensory areas in the cerebral cortex. Distorted appearance is due to certain body regions being more richly innervated and thus having ↑ cortical representation.



Cerebral arteries—cortical distribution



Watershed zones




Cortical border zones occur between anterior and middle cerebral arteries and posterior and middle cerebral arteries (blue areas in **A**). Internal border zones occur between the superficial and deep vascular territories of the middle cerebral artery (red areas in **A**).

Infarct due to severe hypoperfusion → proximal upper and lower extremity weakness (“man-in-the-barrel syndrome”), higher order visual dysfunction (if posterior cerebral/middle cerebral cortical border zone stroke).


Circle of Willis

System of anastomoses between anterior and posterior blood supplies to brain.




Dural venous sinuses


Large venous channels **A** that run through the periosteal and meningeal layers of the dura mater. Drain blood from cerebral veins (arrow) and receive CSF from arachnoid granulations. Empty into internal jugular vein.



Venous sinus thrombosis—presents with signs/symptoms of ↑ ICP (eg, headache, seizures, papilledema, focal neurologic deficits). May lead to venous hemorrhage. Associated with hypercoagulable states (eg, pregnancy, OCP use, factor V Leiden).



Ventricular system




Lateral ventricles → 3rd ventricle via right and left interventricular foramina of Monro.
 3rd ventricle → 4th ventricle via cerebral aqueduct of Sylvius.

4th ventricle → subarachnoid space via:

- Foramina of **L**uschka = **L**ateral.
- Foramen of **M**agendie = **M**edial.


CSF made by choroid plexuses located in the lateral and fourth ventricles. Travels to subarachnoid space via foramina of Luschka and Magendie, is reabsorbed by arachnoid granulations, and then drains into dural venous sinuses.

Brain stem—ventral view



- 4 CN are above pons (I, II, III, IV).
 - 4 CN exit the pons (V, VI, VII, VIII).
 - 4 CN are in medulla (IX, X, XI, XII).
 - 4 CN nuclei are medial (III, IV, VI, XII).
- “Factors of 12, except 1 and 2.”

Brain stem—dorsal view (cerebellum removed)



Pineal gland—melatonin secretion, circadian rhythms.
 Superior colliculi—direct eye movements to stimuli (noise/movements) or objects of interest.
 Inferior colliculi—auditory.
 Your eyes are **above** your ears, and the superior colliculus (visual) is **above** the inferior colliculus (auditory).

Cranial nerve nuclei

Located in tegmentum portion of brain stem (between dorsal and ventral portions):


- Midbrain—nuclei of CN III, IV
- Pons—nuclei of CN V, VI, VII, VIII
- Medulla—nuclei of CN IX, X, XII
- Spinal cord—nucleus of CN XI

Lateral nuclei = sensory (a**L**ar plate).

—**S**ulcus limitans—

Medial nuclei = **M**otor (basal plate).

Cranial nerve and vessel pathways



Divisions of CN V exit owing to **S**tanding **R**oom **O**nly



Cranial nerves

NERVE	CN	FUNCTION	TYPE	MNEMONIC
Olfactory	I	Smell (only CN without thalamic relay to cortex)	Sensory	Some
Optic	II	Sight	Sensory	Say
Oculomotor	III	Eye movement (SR, IR, MR, IO), pupillary constriction (sphincter pupillae: Edinger-Westphal nucleus, muscarinic receptors), accommodation, eyelid opening (levator palpebrae)	Motor	Marry
Trochlear	IV	Eye movement (SO)	Motor	Money
Trigeminal	V	Mastication, facial sensation (ophthalmic, maxillary, mandibular divisions), somatosensation from anterior 2/3 of tongue, dampening of loud noises (tensor tympani)	Both	But
Abducens	VI	Eye movement (LR)	Motor	My
Facial	VII	Facial movement, taste from anterior 2/3 of tongue (chorda tympani), lacrimation, salivation (submandibular and sublingual glands are innervated by CN seven), eye closing (orbicularis oculi), auditory volume modulation (stapedius)	Both	Brother
Vestibulocochlear	VIII	Hearing, balance	Sensory	Says
Glossopharyngeal	IX	Taste and sensation from posterior 1/3 of tongue, swallowing, salivation (parotid gland), monitoring carotid body and sinus chemo- and baroreceptors, and elevation of pharynx/larynx (stylopharyngeus)	Both	Big
Vagus	X	Taste from supraglottic region, swallowing, soft palate elevation, midline uvula, talking, cough reflex, parasympathetics to thoracoabdominal viscera, monitoring aortic arch chemo- and baroreceptors	Both	Brains
Accessory	XI	Head turning, shoulder shrugging (SCM, trapezius)	Motor	Matter
Hypoglossal	XII	Tongue movement	Motor	Most

Vagal nuclei

NUCLEUS	FUNCTION	CRANIAL NERVES
Nucleus tractus Solitarius	Visceral Sensory information (eg, taste, baroreceptors, gut distention)	VII, IX, X
Nucleus aMbiguus	Motor innervation of pharynx, larynx, upper esophagus (eg, swallowing, palate elevation)	IX, X, XI (cranial portion)
Dorsal motor nucleus	Sends autonomic (parasympathetic) fibers to heart, lungs, upper GI	X

Cranial nerve reflexes

REFLEX	AFFERENT	EFFERENT
Corneal	V ₁ ophthalmic (nasociliary branch)	Bilateral VII (temporal branch—orbicularis oculi)
Lacrimation	V ₁ (loss of reflex does not preclude emotional tears)	VII
Jaw jerk	V ₃ (sensory—muscle spindle from masseter)	V ₃ (motor—masseter)
Pupillary	II	III
Gag	IX	X
Cough	X	X

Mastication muscles

3 muscles close jaw: **M**asseter, **tM**poralis, **M**edial pterygoid. 1 opens: **L**ateral pterygoid. All are innervated by trigeminal nerve (V₃).

M's Munch.

Lateral **L**owers (when speaking of pterygoids with respect to jaw motion).

“It takes more muscle to keep your mouth shut.”

Spinal nerves

There are 31 pairs of spinal nerves: 8 cervical, 12 thoracic, 5 lumbar, 5 sacral, 1 coccygeal.

Nerves C1–C7 exit above the corresponding vertebrae (eg, C3 exits above the 3rd cervical vertebra).

C8 spinal nerve exits below C7 and above T1. All other nerves exit below (eg, L2 exits below the 2nd lumbar vertebra).


Spinal cord—lower extent

In adults, spinal cord ends at lower border of L1–L2 vertebrae. **S**ubarachnoid **S**pace (which contains the CSF) extends to lower border of **S2** vertebra. Lumbar puncture is usually performed between L3–L4 or L4–L5 (level of cauda equina).

Goal of lumbar puncture is to obtain sample of CSF without damaging spinal cord. To **keep** the cord **alive**, keep the spinal needle between **L3** and **L5**.


Needle passes through:

- ① skin
- ② fascia and fat
- ③ supraspinous ligament
- ④ interspinous ligament
- ⑤ ligamentum flavum
- ⑥ epidural space
(epidural anesthesia needle stops here)
- ⑦ dura mater
- ⑧ arachnoid mater
- ⑨ subarachnoid space
(CSF collection occurs here)




Spinal cord and associated tracts

Legs (**L**umbosacral) are **L**ateral in **L**ateral corticospinal, spinothalamic tracts. Thoracic spinal cord section in **A**.
Dorsal columns are organized as you are, with hands at sides. “Arms outside, legs inside.”



Spinal tract anatomy and functions Ascending tracts synapse and then cross.

TRACT	FUNCTION	1ST-ORDER NEURON	SYNAPSE 1	2ND-ORDER NEURON	SYNAPSE 2 + PROJECTIONS
Ascending tracts					
Dorsal column	Pressure, vibration, fine touch, proprioception	Sensory nerve ending → bypasses pseudounipolar cell body in dorsal root ganglion → enters spinal cord → ascends ipsilaterally in dorsal columns	Nucleus gracilis, nucleus cuneatus (ipsilateral medulla)	Decussates in medulla → ascends contralaterally as the medial lemniscus	VPL (thalamus) → sensory cortex
Spinothalamic tract	Lateral: pain, temperature Anterior: crude touch, pressure	Sensory nerve ending (A δ and C fibers) → bypasses pseudounipolar cell body in dorsal root ganglion → enters spinal cord	Ipsilateral gray matter (spinal cord)	Decussates in spinal cord as the anterior white commissure → ascends contralaterally	
Descending tract					
Lateral corticospinal tract	Voluntary movement of contralateral limbs	UMN: cell body in 1° motor cortex → descends ipsilaterally (through posterior limb of internal capsule and cerebral peduncle), most fibers decussate at caudal medulla (pyramidal decussation) → descends contralaterally	Cell body of anterior horn (spinal cord)	LMN: leaves spinal cord	NMJ → muscle fibers

Clinical reflexes

Reflexes count up in order (main nerve root in bold):

Achilles reflex = S1, S2 (“buckle my shoe”)

Patellar reflex = L2-L4 (“kick the door”)

Biceps and brachioradialis reflexes = C5, C6
 (“pick up sticks”)

Triceps reflex = C6, C7, C8 (“lay them straight”)

Additional reflexes:

Cremasteric reflex = L1, L2 (“testicles move”)

Anal wink reflex = S3, S4 (“winks galore”)


Primitive reflexes

CNS reflexes that are present in a healthy infant, but are absent in a neurologically intact adult. Normally disappear within 1st year of life. These primitive reflexes are inhibited by a mature/developing frontal lobe. They may reemerge in adults following frontal lobe lesions → loss of inhibition of these reflexes.

Moro reflex	“Hang on for life” reflex—abduct/extend arms when startled, and then draw together
Rooting reflex	Movement of head toward one side if cheek or mouth is stroked (nipple seeking)
Sucking reflex	Sucking response when roof of mouth is touched
Palmar reflex	Curling of fingers if palm is stroked
Plantar reflex	Dorsiflexion of large toe and fanning of other toes with plantar stimulation Babinski sign—presence of this reflex in an adult, which may signify a UMN lesion
Galant reflex	Stroking along one side of the spine while newborn is in ventral suspension (face down) causes lateral flexion of lower body toward stimulated side

Landmark dermatomes

DERMATOME	CHARACTERISTICS
C2	Posterior half of skull
C3	High turtleneck shirt Diaphragm and gallbladder pain referred to the right shoulder via phrenic nerve C3, 4, 5 keeps the diaphragm alive
C4	Low-collar shirt
C6	Includes thumbs Thumbs up sign on left hand looks like a 6
T4	At the nipple T4 at the teat pore
T7	At the xiphoid process 7 letters in xiphoid
T10	At the umbilicus (belly butten) Point of referred pain in early appendicitis
L1	At the Inguinal Ligament
L4	Includes the kneecaps Down on ALL 4 's
S2, S3, S4	Sensation of penile and anal zones S2, 3, 4 keep the penis off the floor



▶ NEUROLOGY—PATHOLOGY


Common brain lesions

AREA OF LESION	CONSEQUENCE	EXAMPLES/COMMENTS
Frontal lobe	Disinhibition and deficits in concentration, orientation, judgment; may have reemergence of primitive reflexes	
Frontal eye fields	Destructive lesions (eg, MCA stroke): eyes look toward brain lesion (ie, away from side of hemiplegia)	
Paramedian pontine reticular formation	Eyes look away from brain lesion (ie, toward side of hemiplegia)	
Medial longitudinal fasciculus	Internuclear ophthalmoplegia (impaired adduction of ipsilateral eye; nystagmus of contralateral eye with abduction)	Multiple sclerosis
Dominant parietal cortex	Agraphia, acalculia, finger agnosia, left-right disorientation	Gerstmann syndrome
Nondominant parietal cortex	Agnosia of the contralateral side of the world	Hemispatial neglect syndrome
Hippocampus (bilateral)	Anterograde amnesia—inability to make new memories	
Basal ganglia	May result in tremor at rest, chorea, athetosis	Parkinson disease, Huntington disease, Wilson disease
Subthalamic nucleus	Contralateral hemiballismus	
Mammillary bodies (bilateral)	Wernicke-Korsakoff syndrome —Confusion, Ataxia, Nystagmus, Ophthalmoplegia, memory loss (anterograde and retrograde amnesia), confabulation, personality changes	Wernicke problems come in a CAN O' beer and other conditions associated with thiamine deficiency
Amygdala (bilateral)	Klüver-Bucy syndrome —disinhibited behavior (eg, hyperphagia, hypersexuality, hyperorality)	HSV-1 encephalitis
Dorsal midbrain	Parinaud syndrome —vertical gaze palsy, pupillary light-near dissociation, lid retraction, convergence-retraction nystagmus	Stroke, hydrocephalus, pinealoma
Reticular activating system (midbrain)	Reduced levels of arousal and wakefulness	Coma
Cerebellar hemisphere	Intention tremor, limb ataxia, loss of balance; damage to cerebellum → ipsilateral deficits; fall toward side of lesion	Cerebellar hemispheres are laterally located— affect lateral limbs
Cerebellar vermis	Truncal ataxia (wide-based, “drunken sailor” gait), nystagmus	Vermis is centrally located—affects central body Degeneration associated with chronic alcohol use
Red nucleus (midbrain)	Decorticate (flexor) posturing—lesion above red nucleus, presents with flexion of upper extremities and extension of lower extremities Decerebrate (extensor) posturing—lesion at or below red nucleus, presents with extension of upper and lower extremities	Worse prognosis with decerebrate posturing In decorticate posturing, your hands are near the cor (heart)

Ischemic brain disease/stroke

Irreversible neuronal injury begins after 5 minutes of hypoxia. Most **vulnerable: hippocampus, neocortex, cerebellum (Purkinje cells), watershed areas** (“**vulnerable hippos need pure water**”). Stroke imaging: noncontrast CT to exclude hemorrhage (before tPA can be given). CT detects ischemic changes in 6–24 hr. Diffusion-weighted MRI can detect ischemia within 3–30 min.

TIME SINCE ISCHEMIC EVENT	12–24 HOURS	24–72 HOURS	3–5 DAYS	1–2 WEEKS	> 2 WEEKS
Histologic features	Eosinophilic cytoplasm + pyknotic nuclei (red neurons)	Necrosis + neutrophils	Macrophages (microglia)	Reactive gliosis (astrocytes) + vascular proliferation	Glial scar

Ischemic stroke

Acute blockage of vessels → disruption of blood flow and subsequent ischemia → infarction → liquefactive necrosis.


3 types:

- Thrombotic—due to a clot forming directly at site of infarction (commonly the MCA **A**), usually over a ruptured atherosclerotic plaque.
- Embolic—embolus from another part of the body obstructs vessel. Can affect multiple vascular territories. Examples: atrial fibrillation, carotid artery stenosis, DVT with patent foramen ovale, infective endocarditis.
- Hypoxic—due to hypoperfusion or hypoxemia. Common during cardiovascular surgeries, tends to affect watershed areas.

Treatment: tPA (if within 3–4.5 hr of onset and no hemorrhage/risk of hemorrhage) and/or thrombectomy (if large artery occlusion). Reduce risk with medical therapy (eg, aspirin, clopidogrel); optimum control of blood pressure, blood sugars, lipids; smoking cessation; and treat conditions that ↑ risk (eg, atrial fibrillation, carotid artery stenosis).

Transient ischemic attack

Brief, reversible episode of focal neurologic dysfunction without acute infarction (⊖ MRI), with the majority resolving in < 15 minutes; ischemia (eg, embolus, small vessel stenosis).

Neonatal intraventricular hemorrhage


Bleeding into ventricles (arrow in coronal transcranial ultrasound **A** shows blood in right intraventricular space, extending into periventricular white matter). Increased risk in premature and low-birth-weight infants. Originates in germinal matrix, a highly vascularized layer within the subventricular zone. Due to reduced glial fiber support and impaired autoregulation of BP in premature infants. Can present with altered level of consciousness, bulging fontanelle, hypotension, seizures, coma.

Intracranial hemorrhage**Epidural hematoma**

Rupture of middle meningeal artery (branch of maxillary artery), often 2° to skull fracture (circle in **A**) involving the pterion (thinnest area of the lateral skull). Might present with transient loss of consciousness → recovery (“lucid interval”) → rapid deterioration due to hematoma expansion.


Scalp hematoma (arrows in **A**) and rapid intracranial expansion (arrows in **B**) under systemic arterial pressure → transtentorial herniation, CN III palsy.

CT shows biconvex (lentiform), hyperdense blood collection **B** not crossing suture lines.

**Subdural hematoma**

Rupture of bridging veins. Can be acute (traumatic, high-energy impact → hyperdense on CT) or chronic (associated with mild trauma, cerebral atrophy, elderly, alcoholism → hypodense on CT). Also seen in shaken babies. Predisposing factors: brain atrophy, trauma.

Crescent-shaped hemorrhage (red arrows in **C** and **D**) that **crosses suture lines**. Can cause midline shift (yellow arrow in **C**), findings of “acute on chronic” hemorrhage (blue arrows in **D**).

**Subarachnoid hemorrhage**


Bleeding **E F** due to trauma, or rupture of an aneurysm (such as a saccular aneurysm **E**) or arteriovenous malformation. Rapid time course. Patients complain of “worst headache of my life.” Bloody or yellow (xanthochromic) lumbar puncture.

Vasospasm can occur due to blood breakdown or rebleed 3–10 days after hemorrhage → ischemic infarct; nimodipine used to prevent/reduce vasospasm. ↑ risk of developing communicating and/or obstructive hydrocephalus.

**Intraparenchymal hemorrhage**

Most commonly caused by systemic hypertension. Also seen with amyloid angiopathy (recurrent lobar hemorrhagic stroke in elderly), vasculitis, neoplasm. May be 2° to reperfusion injury in ischemic stroke.

Hypertensive hemorrhages (Charcot-Bouchard microaneurysm) most often occur in putamen of basal ganglia (lenticulostriate vessels **G**), followed by thalamus, pons, and cerebellum **H**.




Effects of strokes


ARTERY	AREA OF LESION	SYMPTOMS	NOTES
Anterior circulation			
Middle cerebral artery	Motor and sensory cortices A —upper limb and face. Temporal lobe (Wernicke area); frontal lobe (Broca area).	Contralateral paralysis and sensory loss—face and upper limb. Aphasia if in dominant (usually left) hemisphere. Hemineglect if lesion affects nondominant (usually right) hemisphere.	Wernicke aphasia is associated with right superior quadrant visual field defect due to temporal lobe involvement.
Anterior cerebral artery	Motor and sensory cortices—lower limb.	Contralateral paralysis and sensory loss—lower limb, urinary incontinence.	
Lenticulo-striate artery	Striatum, internal capsule.	Contralateral paralysis. Absence of cortical signs (eg, neglect, aphasia, visual field loss).	Pure motor stroke. Common location of lacunar infarcts B , due to hyaline arteriosclerosis (lipohyalinosis) 2° to unmanaged hypertension.
Posterior circulation			
Anterior spinal artery	Corticospinal tract. Medial lemniscus. Caudal medulla—hypoglossal nerve.	Contralateral paralysis—upper and lower limbs. ↓ contralateral proprioception. Ipsilateral hypoglossal dysfunction (tongue deviates ipsilaterally).	Medial medullary syndrome —caused by infarct of paramedian branches of ASA and/or vertebral arteries.
Posterior inferior cerebellar artery	Lateral medulla: Nucleus ambiguus (CN IX, X, XI) Vestibular nuclei Lateral spinothalamic tract, spinal trigeminal nucleus Sympathetic fibers Inferior cerebellar peduncle	Dysphagia, hoarseness, ↓ gag reflex, hiccups. Vomiting, vertigo, nystagmus ↓ pain and temperature sensation from contralateral body, ipsilateral face. Ipsilateral Horner syndrome. Ipsilateral ataxia, dysmetria.	Lateral medullary (Wallenberg) syndrome. Nucleus ambiguus effects are specific to PICA lesions C . “Don’t pick a (PICA) horse (hoarseness) that can’t eat (dysphagia).” Also supplies inferior cerebellar peduncle (part of cerebellum).
Anterior inferior cerebellar artery	Lateral pons: Facial nucleus Vestibular nuclei Spinothalamic tract, spinal trigeminal nucleus Sympathetic fibers Middle and inferior cerebellar peduncles Labyrinthine artery	Paralysis of face (LMN lesion vs UMN lesion in cortical stroke), ↓ lacrimation, ↓ salivation, ↓ taste from anterior 2/3 of tongue. Vomiting, vertigo, nystagmus ↓ pain and temperature sensation from contralateral body, ipsilateral face. Ipsilateral Horner syndrome. Ipsilateral ataxia, dysmetria. Ipsilateral sensorineural deafness, vertigo.	Lateral pontine syndrome. Facial nucleus effects are specific to AICA lesions. “ Facial droop means AICA’s pooped .” Also supplies middle and inferior cerebellar peduncles (part of cerebellum).

Effects of strokes (continued)

ARTERY	AREA OF LESION	SYMPTOMS	NOTES
Basilar artery	Pons, medulla, lower midbrain. Corticospinal and corticobulbar tracts. Ocular cranial nerve nuclei, paramedian pontine reticular formation.	If RAS spared, consciousness is preserved. Quadriplegia; loss of voluntary facial, mouth, and tongue movements. Loss of horizontal, but not vertical, eye movements.	Locked-in syndrome (locked in the basement).
Posterior cerebral artery	Occipital lobe D .	Contralateral hemianopia with macular sparing; alexia without agraphia (dominant hemisphere).	

**Central poststroke pain syndrome**

Neuropathic pain due to thalamic lesions. Initial paresthesias followed in weeks to months by allodynia (ordinarily painless stimuli cause pain) and dysesthesia (altered sensation) on the contralateral side. Occurs in 10% of stroke patients.


Diffuse axonal injury

Caused by traumatic shearing forces during rapid acceleration and/or deceleration of the brain (eg, motor vehicle accident). Usually results in devastating neurologic injury, often causing coma or persistent vegetative state. MRI **A** shows multiple lesions (punctate hemorrhages) involving the white matter tracts.

Aphasia

Aphasia—higher-order language deficit (inability to understand/produce/use language appropriately); caused by pathology in dominant cerebral hemisphere (usually left).


Dysarthria—motor inability to produce speech (movement deficit).



TYPE	COMMENTS
Broca (expressive)	Broca area in inferior frontal gyrus of frontal lobe. Patient appears frustrated, insight intact. B roca = B roken B oca (<i>boca</i> = mouth in Spanish).
Wernicke (receptive)	Wernicke area in superior temporal gyrus of temporal lobe. Patients do not have insight. W ernicke is a W ord salad and makes no sense.
Conduction	Can be caused by damage to ar C uate fasciculus.
Global	Broca and Wernicke areas affected.
Transcortical motor	Affects frontal lobe around Broca area, but Broca area is spared.
Transcortical sensory	Affects temporal lobe around Wernicke area, but Wernicke area is spared.
Transcortical mixed	Broca and Wernicke areas and arcuate fasciculus remain intact; surrounding watershed areas affected.

Aneurysms

Abnormal dilation of an artery due to weakening of vessel wall.

Saccular aneurysm

Also called berry aneurysm **A**. Occurs at bifurcations in the circle of Willis. Most common site is junction of ACom and ACA. Associated with ADPKD, Ehlers-Danlos syndrome. Other risk factors: advanced age, hypertension, smoking, race (↑ risk in African-Americans).

Usually clinically silent until rupture (most common complication) → subarachnoid hemorrhage (“worst headache of my life” or “thunderclap headache”) → focal neurologic deficits. Can also cause symptoms via direct compression of surrounding structures by growing aneurysm.

- ACom—compression → bitemporal hemianopia (compression of optic chiasm); visual acuity deficits; rupture → ischemia in ACA distribution → contralateral lower extremity hemiparesis, sensory deficits.
- MCA—rupture → ischemia in MCA distribution → contralateral upper extremity and lower facial hemiparesis, sensory deficits.
- PCom—compression → ipsilateral CN III palsy → mydriasis (“blown pupil”); may also see ptosis, “down and out” eye.

Charcot-Bouchard microaneurysm

Common, associated with chronic hypertension; affects small vessels (eg, lenticulostriate arteries in basal ganglia, thalamus) and can cause hemorrhagic intraparenchymal strokes. Not visible on angiography.

Seizures

Characterized by synchronized, high-frequency neuronal firing. Variety of forms.

Partial (focal) seizures

Affect single area of the brain. Most commonly originate in medial temporal lobe. Types:

- **Simple partial** (consciousness intact)—motor, sensory, autonomic, psychic
- **Complex partial** (impaired consciousness, automatisms)

Generalized seizures

Diffuse. Types:


- **Absence** (petit mal)—3 Hz spike-and-wave discharges, no postictal confusion, blank stare
- **Myoclonic**—quick, repetitive jerks
- **Tonic-clonic** (grand mal)—alternating stiffening and movement, postictal confusion, urinary incontinence, tongue biting
- **Tonic**—stiffening
- **Atonic**—“drop” seizures (falls to floor); commonly mistaken for fainting

Epilepsy—disorder of recurrent, unprovoked seizures (febrile seizures are not epilepsy).

Status epilepticus—continuous (≥ 5 min) or recurring seizures that may result in brain injury.

Causes of seizures by age:

- Children—genetic, infection (febrile), trauma, congenital, metabolic
- Adults—tumor, trauma, stroke, infection
- Elderly—stroke, tumor, trauma, metabolic, infection






Fever vs heat stroke

	Fever	Heat stroke
PATHOPHYSIOLOGY	Cytokine activation during inflammation (eg, infection)	Inability of body to dissipate heat (eg, exertion)
TEMPERATURE	Usually < 40 °C	Usually > 40 °C
COMPLICATIONS	Febrile seizure (benign, usually self-limiting)	CNS dysfunction (eg, confusion), end-organ damage, acute respiratory distress syndrome, rhabdomyolysis
MANAGEMENT	Acetaminophen or ibuprofen for comfort (does not prevent future febrile seizures), antibiotic therapy if indicated	Rapid external cooling, rehydration and electrolyte correction

Headaches

Pain due to irritation of structures such as the dura, cranial nerves, or extracranial structures. More common in females, except cluster headaches.

CLASSIFICATION	LOCALIZATION	DURATION	DESCRIPTION	TREATMENT
Cluster^a 	Unilateral	15 min–3 hr; repetitive	Excruciating periorbital pain (“suicide headache”) with lacrimation and rhinorrhea. May present with Horner syndrome. More common in males.	Acute: sumatriptan, 100% O ₂ . Prophylaxis: verapamil.
Migraine 	Unilateral	4–72 hr	Pulsating pain with nausea, photophobia, or phonophobia. May have “aura.” Due to irritation of CN V, meninges, or blood vessels (release of vasoactive neuropeptides [eg, substance P, calcitonin gene-related peptide]).	Acute: NSAIDs, triptans, dihydroergotamine. Prophylaxis: lifestyle changes (eg, sleep, exercise, diet), β-blockers, amitriptyline, topiramate, valproate, botulinum toxin, anti-CGRP monoclonal antibodies. POUND —Pulsatile, One-day duration, Unilateral, Nausea, Disabling.
Tension 	Bilateral	> 30 min (typically 4–6 hr); constant	Steady, “band-like” pain. No photophobia or phonophobia. No aura.	Acute: analgesics, NSAIDs, acetaminophen. Prophylaxis: TCAs (eg, amitriptyline), behavioral therapy.

Other causes of headache include subarachnoid hemorrhage (“worst headache of my life”), meningitis, hydrocephalus, neoplasia, giant cell (temporal) arteritis.

^aCompare with **trigeminal neuralgia**, which produces repetitive, unilateral, shooting/shock-like pain in the distribution of CN V. Triggered by chewing, talking, touching certain parts of the face. Lasts (typically) for seconds to minutes, but episodes often increase in intensity and frequency over time. First-line therapy: carbamazepine.

Movement disorders

DISORDER	PRESENTATION	CHARACTERISTIC LESION	NOTES
Akathisia	Restlessness and intense urge to move.		Can be seen with neuroleptic use or as a side effect of Parkinson treatment.
Asterixis	Extension of wrists causes “flapping” motion.		Associated with hepatic encephalopathy, Wilson disease, and other metabolic derangements.
Athetosis	Slow, snake-like, writhing movements; especially seen in the fingers.	Basal ganglia.	Seen in Huntington disease.
Chorea	Sudden, jerky, purposeless movements.	Basal ganglia.	<i>Chorea</i> = dancing. Seen in Huntington disease and in acute rheumatic fever (Sydenham chorea).
Dystonia	Sustained, involuntary muscle contractions.		Writer’s cramp, blepharospasm, torticollis. Treatment: botulinum toxin injection.
Essential tremor	High-frequency tremor with sustained posture (eg, outstretched arms), worsened with movement or when anxious.		Often familial. Patients often self-medicate with alcohol, which ↓ tremor amplitude. Treatment: nonselective β-blockers (eg, propranolol), primidone.
Intention tremor	Slow, zigzag motion when pointing/extending toward a target.	Cerebellar dysfunction.	
Resting tremor	Uncontrolled movement of distal appendages (most noticeable in hands); tremor alleviated by intentional movement.	Substantia nigra (P arkinson disease).	Occurs at rest; “pill-rolling tremor” of Parkinson disease. When you park your car, it is at rest .
Hemiballismus	Sudden, wild flailing of one side of the body.	Contralateral subthalamic nucleus (eg, lacunar stroke).	Pronounce “ H alf-of-body b allistic.”
Myoclonus	Sudden, brief, uncontrolled muscle contraction.		Jerks; hiccups; common in metabolic abnormalities such as renal and liver failure.
Restless legs syndrome	Worse at rest/nighttime. Relieved by movement.		Associated with iron deficiency, CKD. Treatment: dopamine agonists (pramipexole, ropinirole).


Neurodegenerative disorders

↓ in cognitive ability, memory, or function with intact consciousness.
Must rule out depression as cause of dementia (called pseudodementia). Other reversible causes of dementia: hypothyroidism, vitamin B₁₂ deficiency, neurosyphilis, normal pressure hydrocephalus.

DISEASE	DESCRIPTION	HISTOLOGIC/GROSS FINDINGS
Parkinson disease	<p>Parkinson TRAPSS your body:</p> <ul style="list-style-type: none"> Tremor (pill-rolling tremor at rest) Rigidity (cogwheel) Akinesia (or bradykinesia) Postural instability Shuffling gait Small handwriting (micrographia) <p>MPTP, a contaminant in illegal drugs, is metabolized to MPP⁺, which is toxic to substantia nigra.</p>	<p>Loss of dopaminergic neurons (ie, depigmentation) of substantia nigra pars compacta.</p> <p>Lewy bodies: composed of α-synuclein (intracellular eosinophilic inclusions A).</p>
Huntington disease	<p>Autosomal dominant trinucleotide (CAG)_n repeat expansion in the huntingtin (HTT) gene on chromosome 4 (4 letters). Symptoms manifest between ages 20 and 50: chorea, athetosis, aggression, depression, dementia (sometimes initially mistaken for substance abuse).</p> <p>Anticipation results from expansion of CAG repeats. Caudate loses ACh and GABA.</p>	<p>Atrophy of caudate and putamen with ex vacuo ventriculomegaly.</p> <p>↑ dopamine, ↓ GABA, ↓ ACh in brain. Neuronal death via NMDA-R binding and glutamate excitotoxicity.</p>
Alzheimer disease	<p>Most common cause of dementia in elderly. Down syndrome patients have ↑ risk of developing Alzheimer disease, as APP is located on chromosome 21.</p> <p>↓ ACh.</p> <p>Associated with the following altered proteins:</p> <ul style="list-style-type: none"> ▪ ApoE-2: ↓ risk of sporadic form ▪ ApoE-4: ↑ risk of sporadic form ▪ APP, presenilin-1, presenilin-2: familial forms (10%) with earlier onset 	<p>Widespread cortical atrophy (normal cortex B; cortex in Alzheimer disease C), especially hippocampus (arrows in B and C). Narrowing of gyri and widening of sulci.</p> <p>Senile plaques D in gray matter: extracellular β-amyloid core; may cause amyloid angiopathy → intracranial hemorrhage; Aβ (amyloid-β) synthesized by cleaving amyloid precursor protein (APP).</p> <p>Neurofibrillary tangles E: intracellular, hyperphosphorylated tau protein = insoluble cytoskeletal elements; number of tangles correlates with degree of dementia.</p> <p>Hirano bodies—intracellular eosinophilic proteinaceous rods in hippocampus.</p>
Frontotemporal dementia	<p>Formerly called Pick disease. Early changes in personality and behavior (behavioral variant), or aphasia (primary progressive aphasia). May have associated movement disorders.</p>	<p>Frontotemporal lobe degeneration F.</p> <p>Inclusions of hyperphosphorylated tau (round Pick bodies G) or ubiquitinated TDP-43.</p>

Neurodegenerative disorders (continued)

DISEASE	DESCRIPTION	HISTOLOGIC/GROSS FINDINGS
Lewy body dementia	Visual hallucinations (“haLewycinations”), dementia with fluctuating cognition/alertness, REM sleep behavior disorder, and parkinsonism. Called Lewy body dementia if cognitive and motor symptom onset < 1 year apart, otherwise considered dementia 2° to Parkinson disease.	Intracellular Lewy bodies A primarily in cortex.
Vascular dementia	Result of multiple arterial infarcts and/or chronic ischemia. Step-wise decline in cognitive ability with late-onset memory impairment. 2nd most common cause of dementia in elderly.	MRI or CT shows multiple cortical and/or subcortical infarcts.
Creutzfeldt-Jakob disease	Rapidly progressive (weeks to months) dementia with myoclonus (“startle myoclonus”) and ataxia. Commonly see periodic sharp waves on EEG and ↑ 14-3-3 protein in CSF.	Spongiform cortex (vacuolization without inflammation). Prions (PrP ^c → PrP ^{sc} sheet [β-pleated sheet resistant to proteases]) H .

**Idiopathic intracranial hypertension**

Also called pseudotumor cerebri. ↑ ICP with no obvious findings on imaging. Risk factors include **female** sex, **T**etracyclines, **O**besity, vitamin **A** excess, **D**anazol (**f**emale **T**OAD). Associated with cerebral venous sinus stenosis. Findings: headache, tinnitus, diplopia (usually from CN VI palsy), no change in mental status. Impaired optic nerve axoplasmic flow → papilledema. Visual field testing shows enlarged blind spot and peripheral constriction. Lumbar puncture reveals ↑ opening pressure and provides temporary headache relief.

Treatment: weight loss, acetazolamide, invasive procedures for refractory cases (eg, CSF shunt placement, optic nerve sheath fenestration surgery for visual loss).

Hydrocephalus	↑ CSF volume → ventricular dilation +/- ↑ ICP.
Communicating	
Communicating hydrocephalus	↓ CSF absorption by arachnoid granulations (eg, arachnoid scarring post-meningitis) → ↑ ICP, papilledema, herniation.
Normal pressure hydrocephalus	Affects the elderly; idiopathic; CSF pressure elevated only episodically; does not result in increased subarachnoid space volume. Expansion of ventricles A distorts the fibers of the corona radiata → triad of urinary incontinence, gait apraxia (magnetic gait), and cognitive dysfunction . “ Wet, wobbly, and wacky. ” Symptoms potentially reversible with CSF drainage via lumbar puncture or shunt placement.
Noncommunicating (obstructive)	
Noncommunicating hydrocephalus	Caused by structural blockage of CSF circulation within ventricular system (eg, stenosis of aqueduct of Sylvius, colloid cyst blocking foramen of Monro, tumor B).
Hydrocephalus mimics	
Ex vacuo ventriculomegaly	Appearance of ↑ CSF on imaging C , but is actually due to ↓ brain tissue and neuronal atrophy (eg, Alzheimer disease, advanced HIV, Pick disease, Huntington disease). ICP is normal; NPH triad is not seen.




Multiple sclerosis

Autoimmune inflammation and demyelination of CNS (brain and spinal cord) with subsequent axonal damage. Can present with:

- Acute optic neuritis (painful unilateral visual loss associated with Marcus Gunn pupil)
- Brain stem/cerebellar syndromes (eg, diplopia, ataxia, scanning speech, intention tremor, nystagmus/INO [bilateral > unilateral])
- Pyramidal tract demyelination (eg, weakness, spasticity)
- Spinal cord syndromes (eg, electric shock-like sensation along cervical spine on neck flexion, neurogenic bladder, paraparesis, sensory manifestations affecting the trunk or one or more extremity)

Symptoms may exacerbate with increased body temperature (eg, hot bath, exercise). Relapsing and remitting is most common clinical course. Most often affects women in their 20s and 30s; more common in individuals living farther from equator and with low serum vitamin D levels.

FINDINGS


↑ IgG level and myelin basic protein in CSF. Oligoclonal bands are diagnostic. MRI is gold standard. Periventricular plaques **A** (areas of oligodendrocyte loss and reactive gliosis). Multiple white matter lesions disseminated in space and time.

TREATMENT

Stop relapses and halt/slow progression with disease-modifying therapies (eg, β -interferon, glatiramer, natalizumab). Treat acute flares with IV steroids. Symptomatic treatment for neurogenic bladder (catheterization, muscarinic antagonists), spasticity (baclofen, GABA_B receptor agonists), pain (TCAs, anticonvulsants).

Other demyelinating and dysmyelinating disorders

Osmotic demyelination syndrome



Also called central pontine myelinolysis. Massive axonal demyelination in pontine white matter **A** 2° to rapid osmotic changes, most commonly iatrogenic correction of hyponatremia but also rapid shifts of other osmolytes (eg, glucose). Acute paralysis, dysarthria, dysphagia, diplopia, loss of consciousness. Can cause “locked-in syndrome.”

Correcting serum Na⁺ too fast:

- “From low to high, your pons will die” (osmotic demyelination syndrome)
- “From high to low, your brains will blow” (cerebral edema/herniation)

Acute inflammatory demyelinating polyradiculopathy

Most common subtype of **Guillain-Barré syndrome**. Autoimmune condition that destroys Schwann cells via inflammation and demyelination of motor fibers, sensory fibers, peripheral nerves (including CN III-XII). Likely facilitated by molecular mimicry and triggered by inoculations or stress. Despite association with infections (eg, *Campylobacter jejuni*, viruses [eg, Zika]), no definitive causal link to any pathogen. Results in symmetric ascending muscle weakness/paralysis and depressed/absent DTRs beginning in lower extremities. Facial paralysis (usually bilateral) and respiratory failure are common. May see autonomic dysregulation (eg, cardiac irregularities, hypertension, hypotension) or sensory abnormalities. Almost all patients survive; majority recover completely after weeks to months. ↑ CSF protein with normal cell count (albuminocytologic dissociation). Respiratory support is critical until recovery. Disease-modifying treatment: plasmapheresis or IV immunoglobulins. No role for steroids.

Acute disseminated (postinfectious) encephalomyelitis

Multifocal inflammation and demyelination after infection or vaccination. Presents with rapidly progressive multifocal neurologic symptoms, altered mental status.

Charcot-Marie-Tooth disease

Also called hereditary motor and sensory neuropathy. Group of progressive hereditary nerve disorders related to the defective production of proteins involved in the structure and function of peripheral nerves or the myelin sheath. Typically autosomal dominant and associated with foot deformities (eg, pes cavus, hammer toe), lower extremity weakness (eg, foot drop), and sensory deficits. Most common type, CMT1A, is caused by *PMP22* gene duplication.

Progressive multifocal leukoencephalopathy




Demyelination of CNS **B** due to destruction of oligodendrocytes (2° to reactivation of latent JC virus infection). Seen in 2–4% of patients with AIDS. Rapidly progressive, usually fatal. Predominantly involves parietal and occipital areas; visual symptoms are common. ↑ risk associated with natalizumab.

Other disorders

Krabbe disease, metachromatic leukodystrophy, adrenoleukodystrophy.


Neurocutaneous disorders

DISORDER	GENETICS	PRESENTATION	NOTES
Sturge-Weber syndrome	Congenital nonhereditary anomaly of neural crest derivatives. Somatic mosaicism of an activating mutation in one copy of the <i>GNAQ</i> gene.	Capillary vascular malformation → port-wine stain A (nevus flammeus or non-neoplastic birthmark) in CN V ₁ /V ₂ distribution; ipsilateral leptomeningeal angioma B → seizures/epilepsy; intellectual disability; episcleral hemangioma → ↑ IOP → early-onset glaucoma.	Also called encephalotrigeminal angiomatosis. SSTURGG E-Weber: Sporadic, port-wine Stain , T ram track calcifications (opposing gyri), U nilateral, intellectual disability (R etardation), G laucoma, GNAQ gene, E pilepsy.
Tuberous sclerosis	AD, variable expression. Mutation in tumor suppressor genes <i>TSC1</i> on chromosome 9 (hamartin), <i>TSC2</i> on chromosome 16 (tuberin).	Hamartomas in CNS and skin, A ngiofibromas C , M itral regurgitation, A sh-leaf spots D , cardiac R habdomyoma, (T uberous sclerosis), autosomal d ominant; M ental retardation (intellectual disability), renal A ngiomyolipoma E , S eizures, S hagreen patches.	HAMARTOMASS . ↑ incidence of S ubependymal giant cell astrocytomas and unguinal fibromas.
Neurofibromatosis type I	AD, 100% penetrance. Mutation in <i>NF1</i> tumor suppressor gene on chromosome 17 (encodes neurofibromin, a negative RAS regulator).	C afé-au-lait spots F , I ntellectual disability, C utaneous neurofibromas G , L isch nodules (pigmented iris hamartomas H), O ptic gliomas, P heochromocytomas, S eizures/focal neurologic S igns (often from meningioma), bone lesions (eg, sphenoid dysplasia).	Also called von Recklinghausen disease. 17 letters in “von Recklinghausen.” CICLOPSS .
Neurofibromatosis type II	AD. Mutation in <i>NF2</i> tumor suppressor gene (merlin) on chromosome 22.	Bilateral vestibular schwannomas, juvenile cataracts, meningiomas, ependymomas.	NF2 affects 2 ears, 2 eyes.
von Hippel-Lindau disease	AD. Deletion of <i>VHL</i> gene on chromosome 3p. pVHL ubiquitinates hypoxia-inducible factor 1 α .	H emangioblastomas (high vascularity with hyperchromatic nuclei I) in retina, brain stem, cerebellum, spine J ; A ngiomatosis; bilateral R enal cell carcinomas; P heochromocytomas.	Numerous tumors, benign and malignant. VHL = 3 letters. HARP .




Adult primary brain tumors

TUMOR	DESCRIPTION	HISTOLOGY
Glioblastoma multiforme	Grade IV astrocytoma. Common, highly malignant 1° brain tumor with ~ 1-year median survival. Found in cerebral hemispheres. Can cross corpus callosum (“butterfly glioma” A).	Astrocyte origin, GFAP ⊕. “Pseudopalisading” pleomorphic tumor cells B border central areas of necrosis, hemorrhage, and/or microvascular proliferation.
Oligodendroglioma	Relatively rare, slow growing. Most often in frontal lobes C . Often calcified.	Oligodendrocyte origin. “Fried egg” cells—round nuclei with clear cytoplasm D . “Chicken-wire” capillary pattern.
Meningioma	Common, typically benign. Females > males. Most often occurs near surfaces of brain and in parasagittal region. Extra-axial (external to brain parenchyma) and may have a dural attachment (“tail” E). Often asymptomatic; may present with seizures or focal neurologic signs. Resection and/or radiosurgery.	Arachnoid cell origin. Spindle cells concentrically arranged in a whorled pattern F ; psammoma bodies (laminated calcifications).
Hemangioblastoma	Most often cerebellar G . Associated with von Hippel-Lindau syndrome when found with retinal angiomas. Can produce erythropoietin → 2° polycythemia.	Blood vessel origin. Closely arranged, thin-walled capillaries with minimal intervening parenchyma H .




Adult primary brain tumors (continued)

TUMOR	DESCRIPTION	HISTOLOGY
Pituitary adenoma	<p>May be nonfunctioning (silent) or hyperfunctioning (hormone-producing). Nonfunctional tumors present with mass effect (eg, bitemporal hemianopia [due to pressure on optic chiasm I]). Pituitary apoplexy → hyper- or hypopituitarism.</p> <p>Prolactinoma classically presents as galactorrhea, amenorrhea, ↓ bone density due to suppression of estrogen in women and as ↓ libido, infertility in men.</p> <p>Treatment: dopamine agonists (eg, bromocriptine, cabergoline), transsphenoidal resection.</p>	<p>Hyperplasia of only one type of endocrine cells found in pituitary. Most commonly from lactotrophs (prolactin) J → hyperprolactinemia. Less commonly, from somatotrophs (GH) → acromegaly, gigantism; corticotrophs (ACTH) → Cushing disease. Rarely, from thyrotrophs (TSH), gonadotrophs (FSH, LH).</p>
Schwannoma	<p>Classically at the cerebellopontine angle K, benign, involving CNs V, VII, and VIII, but can be along any peripheral nerve. Often localized to CN VIII in internal acoustic meatus → vestibular schwannoma (can present as hearing loss and tinnitus). Bilateral vestibular schwannomas found in NF-2.</p> <p>Resection or stereotactic radiosurgery.</p>	<p>Schwann cell origin, S-100 ⊕. Biphasic, dense, hypercellular areas containing spindle cells alternating with hypocellular, myxoid areas L.</p>




Childhood primary brain tumors

TUMOR	DESCRIPTION	HISTOLOGY
Pilocytic astrocytoma	Low-grade astrocytoma. Most common 1° brain tumor in childhood. Usually well circumscribed. In children, most often found in posterior fossa A (eg, cerebellum). May be supratentorial. Benign; good prognosis.	Astrocyte origin, GFAP ⊕. Rosenthal fibers—eosinophilic, corkscrew fibers B . Cystic + solid (gross).
Medulloblastoma	Most common malignant brain tumor in childhood. Commonly involves cerebellum C . Can compress 4th ventricle, causing noncommunicating hydrocephalus → headaches, papilledema. Can involve the cerebellar vermis → truncal ataxia. Can send “drop metastases” to spinal cord.	Form of primitive neuroectodermal tumor (PNET). Homer-Wright rosettes, small blue cells D . Synaptophysin ⊕.
Ependymoma	Most commonly found in 4th ventricle E . Can cause hydrocephalus. Poor prognosis.	Ependymal cell origin. Characteristic perivascular pseudorosettes F . Rod-shaped blepharoplasts (basal ciliary bodies) found near the nucleus.
Craniopharyngioma	Most common childhood supratentorial tumor. May be confused with pituitary adenoma (both cause bitemporal hemianopia).	Derived from remnants of Rathke pouch (ectoderm). Calcification is common G H . Cholesterol crystals found in “motor oil”-like fluid within tumor.
Pinealoma	Tumor of pineal gland. Can cause Parinaud syndrome (compression of tectum → vertical gaze palsy); obstructive hydrocephalus (compression of cerebral aqueduct); precocious puberty in males (hCG production).	Similar to germ cell tumors (eg, testicular seminoma).



Herniation syndromes



- 1** Cingulate (subfalcine) herniation under falx cerebri Can compress anterior cerebral artery.

- 2** Central/downward transtentorial herniation Caudal displacement of brain stem → rupture of paramedian basilar artery branches → Duret hemorrhages. Usually fatal.








- 3** Uncal transtentorial herniation Uncus = medial temporal lobe. Early herniation → ipsilateral blown pupil (unilateral CN III compression), contralateral hemiparesis. Late herniation → coma, Kernohan phenomenon (misleading contralateral blown pupil and ipsilateral hemiparesis due to contralateral compression against Kernohan notch).

- 4** Cerebellar tonsillar herniation into the foramen magnum Coma and death result when these herniations compress the brain stem.

Motor neuron signs

SIGN	UMN LESION	LMN LESION	COMMENTS
Weakness	+	+	Lower motor neuron = everything lowered (less muscle mass, ↓ muscle tone, ↓ reflexes, downgoing toes)
Atrophy	–	+	
Fasciculations	–	+	Upper motor neuron = everything up (tone, DTRs, toes)
Reflexes	↑	↓	
Tone	↑	↓	Fasciculations = muscle twitching Positive Babinski is normal in infants
Babinski	+	–	
Spastic paresis	+	–	
Flaccid paralysis	–	+	
Clasp knife spasticity	+	–	


Spinal lesions

AREA AFFECTED	DISEASE	CHARACTERISTICS
	Spinal muscular atrophy	Congenital degeneration of anterior horns of spinal cord. LMN symptoms only, symmetric weakness. “ F loppy baby” with marked hypotonia (F laccid paralysis) and tongue F asciculations. Autosomal recessive mutation in <i>SMN1</i> → defective snRNP assembly. SMA type 1 is called W erdnig- H offmann disease.
	Amyotrophic lateral sclerosis	Also called L ou Gehrig disease. Combined UMN (corticobulbar/corticospinal) and LMN (medullary and spinal cord) degeneration. No sensory or bowel/bladder deficits. Can be caused by defect in superoxide dismutase 1. LMN deficits: flaccid limb weakness, fasciculations, atrophy, bulbar palsy (dysarthria, dysphagia, tongue atrophy). UMN deficits: spastic limb weakness, hyperreflexia, clonus, pseudobulbar palsy (dysarthria, dysphagia, emotional lability). Fatal. Treatment: “ri L ouzole”.
	Complete occlusion of anterior spinal artery	Sparses dorsal columns and Lissauer tract; mid-thoracic ASA territory is watershed area, as artery of Adamkiewicz supplies ASA below T8. Can be caused by aortic aneurysm repair. Presents with UMN deficit below the lesion (corticospinal tract), LMN deficit at the level of the lesion (anterior horn), and loss of pain and temperature sensation below the lesion (spinothalamic tract).
	Tabes dorsalis	Caused by 3° syphilis. Results from degeneration/demyelination of dorsal columns and roots → progressive sensory ataxia (impaired proprioception → poor coordination). ⊕ Romberg sign and absent DTRs. Associated with Charcot joints, shooting pain, Argyll Robertson pupils.
	Syringomyelia	Syrinx expands and damages anterior white commissure of spinothalamic tract (2nd-order neurons) → bilateral symmetric loss of pain and temperature sensation in cape-like distribution. Seen with Chiari I malformation. Can affect other tracts.
	Vitamin B₁₂ deficiency	Subacute combined degeneration (S CD)—demyelination of S pinocerebellar tracts, lateral C orticospinal tracts, and D orsal columns. Ataxic gait, paresthesia, impaired position/vibration sense, UMN symptoms.
	Cauda equina syndrome	Compression of spinal roots L2 and below, often due to intervertebral disc herniation or tumor. Radicular pain, absent knee and ankle reflexes, loss of bladder and anal sphincter control, saddle anesthesia.

Poliomyelitis

Caused by poliovirus (fecal-oral transmission). Replicates in oropharynx and small intestine before spreading via bloodstream to CNS. Infection causes destruction of cells in anterior horn of spinal cord (LMN death).


Signs of LMN lesion: asymmetric weakness (vs symmetric weakness in spinal muscular atrophy), hypotonia, flaccid paralysis, fasciculations, hyporeflexia, muscle atrophy. Respiratory muscle involvement leads to respiratory failure. Signs of infection: malaise, headache, fever, nausea, etc. CSF shows ↑ WBCs (lymphocytic pleocytosis) and slight ↑ of protein (with no change in CSF glucose). Virus recovered from stool or throat.

Brown-Séquard syndrome

Hemisection of spinal cord. Findings:

- 1 Ipsilateral loss of all sensation **at** level of lesion
- 2 Ipsilateral LMN signs (eg, flaccid paralysis) **at** level of lesion
- 3 Ipsilateral UMN signs **below** level of lesion (due to corticospinal tract damage)
- 4 Ipsilateral loss of proprioception, vibration, light (2-point discrimination) touch, and tactile sense **below** level of lesion (due to dorsal column damage)
- 5 Contralateral loss of pain, temperature, and crude (non-discriminative) touch **below** level of lesion (due to spinothalamic tract damage)

If lesion occurs above T1, patient may present with ipsilateral Horner syndrome due to damage of oculosympathetic pathway.

**Friedreich ataxia**

Autosomal recessive trinucleotide repeat disorder (GAA)_n on chromosome 9 in gene that encodes frataxin (iron-binding protein). Leads to impairment in mitochondrial functioning. Degeneration of lateral corticospinal tract (spastic paralysis), spinocerebellar tract (ataxia), dorsal columns (↓ vibratory sense, proprioception), and dorsal root ganglia (loss of DTRs). **Staggering** gait, frequent **falling**, nystagmus, dysarthria, pes cavus, hammer toes, **diabetes** mellitus, **hypertrophic cardiomyopathy** (cause of death). Presents in childhood with kyphoscoliosis **A B**.

Friedreich is **F**ratastic (**frataxin**): he's your favorite **frat** brother, always **staggering** and **falling** but has a **sweet, big heart**. Ataxic **GAA**it.



Common cranial nerve lesions


CN V motor lesion	Jaw deviates toward side of lesion due to unopposed force from the opposite pterygoid muscle.
CN X lesion	Uvula deviates away from side of lesion. Weak side collapses and uvula points away.
CN XI lesion	Weakness turning head to contralateral side of lesion (SCM). Shoulder droop on side of lesion (trapezius). The left SCM contracts to help turn the head to the right.
CN XII lesion	LMN lesion. Tongue deviates toward side of lesion (“lick your wounds”) due to weakened tongue muscles on affected side.

Facial nerve lesions



Bell palsy is the most common cause of peripheral facial palsy **A**. Usually develops after HSV reactivation. Treatment: corticosteroids +/- acyclovir. Most patients gradually recover function, but aberrant regeneration can occur. Other causes of peripheral facial palsy include Lyme disease, herpes zoster (Ramsay Hunt syndrome), sarcoidosis, tumors (eg, parotid gland), diabetes mellitus.

	Upper motor neuron lesion	Lower motor neuron lesion
LESION LOCATION	Motor cortex, connection from motor cortex to facial nucleus in pons	Facial nucleus, anywhere along CN VII
AFFECTED SIDE	Contralateral	Ipsilateral
MUSCLES INVOLVED	Lower muscles of facial expression	Upper and lower muscles of facial expression
FOREHEAD INVOLVED?	Spared, due to bilateral UMN innervation	Affected
OTHER SYMPTOMS	None	Incomplete eye closure (dry eyes, corneal ulceration), hyperacusis, loss of taste sensation to anterior tongue









▶ NEUROLOGY—OTOLOGY

Auditory physiology

Outer ear	Visible portion of ear (pinna), includes auditory canal and tympanic membrane. Transfers sound waves via vibration of tympanic membrane.
Middle ear	Air-filled space with three bones called the ossicles (malleus, incus, stapes). Ossicles conduct and amplify sound from tympanic membrane to inner ear.
Inner ear	Snail-shaped, fluid-filled cochlea. Contains basilar membrane that vibrates 2° to sound waves. Vibration transduced via specialized hair cells → auditory nerve signaling → brain stem. Each frequency leads to vibration at specific location on basilar membrane (tonotopy): <ul style="list-style-type: none"> ▪ Low frequency heard at apex near helicotrema (wide and flexible). ▪ High frequency heard best at base of cochlea (thin and rigid).

Diagnosing hearing loss


	Normal	Conductive	Sensorineural
Weber test Tuning fork on vertex of skull	 No localization	 Localizes to affected ear ↓ transmission of background noise	 Localizes to unaffected ear ↓ transmission of all sound
Rinne test Tuning fork in front of ear (air conduction, AC), Tuning fork on mastoid process (bone conduction, BC)	 AC > BC	 BC > AC	 AC > BC

Types of hearing loss

Noise-induced hearing loss	Damage to stereociliated cells in organ of Corti. Loss of high-frequency hearing first. Sudden extremely loud noises can produce hearing loss due to tympanic membrane rupture.
Presbycusis	Aging -related progressive bilateral/symmetric sensorineural hearing loss (often of higher frequencies) due to destruction of hair cells at the cochlear base (preserved low-frequency hearing at apex).

Cholesteatoma

Overgrowth of desquamated keratin debris within the middle ear space (A, arrows); may erode ossicles, mastoid air cells → conductive hearing loss. Often presents with painless otorrhea.



Vertigo

Sensation of spinning while actually stationary. Subtype of “dizziness,” but distinct from “lightheadedness.”

Peripheral vertigo



More common. Inner ear etiology (eg, semicircular canal debris, vestibular nerve infection, Ménière disease [triad: sensorineural hearing loss, vertigo, tinnitus; endolymphatic hydrops → ↑ endolymph within the inner ear], benign paroxysmal positional vertigo [BPPV]). Treatment: antihistamines, anticholinergics, antiemetics (symptomatic relief); low-salt diet +/- diuretics (Ménière disease); Epley maneuver (BPPV).

Central vertigo


Brain stem or cerebellar lesion (eg, stroke affecting vestibular nuclei, demyelinating disease, or posterior fossa tumor). Findings: directional or purely vertical nystagmus, skew deviation (vertical misalignment of the eyes), diplopia, dysmetria. Focal neurologic findings.

▶ NEUROLOGY—OPHTHALMOLOGY

Normal eye anatomy



Conjunctivitis



Inflammation of the conjunctiva → red eye **A**.

Allergic—itchy eyes, bilateral.

Bacterial—pus; treat with antibiotics.


Viral—most common, often adenovirus; sparse mucous discharge, swollen preauricular node, ↑ lacrimation; self-resolving.

Refractive errors	Common cause of impaired vision, correctable with glasses.
Hyperopia	Also called “farsightedness.” Eye too short for refractive power of cornea and lens → light focused behind retina. Correct with convex (converging) lenses.
Myopia	Also called “nearsightedness.” Eye too long for refractive power of cornea and lens → light focused in front of retina. Correct with concave (diverging) lenses.
Astigmatism	Abnormal curvature of cornea → different refractive power at different axes. Correct with cylindrical lens.

Presbyopia	Aging-related impaired accommodation (focusing on near objects), primarily due to ↓ lens elasticity, changes in lens curvature, ↓ strength of the ciliary muscle. Patients often need “reading glasses” (magnifiers).
-------------------	---

Cataract

Painless, often bilateral, opacification of lens **A**, often resulting in glare and ↓ vision, especially at night. Acquired risk factors: ↑ age, smoking, excessive alcohol use, excessive sunlight, prolonged corticosteroid use, diabetes mellitus, trauma, infection. Congenital risk factors: classic galactosemia, galactokinase deficiency, trisomies (13, 18, 21), TORCH infections (eg, rubella), Marfan syndrome, Alport syndrome, myotonic dystrophy, neurofibromatosis 2.

Aqueous humor pathway

Glaucoma

Optic disc atrophy with characteristic cupping (normal **A** versus thinning of outer rim of optic nerve head **B**), usually with elevated intraocular pressure (IOP) and progressive peripheral visual field loss if untreated. Treatment is through pharmacologic or surgical lowering of IOP.

Open-angle glaucoma



Associated with ↑ age, African-American race, family history. Painless, more common in US. Primary—cause unclear. Secondary—blocked trabecular meshwork from WBCs (eg, uveitis), RBCs (eg, vitreous hemorrhage), retinal elements (eg, retinal detachment).

Closed- or narrow-angle glaucoma

Primary—enlargement or anterior movement of lens against central iris (pupil margin) → obstruction of normal aqueous flow through pupil → fluid builds up behind iris, pushing peripheral iris against cornea **C** and impeding flow through trabecular meshwork. Secondary—hypoxia from retinal disease (eg, diabetes mellitus, vein occlusion) induces vasoproliferation in iris that contracts angle.

Chronic closure—often asymptomatic with damage to optic nerve and peripheral vision.

Acute closure—true ophthalmic emergency. ↑ IOP pushes iris forward → angle closes abruptly. Very painful, red eye **D**, sudden vision loss, halos around lights, frontal headache, fixed and mid-dilated pupil, nausea and vomiting. Mydriatic agents contraindicated.


**Uveitis**

Inflammation of uvea; specific name based on location within affected eye. Anterior uveitis: iritis; posterior uveitis: choroiditis and/or retinitis. May have hypopyon (accumulation of pus in anterior chamber **A**) or conjunctival redness. Associated with systemic inflammatory disorders (eg, sarcoidosis, rheumatoid arthritis, juvenile idiopathic arthritis, HLA-B27-associated conditions).

Age-related macular degeneration

Degeneration of macula (central area of retina). Causes distortion (metamorphopsia) and eventual loss of central vision (scotomas).

- **Dry** (nonexudative, > 80%)—**D**eposition of yellowish extracellular material (“**D**rusen”) in between Bruch membrane and retinal pigment epithelium **A** with gradual ↓ in vision. Prevent progression with multivitamin and antioxidant supplements.
- **Wet** (exudative, 10–15%)—rapid loss of vision due to bleeding 2° to choroidal neovascularization. Treat with anti-VEGF (vascular endothelial growth factor) injections (eg, bevacizumab, ranibizumab).

Diabetic retinopathy

Retinal damage due to chronic hyperglycemia. Two types:


- Nonproliferative—damaged capillaries leak blood → lipids and fluid seep into retina → hemorrhages (arrows in **A**) and macular edema. Treatment: blood sugar control.
- Proliferative—chronic hypoxia results in new blood vessel formation with resultant traction on retina → retinal detachment. Treatment: anti-VEGF injections, peripheral retinal photocoagulation, surgery.

Hypertensive retinopathy

Retinal damage due to chronic uncontrolled HTN.


Flame-shaped retinal hemorrhages, arteriovenous nicking, microaneurysms, macular star (exudate, red arrow in **A**), cotton-wool spots (blue arrow in **A**). Presence of papilledema requires immediate lowering of BP.

Associated with ↑ risk of stroke, CAD, kidney disease.

Retinal vein occlusion

Blockage of central or branch retinal vein due to compression from nearby arterial atherosclerosis.

Retinal hemorrhage and venous engorgement (“blood and thunder appearance”; arrows in **A**), edema in affected area.


Retinal detachment

Separation of neurosensory layer of retina (photoreceptor layer with rods and cones) from outermost pigmented epithelium (normally shields excess light, supports retina) → degeneration of photoreceptors → vision loss. May be 2° to retinal breaks, diabetic traction, inflammatory effusions. Visualized on fundoscopy as crinkling of retinal tissue **A** and changes in vessel direction.

Breaks more common in patients with high myopia and/or history of head trauma. Often preceded by posterior vitreous detachment (“flashes” and “floaters”) and eventual monocular loss of vision like a “curtain drawn down.” Surgical emergency.


Central retinal artery occlusion

Acute, painless monocular vision loss. Retina cloudy with attenuated vessels and “cherry-red” spot at fovea (center of macula) **A**. Evaluate for embolic source (eg, carotid artery atherosclerosis, cardiac vegetations, patent foramen ovale).




Retinitis pigmentosa

Inherited progressive retinal degeneration. Nyctalopia (night blindness) → peripheral vision loss. Bone spicule-shaped deposits **A**.



Papilledema

Optic disc swelling (usually bilateral) due to ↑ ICP (eg, 2° to mass effect). Enlarged blind spot and elevated optic disc with blurred margins **A**.



Leukocoria

Loss (whitening) of the red reflex. Important causes in children include retinoblastoma **A**, congenital cataract, toxocariasis.



Pupillary control**Miosis**

Constriction, parasympathetic:


- 1st neuron: Edinger-Westphal nucleus to ciliary ganglion via CN III
- 2nd neuron: short ciliary nerves to sphincter pupillae muscles

Short ciliary nerves **shorten** the pupil diameter.

Pupillary light reflex

Light in either retina sends a signal via CN II to pretectal nuclei (dashed lines in image) in midbrain that activates bilateral Edinger-Westphal nuclei; pupils constrict bilaterally (direct and consensual reflex).

Result: illumination of 1 eye results in bilateral pupillary constriction.

**Mydriasis**

Dilation, sympathetic:

- 1st neuron: hypothalamus to ciliospinal center of Budge (C8–T2)
- 2nd neuron: exit at T1 to superior cervical ganglion (travels along cervical sympathetic chain near lung apex, subclavian vessels)
- 3rd neuron: plexus along internal carotid, through cavernous sinus; enters orbit as long ciliary nerve to pupillary dilator muscles. Sympathetic fibers also innervate smooth muscle of eyelids (minor retractors) and sweat glands of forehead and face.

Long ciliary nerves make the pupil diameter **longer**.

Marcus Gunn pupil

Also called relative afferent pupillary defect (RAPD). When the light shines into a normal eye, constriction of the ipsilateral (direct reflex) and contralateral eye (consensual reflex) is observed. When the light is then swung to the affected eye, both pupils dilate instead of constrict due to impaired conduction of light signal along the injured optic nerve. Associated with optic neuritis, early multiple sclerosis.

Horner syndrome


Sympathetic denervation of face →:

- **P**tos (slight drooping of eyelid: superior tarsal muscle)
- **A**nhidrosis (absence of sweating) and flushing of affected side of face
- **M**iosis (pupil constriction)


Associated with lesions along the sympathetic chain:

- 1st neuron: pontine hemorrhage, lateral medullary syndrome, spinal cord lesion above T1 (eg, Brown-Séquard syndrome, late-stage syringomyelia)
- 2nd neuron: stellate ganglion compression by Pancoast tumor
- 3rd neuron: carotid dissection (painful)


PAM is **horny** (**H**orner).



Ocular motility



CN **VI** innervates the **L**ateral **R**ectus.
 CN **IV** innervates the **S**uperior **O**blique.
 CN **III** innervates the **R**est.
 The “chemical formula” **LR₆SO₄R₃**.




Obliques go **O**pposite (left SO and IO tested with patient looking right).
IOU: **IO** tested looking **U**p.

CN III, IV, VI palsies

CN III damage

CN III has both motor (central) and parasympathetic (peripheral) components. Common causes include:

- Ischemia → pupil sparing (motor fibers affected more than parasympathetic fibers)
- Uncal herniation → coma
- PCom aneurysm → sudden-onset headache
- Cavernous sinus thrombosis → proptosis, involvement of CNs IV, V₁/V₂, VI
- Midbrain stroke → contralateral hemiplegia



Motor output to extraocular muscles—affected primarily by vascular disease (eg, diabetes mellitus: glucose → sorbitol) due to ↓ diffusion of oxygen and nutrients to the interior fibers from compromised vasculature that resides on outside of nerve. Signs: ptosis, “down-and-out” gaze.


Parasympathetic output—fibers on the periphery are first affected by compression (eg, PCom aneurysm, uncal herniation). Signs: diminished or absent pupillary light reflex, “blown pupil” often with “down-and-out” gaze **A**.

Motor = **M**iddle (central)
Parasympathetic = **P**eripheral



CN IV damage

Pupil is higher in the affected eye **B**. Characteristic head tilt to contralateral/unaffected side to compensate for lack of intorsion in affected eye. Can't see the **floor** with CN **IV** damage (eg, difficulty going down stairs, reading).



CN VI damage


Affected eye unable to abduct and is displaced medially in primary position of gaze **C**.



Visual field defects

1. Right anopia (monocular vision loss)
2. Bitemporal hemianopia (pituitary lesion, chiasm)
3. Left homonymous hemianopia
4. Left upper quadrantanopia (right temporal lesion, MCA)
5. Left lower quadrantanopia (right parietal lesion, MCA)
6. Left hemianopia with macular sparing (right occipital lesion, PCA)
7. Central scotoma (eg, macular degeneration)

Meyer **L**oop—**L**ower retina; **L**oops around inferior horn of **L**ateral ventricle.
 Dorsal optic radiation—superior retina; takes shortest path via internal capsule.




Note: When an image hits 1° visual cortex, it is upside down and left-right reversed.

Cavernous sinus

Collection of venous sinuses on either side of pituitary. Blood from eye and superficial cortex → cavernous sinus → internal jugular vein.

CNs III, IV, V₁, V₂, and VI plus postganglionic sympathetic pupillary fibers en route to orbit all pass through cavernous sinus. Cavernous portion of internal carotid artery is also here.

Cavernous sinus syndrome—presents with variable ophthalmoplegia, ↓ corneal sensation, Horner syndrome and occasional decreased maxillary sensation. 2° to pituitary tumor mass effect, carotid-cavernous fistula, or cavernous sinus thrombosis related to infection.



Internuclear ophthalmoplegia

Medial longitudinal fasciculus (MLF): pair of tracts that allows for crosstalk between CN VI and CN III nuclei. Coordinates both eyes to move in same horizontal direction. Highly myelinated (must communicate quickly so eyes move at same time). Lesions may be unilateral or bilateral (latter classically seen in multiple sclerosis, stroke).



Lesion in MLF = internuclear ophthalmoplegia (INO), a conjugate horizontal gaze palsy. Lack of communication such that when CN VI nucleus activates ipsilateral lateral rectus, contralateral CN III nucleus does not stimulate medial rectus to contract. Abducting eye displays nystagmus (CN VI overfires to stimulate CN III). Convergence normal.

MLF in MS.

When looking left, the left nucleus of CN VI fires, which contracts the left lateral rectus and stimulates the contralateral (right) nucleus of CN III via the right MLF to contract the right medial rectus.

Directional term (eg, right INO, left INO) refers to the eye that is unable to adduct.

INO = Ipsilateral adduction failure, Nystagmus **O**pposite.



ⓧ


▶ NEUROLOGY—PHARMACOLOGY

Epilepsy therapy

	PARTIAL (FOCAL)	GENERALIZED			MECHANISM	SIDE EFFECTS	NOTES
		TONIC-CLONIC	ABSENCE	STATUS EPILEPTICUS			
Benzodiazepines				** ✓	↑ GABA _A action	Sedation, tolerance, dependence, respiratory depression	Also for eclampsia seizures (1st line is MgSO ₄)
Carbamazepine	* ✓	✓			Blocks Na ⁺ channels	Diplopia, ataxia, blood dyscrasias (agranulocytosis, aplastic anemia), liver toxicity, teratogenesis (cleft lip/palate, spina bifida), induction of cytochrome P-450, SIADH, SJS	1st line for trigeminal neuralgia
Ethosuximide			* ✓		Blocks thalamic T-type Ca ²⁺ channels	EFGHIJ —Ethosuximide causes F atigue, G I distress, H eadache, I tching (and urticaria), S J	S ucks to have S ilent (absence) S eizures
Gabapentin	✓				Primarily inhibits high-voltage-activated Ca ²⁺ channels; designed as GABA analog	Sedation, ataxia	Also used for peripheral neuropathy, postherpetic neuralgia
Lamotrigine	✓	✓	✓		Blocks voltage-gated Na ⁺ channels, inhibits the release of glutamate	SJS (must be titrated slowly), hemophagocytic lymphohistiocytosis (black box warning)	
Levetiracetam	✓	✓			SV2A receptor blocker; may modulate GABA and glutamate release, inhibit voltage-gated Ca ²⁺ channels	Neuropsychiatric symptoms (eg, personality change), fatigue, drowsiness, headache	
Phenobarbital	✓	✓		✓	↑ GABA _A action	Sedation, tolerance, dependence, induction of cytochrome P-450, cardiorespiratory depression	1st line in neonates (“phenobabytal”)
Phenytoin, fosphenytoin	✓	* ✓		*** ✓	Blocks Na ⁺ channels; zero-order kinetics	PHENYTOIN : cytochrome P-450 induction, H irsutism, E nlarged gums, N ystagmus, Y ellow-brown skin, T eratogenicity (fetal hydantoin syndrome), O steopenia, I nhibited folate absorption, N europathy. Rare: SJS, DRESS syndrome, SLE-like syndrome. Toxicity leads to diplopia, ataxia, sedation.	
Topiramate	✓	✓			Blocks Na ⁺ channels, ↑ GABA action	S edation, s low cognition, kidney s tones, s kinny (weight loss), s ight threatened (glaucoma), s peech (word-finding) difficulties	Also used for migraine prophylaxis
Valproic acid	✓	* ✓	✓		↑ Na ⁺ channel inactivation, ↑ GABA concentration by inhibiting GABA transaminase	GI distress, rare but fatal hepatotoxicity (measure LFTs), pancreatitis, neural tube defects, tremor, weight gain, contraindicated in pregnancy	Also used for myoclonic seizures, bipolar disorder, migraine prophylaxis
Vigabatrin	✓				↑ GABA. Irreversible GABA transaminase inhibitor	Permanent visual loss (black box warning)	V ision gone a ll b ad with V igabatrin

* = Common use, ** = 1st line for acute, *** = 1st line for recurrent seizure prophylaxis.

Epilepsy therapy (continued)



PK

Barbiturates	Phenobarbital, pentobarbital, thiopental, secobarbital.
MECHANISM	Facilitate GABA _A action by ↑ duration of Cl ⁻ channel opening, thus ↓ neuron firing (barbiturates ↑ duration).
CLINICAL USE	Sedative for anxiety, seizures, insomnia, induction of anesthesia (thiopental).
ADVERSE EFFECTS	Respiratory and cardiovascular depression (can be fatal); CNS depression (can be exacerbated by alcohol use); dependence; drug interactions (induces cytochrome P-450). Overdose treatment is supportive (assist respiration and maintain BP). Contraindicated in porphyria.
Benzodiazepines	Diazepam, lorazepam, triazolam, temazepam, oxazepam, midazolam, chlordiazepoxide, alprazolam.
MECHANISM	Facilitate GABA _A action by ↑ frequency of Cl ⁻ channel opening (“ fren zodiazepines” ↑ f requency). ↓ REM sleep. Most have long half-lives and active metabolites (exceptions [ATOM]: A lprazolam, T riazolam, O xazepam, and M idazolam are short acting → higher addictive potential).
CLINICAL USE	Anxiety, panic disorder, spasticity, status epilepticus (lorazepam, diazepam, midazolam), eclampsia, detoxification (especially alcohol withdrawal–DTs), night terrors, sleepwalking, general anesthetic (amnesia, muscle relaxation), hypnotic (insomnia). L orazepam, O xazepam, and T emazepam can be used for those with liver disease who drink a LOT due to minimal first-pass metabolism.
ADVERSE EFFECTS	Dependence, additive CNS depression effects with alcohol and barbiturates (all bind the GABA _A receptor). Less risk of respiratory depression and coma than with barbiturates. Treat overdose with flumazenil (competitive antagonist at GABA benzodiazepine receptor). Can precipitate seizures by causing acute benzodiazepine withdrawal.
Nonbenzodiazepine hypnotics	Z olpidem, Z aleplon, esZ opiclone. “These ZZZ s put you to sleep.”
MECHANISM	Act via the BZ ₁ subtype of the GABA receptor. Effects reversed by flumazenil. Sleep cycle less affected as compared with benzodiazepine hypnotics.
CLINICAL USE	Insomnia.
ADVERSE EFFECTS	Ataxia, headaches, confusion. Short duration because of rapid metabolism by liver enzymes. Unlike older sedative-hypnotics, cause only modest day-after psychomotor depression and few amnestic effects. ↓ dependence risk than benzodiazepines.

Suvorexant

MECHANISM	Orexin (hypocretin) receptor antagonist.	Suvorexant is an orexin antagonist .
CLINICAL USE	Insomnia.	
ADVERSE EFFECTS	CNS depression (somnolence), headache, abnormal sleep-related activities. Contraindications: narcolepsy, combination with strong CYP3A4 inhibitors. Not recommended in patients with liver disease. Limited physical dependence or abuse potential.	

Ramelteon

MECHANISM	Melatonin receptor agonist; binds MT ₁ and MT ₂ in suprachiasmatic nucleus.	Ramelteon is a melatonin receptor agonist .
CLINICAL USE	Insomnia.	
ADVERSE EFFECTS	Dizziness, nausea, fatigue, headache. No dependence (not a controlled substance).	

Triptans**Sumatriptan**


MECHANISM	5-HT _{1B/1D} agonists. Inhibit trigeminal nerve activation, prevent vasoactive peptide release, induce vasoconstriction.	A sumo wrestler trips and falls on his head .
CLINICAL USE	Acute migraine, cluster headache attacks.	
ADVERSE EFFECTS	Coronary vasospasm (contraindicated in patients with CAD or vasospastic angina), mild paresthesia, serotonin syndrome (in combination with other 5-HT agonists).	

Parkinson disease therapy

Parkinsonism is due to loss of dopaminergic neurons and excess cholinergic activity.

Bromocriptine, **A**mantadine, **L**evodopa (with carbidopa), **S**elegiline (and COMT inhibitors), **A**ntimuscarinics (**BALSA**).

STRATEGY	AGENTS
Dopamine agonists	Ergot— B romocriptine. Non-ergot (preferred)—pramipexole, ropinirole; toxicity includes nausea, impulse control disorder (eg, gambling), postural hypotension, hallucinations, confusion.
↑ dopamine availability	A mantadine (↑ dopamine release and ↓ dopamine reuptake); toxicity = peripheral edema, livedo reticularis, ataxia.
↑ L-DOPA availability	Agents prevent peripheral (pre-BBB) L-DOPA degradation → ↑ L-DOPA entering CNS → ↑ central L-DOPA available for conversion to dopamine. <ul style="list-style-type: none"> ▪ Levodopa (L-DOPA)/carbidopa—carbidopa blocks peripheral conversion of L-DOPA to dopamine by inhibiting DOPA decarboxylase. Also reduces side effects of peripheral L-DOPA conversion into dopamine (eg, nausea, vomiting). ▪ Entacapone and tolcapone prevent peripheral L-DOPA degradation to 3-O-methyldopa (3-OMD) by inhibiting COMT. Used in conjunction with levodopa.
Prevent dopamine breakdown	Agents act centrally (post-BBB) to inhibit breakdown of dopamine. <ul style="list-style-type: none"> ▪ Selegiline, rasagiline—block conversion of dopamine into DOPAC by selectively inhibiting MAO-B. ▪ Tolcapone—crosses BBB and blocks conversion of dopamine to 3-methoxytyramine (3-MT) in the brain by inhibiting central COMT.
Curb excess cholinergic activity	B enzotropine, trihexyphenidyl (A ntimuscarinic; improves tremor and rigidity but has little effect on bradykinesia in P arkinson disease). P ark your Mercedes- B enz.



Carbidopa/levodopa

MECHANISM	↑ dopamine in brain. Unlike dopamine, L-DOPA can cross blood-brain barrier and is converted by dopa decarboxylase in the CNS to dopamine. Carbidopa, a peripheral DOPA decarboxylase inhibitor, is given with L-DOPA to ↑ bioavailability of L-DOPA in the brain and to limit peripheral side effects.
CLINICAL USE	Parkinson disease.
ADVERSE EFFECTS	Nausea, hallucinations, postural hypotension. With progressive disease, L-DOPA can lead to “on-off” phenomenon with improved mobility during “on” periods, then impaired motor function during “off” periods when patient responds poorly to L-DOPA or medication wears off.

Selegiline, rasagiline

MECHANISM	Selectively inhibit MAO-B (metabolize dopamine) → ↑ dopamine availability. Selegiline selectively inhibits MAO-B and is more commonly found in the Brain than in the periphery.
CLINICAL USE	Adjunctive agent to L-DOPA in treatment of Parkinson disease.
ADVERSE EFFECTS	May enhance adverse effects of L-DOPA.

Neurodegenerative disease therapy

DISEASE	AGENT	MECHANISM	NOTES
Alzheimer disease	Donepezil, rivastigmine, galantamine	AChE inhibitor	1st-line treatment Adverse effects: nausea, dizziness, insomnia Dona Riva dances at the gala
	Memantine	NMDA receptor antagonist; helps prevent excitotoxicity (mediated by Ca ²⁺)	Used for moderate to advanced dementia Adverse effects: dizziness, confusion, hallucinations
Amyotrophic lateral sclerosis	Riluzole	↓ neuron glutamate excitotoxicity	↑ survival Treat Lou Gehrig disease with rilouzole
Huntington disease	Tetrabenazine	Inhibit vesicular monoamine transporter (VMAT) → ↓ dopamine vesicle packaging and release	May be used for Huntington chorea and tardive dyskinesia

Anesthetics—general principles

CNS drugs must be lipid soluble (cross the blood-brain barrier) or be actively transported.
Drugs with ↓ solubility in blood = rapid induction and recovery times.

Drugs with ↑ solubility in lipids = ↑ potency = $\frac{1}{\text{MAC}}$

MAC = **M**inimum **A**lveolar **C**oncentration (of inhaled anesthetic) required to prevent 50% of subjects from moving in response to noxious stimulus (eg, skin incision).

Examples: nitrous oxide (N₂O) has ↓ blood and lipid solubility, and thus fast induction and low potency. Halothane has ↑ lipid and blood solubility, and thus high potency and slow induction.

Inhaled anesthetics	Desflurane, halothane, enflurane, isoflurane, sevoflurane, methoxyflurane, N ₂ O.
MECHANISM	Mechanism unknown.
EFFECTS	Myocardial depression, respiratory depression, postoperative nausea/vomiting, ↑ cerebral blood flow, ↓ cerebral metabolic demand.
ADVERSE EFFECTS	Hepatotoxicity (halothane), nephrotoxicity (methoxyflurane), proconvulsant (enflurane, epileptogenic), expansion of trapped gas in a body cavity (N ₂ O). Malignant hyperthermia —rare, life-threatening condition in which inhaled anesthetics or succinylcholine induce severe muscle contractions and hyperthermia. Susceptibility is often inherited as autosomal dominant with variable penetrance. Mutations in voltage-sensitive ryanodine receptor (RYR1 gene) cause ↑ Ca ²⁺ release from sarcoplasmic reticulum. Treatment: dantrolene (a ryanodine receptor antagonist).

Intravenous anesthetics

AGENT	MECHANISM	ANESTHESIA USE	NOTES
Thiopental	Facilitates GABA _A (barbiturate)	Anesthesia induction, short surgical procedures	↓ cerebral blood flow. High lipid solubility Effect terminated by rapid redistribution into tissue, fat
Midazolam	Facilitates GABA _A (benzodiazepine)	Procedural sedation (eg, endoscopy), anesthesia induction	May cause severe postoperative respiratory depression, ↓ BP, anterograde amnesia
Propofol	Potentiates GABA _A	Rapid anesthesia induction, short procedures, ICU sedation	May cause respiratory depression, hypotension
Ketamine	NMDA receptor antagonist	Dissociative anesthesia Sympathomimetic	↑ cerebral blood flow Emergence reaction possible with disorientation, hallucination, vivid dreams

Local anesthetics

	Esters—procaine, tetracaine, benzocaine, chlorprocaine. Amides—lidocaine, mepivacaine, bupivacaine, ropivacaine (amides have 2 I's in name).
MECHANISM	Block Na ⁺ channels by binding to specific receptors on inner portion of channel. Most effective in rapidly firing neurons. 3° amine local anesthetics penetrate membrane in uncharged form, then bind to ion channels as charged form. Can be given with vasoconstrictors (usually epinephrine) to enhance local action—↓ bleeding, ↑ anesthesia by ↓ systemic concentration. In infected (acidic) tissue, alkaline anesthetics are charged and cannot penetrate membrane effectively → need more anesthetic. Order of nerve blockade: small-diameter fibers > large diameter. Myelinated fibers > unmyelinated fibers. Overall, size factor predominates over myelination such that small myelinated fibers > small unmyelinated fibers > large myelinated fibers > large unmyelinated fibers. Order of loss: (1) pain, (2) temperature, (3) touch, (4) pressure.
CLINICAL USE	Minor surgical procedures, spinal anesthesia. If allergic to esters, give amides.
ADVERSE EFFECTS	CNS excitation, severe cardiovascular toxicity (bupivacaine), hypertension, hypotension, arrhythmias (cocaine), methemoglobinemia (benzocaine).

Neuromuscular blocking drugs	Muscle paralysis in surgery or mechanical ventilation. Selective for Nm nicotinic receptors at neuromuscular junction but not autonomic Nn receptors.
Depolarizing neuromuscular blocking drugs	Succinylcholine—strong ACh receptor agonist; produces sustained depolarization and prevents muscle contraction. Reversal of blockade: <ul style="list-style-type: none"> Phase I (prolonged depolarization)—no antidote. Block potentiated by cholinesterase inhibitors. Phase II (repolarized but blocked; ACh receptors are available, but desensitized)—may be reversed with cholinesterase inhibitors. Complications include hypercalcemia, hyperkalemia, malignant hyperthermia.
Nondepolarizing neuromuscular blocking drugs	Atracurium, cisatracurium, pancuronium, rocuronium, tubocurarine, vecuronium—competitive ACh antagonist. Reversal of blockade—cholinesterase inhibitors (eg, neostigmine, edrophonium) are given with anticholinergics (eg, atrophine, glycopyrrolate) to prevent muscarinic effects, such as bradycardia.

Spasmolytics, antispasmodics

DRUG	MECHANISM	CLINICAL USE	NOTES
Baclofen	GABA _B receptor agonist in spinal cord.	Muscle spasticity, dystonia, multiple sclerosis.	Acts on the back (spinal cord).
Cyclobenzaprine	Acts within CNS, mainly at the brain stem.	Muscle spasticity.	C entrally acting. Structurally related to TCAs. May cause anticholinergic side effects, sedation.
Dantrolene	Prevents release of Ca ²⁺ from sarcoplasmic reticulum of skeletal muscle by inhibiting the ryanodine receptor.	Malignant hyperthermia (toxicity of inhaled anesthetics and succinylcholine) and neuroleptic malignant syndrome (toxicity of antipsychotic drugs).	Acts D irectly on muscle.
Tizanidine	α ₂ agonist, acts centrally.	Muscle spasticity, multiple sclerosis, ALS, cerebral palsy.	

Opioid analgesics

MECHANISM	Act as agonists at opioid receptors (μ = β-endorphin, δ = enkephalin, κ = dynorphin) to modulate synaptic transmission—close presynaptic Ca ²⁺ channels, open postsynaptic K ⁺ channels → ↓ synaptic transmission. Inhibit release of ACh, norepinephrine, 5-HT, glutamate, substance P.
EFFICACY	Full agonist: morphine, heroin, meperidine, methadone, codeine, fentanyl. Partial agonist: buprenorphine. Mixed agonist/antagonist: nalbuphine, pentazocine, butorphanol. Antagonist: naloxone, naltrexone, methyl naltrexone.
CLINICAL USE	Moderate to severe or refractory pain, diarrhea (loperamide, diphenoxylate), acute pulmonary edema, maintenance programs for heroin addicts (methadone, buprenorphine + naloxone).
ADVERSE EFFECTS	Nausea, vomiting, pruritus, addiction, respiratory depression, constipation, sphincter of Oddi spasm, miosis (except meperidine → mydriasis), additive CNS depression with other drugs. Tolerance does not develop to miosis and constipation. Treat toxicity with naloxone (competitive opioid receptor antagonist) and prevent relapse with naltrexone once detoxified.

Mixed agonist and antagonist opioid analgesics

DRUG	MECHANISM	CLINICAL USE	NOTES
Pentazocine	κ -opioid receptor agonist and μ -opioid receptor weak antagonist or partial agonist.	Analgesia for moderate to severe pain.	Can cause opioid withdrawal symptoms if patient is also taking full opioid agonist (due to competition for opioid receptors).
Butorphanol	κ -opioid receptor agonist and μ -opioid receptor partial agonist.	Severe pain (eg, migraine, labor).	Causes less respiratory depression than full opioid agonists. Use with full opioid agonist can precipitate withdrawal. Not easily reversed with naloxone.

Tramadol

MECHANISM	Very weak opioid agonist; also inhibits the reuptake of norepinephrine and serotonin.	Tramadol is a S light opioid agonist, and a S erotonin and norepinephrine reuptake inhibitor. It is used for S tubborn pain, but can lower S eizure threshold, and may cause S erotonin S yndrome.
CLINICAL USE	Chronic pain.	
ADVERSE EFFECTS	Similar to opioids; decreases seizure threshold; serotonin syndrome.	

Glaucoma therapy

↓ IOP via ↓ amount of aqueous humor (inhibit synthesis/secretion or ↑ drainage).
BAD humor may not be **P**olitically **C**orrect.

DRUG CLASS	EXAMPLES	MECHANISM	ADVERSE EFFECTS
β-blockers	Timolol, betaxolol, carteolol	↓ aqueous humor synthesis	No pupillary or vision changes
α-agonists	Epinephrine (α_1), apraclonidine, brimonidine (α_2)	↓ aqueous humor synthesis via vasoconstriction (epinephrine) ↓ aqueous humor synthesis (apraclonidine, brimonidine)	Mydriasis (α_1); do not use in closed-angle glaucoma Blurry vision, ocular hyperemia, foreign body sensation, ocular allergic reactions, ocular pruritus
Diuretics	Acetazolamide	↓ aqueous humor synthesis via inhibition of carbonic anhydrase	No pupillary or vision changes
Prostaglandins	Bimatoprost, latanoprost (PGF _{2α})	↑ outflow of aqueous humor via ↓ resistance of flow through uveoscleral pathway	Darkens color of iris (browning), eyelash growth
Cholinomimetics (M₃)	Direct: pilocarpine, carbachol Indirect: physostigmine, echothiophate	↑ outflow of aqueous humor via contraction of ciliary muscle and opening of trabecular meshwork Use pilocarpine in acute angle closure glaucoma—very effective at opening meshwork into canal of Schlemm	Miosis (contraction of pupillary sphincter muscles) and cyclospasm (contraction of ciliary muscle)

HIGH-YIELD PRINCIPLES IN

Psychiatry

“Words of comfort, skillfully administered, are the oldest therapy known to man.”

—Louis Nizer

“All men should strive to learn before they die what they are running from, and to, and why.”

—James Thurber

“The sorrow which has no vent in tears may make other organs weep.”

—Henry Maudsley

“It’s no use going back to yesterday, because I was a different person then.”

—Lewis Carroll, *Alice in Wonderland*

This chapter encompasses overlapping areas in psychiatry, psychology, sociology, and psychopharmacology. High-yield topics include schizophrenia, mood disorders, eating disorders, personality disorders, somatic symptom disorders, substance abuse, and antipsychotic agents. Know the DSM-5 criteria for diagnosing common psychiatric disorders.

▶ Psychology	554
▶ Pathology	556
▶ Pharmacology	572

▶ PSYCHIATRY—PSYCHOLOGY

Classical conditioning	Learning in which a natural response (salivation) is elicited by a conditioned, or learned, stimulus (bell) that previously was presented in conjunction with an unconditioned stimulus (food).	Usually elicits involuntary responses. Pavlov's classical experiments with dogs—ringing the bell provoked salivation.									
Operant conditioning	Learning in which a particular action is elicited because it produces a punishment or reward. Usually elicits voluntary responses.										
Reinforcement	Target behavior (response) is followed by desired reward (positive reinforcement) or removal of aversive stimulus (negative reinforcement).	Skinner operant conditioning quadrants: <table border="1"> <thead> <tr> <th></th> <th>Increase behavior</th> <th>Decrease behavior</th> </tr> </thead> <tbody> <tr> <th>Add a stimulus</th> <td>Positive reinforcement</td> <td>Positive punishment</td> </tr> <tr> <th>Remove a stimulus</th> <td>Negative reinforcement</td> <td>Negative punishment</td> </tr> </tbody> </table>		Increase behavior	Decrease behavior	Add a stimulus	Positive reinforcement	Positive punishment	Remove a stimulus	Negative reinforcement	Negative punishment
	Increase behavior		Decrease behavior								
Add a stimulus	Positive reinforcement		Positive punishment								
Remove a stimulus	Negative reinforcement		Negative punishment								
Punishment	Repeated application of aversive stimulus (positive punishment) or removal of desired reward (negative punishment) to extinguish unwanted behavior.										
Extinction	Discontinuation of reinforcement (positive or negative) eventually eliminates behavior. Can occur in operant or classical conditioning.										
Transference and countertransference											
Transference	Patient projects feelings about formative or other important persons onto physician (eg, psychiatrist is seen as parent).										
Countertransference	Doctor projects feelings about formative or other important persons onto patient (eg, patient reminds physician of younger sibling).										
Ego defenses											
	Thoughts and behaviors (voluntary or involuntary) used to resolve conflict and prevent undesirable feelings (eg, anxiety, depression).										
IMMATURE DEFENSES	DESCRIPTION	EXAMPLE									
Acting out	Subconsciously coping with stressors or emotional conflict using actions rather than reflections or feelings.	A patient skips therapy appointments after deep discomfort from dealing with his past.									
Denial	Avoiding the awareness of some painful reality.	A patient with cancer plans a full-time work schedule despite being warned of significant fatigue during chemotherapy.									
Displacement	Redirection of emotions or impulses to a neutral person or object (vs projection).	After being reprimanded by her principal, a frustrated teacher returns home and criticizes her husband's cooking instead of confronting the principal directly.									
Dissociation	Temporary, drastic change in personality, memory, consciousness, or motor behavior to avoid emotional stress. Patient has incomplete or no memory of traumatic event.	A victim of sexual abuse suddenly appears numb and detached when she is exposed to her abuser.									

Ego defenses (continued)

IMMATURE DEFENSES	DESCRIPTION	EXAMPLE
Fixation	Partially remaining at a more childish level of development (vs regression).	A surgeon throws a tantrum in the operating room because the last case ran very late.
Idealization	Expressing extremely positive thoughts of self and others while ignoring negative thoughts.	A patient boasts about his physician and his accomplishments while ignoring any flaws.
Identification	Largely unconscious assumption of the characteristics, qualities, or traits of another person or group.	A resident starts putting his stethoscope in his pocket like his favorite attending, instead of wearing it around his neck like before.
Intellectualization	Using facts and logic to emotionally distance oneself from a stressful situation.	A patient diagnosed with cancer discusses the pathophysiology of the disease.
Isolation (of affect)	Separating feelings from ideas and events.	Describing murder in graphic detail with no emotional response.
Passive aggression	Demonstrating hostile feelings in a nonconfrontational manner; showing indirect opposition.	A disgruntled employee is repeatedly late to work, but won't admit it is a way to get back at the manager.
Projection	Attributing an unacceptable internal impulse to an external source (vs displacement).	A man who wants to cheat on his wife accuses his wife of being unfaithful.
Rationalization	Asserting plausible explanations for events that actually occurred for other reasons, usually to avoid self-blame.	A man who was recently fired claims that the job was not important anyway.
Reaction formation	Replacing a warded-off idea or feeling with an emphasis on its opposite (vs sublimation).	A stepmother treats a child she resents with excessive nurturing and overprotection.
Regression	Involuntarily turning back the maturational clock to behaviors previously demonstrated under stress (vs fixation).	A previously toilet-trained child begins bedwetting again following the birth of a sibling.
Repression	Involuntarily withholding an idea or feeling from conscious awareness (vs suppression).	A 20-year-old does not remember going to counseling during his parents' divorce 10 years earlier.
Splitting	Believing that people are either all good or all bad at different times due to intolerance of ambiguity. Common in borderline personality disorder.	A patient says that all the nurses are cold and insensitive, but the doctors are warm and friendly.
MATURE DEFENSES		
Sublimation	Replacing an unacceptable wish with a course of action that is similar to the wish but socially acceptable (vs reaction formation).	A teenager's aggression toward his parents because of their high expectations is channeled into excelling in sports.
Altruism	Alleviating negative feelings via unsolicited generosity, which provides gratification (vs reaction formation).	A mafia boss makes a large donation to charity.
Suppression	Intentionally withholding an idea or feeling from conscious awareness (vs repression); temporary.	An athlete focuses on other tasks to prevent worrying about an important upcoming match.
Humor	Lightheartedly expressing uncomfortable feelings to shift the internal focus away from the distress.	A nervous medical student jokes about the boards.

Mature adults wear a **SASH**.

▶ PSYCHIATRY—PATHOLOGY

Infant deprivation effects

Long-term deprivation of affection results in:

- Failure to thrive
- Poor language/socialization skills
- Lack of basic trust
- Reactive attachment disorder (infant withdrawn/unresponsive to comfort)
- Disinhibited social engagement (child indiscriminately attaches to strangers)

Deprivation for > 6 months can lead to irreversible changes.

Severe deprivation can result in infant death.

Child abuse

	Physical abuse	Sexual abuse	Emotional abuse
SIGNS	Fractures, bruises, or burns. Injuries often in different stages of healing or in patterns resembling possible implements of injury. Includes abusive head trauma (shaken baby syndrome), characterized by subdural hematomas or retinal hemorrhages. Caregivers may delay seeking medical attention for the child or provide explanations inconsistent with the child's developmental stage or pattern of injury.	STIs, UTIs, and genital, anal, or oral trauma. Most often, there are no physical signs; sexual abuse should not be excluded from a differential diagnosis in the absence of physical trauma. Children often exhibit sexual knowledge or behavior incongruent with their age.	Babies or young children may lack a bond with the caregiver but are overly affectionate with less familiar adults. They may be aggressive toward children and animals or unusually anxious. Older children are often emotionally labile and prone to angry outbursts. They may distance themselves from caregivers and other children. They can experience vague somatic symptoms for which a medical cause cannot be found.
EPIDEMIOLOGY	40% of deaths related to child abuse or neglect occur in children < 1 year old.	Peak incidence 9–12 years old.	~80% of young adult victims of child emotional abuse meet the criteria for ≥ 1 psychiatric illness by age 21.

Child neglect

Failure to provide a child with adequate food, shelter, supervision, education, and/or affection. Most common form of child maltreatment. Signs: poor hygiene, malnutrition, withdrawal, impaired social/emotional development, failure to thrive. As with child abuse, suspected child neglect must be reported to local child protective services.

Vulnerable child syndrome

Parents perceive the child as especially susceptible to illness or injury (vs factitious disorder imposed on another). Usually follows a serious illness or life-threatening event. Can result in missed school or overuse of medical services.

Childhood and early-onset disorders

Attention-deficit hyperactivity disorder	Onset before age 12. ≥ 6 months of limited attention span and/or poor impulse control. Characterized by hyperactivity, impulsivity, and/or inattention in ≥ 2 settings (eg, school, home, places of worship). Normal intelligence, but commonly coexists with difficulties in school. Often persists into adulthood. Commonly coexists with oppositional defiant disorder. Treatment: stimulants (eg, methylphenidate) +/- behavioral therapy; alternatives include atomoxetine, guanfacine, clonidine.
Autism spectrum disorder	Onset in early childhood. Social and communication deficits, repetitive/ritualized behaviors, restricted interests. May be accompanied by intellectual disability and/or above average abilities in specific skills (eg, music). More common in boys. Associated with \uparrow head and/or brain size.
Conduct disorder	Repetitive, pervasive behavior violating societal norms or the basic rights of others (eg, aggression toward people and animals, destruction of property, theft). After age 18, often reclassified as antisocial personality disorder. Treatment: psychotherapy (eg, cognitive behavioral therapy [CBT]).
Disruptive mood dysregulation disorder	Onset before age 10. Severe, recurrent temper outbursts out of proportion to situation. Child is constantly angry and irritable between outbursts. Treatment: CBT, stimulants, antipsychotics.
Intellectual disability	Global cognitive deficits (vs specific learning disorder) that affect reasoning, memory, abstract thinking, judgment, language, learning. Adaptive functioning is impaired, leading to major difficulties with education, employment, communication, socialization, independence. Treatment: psychotherapy, occupational therapy, special education.
Oppositional defiant disorder	Enduring pattern of anger and irritability with argumentative, vindictive, and defiant behavior toward authority figures. Treatment: psychotherapy (eg, CBT).
Selective mutism	Onset before age 5. Anxiety disorder lasting ≥ 1 month involving refraining from speech in certain situations despite speaking in other, usually more comfortable situations. Development (eg, speech and language) not typically impaired. Interferes with social, academic, and occupational tasks. Commonly coexists with social anxiety disorder. Treatment: behavioral, family, and play therapy; SSRIs.
Separation anxiety disorder	Overwhelming fear of separation from home or attachment figure lasting ≥ 4 weeks. Can be normal behavior up to age 3–4. May lead to factitious physical complaints to avoid school. Treatment: CBT, play therapy, family therapy.
Specific learning disorder	Onset during school-age years. Inability to acquire or use information from a specific subject (eg, math, reading, writing) near age-expected proficiency for ≥ 6 months despite focused intervention. General functioning and intelligence are normal (vs intellectual disability). Treatment: academic support, counseling, extracurricular activities.
Tourette syndrome	Onset before age 18. Sudden, recurrent, nonrhythmic, stereotyped motor and vocal tics that persist for > 1 year. Coprolalia (involuntary obscene speech) found in some patients. Associated with OCD and ADHD. Treatment: psychoeducation, behavioral therapy. For intractable and distressing tics, high-potency antipsychotics (eg, haloperidol, fluphenazine), tetrabenazine, α_2 -agonists (eg, guanfacine, clonidine), or atypical antipsychotics.

Orientation

Patients' ability to know the date and time, where they are, and who they are (order of loss: time \rightarrow place \rightarrow person). Common causes of loss of orientation: alcohol, drugs, fluid/electrolyte imbalance, head trauma, hypoglycemia, infection, nutritional deficiencies, hypoxia.

Amnesias

Retrograde amnesia	Inability to remember things that occurred before a CNS insult.
Anterograde amnesia	Inability to remember things that occurred after a CNS insult (↓ acquisition of new memory).
Korsakoff syndrome	Amnesia (anterograde > retrograde) and disorientation caused by vitamin B ₁ deficiency. Associated with disruption and destruction of the limbic system, especially mammillary bodies and anterior thalamus. Seen in alcoholics as a late neuropsychiatric manifestation of Wernicke encephalopathy. Confabulations are characteristic.

Dissociative disorders

Depersonalization/derealization disorder	Persistent feelings of detachment or estrangement from one's own body, thoughts, perceptions, and actions (depersonalization) or one's environment (derealization). Intact reality testing (vs psychosis).
Dissociative amnesia	Inability to recall important personal information, usually following severe trauma or stress. May be accompanied by dissociative fugue (abrupt, unexpected travelling away from home).
Dissociative identity disorder	Formerly called multiple personality disorder. Presence of ≥ 2 distinct identities or personality states. More common in women. Associated with history of sexual abuse, PTSD, depression, substance abuse, borderline personality, somatic symptom disorders.

Delirium

“Waxing and waning” level of consciousness with acute onset, ↓ attention span, ↓ level of arousal. Characterized by disorganized thinking, hallucinations (often visual), misperceptions (eg, illusions), disturbance in sleep-wake cycle, cognitive dysfunction, agitation. Reversible.

Usually 2° to other identifiable illness (eg, CNS disease, infection, trauma, substance abuse/withdrawal, metabolic/electrolyte disturbances, hemorrhage, urinary/fecal retention), or medications (eg, anticholinergics), especially in the elderly.

Most common presentation of altered mental status in inpatient setting, especially in the ICU or during prolonged hospital stays. EEG may show diffuse background rhythm slowing.

Delirium = changes in **sensorium**.

Treatment: identification and management of underlying condition. Orientation protocols (eg, keeping a clock or calendar nearby), ↓ sleep disturbances, and ↑ cognitive stimulation to manage symptoms.

Antipsychotics as needed. Avoid unnecessary restraints and drugs that may worsen delirium (eg, anticholinergics, benzodiazepines, opioids).

Psychosis	Distorted perception of reality characterized by delusions, hallucinations, and/or disorganized thought/speech. Can occur in patients with medical illness, psychiatric illness, or both.
Delusions	False, fixed, idiosyncratic beliefs that persist despite evidence to the contrary and are not typical of a patient's culture or religion (eg, a patient who believes that others are reading his thoughts). Types include erotomanic, grandiose, jealous, persecutory, somatic, mixed, and unspecified.
Disorganized thought	Speech may be incoherent ("word salad"), tangential, or derailed ("loose associations").
Hallucinations	Perceptions in the absence of external stimuli (eg, seeing a light that is not actually present). Contrast with misperceptions (eg, illusions) of real external stimuli. Types include: <ul style="list-style-type: none"> ▪ Auditory—more commonly due to psychiatric illness (eg, schizophrenia) than medical illness. ▪ Visual—more commonly due to medical illness (eg, drug intoxication, delirium) than psychiatric illness. ▪ Tactile—common in alcohol withdrawal and stimulant use (eg, "cocaine crawlies," a type of delusional parasitosis). ▪ Olfactory—often occur as an aura of temporal lobe epilepsy (eg, burning rubber) and in brain tumors. ▪ Gustatory—rare, but seen in epilepsy. ▪ Hypnagogic—occurs while going to sleep. Sometimes seen in narcolepsy. ▪ Hypnopompic—occurs while waking from sleep ("get pumped up in the morning"). Sometimes seen in narcolepsy.

Schizophrenia spectrum disorders

Schizophrenia

Chronic illness causing profound functional impairment. Symptom categories include:

- Positive—hallucinations, delusions, unusual thought processes, disorganized speech, bizarre behavior
- Negative—flat or blunted affect, apathy, anhedonia, alogia, social withdrawal
- Cognitive—reduced ability to understand or make plans, diminished working memory, inattention

Diagnosis requires ≥ 2 of the following active symptoms, including ≥ 1 from symptoms #1–3:

1. Delusions
2. Hallucinations, often auditory
3. Disorganized speech
4. Disorganized or catatonic behavior
5. Negative symptoms

Requires ≥ 1 month of active symptoms over the past 6 months; onset ≥ 6 months prior to diagnosis.

Brief psychotic disorder— ≥ 1 positive symptom(s) lasting < 1 month, usually stress-related.

Schizophreniform disorder— ≥ 2 symptoms lasting 1–6 months.

Associated with altered dopaminergic activity, \uparrow serotonergic activity, and \downarrow dendritic branching. Ventriculomegaly on brain imaging. Lifetime prevalence—1.5% (males $>$ females). Presents earlier in men (late teens to early 20s) than in women (late 20s to early 30s). \uparrow suicide risk.

Heavy cannabis use in adolescence is associated with \uparrow incidence and worsened course of psychotic, mood, and anxiety disorders.

Treatment: atypical antipsychotics (eg, risperidone) are first line.

Negative symptoms often persist after treatment, despite resolution of positive symptoms.

Schizoaffective disorder

Shares symptoms with both schizophrenia and mood disorders (major depressive or bipolar disorder). To differentiate from a mood disorder with psychotic features, patient must have > 2 weeks of psychotic symptoms without a manic or depressive episode.

Delusional disorder

≥ 1 delusion(s) lasting > 1 month, but without a mood disorder or other psychotic symptoms. Daily functioning, including socialization, may be impacted by the pathological, fixed belief but is otherwise unaffected. Can be shared by individuals in close relationships (folie à deux).

Schizotypal personality disorder

Cluster A personality disorder that also falls on the schizophrenia spectrum. May include brief psychotic episodes (eg, delusions) that are less frequent and severe than in schizophrenia.

Mood disorder

Characterized by an abnormal range of moods or internal emotional states and loss of control over them. Severity of moods causes distress and impairment in social and occupational functioning. Includes major depressive, bipolar, dysthymic, and cyclothymic disorders. Episodic superimposed psychotic features (delusions, hallucinations, disorganized speech/behavior) may be present.

Manic episode

Distinct period of abnormally and persistently elevated, expansive, or irritable mood and \uparrow activity or energy lasting ≥ 1 week. Diagnosis requires hospitalization or marked functional impairment with ≥ 3 of the following (manics **DIG FAST**):

- **D**istractibility
- **I**mpulsivity/**I**ndiscretion—seeks pleasure without regard to consequences (hedonistic)
- **G**randiosity—inflated self-esteem
- **F**light of ideas—racing thoughts
- \uparrow goal-directed **A**ctivity/psychomotor **A**gitation
- \downarrow need for **S**leep
- **T**alkativeness or pressured speech

Hypomanic episode	Similar to a manic episode except mood disturbance is not severe enough to cause marked impairment in social and/or occupational functioning or to necessitate hospitalization. Abnormally ↑ activity or energy usually present. No psychotic features. Lasts ≥ 4 consecutive days.
Bipolar disorder	<p>Bipolar I—≥ 1 manic episode +/- a hypomanic or depressive episode (may be separated by any length of time).</p> <p>Bipolar II—a hypomanic and a depressive episode (no history of manic episodes). Patient's mood and functioning usually normalize between episodes. Use of antidepressants can destabilize mood. High suicide risk. Treatment: mood stabilizers (eg, lithium, valproic acid, carbamazepine, lamotrigine), atypical antipsychotics.</p> <p>Cyclothymic disorder—milder form of bipolar disorder fluctuating between mild depressive and hypomanic symptoms. Must last ≥ 2 years with symptoms present at least half of the time, with any remission lasting ≤ 2 months.</p>
Major depressive disorder	<p>Recurrent episodes lasting ≥ 2 weeks characterized by ≥ 5 of 9 diagnostic symptoms (must include depressed mood or anhedonia) (DIGS SPACE):</p> <ul style="list-style-type: none"> ▪ Depressed mood (or irritability in children) ▪ ↓ Interest (anhedonia) ▪ Guilt or feelings of worthlessness ▪ Sleep disturbances ▪ Suicidal ideation ▪ Psychemotor retardation or agitation ▪ Appetite/weight changes ▪ ↓ Concentration ▪ ↓ Energy <p>Screen for previous manic or hypomanic episodes to rule out bipolar disorder. Treatment: CBT and SSRIs are first line. Also SNRIs, mirtazapine, bupropion, electroconvulsive therapy (ECT).</p>
MDD with psychotic features	MDD + hallucinations or delusions. Psychotic features are typically mood congruent (eg, depressive themes of inadequacy, guilt, punishment, nihilism, disease, or death) and occur only in the context of major depressive episode (vs schizoaffective disorder). Treatment: antidepressant with atypical antipsychotic, ECT.
Persistent depressive disorder (dysthymia)	Often milder than MDD; ≥ 2 depressive symptoms lasting ≥ 2 years (≥ 1 year in children), with any remission lasting ≤ 2 months.
MDD with seasonal pattern	Formerly called seasonal affective disorder. Major depressive episodes occurring only during a particular season (usually winter) in ≥ 2 consecutive years and in most years across a lifetime. Atypical symptoms common.
Depression with atypical features	Characterized by mood reactivity (transient improvement in response to a positive event), hypersomnia, hyperphagia, leaden paralysis (heavy feeling in arms and legs), long-standing interpersonal rejection sensitivity. Most common subtype of depression. Treatment: CBT and SSRIs are first line. MAO inhibitors (MAOIs) are effective but not first line because of their risk profile.

Peripartum mood disturbances	Onset during or shortly after pregnancy or within 4 weeks of delivery. ↑ risk with history of mood disorders.
Maternal (postpartum) blues	50–85% incidence rate. Characterized by depressed affect, tearfulness, and fatigue starting 2–3 days after delivery. Usually resolves within 2 weeks. Treatment: supportive. Follow up to assess for possible MDD with peripartum onset.
MDD with peripartum onset	10–15% incidence rate. Formerly called postpartum depression. Meets MDD criteria with onset no later than 1 year after delivery. Treatment: CBT and SSRIs are first line.
Postpartum psychosis	0.1–0.2% incidence rate. Characterized by mood-congruent delusions, hallucinations, and thoughts of harming the baby or self. Risk factors include first pregnancy, family history, bipolar disorder, psychotic disorder, recent medication change. Treatment: hospitalization and initiation of atypical antipsychotic; if insufficient, ECT may be used.

Grief	<p>The five stages of grief per the Kübler-Ross model are denial, anger, bargaining, depression, and acceptance (may occur in any order). Other normal grief symptoms include shock, guilt, sadness, anxiety, yearning, and somatic symptoms that usually occur in waves. Simple hallucinations of the deceased person are common (eg, hearing the deceased speaking). Any thoughts of dying are limited to joining the deceased (vs complicated grief). Duration varies widely; usually resolves within 6–12 months.</p> <p>Persistent complex bereavement disorder involves obsessive preoccupation with the deceased and causes functional impairment, lasting at least 12 months (6 months in children). Can also meet criteria for major depressive episode.</p>
--------------	---

Electroconvulsive therapy	Rapid-acting method to treat refractory depression, depression with psychotic symptoms, catatonia, and acute suicidality. Induces tonic-clonic seizure under anesthesia and neuromuscular blockade. Adverse effects include disorientation, headache, partial anterograde/retrograde amnesia usually resolving in 6 months. No absolute contraindications. Safe in pregnant and elderly individuals.
----------------------------------	--

Risk factors for suicide completion	<p>Sex (male)</p> <p>Age (young adult or elderly)</p> <p>Depression</p> <p>Previous attempt (highest risk factor)</p> <p>Ethanol or drug use</p> <p>Rational thinking loss (psychosis)</p> <p>Sickness (medical illness)</p> <p>Organized plan</p> <p>No spouse or other social support</p> <p>Stated future intent</p>	<p>SAD PERSONS are more likely to complete suicide.</p> <p>Most common method in US is firearms; access to guns ↑ risk of suicide completion.</p> <p>Women try more often; men complete more often.</p> <p>Other risk factors include recent psychiatric hospitalization and family history of completed suicide.</p>
--	---	--

Anxiety disorders	Inappropriate experiences of fear/worry and their physical manifestations incongruent with the magnitude of the stressors. Symptoms are not attributable to another psychiatric disorder, medical condition (eg, hyperthyroidism), or substance abuse. Includes panic disorder, phobias, generalized anxiety disorder, and selective mutism.
--------------------------	--

Panic disorder

Recurrent panic attacks involving intense fear and discomfort +/- a known trigger. Attacks typically peak in 10 minutes with ≥ 4 of the following: palpitations, paresthesias, depersonalization or derealization, abdominal pain, nausea, intense fear of dying, intense fear of losing control, lightheadedness, chest pain, chills, choking, sweating, shaking, shortness of breath. Strong genetic component. ↑ risk of suicide.

Diagnosis requires attack followed by ≥ 1 month of ≥ 1 of the following:

- Persistent concern of additional attacks
- Worrying about consequences of attack
- Behavioral change related to attacks

Symptoms are systemic manifestations of fear.

Treatment: CBT, SSRIs, and venlafaxine are first line. Benzodiazepines occasionally used in acute setting.

Phobias

Severe, persistent (≥ 6 months) fear or anxiety due to presence or anticipation of a specific object or situation. Person often recognizes fear is excessive. Treatment: CBT with exposure therapy.

Social anxiety disorder—exaggerated fear of embarrassment in social situations (eg, public speaking, using public restrooms). Treatment: CBT, SSRIs, venlafaxine. For performance type (eg, anxiety restricted to public speaking), use β -blockers or benzodiazepines as needed.

Agoraphobia—irrational fear/anxiety while facing or anticipating ≥ 2 specific situations (eg, open/closed spaces, lines, crowds, public transport). If severe, patients may refuse to leave their homes. Associated with panic disorder. Treatment: CBT, SSRIs.

Generalized anxiety disorder

Excessive anxiety and worry about different aspects of daily life (eg, work, school, children) for most days of ≥ 6 months. Associated with ≥ 3 of the following for adults (≥ 1 for kids): restlessness, irritability, sleep disturbance, fatigue, muscle tension, difficulty concentrating. Treatment: CBT, SSRIs, SNRIs are first line. Bupropion, TCAs, benzodiazepines are second line.

Obsessive-compulsive disorders

Obsessions (recurring intrusive thoughts, feelings, or sensations) that cause severe distress, relieved in part by compulsions (performance of repetitive, often time-consuming actions). Ego-dystonic: behavior inconsistent with one's beliefs and attitudes (vs obsessive-compulsive personality disorder, ego-syntonic). Associated with Tourette syndrome. Treatment: CBT and SSRIs; clomipramine and venlafaxine are second line.

Body dysmorphic disorder—preoccupation with minor or imagined defects in appearance.

Causes significant emotional distress and repetitive appearance-related behaviors (eg, mirror checking, excessive grooming). Common in eating disorders. Treatment: CBT.

Trichotillomania

Compulsively pulling out one's hair. Causes significant distress and persists despite attempts to stop. Presents with areas of thinning hair or baldness on any area of the body, most commonly the scalp

A. Incidence highest in childhood but spans all ages. Treatment: psychotherapy.


Trauma and stress-related disorders

Adjustment disorder Emotional or behavioral symptoms (eg, anxiety, outbursts) that occur within 3 months of an identifiable psychosocial stressor (eg, divorce, illness) lasting < 6 months once the stressor has ended. If symptoms persist > 6 months after stressor ends, it is GAD. Symptoms do not meet criteria for MDD. Treatment: CBT is first line; antidepressants and anxiolytics may be considered.

Post-traumatic stress disorder Experiencing, or discovering that a loved one has experienced, a life-threatening situation (eg, serious injury, rape, witnessing death) → persistent **H**yperarousal, **A**voidance of associated stimuli, intrusive **R**e-experiencing of the event (eg, nightmares, flashbacks), changes in cognition or mood (eg, fear, horror, **D**istress) (having PTSD is **HARD**). Disturbance lasts > 1 month with significant distress or impaired functioning. Treatment: CBT, SSRIs, and venlafaxine are first line. Prazosin can reduce nightmares.

Acute stress disorder—lasts between 3 days and 1 month. Treatment: CBT; pharmacotherapy is usually not indicated.

Diagnostic criteria by symptom duration



Personality

Personality trait	An enduring, repetitive pattern of perceiving, relating to, and thinking about the environment and oneself.	
Personality disorder	Inflexible, maladaptive, and rigidly pervasive pattern of behavior causing subjective distress and/or impaired functioning; person is usually not aware of problem (ego-syntonic). Usually presents by early adulthood. Three clusters: A, B, C ; remember as Weird, Wild, and Worried , respectively, based on symptoms.	
Cluster A personality disorders	Odd or eccentric; inability to develop meaningful social relationships. No psychosis; genetic association with schizophrenia.	Cluster A : Accusatory, Aloof, Awkward . “ Weird .”
Paranoid	Pervasive distrust (Accusatory), suspiciousness, hypervigilance, and a profoundly cynical view of the world.	
Schizoid	Voluntary social withdrawal (Aloof), limited emotional expression, content with social isolation (vs avoidant).	
Schizotypal	Eccentric appearance, odd beliefs or magical thinking, interpersonal Awkwardness .	Included on the schizophrenia spectrum. Pronounce schizo- type -al: odd-type thoughts.
Cluster B personality disorders	Dramatic, emotional, or erratic; genetic association with mood disorders and substance abuse.	Cluster B : Bad, Borderline, flamBoyant , must be the Best . “ Wild .”
Antisocial	Disregard for the rights of others with lack of remorse. Involves criminality, impulsivity, hostility, and manipulation. Males > females. Must be ≥ 18 years old with evidence of conduct disorder onset before age 15. Diagnosis is conduct disorder if < 18 years old.	Antisocial = sociopath . Bad .
Borderline	Unstable mood and interpersonal relationships, fear of abandonment, impulsivity, self-mutilation, suicidality, sense of emotional emptiness. Females > males. Splitting is a major defense mechanism.	Treatment: dialectical behavior therapy. Borderline .
Histrionic	Attention-seeking, dramatic speech and emotional expression, shallow and labile emotions, sexually provocative. May use physical appearance to draw attention.	FlamBoyant .
Narcissistic	Grandiosity, sense of entitlement; lacks empathy and requires excessive admiration; often demands the “best” and reacts to criticism with rage and/or defensiveness. Fragile self-esteem. Often envious of others.	Must be the Best .

Cluster C personality disorders	Anxious or fearful; genetic association with anxiety disorders.	Cluster C : C owardly, obsessive- C ompulsive, C lingy. “ W orried.”
Avoidant	Hypersensitive to rejection and criticism, socially inhibited, timid, feelings of inadequacy, desires relationships with others (vs schizoid).	C owardly.
Obsessive-Compulsive	Preoccupation with order, perfectionism, and control; ego-syntonic: behavior consistent with one’s own beliefs and attitudes (vs OCD).	
Dependent	Excessive need for support, low self-confidence. Patients often get stuck in abusive relationships.	Submissive and C lingy.
Malingering	Symptoms are intentional , motivation is intentional . Patient consciously fakes, profoundly exaggerates, or claims to have a disorder in order to attain a specific 2° (external) gain (eg, avoiding work, obtaining compensation). Poor compliance with treatment or follow-up of diagnostic tests. Complaints cease after gain (vs factitious disorder).	
Factitious disorders	Symptoms are intentional , motivation is unconscious . Patient consciously creates physical and/or psychological symptoms in order to assume “sick role” and to get medical attention and sympathy (1° [internal] gain).	
Factitious disorder imposed on self	Formerly called Munchausen syndrome. Chronic factitious disorder with predominantly physical signs and symptoms. Characterized by a history of multiple hospital admissions and willingness to undergo invasive procedures. More common in women and healthcare workers.	
Factitious disorder imposed on another	Formerly called Munchausen syndrome by proxy. Illness in a child or elderly patient is caused or fabricated by the caregiver. Motivation is to assume a sick role by proxy. Form of child/elder abuse.	
Somatic symptom and related disorders	Symptoms are unconscious , motivation is unconscious . Category of disorders characterized by physical symptoms causing significant distress and impairment. Symptoms not intentionally produced or feigned.	
Somatic symptom disorder	≥ 1 bodily complaints (eg, abdominal pain, fatigue) lasting months to years. Associated with excessive, persistent thoughts and anxiety about symptoms. May co-occur with medical illness. Treatment: regular office visits with the same physician in combination with psychotherapy.	
Conversion disorder	Also called functional neurologic symptom disorder. Loss of sensory or motor function (eg, paralysis, blindness, mutism), often following an acute stressor; patient may be aware of but indifferent toward symptoms (<i>la belle indifférence</i>); more common in females, adolescents, and young adults.	
Illness anxiety disorder	Preoccupation with acquiring or having a serious illness, often despite medical evaluation and reassurance; minimal to no somatic symptoms.	

Eating disorders	Most common in young women.
Anorexia nervosa	Intense fear of weight gain, overvaluation of thinness, and body image distortion leading to calorie restriction and severe weight loss resulting in inappropriately low body weight. Binge-eating/purging type —recurring purging behaviors (eg, laxative or diuretic abuse, self-induced vomiting) or binge eating over the last 3 months. Restricting type —primary disordered behaviors include dieting, fasting, and/or over-exercising. No recurring purging behaviors or binge eating over the last 3 months. Refeeding syndrome —often occurs in significantly malnourished patients with sudden ↑ calorie intake → ↑ insulin → ↓ PO ₄ ³⁻ , ↓ K ⁺ , ↓ Mg ²⁺ → cardiac complications, rhabdomyolysis, seizures. Treatment: psychotherapy, nutritional rehabilitation, antidepressants (eg, SSRIs).
Bulimia nervosa	Recurring episodes of binge eating with compensatory purging behaviors at least weekly over the last 3 months. BMI often normal or slightly overweight (vs anorexia). Associated with parotid gland hypertrophy (may see ↑ serum amylase), enamel erosion, Mallory-Weiss syndrome, electrolyte disturbances (eg, ↓ K ⁺ , ↓ Cl ⁻), metabolic alkalosis, dorsal hand calluses from induced vomiting (Russell sign). Treatment: psychotherapy, nutritional rehabilitation, antidepressants (eg, SSRIs). Bupropion is contraindicated due to seizure risk.
Binge-eating disorder	Recurring episodes of binge eating without purging behaviors at least weekly over the last 3 months. ↑ diabetes risk. Most common eating disorder in adults. Treatment: psychotherapy (first line); SSRIs; lisdexamfetamine.
Pica	Recurring episodes of eating non-food substances (eg, dirt, hair, paint chips) over ≥ 1 month that are not culturally or developmentally recognized as normal. May provide temporary emotional relief. Common in children and during pregnancy. Associated with malnutrition, iron deficiency anemia, developmental disabilities, emotional trauma. Treatment: psychotherapy and nutritional rehabilitation (first line); SSRIs (second line).
Gender dysphoria	Significant incongruence between one's experienced gender and the gender assigned at birth, lasting > 6 months and leading to persistent distress. Individuals may self-identify as another gender, pursue surgery or hormone treatment to rid self of primary/secondary sex characteristics, and/or live as another gender. Gender nonconformity itself is not a mental disorder. Transgender —desiring and often making lifestyle changes to live as a different gender . Medical interventions (eg, hormone therapy, sex reassignment surgery) may be utilized during the transition to enable the individual's appearance to match their gender identity. Transvestism —deriving pleasure from wearing clothes (eg, a vest) of the opposite sex (cross-dressing). Transvestic disorder —transvestism that causes significant distress/functional impairment. It is a paraphilia (psychosexual disorder), not part of gender dysphoria.
Sexual dysfunction	Includes sexual desire disorders (hypoactive sexual desire or sexual aversion), sexual arousal disorders (erectile dysfunction), orgasmic disorders (anorgasmia, premature ejaculation), sexual pain disorders (dyspareunia, vaginismus). Differential diagnosis includes (PENIS): <ul style="list-style-type: none"> ▪ P psychological (if nighttime erections still occur) ▪ Endocrine (eg, diabetes, low testosterone) ▪ Neurogenic (eg, postoperative, spinal cord injury) ▪ Insufficient blood flow (eg, atherosclerosis) ▪ Substances (eg, antihypertensives, antidepressants, ethanol)

Sleep terror disorder

Periods of inconsolable terror with screaming in the middle of the night. Most common in children. Occurs during slow-wave/deep (stage N3) non-REM sleep with no memory of the arousal episode, as opposed to nightmares that occur during **REM** sleep (**re**mbering a scary dream). Triggers include emotional stress, fever, and lack of sleep. Usually self limited.

Enuresis

Nighttime urinary incontinence ≥ 2 times/week for ≥ 3 months in person > 5 years old. First-line treatment: behavioral modification (eg, scheduled voids, nighttime fluid restriction) and positive reinforcement. For refractory cases: bedwetting alarm, oral desmopressin (ADH analog; preferred over imipramine due to fewer side effects).

Narcolepsy

Excessive daytime sleepiness (despite awakening well-rested) with recurrent episodes of rapid-onset, overwhelming sleepiness ≥ 3 times/week for the last 3 months. Due to \downarrow orexin (hypocretin) production in lateral hypothalamus and dysregulated sleep-wake cycles. Associated with:

- Hypnagogic (just before **g**oing to sleep) or hypnopompic (just before awakening; get **p**omped up in the morning) hallucinations.
- Nocturnal and narcoleptic sleep episodes that start with REM sleep (sleep paralysis).
- Cataplexy (loss of all muscle tone following strong emotional stimulus, such as laughter).

Treatment: good sleep hygiene (scheduled naps, regular sleep schedule), daytime stimulants (eg, amphetamines, modafinil) and/or nighttime sodium oxybate (GHB).


Substance use disorder

Maladaptive pattern of substance use involving ≥ 2 of the following in the past year:

- Tolerance
- Withdrawal
- Intense, distracting cravings
- Using more, or longer, than intended
- Persistent desire but inability to cut down
- Time-consuming substance acquisition, use, or recovery
- Impaired functioning at work, school, or home
- Social or interpersonal conflicts
- Reduced recreational activities
- > 1 episode of use involving danger (eg, unsafe sex, driving while impaired)
- Continued use despite awareness of harm

Stages of change in overcoming addiction


1. **Precontemplation**—denying problem
2. **Contemplation**—acknowledging problem, but unwilling to change
3. **Preparation/determination**—preparing for behavioral changes
4. **Action/willpower**—changing behaviors
5. **Maintenance**—maintaining changes
6. **Relapse**—(if applicable) returning to old behaviors and abandoning changes



Psychiatric emergencies

	CAUSE	MANIFESTATION	TREATMENT
Serotonin syndrome	Any drug that ↑ 5-HT. Psychiatric drugs: MAOIs, SSRIs, SNRIs, TCAs, vilazodone, vortioxetine, bupirone Nonpsychiatric drugs: tramadol, ondansetron, triptans, linezolid, MDMA, dextromethorphan, meperidine, St. John's wort	3 A's: ↑ A ctivity (neuromuscular; eg, clonus, hyperreflexia, hypertonia, tremor, seizure), A utonomic instability (eg, hyperthermia, diaphoresis, diarrhea), A ltered mental status	Cyproheptadine (5-HT ₂ receptor antagonist)
Hypertensive crisis	Eating tyramine-rich foods (eg, aged cheeses, cured meats, wine, chocolate) while taking MAOIs	Hypertensive crisis (tyramine displaces other neurotransmitters [eg, NE] in the synaptic cleft → ↑ sympathetic stimulation)	Phentolamine
Neuroleptic malignant syndrome	Antipsychotics (typical > atypical) + genetic predisposition	Malignant FEVER: M yoalbuminuria, F ever, E ncephalopathy, V itals unstable, ↑ E nzymes (eg, CK), muscle R igidity ("lead pipe")	Dantrolene, dopamine agonist (eg, bromocriptine), discontinue causative agent
Delirium tremens	Alcohol withdrawal; occurs 2–4 days after last drink Classically seen in hospital setting when inpatient cannot drink	Altered mental status, hallucinations, autonomic hyperactivity, anxiety, seizures, tremors, psychomotor agitation, insomnia, nausea	Benzodiazepines (eg, chlordiazepoxide, lorazepam, diazepam)
Acute dystonia	Typical antipsychotics, anticonvulsants (eg, carbamazepine), metoclopramide	Sudden onset of muscle spasms, stiffness, and/or oculogyric crisis occurring hours to days after medication use; can lead to laryngospasm requiring intubation	Benzotropine or diphenhydramine
Lithium toxicity	↑ lithium dosage, ↓ renal elimination (eg, acute kidney injury), medications affecting clearance (eg, ACE inhibitors, thiazide diuretics, NSAIDs). Narrow therapeutic window.	Nausea, vomiting, slurred speech, hyperreflexia, seizures, ataxia, nephrogenic diabetes insipidus	Discontinue lithium, hydrate aggressively with isotonic sodium chloride, consider hemodialysis
Tricyclic antidepressant toxicity	TCA overdose	Respiratory depression, hyperpyrexia, prolonged QT Tri-CyCliC's: C onvulsions, C oma, C ardiotoxicity (arrhythmia due to Na ⁺ channel inhibition)	Supportive treatment, monitor ECG, NaHCO ₃ (prevents arrhythmia), activated charcoal

Psychoactive drug intoxication and withdrawal

DRUG	INTOXICATION	WITHDRAWAL
Depressants		
	Nonspecific: mood elevation, ↓ anxiety, sedation, behavioral disinhibition, respiratory depression.	Nonspecific: anxiety, tremor, seizures, insomnia.
Alcohol	Emotional lability, slurred speech, ataxia, coma, blackouts. Serum γ -glutamyltransferase (GGT)—sensitive indicator of alcohol use. AST value is 2×ALT value (“ ToAST 2 AL cohol”). Treatment: benzodiazepines.	 <p>Timeline of alcohol withdrawal symptoms:</p> <ul style="list-style-type: none"> 0-36 hours: Tremors, insomnia, diaphoresis, agitation, GI upset 12-48 hours: Alcoholic hallucinosis (usually visual), Withdrawal seizures 48-96 hours: Delirium tremens
Barbiturates	Low safety margin, marked respiratory depression. Treatment: symptom management (eg, assist respiration, ↑ BP).	Delirium, life-threatening cardiovascular collapse.
Benzodiazepines	Greater safety margin. Ataxia, minor respiratory depression. Treatment: flumazenil (benzodiazepine receptor antagonist, but rarely used as it can precipitate seizures).	Sleep disturbance, depression.
Opioids	Euphoria, respiratory and CNS depression, ↓ gag reflex, pupillary constriction (pinpoint pupils), seizures. Most common cause of drug overdose death. Treatment: naloxone.	Sweating, dilated pupils, piloerection (“cold turkey”), rhinorrhea, lacrimation, yawning, nausea, stomach cramps, diarrhea (“flu-like” symptoms). Treatment: symptom management, methadone, buprenorphine.
Inhalants	Disinhibition, euphoria, slurred speech, disturbed gait, disorientation, drowsiness.	Irritability, dysphoria, sleep disturbance, headache.
Stimulants		
	Nonspecific: mood elevation, ↓ appetite, psychomotor agitation, insomnia, cardiac arrhythmias, tachycardia, anxiety.	Nonspecific: post-use “crash,” including depression, lethargy, ↑ appetite, sleep disturbance, vivid nightmares.
Amphetamines	Euphoria, grandiosity, pupillary dilation, prolonged wakefulness, hyperalertness, hypertension, paranoia, fever, fractured teeth. Skin excoriations with methamphetamine use. Severe: cardiac arrest, seizures. Treatment: benzodiazepines for agitation and seizures.	
Caffeine	Palpitation, agitation, tremor, insomnia.	Headache, difficulty concentrating, flu-like symptoms.

Psychoactive drug intoxication and withdrawal (continued)

DRUG	INTOXICATION	WITHDRAWAL
Cocaine	Impaired judgment, pupillary dilation, hallucinations (including tactile), paranoia, angina, sudden cardiac death. Chronic use may lead to perforated nasal septum due to vasoconstriction and resulting ischemic necrosis. Treatment: benzodiazepines; consider mixed α -/ β -blocker (eg, labetalol) for hypertension and tachycardia. Pure β -blocker usage is controversial as a first-line therapy.	
Nicotine	Restlessness.	Irritability, anxiety, restlessness, ↓ concentration, ↑ appetite/weight. Treatment: nicotine patch, gum, or lozenges; bupropion/varenicline.
Hallucinogens		
Lysergic acid diethylamide	Perceptual distortion (visual, auditory), depersonalization, anxiety, paranoia, psychosis, flashbacks (usually nondisturbing).	
Marijuana (cannabinoid)	Euphoria, anxiety, paranoid delusions, perception of slowed time, impaired judgment, social withdrawal, ↑ appetite, dry mouth, conjunctival injection, hallucinations. Pharmaceutical form is dronabinol : used as antiemetic (chemotherapy) and appetite stimulant (in AIDS).	Irritability, anxiety, depression, insomnia, restlessness, ↓ appetite.
MDMA (ecstasy)	Hallucinogenic stimulant: euphoria, hallucinations, disinhibition, hyperactivity, ↑ thirst, bruxism, distorted sensory and time perception. Life-threatening effects include hypertension, tachycardia, hyperthermia, hyponatremia, serotonin syndrome.	Depression, fatigue, change in appetite, difficulty concentrating, anxiety.
Phencyclidine	Violence, impulsivity, psychomotor agitation, nystagmus, tachycardia, hypertension, analgesia, psychosis, delirium, seizures. Trauma is most common complication.	
Alcohol use disorder	Physiologic tolerance and dependence on alcohol with symptoms of withdrawal when intake is interrupted. Complications: vitamin B ₁ (thiamine) deficiency, alcoholic cirrhosis, hepatitis, pancreatitis, peripheral neuropathy, testicular atrophy. Treatment: naltrexone (reduces cravings), acamprosate, disulfiram (to condition the patient to abstain from alcohol use). Support groups such as Alcoholics Anonymous are helpful in sustaining abstinence and supporting patient and family.	
Wernicke-Korsakoff syndrome	Results from vitamin B ₁ deficiency. Symptoms can be precipitated by administering dextrose before vitamin B ₁ . Triad of confusion, ophthalmoplegia, ataxia (Wernicke encephalopathy). May progress to irreversible memory loss, confabulation, personality change (Korsakoff syndrome). Treatment: IV vitamin B ₁ (before dextrose).	

▶ PSYCHIATRY—PHARMACOLOGY

Psychotherapy

Behavioral therapy	Teaches patients how to identify and change maladaptive behaviors or reactions to stimuli. Examples include systematic desensitization for treatment of phobia.
Cognitive behavioral therapy	Teaches patients to recognize distortions in their thought processes, develop constructive coping skills, and ↓ maladaptive coping behaviors → greater emotional control and tolerance of distress. Examples include recognizing triggers for alcohol consumption.
Dialectical behavioral therapy	Designed for use in borderline personality disorder, but can be used in other psychiatric conditions as well (eg, depression).
Interpersonal therapy	Focused on improving interpersonal relationships and communication skills.
Supportive therapy	Utilizes empathy to help individuals during a time of hardship to maintain optimism or hope.

Preferred medications for selected psychiatric conditions

PSYCHIATRIC CONDITION	PREFERRED DRUGS
ADHD	Stimulants (methylphenidate, amphetamines)
Alcohol withdrawal	Benzodiazepines (eg, chlordiazepoxide, lorazepam, diazepam)
Bipolar disorder	Lithium, valproic acid, carbamazepine, lamotrigine, atypical antipsychotics
Bulimia nervosa	SSRIs
Depression	SSRIs
Generalized anxiety disorder	SSRIs, SNRIs
Obsessive-compulsive disorder	SSRIs, venlafaxine, clomipramine
Panic disorder	SSRIs, venlafaxine, benzodiazepines
PTSD	SSRIs, venlafaxine
Schizophrenia	Atypical antipsychotics
Social anxiety disorder	SSRIs, venlafaxine
Tourette syndrome	Performance only: β-blockers, benzodiazepines Antipsychotics (eg, fluphenazine, risperidone), tetrabenazine

Central nervous system stimulants

Methylphenidate, dextroamphetamine, methamphetamine, lisdexamfetamine.

MECHANISM	↑ catecholamines in the synaptic cleft, especially norepinephrine and dopamine.
CLINICAL USE	ADHD, narcolepsy, binge-eating disorder.
ADVERSE EFFECTS	Nervousness, agitation, anxiety, insomnia, anorexia, tachycardia, hypertension, weight loss, tics, bruxism.

Typical antipsychotics	Haloperidol, pimozide, trifluoperazine, fluphenazine, thioridazine, chlorpromazine.	
MECHANISM	Block dopamine D ₂ receptor (↑ cAMP).	
CLINICAL USE	Schizophrenia (1° positive symptoms), psychosis, bipolar disorder, delirium, Tourette syndrome, Huntington disease, OCD. Use with caution in dementia.	
POTENCY	<p>High potency: Haloperidol, Trifluoperazine, Fluphenazine (Hal Tries to Fly High)—more neurologic side effects (eg, extrapyramidal symptoms [EPS]).</p> <p>Low potency: Chlorpromazine, Thioridazine (Cheating Thieves are low)—more anticholinergic, antihistamine, α₁-blockade effects.</p>	
ADVERSE EFFECTS	<p>Lipid soluble → stored in body fat → slow to be removed from body.</p> <p>Endocrine: dopamine receptor antagonism → hyperprolactinemia → galactorrhea, oligomenorrhea, gynecomastia.</p> <p>Metabolic: dyslipidemia, weight gain, hyperglycemia.</p> <p>Antimuscarinic: dry mouth, constipation.</p> <p>Antihistamine: sedation.</p> <p>α₁-blockade: orthostatic hypotension.</p> <p>Cardiac: QT prolongation.</p> <p>Ophthalmologic: Chlorpromazine—Corneal deposits; Thioridazine—reTinal deposits.</p> <p>Neuroleptic malignant syndrome.</p> <p>Extrapyramidal symptoms— ADAPT:</p> <ul style="list-style-type: none"> ▪ Hours to days: Acute Dystonia (muscle spasm, stiffness, oculogyric crisis). Treatment: benztropine, diphenhydramine. ▪ Days to months: <ul style="list-style-type: none"> ▪ Akathisia (restlessness). Treatment: β-blockers, benztropine, benzodiazepines. ▪ Parkinsonism (bradykinesia). Treatment: benztropine, amantadine. ▪ Months to years: Tardive dyskinesia (chorea, especially orofacial). Treatment: atypical antipsychotics (eg, clozapine), valbenazine, deutetrabenazine. 	
Atypical antipsychotics	Aripiprazole, asenapine, clozapine, olanzapine, quetiapine, iloperidone, paliperidone, risperidone, lurasidone, ziprasidone.	
MECHANISM	Not completely understood. Most are 5-HT ₂ and D ₂ antagonists; aripiprazole is a D ₂ partial agonist. Varied effects on α and H ₁ receptors.	
CLINICAL USE	Schizophrenia—both positive and negative symptoms. Also used for bipolar disorder, OCD, anxiety disorders, depression, mania, Tourette syndrome.	Use clozapine for treatment-resistant schizophrenia or schizoaffective disorder and for suicidality in schizophrenia.
ADVERSE EFFECTS	<p>All—prolonged QT, fewer EPS and anticholinergic side effects than typical antipsychotics.</p> <p>“-apines”—metabolic syndrome (weight gain, diabetes, dyslipidemia).</p> <p>Clozapine—agranulocytosis (monitor WBCs frequently) and seizures (dose related).</p> <p>Risperidone—hyperprolactinemia (amenorrhea, galactorrhea, gynecomastia).</p>	<p>Olanzapine, clozapine → Obesity</p> <p>Must watch bone marrow clozely with clozapine.</p>

Lithium

MECHANISM	Not established; possibly related to inhibition of phosphoinositol cascade.
CLINICAL USE	Mood stabilizer for bipolar disorder; treats acute manic episodes and prevents relapse.
ADVERSE EFFECTS	Tremor, thyroid abnormalities (eg, hypothyroidism), polyuria (causes nephrogenic diabetes insipidus), teratogenesis. Causes Ebstein anomaly in newborn if taken by pregnant mother. Narrow therapeutic window requires close monitoring of serum levels. Almost exclusively excreted by kidneys; most is reabsorbed at PCT via Na ⁺ channels. Thiazides, NSAIDs, and other drugs affecting clearance are implicated in lithium toxicity.

LiTHIUM:


- L**ow **T**hyroid (hypothyroidism)
- H**ead (Ebstein anomaly)
- I**nsipidus (nephrogenic diabetes insipidus)
- U**nwanted **M**ovements (tremor)

Buspirone

MECHANISM	Stimulates 5-HT _{1A} receptors.
CLINICAL USE	Generalized anxiety disorder. Does not cause sedation, addiction, or tolerance. Begins to take effect after 1–2 weeks. Does not interact with alcohol (vs barbiturates, benzodiazepines).

I get **anxious** if the **bus** doesn't arrive at **one**, so I take **buspirone**.

Antidepressants



Selective serotonin reuptake inhibitors

	Fluoxetine, fluvoxamine, paroxetine, sertraline, escitalopram, citalopram.	
MECHANISM	Inhibit 5-HT reuptake.	It normally takes 4–8 weeks for antidepressants to show appreciable effect.
CLINICAL USE	Depression, generalized anxiety disorder, panic disorder, OCD, bulimia, binge-eating disorder, social anxiety disorder, PTSD, premature ejaculation, premenstrual dysphoric disorder.	
ADVERSE EFFECTS	Fewer than TCAs. Serotonin syndrome, GI distress, SIADH, sexual dysfunction (anorgasmia, ↓ libido).	

Serotonin-norepinephrine reuptake inhibitors

	Venlafaxine, desvenlafaxine, duloxetine, levomilnacipran, milnacipran.
MECHANISM	Inhibit 5-HT and NE reuptake.
CLINICAL USE	Depression, generalized anxiety disorder, diabetic neuropathy. Venlafaxine is also indicated for social anxiety disorder, panic disorder, PTSD, OCD. Duloxetine and milnacipran are also indicated for fibromyalgia.
ADVERSE EFFECTS	↑ BP, stimulant effects, sedation, nausea.

Tricyclic antidepressants

	Amitriptyline, nortriptyline, imipramine, desipramine, clomipramine, doxepin, amoxapine.
MECHANISM	TCAs inhibit 5-HT and NE reuptake.
CLINICAL USE	MDD, peripheral neuropathy, chronic neuropathic pain, migraine prophylaxis, OCD (clomipramine), nocturnal enuresis (imipramine, although adverse effects may limit use).
ADVERSE EFFECTS	Sedation, α_1 -blocking effects including postural hypotension, and atropine-like (anticholinergic) side effects (tachycardia, urinary retention, dry mouth). 3° TCAs (amitriptyline) have more anticholinergic effects than 2° TCAs (nortriptyline). Can prolong QT interval. Tri-CyCliC's: Convulsions, Coma, Cardiotoxicity (arrhythmia due to Na ⁺ channel inhibition); also respiratory depression, hyperpyrexia. Confusion and hallucinations are more common in the elderly due to anticholinergic side effects (2° amines [eg, nortriptyline] better tolerated). Treatment: NaHCO ₃ to prevent arrhythmia.

Monoamine oxidase inhibitors

	Tranylcypromine, Phenelzine, Isocarboxazid, Selegiline (selective MAO-B inhibitor). (MAO Takes Pride In Shanghai).
MECHANISM	Nonselective MAO inhibition → ↑ levels of amine neurotransmitters (norepinephrine, 5-HT, dopamine).
CLINICAL USE	Atypical depression, anxiety. Parkinson disease (selegiline).
ADVERSE EFFECTS	CNS stimulation; hypertensive crisis, most notably with ingestion of tyramine. Contraindicated with SSRIs, TCAs, St. John's wort, meperidine, dextromethorphan, linezolid (to avoid precipitating serotonin syndrome). Wait 2 weeks after stopping MAOIs before starting serotonergic drugs or stopping dietary restrictions.

Atypical antidepressants

Bupropion	Inhibits NE and DA reuptake. Also used for smoking cessation. Toxicity: stimulant effects (tachycardia, insomnia), headache, seizures in patients with bulimia and anorexia nervosa. Favorable sexual side effect profile.
Mirtazapine	α_2 -antagonist (\uparrow release of NE and 5-HT), potent 5-HT ₂ and 5-HT ₃ receptor antagonist, and H ₁ antagonist. Toxicity: sedation (which may be desirable in depressed patients with insomnia), \uparrow appetite, weight gain (which may be desirable in underweight patients), dry mouth.
Trazodone	Primarily blocks 5-HT ₂ , α_1 -adrenergic, and H ₁ receptors; also weakly inhibits 5-HT reuptake. Used primarily for insomnia, as high doses are needed for antidepressant effects. Toxicity: sedation, nausea, priapism, postural hypotension. Think tra ZZZ obone due to sedative and male-specific side effects.
Varenicline	Nicotinic ACh receptor partial agonist. Used for smoking cessation. Toxicity: sleep disturbance, depressed mood, suicidal ideation. Varen icline helps nicotine cravings decline .
Vilazodone	Inhibits 5-HT reuptake; 5-HT _{1A} receptor partial agonist. Used for MDD. Toxicity: headache, diarrhea, nausea, anticholinergic effects. May cause serotonin syndrome if taken with other serotonergic agents.
Vortioxetine	Inhibits 5-HT reuptake; 5-HT _{1A} receptor agonist and 5-HT ₃ receptor antagonist. Used for MDD. Toxicity: nausea, sexual dysfunction, sleep disturbances, anticholinergic effects. May cause serotonin syndrome if taken with other serotonergic agents.

Opioid detoxification and relapse prevention

Intravenous drug users at \uparrow risk for hepatitis, HIV, abscesses, bacteremia, right-heart endocarditis.

Methadone	Long-acting oral opiate used for heroin detoxification or long-term maintenance therapy.
Buprenorphine	Sublingual form (partial agonist) used to prevent relapse.
Naloxone	Short-acting opioid antagonist given IM, IV, or as a nasal spray to treat acute opioid overdose, particularly to reverse respiratory and CNS depression.
Naltrexone	Long-acting oral opioid antagonist used after detoxification to prevent relapse. Use nalt rex one for the long trex back to sobriety.

HIGH-YIELD SYSTEMS

Renal

“But I know all about love already. I know precious little still about kidneys.”

—Aldous Huxley, *Antic Hay*

“This too shall pass. Just like a kidney stone.”

—Hunter Madsen

“I drink too much. The last time I gave a urine sample it had an olive in it.”

—Rodney Dangerfield

Being able to understand and apply renal physiology will be critical for the exam. Important topics include electrolyte disorders, acid-base derangements, glomerular disorders (including histopathology), acute and chronic kidney disease, urine casts, diuretics, ACE inhibitors, and AT-II receptor blockers. Renal anomalies associated with various congenital defects are also high-yield associations to think about when evaluating pediatric vignettes.

▶ Embryology	578
▶ Anatomy	580
▶ Physiology	581
▶ Pathology	594
▶ Pharmacology	607

▶ RENAL—EMBRYOLOGY

Kidney embryology



Pronephros—week 4; then degenerates.

Mesonephros—functions as interim kidney for 1st trimester; later contributes to male genital system.

Metanephros—permanent; first appears in 5th week of gestation; nephrogenesis continues through weeks 32–36 of gestation.

- Ureteric bud (metanephric diverticulum)—derived from caudal end of mesonephric duct; gives rise to ureter, pelvises, calyces, collecting ducts; fully canalized by 10th week
- Metanephric mesenchyme (ie, metanephric blastema)—ureteric bud interacts with this tissue; interaction induces differentiation and formation of glomerulus through to distal convoluted tubule (DCT)
- Aberrant interaction between these 2 tissues may result in several congenital malformations of the kidney (eg, renal agenesis, multicystic dysplastic kidney)

Ureteropelvic junction—last to canalize
→ congenital obstruction. Most common cause of prenatal hydronephrosis. Detected by prenatal ultrasound.

**Potter sequence (syndrome)**

Oligohydramnios → compression of developing fetus → limb deformities, facial anomalies (eg, low-set ears and retrognathia **A**, flattened nose), compression of chest and lack of amniotic fluid aspiration into fetal lungs → pulmonary hypoplasia (cause of death).


Causes include ARPKD, obstructive uropathy (eg, posterior urethral valves), bilateral renal agenesis, chronic placental insufficiency.

Babies who can't "Pee" in utero develop Potter sequence.

POTTER sequence associated with:

- P**ulmonary hypoplasia
- O**ligohydramnios (trigger)
- T**wisted face
- T**wisted skin
- E**xtrernity defects
- R**enal failure (in utero)

Horseshoe kidney



Inferior poles of both kidneys fuse abnormally **A**. As they ascend from pelvis during fetal development, horseshoe kidneys get trapped under inferior mesenteric artery and remain low in the abdomen. Kidneys function normally. Associated with hydronephrosis (eg, ureteropelvic junction obstruction), renal stones, infection, ↑ risk of renal cancer.

Higher incidence in chromosomal aneuploidy (eg, Turner syndrome, trisomies 13, 18, 21).



Congenital solitary functioning kidney

Condition of being born with only one functioning kidney. Majority asymptomatic with compensatory hypertrophy of contralateral kidney, but anomalies in contralateral kidney are common. Often diagnosed prenatally via ultrasound.

Unilateral renal agenesis

Ureteric bud fails to develop and induce differentiation of metanephric mesenchyme → complete absence of kidney and ureter.

Multicystic dysplastic kidney


Ureteric bud fails to induce differentiation of metanephric mesenchyme → nonfunctional kidney consisting of cysts and connective tissue. Predominantly nonhereditary and usually unilateral; bilateral leads to Potter sequence.

Duplex collecting system

Bifurcation of ureteric bud before it enters the metanephric blastema creates a Y-shaped bifid ureter. Duplex collecting system can alternatively occur through two ureteric buds reaching and interacting with metanephric blastema. Strongly associated with vesicoureteral reflux and/or ureteral obstruction, ↑ risk for UTIs.


Posterior urethral valves

Membrane remnant in the posterior urethra in males; its persistence can lead to urethral obstruction. Can be diagnosed prenatally by bilateral hydronephrosis and dilated or thick-walled bladder on ultrasound. Most common cause of bladder outlet obstruction in male infants. Associated with oligohydramnios in cases of severe obstruction.




▶ RENAL—ANATOMY

Kidney anatomy and glomerular structure



Cross-section of kidney



Cross-section of glomerulus

Left kidney is taken during living donor transplantation because it has a longer renal vein.


Afferent = **A**rriving.


Efferent = **E**xiting.

Renal blood flow: renal artery → segmental artery → interlobar artery → arcuate artery → interlobular artery → afferent arteriole → glomerulus **A** → efferent arteriole → vasa recta/peritubular capillaries → venous outflow.

Left renal vein receives two additional veins: left suprarenal and left gonadal veins.

Despite high overall renal blood flow, renal medulla receives significantly less blood flow than renal cortex → very sensitive to hypoxia → vulnerable to ischemic damage.



Course of ureters

Course of ureter **A**: arises from renal pelvis, travels under gonadal arteries → **over** common iliac artery → **under** uterine artery/vas deferens (retroperitoneal).

Gynecologic procedures (eg, ligation of uterine or ovarian vessels) may damage ureter → ureteral obstruction or leak.



Bladder contraction compresses the intravesical ureter, preventing urine reflux.

Blood supply to ureter:

- Proximal—renal arteries
- Middle—gonadal artery, aorta, common and internal iliac arteries
- Distal—internal iliac and superior vesical arteries

3 common points of ureteral obstruction: ureteropelvic junction, pelvic inlet, ureterovesical junction.

Water (ureters) flows **over** the iliacs and **under** the bridge (uterine artery or vas deferens).

**► RENAL—PHYSIOLOGY****Fluid compartments**

HIKIN: **H**igh **K**⁺ **I**ntracellularly.

60–40–20 rule (% of body weight for average person):


- 60% total body water
- 40% ICF, mainly composed of K⁺, Mg²⁺, organic phosphates (eg, ATP)
- 20% ECF, mainly composed of Na⁺, Cl⁻, HCO₃⁻, albumin

Plasma volume can be measured by radiolabeling albumin.

Extracellular volume can be measured by inulin or mannitol.

Serum osmolality = 285–295 mOsm/kg H₂O.

Plasma volume = TBW × (1 – Hct).

Glomerular filtration barrier

Responsible for filtration of plasma according to size and charge selectivity.

Composed of:

- Fenestrated capillary endothelium
- Basement membrane with type IV collagen chains and heparan sulfate
- Visceral epithelial layer consisting of podocyte foot processes (FPs) **A**

Charge barrier—all 3 layers contain ⊖ charged glycoproteins that prevent entry of ⊖ charged molecules (eg, albumin).

Size barrier—fenestrated capillary endothelium (prevents entry of > 100 nm molecules/blood cells); podocyte foot processes interpose with glomerular basement membrane (GBM); slit diaphragm (prevents entry of molecules > 50–60 nm).

Renal clearance

$C_x = (U_x V)/P_x$ = volume of plasma from which the substance is completely cleared in the urine per unit time.

If $C_x < \text{GFR}$: net tubular reabsorption and/or not freely filtered.

If $C_x > \text{GFR}$: net tubular secretion of X.

If $C_x = \text{GFR}$: no net secretion or reabsorption.

C_x = clearance of X (mL/min).

U_x = urine concentration of X (eg, mg/mL).

P_x = plasma concentration of X (eg, mg/mL).

V = urine flow rate (mL/min).

Glomerular filtration rate


Inulin clearance can be used to calculate GFR because it is freely filtered and is neither reabsorbed nor secreted.

$$C_{\text{inulin}} = \text{GFR} = U_{\text{inulin}} \times V / P_{\text{inulin}} \\ = K_f [(P_{\text{GC}} - P_{\text{BS}}) - (\pi_{\text{GC}} - \pi_{\text{BS}})]$$

(GC = glomerular capillary; BS = Bowman space; π_{BS} normally equals zero; K_f = filtration coefficient).

Normal GFR \approx 100 mL/min.

Creatinine clearance is an approximate measure of GFR. Slightly overestimates GFR because creatinine is moderately secreted by renal tubules.

**Effective renal plasma flow**

Effective renal plasma flow (eRPF) can be estimated using *para*-aminohippuric acid (PAH) clearance. Between filtration and secretion, there is nearly 100% excretion of all PAH that enters the kidney.

$$e\text{RPF} = U_{\text{PAH}} \times V / P_{\text{PAH}} = C_{\text{PAH}}$$

Renal blood flow (RBF) = RPF/(1 - Hct). Usually 20–25% of cardiac output.

eRPF underestimates true renal plasma flow (RPF) slightly.

Filtration

Filtration fraction (FF) = GFR/RPF.

Normal FF = 20%.


Filtered load (mg/min) = GFR (mL/min) × plasma concentration (mg/mL).

GFR can be estimated with creatinine clearance.

RPF is best estimated with PAH clearance.

Prostaglandins Dilate Afferent arteriole (PDA).

Angiotensin II Constricts Efferent arteriole (ACE).



Changes in glomerular dynamics

	GFR	RPF	FF (GFR/RPF)
Afferent arteriole constriction	↓	↓	—
Efferent arteriole constriction	↑	↓	↑
↑ plasma protein concentration	↓	—	↓
↓ plasma protein concentration	↑	—	↑
Constriction of ureter	↓	—	↓
Dehydration	↓	↓↓	↑

Calculation of reabsorption and secretion rate

Filtered load = $GFR \times P_x$.

Excretion rate = $V \times U_x$.

Reabsorption rate = filtered – excreted.

Secretion rate = excreted – filtered.

Fe_{Na} = fractional excretion of sodium.

$$Fe_{Na} = \frac{Na^+ \text{ excreted}}{Na^+ \text{ filtered}} = \frac{V \times U_{Na}}{GFR \times P_{Na}} = \frac{P_{Cr} \times U_{Na}}{U_{Cr} \times P_{Na}} \text{ where } GFR = \frac{U_{Cr} \times V}{P_{Cr}}$$

Glucose clearance

Glucose at a normal plasma level (range 60–120 mg/dL) is completely reabsorbed in proximal convoluted tubule (PCT) by Na^+ /glucose cotransport.

In adults, at plasma glucose of ~ 200 mg/dL, glucosuria begins (threshold). At rate of ~ 375 mg/min, all transporters are fully saturated (T_m).


Normal pregnancy is associated with ↑ GFR.

With ↑ filtration of all substances, including glucose, the glucose threshold occurs at lower plasma glucose concentrations → glucosuria at normal plasma glucose levels.


Sodium-glucose cotransporter 2 (SGLT2) inhibitors (eg, -flozin drugs) result in glucosuria at plasma concentrations < 200 mg/dL.

Glucosuria is an important clinical clue to diabetes mellitus.

Splay phenomenon— T_m for glucose is reached gradually rather than sharply due to the heterogeneity of nephrons (ie, different T_m points); represented by the portion of the titration curve between threshold and T_m .




Nephron transport physiology




Early PCT—contains brush border. Reabsorbs all glucose and amino acids and most HCO_3^- , Na^+ , Cl^- , PO_4^{3-} , K^+ , H_2O , and uric acid. Isotonic absorption. Generates and secretes NH_3 , which enables the kidney to secrete more H^+ .

PTH— inhibits $\text{Na}^+/\text{PO}_4^{3-}$ cotransport $\rightarrow \uparrow \text{PO}_4^{3-}$ excretion.
 AT II— stimulates Na^+/H^+ exchange $\rightarrow \uparrow \text{Na}^+$, H_2O , and HCO_3^- reabsorption (permitting contraction alkalosis).
 65–80% Na^+ and H_2O reabsorbed.

Thin descending loop of Henle— passively reabsorbs H_2O via medullary hypertonicity (impermeable to Na^+).
 Concentrating segment. Makes urine hypertonic.




Thick ascending loop of Henle— reabsorbs Na^+ , K^+ , and Cl^- . Indirectly induces paracellular reabsorption of Mg^{2+} and Ca^{2+} through \oplus lumen potential generated by K^+ backleak. Impermeable to H_2O . Makes urine less concentrated as it ascends.
 10–20% Na^+ reabsorbed.



Early DCT— reabsorbs Na^+ , Cl^- . Impermeable to H_2O .

Makes urine fully dilute (hypotonic).

PTH— $\uparrow \text{Ca}^{2+}/\text{Na}^+$ exchange $\rightarrow \uparrow \text{Ca}^{2+}$ reabsorption.
 5–10% Na^+ reabsorbed.




Collecting tubule— reabsorbs Na^+ in exchange for secreting K^+ and H^+ (regulated by aldosterone).

Aldosterone— acts on mineralocorticoid receptor \rightarrow mRNA \rightarrow protein synthesis. In principal cells: \uparrow apical K^+ conductance, \uparrow Na^+/K^+ pump, \uparrow epithelial Na^+ channel (ENaC) activity \rightarrow lumen negativity \rightarrow K^+ secretion. In α -intercalated cells: lumen negativity \rightarrow \uparrow H^+ ATPase activity \rightarrow \uparrow H^+ secretion \rightarrow \uparrow $\text{HCO}_3^-/\text{Cl}^-$ exchanger activity.

ADH— acts at V_2 receptor \rightarrow insertion of aquaporin H_2O channels on apical side.

3–5% Na^+ reabsorbed.

Renal tubular defects Order: **Fanconi's BaGeLS**


	DEFECTS	EFFECTS	CAUSES	NOTES
Fanconi syndrome	Generalized reabsorption defect in PCT → ↑ excretion of amino acids, glucose, HCO_3^- , and PO_4^{3-} , and all substances reabsorbed by the PCT	May lead to metabolic acidosis (proximal RTA), hypophosphatemia, osteopenia	Hereditary defects (eg, Wilson disease, tyrosinemia, glycogen storage disease), ischemia, multiple myeloma, nephrotoxins/drugs (eg, ifosfamide, cisplatin), lead poisoning	
Bartter syndrome	Reabsorption defect in thick ascending loop of Henle (affects $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporter)	Metabolic alkalosis, hypokalemia, hypercalciuria	Autosomal recessive	Presents similarly to chronic loop diuretic use
Gitelman syndrome	Reabsorption defect of NaCl in DCT	Metabolic alkalosis, hypomagnesemia, hypokalemia, hypocalciuria	Autosomal recessive	Presents similarly to lifelong thiazide diuretic use Less severe than Bartter syndrome
Liddle syndrome	Gain of function mutation → ↓ Na^+ channel degradation → ↑ Na^+ reabsorption in collecting tubules	Metabolic alkalosis, hypokalemia, hypertension, ↓ aldosterone	Autosomal dominant	Presents similarly to hyperaldosteronism, but aldosterone is nearly undetectable Treatment: amiloride
Syndrome of Apparent Mineralocorticoid Excess	Cortisol activates mineralocorticoid receptors. 11β -HSD converts cortisol to cortisone (inactive on these receptors) Hereditary 11β -HSD deficiency → ↑ cortisol → ↑ mineralocorticoid receptor activity	Metabolic alkalosis, hypokalemia, hypertension ↓ serum aldosterone level; cortisol tries to be the SAME as aldosterone	Autosomal recessive Can acquire disorder from glycyrrhetic acid (present in licorice), which blocks activity of 11β -hydroxysteroid dehydrogenase	Treatment: K^+ -sparing diuretics (↓ mineralocorticoid effects) or corticosteroids (exogenous corticosteroid ↓ endogenous cortisol production → ↓ mineralocorticoid receptor activation)

Relative concentrations along proximal convoluted tubules

[TF/P] > 1
when solute is reabsorbed less quickly than water or when solute is secreted


[TF/P] = 1
when solute and water are reabsorbed at the same rate

[TF/P] < 1
when solute is reabsorbed more quickly than water



Tubular inulin ↑ in concentration (but not amount) along the PCT as a result of water reabsorption. Cl⁻ reabsorption occurs at a slower rate than Na⁺ in early PCT and then matches the rate of Na⁺ reabsorption more distally. Thus, its relative concentration ↑ before it plateaus.

Renin-angiotensin-aldosterone system

**Renin**

Secreted by JG cells in response to ↓ renal perfusion pressure (detected by renal baroreceptors in afferent arteriole), ↑ renal sympathetic discharge (β_1 effect), and ↓ NaCl delivery to macula densa cells.

AT II

Helps maintain blood volume and blood pressure. Affects baroreceptor function; limits reflex bradycardia, which would normally accompany its pressor effects.

ANP, BNP

Released from atria (ANP) and ventricles (BNP) in response to ↑ volume; inhibits renin-angiotensin-aldosterone system; relaxes vascular smooth muscle via cGMP → ↑ GFR, ↓ renin. Dilates afferent arteriole, promotes natriuresis.

ADH

Primarily regulates serum osmolality; also responds to low blood volume states. Stimulates reabsorption of water in collecting ducts. Also stimulates reabsorption of urea in collecting ducts to maximize corticopapillary osmotic gradient.

Aldosterone

Primarily regulates ECF volume and Na⁺ content; ↑ release in ↓ blood volume states. Responds to hyperkalemia by ↑ K⁺ excretion.

Juxtaglomerular apparatus

Consists of mesangial cells, JG cells (modified smooth muscle of afferent arteriole), and the macula densa (NaCl sensor, located at distal end of loop of Henle). JG cells secrete renin in response to ↓ renal blood pressure and ↑ sympathetic tone (β_1). Macula densa cells sense ↓ NaCl delivery to DCT → ↑ renin release → efferent arteriole vasoconstriction → ↑ GFR.

JGA maintains GFR via renin-angiotensin-aldosterone system.

In addition to vasodilatory properties, β -blockers can decrease BP by inhibiting β_1 -receptors of the JGA → ↓ renin release.

Kidney endocrine functions


Erythropoietin

Released by interstitial cells in peritubular capillary bed in response to hypoxia.

Stimulates RBC proliferation in bone marrow. Administered for anemia secondary to chronic kidney disease. ↑ risk of HTN.

Calciferol (vitamin D)

PCT cells convert 25-OH vitamin D₃ to 1,25-(OH)₂ vitamin D₃ (calcitriol, active form).



Prostaglandins

Paracrine secretion vasodilates the afferent arterioles to ↑ RBF.

NSAIDs block renal-protective prostaglandin synthesis → constriction of afferent arteriole and ↓ GFR; this may result in acute kidney injury in low renal blood flow states.

Dopamine

Secreted by PCT cells, promotes natriuresis. At low doses; dilates interlobular arteries, afferent arterioles, efferent arterioles → ↑ RBF, little or no change in GFR. At higher doses; acts as vasoconstrictor.

Hormones acting on kidney

Atrial natriuretic peptide


Secreted in response to ↑ atrial pressure. Causes ↑ GFR and ↑ Na⁺ filtration with no compensatory Na⁺ reabsorption in distal nephron. Net effect: Na⁺ loss and volume loss.

Angiotensin II

Synthesized in response to ↓ BP. Causes efferent arteriole constriction → ↑ GFR and ↑ FF but with compensatory Na⁺ reabsorption in proximal and distal nephron. Net effect: preservation of renal function (↑ FF) in low-volume state with simultaneous Na⁺ reabsorption (both proximal and distal) to maintain circulating volume.

Parathyroid hormone

Secreted in response to ↓ plasma [Ca²⁺], ↑ plasma [PO₄³⁻], or ↓ plasma 1,25-(OH)₂D₃. Causes ↑ [Ca²⁺] reabsorption (DCT), ↓ [PO₄³⁻] reabsorption (PCT), and ↑ 1,25-(OH)₂D₃ production (↑ Ca²⁺ and PO₄³⁻ absorption from gut via vitamin D).



Aldosterone

Secreted in response to ↓ blood volume (via AT II) and ↑ plasma [K⁺]; causes ↑ Na⁺ reabsorption, ↑ K⁺ secretion, ↑ H⁺ secretion.

ADH (vasopressin)

Secreted in response to ↑ plasma osmolarity and ↓ blood volume. Binds to receptors on principal cells, causing ↑ number of aquaporins and ↑ H₂O reabsorption. ↑ reabsorption of urea in collecting ducts to maximize corticopapillary osmotic gradient.



Potassium shifts

SHIFTS K⁺ INTO CELL (CAUSING HYPOKALEMIA)

Hypo-osmolarity

Alkalosis

β-adrenergic agonist (↑ Na⁺/K⁺ ATPase)

Insulin (↑ Na⁺/K⁺ ATPase)

Insulin shifts K⁺ into cells

SHIFTS K⁺ OUT OF CELL (CAUSING HYPERKALEMIA)

Digitalis (blocks Na⁺/K⁺ ATPase)

HyperOsmolarity

Lysis of cells (eg, crush injury, rhabdomyolysis, tumor lysis syndrome)

Acidosis

β-blocker

High blood Sugar (insulin deficiency)

Succinylcholine (↑ risk in burns/muscle trauma)

Hyperkalemia? **DO LAβSS**

Electrolyte disturbances

ELECTROLYTE	LOW SERUM CONCENTRATION	HIGH SERUM CONCENTRATION
Sodium	Nausea, malaise, stupor, coma, seizures	Irritability, stupor, coma
Potassium	U waves and flattened T waves on ECG, arrhythmias, muscle cramps, spasm, weakness	Wide QRS and peaked T waves on ECG, arrhythmias, muscle weakness
Calcium	Tetany, seizures, QT prolongation, twitching (eg, Chvostek sign), spasm (eg, Trousseau sign)	Stones (renal), bones (pain), groans (abdominal pain), thrones (↑ urinary frequency), psychiatric overtones (anxiety, altered mental status)
Magnesium	Tetany, torsades de pointes, hypokalemia, hypocalcemia (when $[Mg^{2+}] < 1.0$ mEq/L)	↓ DTRs, lethargy, bradycardia, hypotension, cardiac arrest, hypocalcemia
Phosphate	Bone loss, osteomalacia (adults), rickets (children)	Renal stones, metastatic calcifications, hypocalcemia

Features of renal disorders

CONDITION	BLOOD PRESSURE	PLASMA RENIN	ALDOSTERONE	SERUM Mg^{2+}	URINE Ca^{2+}
SIADH	—/↑	↓	↓		
Primary hyperaldosteronism	↑	↓	↑		
Renin-secreting tumor	↑	↑	↑		
Bartter syndrome		↑	↑		↑
Gitelman syndrome		↑	↑	↓	↓
Liddle syndrome, syndrome of apparent mineralocorticoid excess	↑	↓	↓		

↑ ↓ = important differentiating feature.

Acid-base physiology

	pH	Pco ₂	[HCO ₃ ⁻]	COMPENSATORY RESPONSE
Metabolic acidosis	↓	↓	↓	Hyperventilation (immediate)
Metabolic alkalosis	↑	↑	↑	Hypoventilation (immediate)
Respiratory acidosis	↓	↑	↑	↑ renal [HCO ₃ ⁻] reabsorption (delayed)
Respiratory alkalosis	↑	↓	↓	↓ renal [HCO ₃ ⁻] reabsorption (delayed)


Key: ↓ ↑ = compensatory response.

Henderson-Hasselbalch equation: $pH = 6.1 + \log \frac{[HCO_3^-]}{0.03 Pco_2}$

Predicted respiratory compensation for a simple metabolic acidosis can be calculated using the Winters formula. If measured Pco₂ > predicted Pco₂ → concomitant respiratory acidosis; if measured Pco₂ < predicted Pco₂ → concomitant respiratory alkalosis:

$Pco_2 = 1.5 [HCO_3^-] + 8 \pm 2$

Acidosis and alkalosis



Renal tubular acidosis

Disorder of the renal tubules that causes normal anion gap (hyperchloremic) metabolic acidosis.

RTA TYPE	DEFECT	URINE PH	SERUM K ⁺	CAUSES	ASSOCIATIONS
Distal renal tubular acidosis (type 1)	Inability of α -intercalated cells to secrete H ⁺ → no new HCO ₃ ⁻ is generated → metabolic acidosis	> 5.5	↓	Amphotericin B toxicity, analgesic nephropathy, congenital anomalies (obstruction) of urinary tract, autoimmune diseases (eg, SLE)	↑ risk for calcium phosphate kidney stones (due to ↑ urine pH and ↑ bone turnover related to buffering)
Proximal renal tubular acidosis (type 2)	Defect in PCT HCO ₃ ⁻ reabsorption → ↑ excretion of HCO ₃ ⁻ in urine → metabolic acidosis Urine can be acidified by α -intercalated cells in collecting duct, but not enough to overcome ↑ HCO ₃ ⁻ excretion	> 5.5 when resorptive threshold for serum HCO ₃ ⁻ exceeded; < 5.5 when HCO ₃ ⁻ depleted below resorptive threshold	↓	Fanconi syndrome, multiple myeloma, carbonic anhydrase inhibitors	↑ risk for hypophosphatemic rickets (in Fanconi syndrome)
Hyperkalemic tubular acidosis (type 4)	Hypoaldosteronism or aldosterone resistance; hyperkalemia → ↓ NH ₃ synthesis in PCT → ↓ NH ₄ ⁺ excretion	< 5.5 (or variable)	↑	↓ aldosterone production (eg, diabetic hyporeninism, ACE inhibitors, ARBs, NSAIDs, heparin, cyclosporine, adrenal insufficiency) or aldosterone resistance (eg, K ⁺ -sparing diuretics, nephropathy due to obstruction, TMP-SMX)	


▶ RENAL—PATHOLOGY

Casts in urine	Presence of casts indicates that hematuria/pyuria is of glomerular or renal tubular origin. Bladder cancer, kidney stones → hematuria, no casts. Acute cystitis → pyuria, no casts.
RBC casts A	Glomerulonephritis, hypertensive emergency.
WBC casts B	Tubulointerstitial inflammation, acute pyelonephritis, transplant rejection.
Granular casts C	Acute tubular necrosis (ATN). Can be “muddy brown” in appearance.
Fatty casts (“oval fat bodies”)	Nephrotic syndrome. Associated with “Maltese cross” sign D .
Waxy casts	End-stage renal disease/chronic kidney disease.
Hyaline casts E	Nonspecific, can be a normal finding. Form via solidification of Tamm–Horsfall mucoprotein (secreted by renal tubular cells).

**Nomenclature of glomerular disorders**


TYPE	CHARACTERISTICS	EXAMPLE
Focal	< 50% of glomeruli are involved	Focal segmental glomerulosclerosis
Diffuse	> 50% of glomeruli are involved	Diffuse proliferative glomerulonephritis
Proliferative	Hypercellular glomeruli	Membranoproliferative glomerulonephritis
Membranous	Thickening of glomerular basement membrane (GBM)	Membranous nephropathy
Primary glomerular disease	1° disease of the kidney specifically impacting the glomeruli	Minimal change disease
Secondary glomerular disease	Systemic disease or disease of another organ system that also impacts the glomeruli	SLE, diabetic nephropathy

Glomerular diseases




TYPE	ETIOLOGY	CLINICAL PRESENTATION	EXAMPLES
Nephritic syndrome	Glomerular inflammation → GBM damage → loss of RBCs into urine → hematuria	Hematuria, RBC casts in urine ↓ GFR → oliguria, azotemia, ↑ renin release, HTN Proteinuria often in the subnephrotic range (< 3.5 g/day) but in severe cases may be in nephrotic range	<ul style="list-style-type: none"> Acute poststreptococcal glomerulonephritis Rapidly progressive glomerulonephritis IgA nephropathy (Berger disease) Alport syndrome Membranoproliferative glomerulonephritis
Nephrotic syndrome	Podocyte damage → impaired charge barrier → proteinuria	Massive proteinuria (> 3.5 g/day) with hypoalbuminemia, edema Frothy urine with fatty casts Associated with hypercoagulable state due to antithrombin III loss in urine and ↑ risk of infection (loss of IgGs in urine and soft tissue compromise by edema)	May be 1° (eg, direct podocyte damage) or 2° (podocyte damage from systemic process): <ul style="list-style-type: none"> Focal segmental glomerulosclerosis (1° or 2°) Minimal change disease (1° or 2°) Membranous nephropathy (1° or 2°) Amyloidosis (2°) Diabetic glomerulonephropathy (2°)
Nephritic-nephrotic syndrome	Severe GBM damage → loss of RBCs into urine + impaired charge barrier → hematuria + proteinuria	Nephrotic-range proteinuria (> 3.5 g/day) and concomitant features of nephrotic syndrome	Can occur with any form of nephritic syndrome, but is most common with: <ul style="list-style-type: none"> Diffuse proliferative glomerulonephritis Membranoproliferative glomerulonephritis

Nephritic syndrome	Nephritic syndrome = Inflammatory process.
Acute poststreptococcal glomerulonephritis	<p>Most frequently seen in children. ~ 2–4 weeks after group A streptococcal infection of pharynx or skin. Resolves spontaneously in most children; may progress to renal insufficiency in adults. Type III hypersensitivity reaction. Presents with peripheral and periorbital edema, tea or cola-colored urine, HTN. ⊕ strep titers/serologies, ↓ complement levels (C3) due to consumption.</p> <ul style="list-style-type: none"> LM—glomeruli enlarged and hypercellular A IF—(“starry sky”) granular appearance (“lumpy-bumpy”) B due to IgG, IgM, and C3 deposition along GBM and mesangium EM—subepithelial IC humps
Rapidly progressive (crescentic) glomerulonephritis	<p>Poor prognosis, rapidly deteriorating renal function (days to weeks).</p> <ul style="list-style-type: none"> LM—crescent moon shape C. Crescents consist of fibrin and plasma proteins (eg, C3b) with glomerular parietal cells, monocytes, macrophages <p>Several disease processes may result in this pattern which may be delineated via IF pattern.</p> <ul style="list-style-type: none"> Linear IF due to antibodies to GBM and alveolar basement membrane: Goodpasture syndrome—hematuria/hemoptysis; type II hypersensitivity reaction. Treatment: plasmapheresis Negative IF/Pauci-immune (no Ig/C3 deposition): granulomatosis with polyangiitis (Wegener)—PR3-ANCA/c-ANCA, eosinophilic granulomatosis with polyangiitis (Churg-Strauss) or Microscopic polyangiitis—MPO-ANCA/p-ANCA Granular IF—PSGN or DPGN
Diffuse proliferative glomerulonephritis	<p>Often due to SLE (think “wire lupus”). DPGN and MPGN often present as nephrotic syndrome and nephritic syndrome concurrently.</p> <ul style="list-style-type: none"> LM—“wire looping” of capillaries D IF—granular; EM—subendothelial, sometimes subepithelial or intramembranous IgG-based ICs often with C3 deposition
IgA nephropathy (Berger disease)	<p>Episodic hematuria that usually occurs concurrently with respiratory or GI tract infections (IgA is secreted by mucosal linings). Renal pathology of IgA vasculitis (HSP).</p> <ul style="list-style-type: none"> LM—mesangial proliferation IF—IgA-based IC deposits in mesangium; EM—mesangial IC deposition
Alport syndrome	<p>Mutation in type IV collagen → thinning and splitting of glomerular basement membrane. Most commonly X-linked dominant. Eye problems (eg, retinopathy, anterior lenticonus), glomerulonephritis, sensorineural deafness; “can’t see, can’t pee, can’t hear a bee.”</p> <ul style="list-style-type: none"> EM—“basket-weave” appearance due to irregular thickening of GBM
Membrano-proliferative glomerulonephritis	<p>MPGN is a nephritic syndrome that often co-presents with nephrotic syndrome. Type I may be 2° to hepatitis B or C infection. May also be idiopathic.</p> <ul style="list-style-type: none"> Subendothelial IC deposits with granular IF <p>Type II is associated with C3 nephritic factor (IgG autoantibody that stabilizes C3 convertase → persistent complement activation → ↓ C3 levels).</p> <ul style="list-style-type: none"> Intramembranous deposits, also called dense deposit disease <p>Both types: mesangial ingrowth → GBM splitting → “tram-track” on H&E and PAS E stains.</p>



Nephrotic syndrome	Nephrotic syndrome—massive proteinuria (> 3.5 g/day)
Minimal change disease	<p>Also known as lipid nephrosis. Most common cause of nephrotic syndrome in children. Often 1° (Idiopathic) and may be triggered by recent Infection, Immunization, Immune stimulus (4 I's of MCD). Rarely, may be 2° to lymphoma (eg, cytokine-mediated damage). 1° disease has excellent response to corticosteroids.</p> <ul style="list-style-type: none"> ▪ LM—Normal glomeruli (lipid may be seen in PCT cells) ▪ IF—⊖ ▪ EM—effacement of podocyte foot processes A
Focal segmental glomerulosclerosis	<p>Most common cause of nephrotic syndrome in African-Americans and Hispanics. Can be 1° (idiopathic) or 2° to other conditions (eg, HIV infection, sickle cell disease, heroin abuse, massive obesity, interferon treatment, or congenital malformations). 1° disease has inconsistent response to steroids. May progress to CKD.</p> <ul style="list-style-type: none"> ▪ LM—segmental sclerosis and hyalinosis B ▪ IF—often ⊖ but may be ⊕ for nonspecific focal deposits of IgM, C3, C1 ▪ EM—effacement of foot processes similar to minimal change disease
Membranous nephropathy	<p>Also known as membranous glomerulonephritis. Can be 1° (eg, antibodies to phospholipase A₂ receptor) or 2° to drugs (eg, NSAIDs, penicillamine, gold), infections (eg, HBV, HCV, syphilis), SLE, or solid tumors. 1° disease has poor response to steroids. May progress to CKD.</p> <ul style="list-style-type: none"> ▪ LM—diffuse capillary and GBM thickening C ▪ IF—granular due to immune complex (IC) deposition ▪ EM—“Spike and dome” appearance of subepithelial deposits
Amyloidosis	<p>Kidney is the most commonly involved organ (systemic amyloidosis). Associated with chronic conditions that predispose to amyloid deposition (eg, AL amyloid, AA amyloid).</p> <ul style="list-style-type: none"> ▪ LM—Congo red stain shows apple-green birefringence under polarized light due to amyloid deposition in the mesangium
Diabetic glomerulonephropathy	<p>Most common cause of ESRD in the United States. Hyperglycemia → nonenzymatic glycation of tissue proteins → mesangial expansion; GBM thickening and ↑ permeability. Hyperfiltration (glomerular HTN and ↑ GFR) → glomerular hypertrophy and glomerular scarring (glomerulosclerosis) → further progression of nephropathy.</p> <ul style="list-style-type: none"> ▪ LM—Mesangial expansion, GBM thickening, eosinophilic nodular glomerulosclerosis (Kimmelstiel-Wilson lesions D)




Kidney stones

Can lead to severe complications such as hydronephrosis, pyelonephritis, and acute kidney injury. Obstructed stone presents with unilateral flank tenderness, colicky pain radiating to groin, hematuria. Treat and prevent by encouraging fluid intake.

CONTENT	PRECIPITATES WITH	X-RAY FINDINGS	CT FINDINGS	URINE CRYSTAL	NOTES
Calcium	Calcium oxalate: hypocitraturia	Radiopaque	Radiopaque	Shaped like envelope A or dumbbell	Calcium stones most common (80%); calcium oxalate more common than calcium phosphate stones. Can result from ethylene glycol (antifreeze) ingestion, vitamin C abuse, hypocitraturia (associated with ↓ urine pH), malabsorption (eg, Crohn disease). Treatment: thiazides, citrate, low-sodium diet.
	Calcium phosphate: ↑ pH	Radiopaque	Radiopaque	Wedge-shaped prism	Treatment: low-sodium diet, thiazides.
Ammonium magnesium phosphate (struvite)	↑ pH	Radiopaque	Radiopaque	Coffin lid B	Account for 15% of stones. Caused by infection with urease ⊕ bugs (eg, <i>Proteus mirabilis</i> , <i>Staphylococcus saprophyticus</i> , <i>Klebsiella</i>) that hydrolyze urea to ammonia → urine alkalization. Commonly form staghorn calculi C . Treatment: eradication of underlying infection, surgical removal of stone.
Uric acid	↓ pH	Radiolucent	Minimally visible	Rhomboid D or rosettes	About 5% of all stones. Risk factors: ↓ urine volume, arid climates, acidic pH. Strong association with hyperuricemia (eg, gout). Often seen in diseases with ↑ cell turnover (eg, leukemia). Treatment: alkalization of urine, allopurinol.
Cystine	↓ pH	Faintly radiopaque	Moderately radiopaque	Hexagonal E	Hereditary (autosomal recessive) condition in which Cystine-reabsorbing PCT transporter loses function, causing cystinuria. Transporter defect also results in poor reabsorption of Ornithine, Lysine, Arginine (COLA). Cystine is poorly soluble, thus stones form in urine. Usually begins in childhood. Can form staghorn calculi. Sodium cyanide nitroprusside test ⊕. “SIXtine” stones have SIX sides. Treatment: low sodium diet, alkalization of urine, chelating agents (eg, penicillamine) if refractory.



Hydronephrosis





Distention/dilation of renal pelvis and calyces **A**. Usually caused by urinary tract obstruction (eg, renal stones, severe BPH, congenital obstructions, cervical cancer, injury to ureter); other causes include retroperitoneal fibrosis, vesicoureteral reflux. Dilation occurs proximal to site of pathology. Serum creatinine becomes elevated if obstruction is bilateral or if patient has an obstructed solitary kidney. Leads to compression and possible atrophy of renal cortex and medulla.

Urinary incontinence

Mixed incontinence has features of both stress and urgency incontinence.

	Stress incontinence	Urgency incontinence	Overflow incontinence
MECHANISM	Outlet incompetence (urethral hypermobility or intrinsic sphincter deficiency) → leak with ↑ intra-abdominal pressure (eg, sneezing, lifting) ⊕ bladder stress test (directly observed leakage from urethra upon coughing or Valsalva maneuver)	Detrusor overactivity → leak with urge to void immediately	Incomplete emptying (detrusor underactivity or outlet obstruction) → leak with overfilling, ↑ postvoid residual on catheterization or ultrasound
ASSOCIATIONS	Obesity, vaginal delivery, prostate surgery	UTI	Polyuria (eg, diabetes), bladder outlet obstruction (eg, BPH), neurogenic bladder (eg, MS)
TREATMENT	Pelvic floor muscle strengthening (Kegel) exercises, weight loss, pessaries	Kegel exercises, bladder training (timed voiding, distraction or relaxation techniques), antimuscarinics (eg, oxybutynin for overactive bladder), mirabegron	Catheterization, relieve obstruction (eg, α-blockers for BPH)



Acute cystitis

Inflammation of urinary bladder. Presents as suprapubic pain, dysuria, urinary frequency, urgency.

Systemic signs (eg, high fever, chills) are usually absent.

Risk factors include female sex (short urethra), sexual intercourse, indwelling catheter, diabetes mellitus, impaired bladder emptying.

Causes:

- *E coli* (most common)
- *Staphylococcus saprophyticus*—seen in sexually active young women (*E coli* is still more common in this group)
- *Klebsiella*
- *Proteus mirabilis*—urine has ammonia scent

Labs: ⊕ leukocyte esterase. ⊕ nitrites (indicate gram ⊖ organisms). Sterile pyuria (pyuria with ⊖ urine cultures) could suggest urethritis by *Neisseria gonorrhoeae* or *Chlamydia trachomatis*.

Treatment: antibiotics (eg, TMP-SMX, nitrofurantoin).

Pyelonephritis**Acute pyelonephritis**

Neutrophils infiltrate renal interstitium **A**. Affects cortex with relative sparing of glomeruli/vessels.

Presents with fevers, flank pain (costovertebral angle tenderness), nausea/vomiting, chills.

Causes include ascending UTI (*E coli* is most common), hematogenous spread to kidney. Presents with WBCs in urine +/- WBC casts. CT would show striated parenchymal enhancement **B**.

Risk factors include indwelling urinary catheter, urinary tract obstruction, vesicoureteral reflux, diabetes mellitus, pregnancy.


Complications include chronic pyelonephritis, renal papillary necrosis, perinephric abscess, urosepsis.

Treatment: antibiotics.

Chronic pyelonephritis

The result of recurrent or inadequately treated episodes of acute pyelonephritis. Typically requires predisposition to infection such as vesicoureteral reflux or chronically obstructing kidney stones. Coarse, asymmetric corticomedullary scarring, blunted calyx. Tubules can contain eosinophilic casts resembling thyroid tissue **C** (thyroidization of kidney).

Xanthogranulomatous pyelonephritis—rare; grossly orange nodules that can mimic tumor nodules; characterized by widespread kidney damage due to granulomatous tissue containing foamy macrophages. Associated with *Proteus* infection.



Acute kidney injury

	Prerenal azotemia	Intrinsic renal failure	Postrenal azotemia
ETIOLOGY	Hypovolemia ↓ cardiac output ↓ effective circulating volume (eg, HF, liver failure)	Tubules and interstitium: ▪ Acute tubular necrosis (ischemia, sepsis, infection, nephrotoxins) ▪ Acute interstitial nephritis Glomerulus: ▪ Acute glomerulonephritis Vascular: ▪ Vasculitis ▪ Malignant hypertension ▪ TTP-HUS	Stones BPH Neoplasm Congenital anomalies
PATHOPHYSIOLOGY	↓ RBF → ↓ GFR → ↑ reabsorption of Na ⁺ /H ₂ O and urea	In ATN, patchy necrosis → debris obstructing tubules and fluid backflow → ↓ GFR In ATN, epithelial/granular casts	Outflow obstruction (bilateral)
URINE OSMOLALITY (mOsm/kg)	>500	<350	<350
URINE Na⁺ (mEq/L)	<20	>40	Varies
FE_{Na}	<1%	>2%	Varies
SERUM BUN/Cr	>20	<15	Varies

Acute interstitial nephritis

Also called tubulointerstitial nephritis. Acute interstitial renal inflammation. Pyuria (classically eosinophils) and azotemia occurring after administration of drugs that act as haptens, inducing hypersensitivity (eg, diuretics, NSAIDs, penicillin derivatives, proton pump inhibitors, rifampin, quinolones, sulfonamides). Less commonly may be 2° to other processes such as systemic infections (eg, *Mycoplasma*) or autoimmune diseases (eg, Sjögren syndrome, SLE, sarcoidosis).

Associated with fever, rash, hematuria, pyuria, and costovertebral angle tenderness, but can be asymptomatic.

Remember these **5 P'S**:

- **P**ee (diuretics)
- **P**ain-free (NSAIDs)
- **P**enicillins and cephalosporins
- **P**roton pump inhibitors
- **R**ifam**P**in
- **S**ulfa drugs

Acute tubular necrosis

Most common cause of acute kidney injury in hospitalized patients. Spontaneously resolves in many cases. Can be fatal, especially during initial oliguric phase. ↑ FE_{Na}.



Key finding: granular casts (often muddy brown in appearance) **A**.

3 stages:

1. Inciting event
2. Maintenance phase—oliguric; lasts 1–3 weeks; risk of hyperkalemia, metabolic acidosis, uremia
3. Recovery phase—polyuric; BUN and serum creatinine fall; risk of hypokalemia and renal wasting of other electrolytes and minerals

Can be caused by ischemic or nephrotoxic injury:

- Ischemic—2° to ↓ renal blood flow (eg, hypotension, shock, sepsis, hemorrhage, HF). Results in death of tubular cells that may slough into tubular lumen **B** (PCT and thick ascending limb are highly susceptible to injury).
- Nephrotoxic—2° to injury resulting from toxic substances (eg, aminoglycosides, radiocontrast agents, lead, cisplatin, ethylene glycol), crush injury (myoglobinuria), hemoglobinuria. Proximal tubules are particularly susceptible to injury.



Diffuse cortical necrosis


Acute generalized cortical infarction of both kidneys. Likely due to a combination of vasospasm and DIC.

Associated with obstetric catastrophes (eg, abruptio placentae), septic shock.

Renal papillary necrosis

Sloughing of necrotic renal papillae **A** → gross hematuria and proteinuria. May be triggered by recent infection or immune stimulus.

Associated with: Sick cell disease or trait, Acute pyelonephritis, Analgesics (NSAIDs), Diabetes mellitus (SAAD papa with papillary necrosis).



Consequences of renal failure

Decline in renal filtration can lead to excess retained nitrogenous waste products and electrolyte disturbances.

Consequences (**MAD HUNGER**):

- **M**etabolic **A**cidosis
- **D**yslipidemia (especially ↑ triglycerides)
- **H**igh potassium
- **U**remia—clinical syndrome marked by:
 - Nausea and anorexia
 - Pericarditis
 - Asterixis
 - Encephalopathy
 - Platelet dysfunction
- **N**a⁺/H₂O retention (HF, pulmonary edema, hypertension)
- **G**rowth retardation and developmental delay
- **E**rythropoietin deficiency (anemia)
- **R**enal osteodystrophy

2 forms of renal failure: acute (eg, ATN) and chronic (eg, hypertension, diabetes mellitus, congenital anomalies).

Incremental reductions in GFR define the stages of chronic kidney disease.

Renal osteodystrophy

Hypocalcemia, hyperphosphatemia, and failure of vitamin D hydroxylation associated with chronic kidney disease → 2° hyperparathyroidism → 3° hyperparathyroidism (if 2° poorly managed). High serum phosphate can bind with Ca²⁺ → tissue deposits → ↓ serum Ca²⁺. ↓ 1,25-(OH)₂D₃ → ↓ intestinal Ca²⁺ absorption. Causes subperiosteal thinning of bones.

Renal cyst disorders**Autosomal dominant polycystic kidney disease**

Numerous cysts in cortex and medulla **A** causing bilateral enlarged kidneys ultimately destroy kidney parenchyma. Presents with flank pain, hematuria, hypertension, urinary infection, progressive renal failure in ~ 50% of individuals.

Mutation in *PKD1* (85% of cases, chromosome 16) or *PKD2* (15% of cases, chromosome 4).

Complications include chronic kidney disease and hypertension (caused by ↑ renin production).

Associated with berry aneurysms, mitral valve prolapse, benign hepatic cysts, diverticulosis.

Treatment: If hypertension or proteinuria develops, treat with ACE inhibitors or ARBs.

Autosomal recessive polycystic kidney disease

Cystic dilation of collecting ducts **B**. Often presents in infancy. Associated with congenital hepatic fibrosis. Significant oliguric renal failure in utero can lead to Potter sequence. Concerns beyond neonatal period include systemic hypertension, progressive renal insufficiency, and portal hypertension from congenital hepatic fibrosis.

Autosomal dominant tubulointerstitial kidney disease

Also called medullary cystic kidney disease. Causes tubulointerstitial fibrosis and progressive renal insufficiency with inability to concentrate urine. Medullary cysts usually not visualized; smaller kidneys on ultrasound. Poor prognosis.

Simple vs complex renal cysts

Simple cysts are filled with ultrafiltrate (anechoic on ultrasound **C**). Very common and account for majority of all renal masses. Found incidentally and typically asymptomatic.

Complex cysts, including those that are septated, enhanced, or have solid components on imaging require follow-up or removal due to risk of renal cell carcinoma.

**Renovascular disease**

Renal impairment due to ischemia from renal artery stenosis or microvascular disease.

↓ renal perfusion (one or both kidneys)

→ ↑ renin → ↑ angiotensin → HTN.

Main causes of renal artery stenosis:

- Atherosclerotic plaques—proximal 1/3 of renal artery, usually in older males, smokers.
- Fibromuscular dysplasia—distal 2/3 of renal artery or segmental branches, usually young or middle-aged females.

Clinically, patients can have refractory HTN with negative family history of HTN, asymmetric renal size, epigastric/flank bruits.

Most common cause of 2° HTN in adults. Other large vessels are often involved.

Renal cell carcinoma

Polygonal clear cells **A** filled with accumulated lipids and carbohydrate. Often golden-yellow **B** due to ↑ lipid content.

Originates from PCT → invades renal vein (may develop varicocele if left sided) → IVC → hematogenous spread → metastasis to lung and bone.


Manifests with hematuria, palpable masses, 2° polycythemia, flank pain, fever, weight loss. Treatment: surgery/ablation for localized disease. Immunotherapy (eg, aldesleukin) or targeted therapy for metastatic disease, rarely curative. Resistant to chemotherapy and radiation therapy.

Most common 1° renal malignancy **C**. Most common in men 50–70 years old, ↑ incidence with smoking and obesity.


Associated with paraneoplastic syndromes, eg, PTHrP, Ectopic EPO, ACTH, Renin (“PEAR”-aneoplastic).

Clear cell (most common subtype) associated with gene deletion on chromosome 3 (sporadic, or inherited as von Hippel-Lindau syndrome).

RCC = 3 letters = chromosome 3.




Renal oncocytoma




Benign epithelial cell tumor arising from collecting ducts (arrows in **A** point to well-circumscribed mass with central scar).

Large eosinophilic cells with abundant mitochondria without perinuclear clearing **B** (vs chromophobe renal cell carcinoma). Presents with painless hematuria, flank pain, abdominal mass.

Often resected to exclude malignancy (eg, renal cell carcinoma).



Nephroblastoma




Also called Wilms tumor. Most common renal malignancy of early childhood (ages 2–4). Contains embryonic glomerular structures. Presents with large, palpable, unilateral flank mass **A** and/or hematuria and possible HTN.

“Loss of function” mutations of tumor suppressor genes *WT1* or *WT2* on chromosome 11.

May be a part of several syndromes:

- **WAGR complex**—Wilms tumor, **A**niridia (absence of iris), **G**enitourinary malformations, mental **R**etardation/intellectual disability (*WT1* deletion)
- **Denys-Drash syndrome**—Wilms tumor, **D**iffuse mesangial sclerosis (early-onset nephrotic syndrome), **D**ysgenesis of gonads (male pseudohermaphroditism), *WT1* mutation
- **Beckwith-Wiedemann syndrome**—Wilms tumor, macroglossia, organomegaly, hemihyperplasia (*WT2* mutation), omphalocele


Urothelial carcinoma of the bladder



Also called transitional cell carcinoma. Most common tumor of urinary tract system (can occur in renal calyces, renal pelvis, ureters, and bladder) **A B**. Can be suggested by painless hematuria (no casts).

Associated with problems in your **Pee SAC**:

- P**henacetin, **S**moking, **A**niline dyes, and **C**yclophosphamide.




Squamous cell carcinoma of the bladder

Chronic irritation of urinary bladder → squamous metaplasia → dysplasia and squamous cell carcinoma.

Risk factors include *Schistosoma haematobium* infection (Middle East), chronic cystitis, smoking, chronic nephrolithiasis. Presents with painless hematuria (no casts).

▶ RENAL—PHARMACOLOGY

Diuretics site of action




Mannitol

MECHANISM	Osmotic diuretic. ↑ tubular fluid osmolarity → ↑ urine flow, ↓ intracranial/intraocular pressure.
CLINICAL USE	Drug overdose, elevated intracranial/intraocular pressure.
ADVERSE EFFECTS	Pulmonary edema, dehydration, hypo- or hypernatremia. Contraindicated in anuria, HF.

Acetazolamide

MECHANISM	Carbonic anhydrase inhibitor. Causes self-limited NaHCO_3 diuresis and \downarrow total body HCO_3^- stores. Alkalinizes urine.
CLINICAL USE	Glaucoma, metabolic alkalosis, altitude sickness, idiopathic intracranial hypertension.




ADVERSE EFFECTS	Proximal renal tubular acidosis, paresthesias, NH_3 toxicity, sulfa allergy, hypokalemia. Promotes calcium phosphate stone formation (insoluble at high pH).
-----------------	---

“Acid”azolamide causes Acidosis.

Loop diuretics

Furosemide, bumetanide, torsemide

MECHANISM	Sulfonamide loop diuretics. Inhibit cotransport system ($\text{Na}^+/\text{K}^+/2\text{Cl}^-$) of thick ascending limb of loop of Henle. Abolish hypertonicity of medulla, preventing concentration of urine. Associated with \uparrow PGE (vasodilatory effect on afferent arteriole); inhibited by NSAIDs. \uparrow Ca^{2+} excretion. L oops L ose Ca^{2+} .
CLINICAL USE	Edematous states (HF, cirrhosis, nephrotic syndrome, pulmonary edema), hypertension, hypercalcemia.



ADVERSE EFFECTS	Ototoxicity, Hypokalemia, Hypomagnesemia, Dehydration, Allergy (sulfa), metabolic Alkalosis, Nephritis (interstitial), Gout.
-----------------	--


OHH DAANG!

Ethacrynic acid

MECHANISM	Nonsulfonamide inhibitor of cotransport system ($\text{Na}^+/\text{K}^+/2\text{Cl}^-$) of thick ascending limb of loop of Henle.
CLINICAL USE	Diuresis in patients allergic to sulfa drugs.
ADVERSE EFFECTS	Similar to furosemide, but more ototoxic.


Loop earrings hurt your ears.

Thiazide diuretics	Hydrochlorothiazide, chlorthalidone, metolazone.
MECHANISM	Inhibit NaCl reabsorption in early DCT → ↓ diluting capacity of nephron. ↓ Ca ²⁺ excretion.
CLINICAL USE	Hypertension, HF, idiopathic hypercalciuria, nephrogenic diabetes insipidus, osteoporosis.
ADVERSE EFFECTS	Hypokalemic metabolic alkalosis, hyponatremia, hyperGlycemia, hyperLipidemia, hyperUricemia, hyperCalcemia. Sulfa allergy.



Potassium-sparing diuretics	Spironolactone, Eplerenone, Amiloride, Triamterene.
MECHANISM	Spironolactone and eplerenone are competitive aldosterone receptor antagonists in cortical collecting tubule. Triamterene and amiloride block Na ⁺ channels at the same part of the tubule.
CLINICAL USE	Hyperaldosteronism, K ⁺ depletion, HF, hepatic ascites (spironolactone), nephrogenic DI (amiloride), antiandrogen.
ADVERSE EFFECTS	Hyperkalemia (can lead to arrhythmias), endocrine effects with spironolactone (eg, gynecomastia, antiandrogen effects).

Keep your SEAT

**Diuretics: electrolyte changes**

Urine NaCl	↑ with all diuretics (strength varies based on potency of diuretic effect). Serum NaCl may decrease as a result.
Urine K⁺	↑ especially with loop and thiazide diuretics. Serum K ⁺ may decrease as a result.
Blood pH	<p>↓ (acidemia): carbonic anhydrase inhibitors: ↓ HCO₃⁻ reabsorption. K⁺ sparing: aldosterone blockade prevents K⁺ secretion and H⁺ secretion. Additionally, hyperkalemia leads to K⁺ entering all cells (via H⁺/K⁺ exchanger) in exchange for H⁺ exiting cells.</p> <p>↑ (alkalemia): loop diuretics and thiazides cause alkalemia through several mechanisms:</p> <ul style="list-style-type: none"> ▪ Volume contraction → ↑ AT II → ↑ Na⁺/H⁺ exchange in PCT → ↑ HCO₃⁻ reabsorption (“contraction alkalosis”) ▪ K⁺ loss leads to K⁺ exiting all cells (via H⁺/K⁺ exchanger) in exchange for H⁺ entering cells ▪ In low K⁺ state, H⁺ (rather than K⁺) is exchanged for Na⁺ in cortical collecting tubule → alkalosis and “paradoxical aciduria”
Urine Ca²⁺	<p>↑ with loop diuretics: ↓ paracellular Ca²⁺ reabsorption → hypocalcemia.</p> <p>↓ with thiazides: enhanced Ca²⁺ reabsorption.</p>

Angiotensin-converting enzyme inhibitors

Captopril, enalapril, lisinopril, ramipril.

MECHANISM	Inhibit ACE → ↓ AT II → ↓ GFR by preventing constriction of efferent arterioles. ↑ renin due to loss of negative feedback. Inhibition of ACE also prevents inactivation of bradykinin, a potent vasodilator.	
CLINICAL USE	Hypertension, HF (↓ mortality), proteinuria, diabetic nephropathy. Prevent unfavorable heart remodeling as a result of chronic hypertension.	In chronic kidney disease (eg, diabetic nephropathy), ↓ intraglomerular pressure, slowing GBM thickening.
ADVERSE EFFECTS	C ough, A ngioedema (both due to ↑ bradykinin; contraindicated in C1 esterase inhibitor deficiency), T eratogen (fetal renal malformations), ↑ C reatinine (↓ GFR), H yperkalemia, and H ypotension. Used with caution in bilateral renal artery stenosis because ACE inhibitors will further ↓ GFR → renal failure.	

Angiotensin II receptor blockers

Losartan, candesartan, valsartan.

MECHANISM	Selectively block binding of angiotensin II to AT ₁ receptor. Effects similar to ACE inhibitors, but ARBs do not increase bradykinin.	
CLINICAL USE	Hypertension, HF, proteinuria, or chronic kidney disease (eg, diabetic nephropathy) with intolerance to ACE inhibitors (eg, cough, angioedema).	
ADVERSE EFFECTS	Hyperkalemia, ↓ GFR, hypotension; teratogen.	

Aliskiren

MECHANISM	Direct renin inhibitor, blocks conversion of angiotensinogen to angiotensin I. Aliskiren Kills Renin.	
CLINICAL USE	Hypertension.	
ADVERSE EFFECTS	Hyperkalemia, ↓ GFR, hypotension, angioedema. Relatively contraindicated in patients already taking ACE inhibitors or ARBs and contraindicated in pregnancy.	

HIGH-YIELD SYSTEMS

Reproductive

“Artificial insemination is when the farmer does it to the cow instead of the bull.”

—Student essay

Make no mistake about why these babies are here - they are here to replace us.

—Jerry Seinfeld

“Whoever called it necking was a poor judge of anatomy.”

—Groucho Marx

“See, the problem is that God gives men a brain and a penis, and only enough blood to run one at a time.”

—Robin Williams

The reproductive system can be intimidating at first but is manageable once you organize the concepts into the pregnancy, endocrinologic, embryologic, and oncologic aspects of reproduction. Study the endocrine and reproductive chapters together, because mastery of the hypothalamic-pituitary-gonadal axis is key to answering questions on ovulation, menstruation, disorders of sexual development, contraception, and many pathologies.

Embryology is a nuanced subject that covers multiple organ systems. Approaching it from a clinical perspective will allow for better understanding. For instance, make the connection between the presentation of DiGeorge syndrome and the 3rd/4th pharyngeal pouch, and between the Müllerian/Wolffian systems and disorders of sexual development.

As for oncology, don't worry about remembering screening or treatment guidelines. It is more important to know how these cancers present (eg, signs and symptoms) and their associated labs, histopathology, and risk factors. In addition, some of the testicular and ovarian cancers have distinct patterns of hCG, AFP, LH, or FSH derangements that serve as helpful clues in exam questions.

▶ Embryology	612
▶ Anatomy	624
▶ Physiology	629
▶ Pathology	638
▶ Pharmacology	655


▶ REPRODUCTIVE—EMBRYOLOGY

Important genes of embryogenesis

GENE	LOCATION	FUNCTION	NOTES
Sonic hedgehog (SHH) gene	Zone of polarizing activity at base of limb buds	Anterior-posterior axis patterning, CNS development	Mutations → holoprosencephaly
Wnt-7 gene	Apical ectodermal ridge at distal end of each limb	Dorsal-ventral axis patterning, limb development	
Fibroblast growth factor (FGF) gene	Apical ectodermal ridge	Limb lengthening (via mitosis of mesoderm)	“Look at that F etus, G rowing F ingers”
Homeobox (Hox) genes	Multiple	Segmental organization in cranial-caudal direction, transcription factor coding	Mutations → appendages in wrong locations. Isotretinoin → ↑ <i>Hox</i> gene expression

Early fetal development

Early embryonic development



Within week 1	hCG secretion begins around the time of implantation of blastocyst.	Blastocyst “sticks” at day 6.
Within week 2	Bilaminar disc (epiblast, hypoblast).	2 weeks = 2 layers.
Within week 3	Gastrulation forms trilaminar embryonic disc. Cells from epiblast invaginate → primitive streak → endoderm, mesoderm, ectoderm. Notochord arises from midline mesoderm; overlying ectoderm becomes neural plate.	3 weeks = 3 layers.
Weeks 3–8 (embryonic period)	Neural tube formed by neuroectoderm and closes by week 4. Organogenesis.	Extremely susceptible to teratogens.
Week 4	Heart begins to beat. Upper and lower limb buds begin to form.	4 weeks = 4 limbs and 4 heart chambers.
Week 6	Fetal cardiac activity visible by transvaginal ultrasound.	
Week 8	Fetal movements start.	Gait at week 8.
Week 10	Genitalia have male/female characteristics.	Tenitalia .

Embryologic derivatives

Ectoderm		External/outer layer
Surface ectoderm	Epidermis; adenohypophysis (from Rathke pouch); lens of eye; epithelial linings of oral cavity, sensory organs of ear, and olfactory epithelium; anal canal below the pectinate line; parotid, sweat, mammary glands.	Craniopharyngioma —benign Rathke pouch tumor with cholesterol crystals, calcifications.
Neural tube	Brain (neurohypophysis, CNS neurons, oligodendrocytes, astrocytes, ependymal cells, pineal gland), retina, spinal cord.	Neuroectoderm—think CNS.
Neural crest	Melanocytes , O dontoblasts, T racheal cartilage, E nterochromaffin cells, L eptomeninges (arachnoid, pia), P NS ganglia (cranial, dorsal root, autonomic), A drenal medulla, S chwann cells, S piral membrane (aorticopulmonary septum), E ndocardial cushions (also derived partially from mesoderm), S kull bones.	MOTEL PASSES Neural crest—think PNS and non-neural structures nearby.
Mesoderm	Muscle, bone, connective tissue, serous linings of body cavities (eg, peritoneum, pericardium, pleura), spleen (develops within foregut mesentery), cardiovascular structures, lymphatics, blood, wall of gut tube, upper vagina, kidneys, adrenal cortex, dermis, testes, ovaries, microglia. Notochord induces ectoderm to form neuroectoderm (neural plate); its only postnatal derivative is the nucleus pulposus of the intervertebral disc.	M iddle/“ m eat” layer. Mesodermal defects = VACTERL : V ertebral defects A nal atresia C ardiac defects T racheo- E sophageal fistula R enal defects L imb defects (bone and muscle)
Endoderm	Gut tube epithelium (including anal canal above the pectinate line), most of urethra and lower vagina (derived from urogenital sinus), luminal epithelial derivatives (eg, lungs, liver, gallbladder, pancreas, eustachian tube, thymus, parathyroid, thyroid follicular and parafollicular [C] cells).	“ E nternal” layer.

Types of errors in morphogenesis

Agenesis	Absent organ due to absent primordial tissue.
Aplasia	Absent organ despite presence of primordial tissue.
Hypoplasia	Incomplete organ development; primordial tissue present.
Disruption	2° breakdown of previously normal tissue or structure (eg, amniotic band syndrome).
Deformation	Extrinsic disruption (eg, multiple gestations → crowding → foot deformities); occurs after embryonic period.
Malformation	Intrinsic disruption; occurs during embryonic period (weeks 3–8).
Sequence	Abnormalities result from a single 1° embryologic event (eg, oligohydramnios → Potter sequence).

Teratogens

Most susceptible in 3rd–8th weeks (embryonic period—organogenesis) of pregnancy. Before week 3, “all-or-none” effects. After week 8, growth and function affected.

TERATOGEN	EFFECTS ON FETUS	NOTES
Medications		
ACE inhibitors	Renal failure, oligohydramnios, hypocalvaria.	
Alkylating agents	Absence of digits, multiple anomalies.	
Aminoglycosides	Ototoxicity.	A mean guy hit the baby in the ear .
Antiepileptic drugs	Neural tube defects, cardiac defects, cleft palate, skeletal abnormalities (eg, phalanx/nail hypoplasia, facial dysmorphism).	High-dose folate supplementation recommended. Most commonly valproate, carbamazepine, phenytoin, phenobarbital.
Diethylstilbestrol (DES)	Vaginal clear cell adenocarcinoma, congenital Müllerian anomalies.	
Fluoroquinolones	Cartilage damage.	
Folate antagonists	Neural tube defects.	Antiepileptics, trimethoprim, methotrexate.
Isotretinoin	Multiple severe birth defects.	Contraception mandatory. Iso TERAT inoin.
Lithium	Ebstein anomaly.	
Methimazole	Aplasia cutis congenita (congenital absence of skin, particularly on scalp).	
Tetracyclines	Discolored teeth, inhibited bone growth.	“ Teeth racyclines.”
Thalidomide	Limb defects (phocomelia, micromelia—“flipper” limbs).	Limb defects with “tha- limb -domide.”
Warfarin	Bone and cartilage deformities (stippled epiphyses, nasal and limb hypoplasia), optic nerve atrophy, fetal cerebral hemorrhage.	Do not wage warfare on the baby; keep it heppy with heparin (does not cross placenta).
Substance abuse		
Alcohol	Fetal alcohol syndrome.	
Cocaine	Low birth weight, preterm birth, IUGR, placental abruption.	Cocaine → vasoconstriction.
Smoking (nicotine, CO)	Low birth weight (leading cause in developed countries), preterm labor, placental problems, IUGR, SIDS, ADHD.	Nicotine → vasoconstriction. CO → impaired O ₂ delivery.
Other		
Iodine (lack or excess)	Congenital goiter or hypothyroidism (cretinism).	
Maternal diabetes	Caudal regression syndrome, cardiac defects (eg, VSD), neural tube defects, macrosomia, neonatal hypoglycemia (due to islet cell hyperplasia), polycythemia.	
Methylmercury	Neurotoxicity.	Highest in swordfish, shark, tilefish, king mackerel.
Vitamin A excess	Extremely high risk for spontaneous abortions and birth defects (cleft palate, cardiac).	
X-rays	Microcephaly, intellectual disability.	Minimized by lead shielding.

Fetal alcohol syndrome

One of the leading preventable causes of intellectual disability in the US. Newborns of mothers who consumed alcohol during any stage of pregnancy have ↑ incidence of congenital abnormalities, including pre- and postnatal developmental retardation, microcephaly, facial abnormalities **A** (eg, smooth philtrum, thin vermilion border, small palpebral fissures), limb dislocation, heart defects. Heart-lung fistulas and holoprosencephaly in most severe form. One mechanism is due to impaired migration of neuronal and glial cells.

Neonatal abstinence syndrome

Complex disorder involving CNS, ANS, and GI systems. Secondary to maternal substance use/abuse (most commonly opioids).

Universal screening for substance abuse is recommended in all pregnant patients.


Newborns may present with uncoordinated sucking reflexes, irritability, high-pitched crying, tremors, tachypnea, sneezing, diarrhea, and possibly seizures.

Treatment (for opiate abuse): methadone, morphine, buprenorphine.

Twinning

Dizygotic (“fraternal”) twins arise from 2 eggs that are separately fertilized by 2 different sperm (always 2 zygotes) and will have 2 separate amniotic sacs and 2 separate placentas (chorions). Monozygotic (“identical”) twins arise from 1 fertilized egg (1 egg + 1 sperm) that splits in early pregnancy. The timing of cleavage determines chorionicity (number of chorions) and amnionicity (number of amnions) (**SCAB**):

- Cleavage 0–4 days: Separate chorion and amnion
- Cleavage 4–8 days: shared Chorion
- Cleavage 8–12 days: shared Amnion
- Cleavage 13+ days: shared Body (conjoined)




Placenta 1° site of nutrient and gas exchange between mother and fetus.

Fetal component

Cytotrophoblast	Inner layer of chorionic villi.	Cytotrophoblast makes Cells.
Syncytiotrophoblast	Outer layer of chorionic villi; synthesizes and secretes hormones, eg, hCG (structurally similar to LH; stimulates corpus luteum to secrete progesterone during first trimester).	Syncytiotrophoblast synthesizes hormones. Lacks MHC-I expression → ↓ chance of attack by maternal immune system.

Maternal component

Decidua basalis Derived from endometrium. Maternal blood in lacunae.





Umbilical cord

Two umbilical arteries return deoxygenated blood from fetal internal iliac arteries to placenta **A**.

One umbilical vein supplies oxygenated blood from placenta to fetus; drains into IVC via liver or via ductus venosus.

Single umbilical artery (2-vessel cord) is associated with congenital and chromosomal anomalies.

Umbilical arteries and vein are derived from allantois.



Urachus

Allantois forms from hindgut and extends into urogenital sinus. Allantois becomes the urachus, a duct between fetal bladder and umbilicus. Failure of urachus to involute can lead to anomalies that may increase risk of infection and/or malignancy (eg, adenocarcinoma) if not treated. Obliterated urachus is represented by the median umbilical ligament after birth, which is covered by median umbilical fold of the peritoneum.

Patent urachus


Total failure of urachus to obliterate → urine discharge from umbilicus.

Urachal cyst

Partial failure of urachus to obliterate; fluid-filled cavity lined with uroepithelium, between umbilicus and bladder. Cyst can become infected and present as painful mass below umbilicus.

Vesicourachal diverticulum

Slight failure of urachus to obliterate → outpouching of bladder.



Vitelline duct



7th week—obliteration of vitelline duct (omphalomesenteric duct), which connects yolk sac to midgut lumen.

Vitelline fistula


Vitelline duct fails to close → meconium discharge from umbilicus.

Meckel diverticulum

Partial closure of vitelline duct, with patent portion attached to ileum (true diverticulum, white arrow in **B**). May be asymptomatic. May have heterotopic gastric and/or pancreatic tissue → melena, hematochezia, abdominal pain.




Aortic arch derivatives	Develop into arterial system.	
1st	Part of max illary artery (branch of external carotid)	1st arch is max imal
2nd	S tapedial artery and hyoid artery	Second = S tapedial
3rd	C ommon C arotid artery and proximal part of internal C arotid artery	C is 3rd letter of alphabet
4th	On left, aortic arch; on right, proximal part of right subclavian artery	4th arch (4 limbs) = systemic
6th	Proximal part of pulmonary arteries and (on left only) ductus arteriosus	6th arch = pulmonary and the pulmonary-to-systemic shunt (ductus arteriosus)



Pharyngeal apparatus Composed of pharyngeal clefts, arches, pouches.
 Pharyngeal **c**lefts—derived from **e**ctoderm. Also called pharyngeal grooves.
 Pharyngeal **a**rches—derived from mesoderm (muscles, arteries) and neural crest (bones, cartilage).
 Pharyngeal **p**ouches—derived from endoderm.

CAP covers outside to inside:
Clefts = **e**ctoderm
Arches = mesoderm + neural crest
Pouches = endoderm



Pharyngeal cleft derivatives

1st cleft develops into external auditory meatus.
 2nd through 4th clefts form temporary cervical sinuses, which are obliterated by proliferation of 2nd arch mesenchyme.
 Persistent cervical sinus → pharyngeal cleft cyst within lateral neck, anterior to sternocleidomastoid muscle (does not move with swallowing, vs thyroglossal duct cyst).

Pharyngeal arch derivatives

ARCH	CARTILAGE	MUSCLES	NERVES ^a	NOTES
1st pharyngeal arch	M axillary process → M axilla, zygo M atic bone M andibular process → M eckel cartilage → M andible, M alleus and incus, spheno M andibular ligament	M uscles of M astication (temporalis, M asseter, lateral and M edial pterygoids), M ylorhyoid, anterior belly of digastric, tensor tympani, anterior 2/3 of tongue, tensor veli palatini	CN V ₃ chew	P ierre Robin sequence—micrognathia, glossoptosis, cleft palate, airway obstruction T reacher Collins s ndrome—autosomal dominant neural crest dysfunction → craniofacial abnormalities (eg, zygomatic bone and mandibular hypoplasia), hearing loss, airway compromise
2nd pharyngeal arch	Reichert cartilage: S tapes, S tyleoid process, le S Ser horn of hyoid, S tyleohyoid ligament	Muscles of facial expression, S tapedius, S tyleohyoid, platy S ma, posterior belly of digastric	CN VII (facial expression) smile	
3rd pharyngeal arch	Greater horn of hyoid	Stylopharyngeus (think of stylo pharyngeus innervated by glosso pharyngeal nerve)	CN IX (styo -pharyngeus) swallow stylishly	
4th and 6th pharyngeal arches	A rytenoids, C ricoid, C orniculate, C uneiform, T hyroid (used to sing and ACCCT)	4th arch: most pharyngeal constrictors; cricothyroid, levator veli palatini 6th arch: all intrinsic muscles of larynx except cricothyroid	4th arch: CN X (superior laryngeal branch) simply swallow 6th arch: CN X (recurrent/inferior laryngeal branch) speak	Arches 3 and 4 form posterior 1/3 of tongue Arch 5 makes no major developmental contributions

^aSensory and motor nerves are not pharyngeal arch derivatives. They grow into the arches and are derived from neural crest (sensory) and neuroectoderm (motor).

When at the restaurant of the golden **arches**, children tend to first **chew** (1), then **smile** (2), then **swallow stylishly** (3) or **simply swallow** (4), and then **speak** (6).

Pharyngeal pouch derivatives

POUCH	DERIVATIVES	NOTES	MNEMONIC
1st pharyngeal pouch	Middle ear cavity, eustachian tube, mastoid air cells	1st pouch contributes to endoderm-lined structures of ear	Ear, tonsils, bottom-to-top: 1 (ear) 2 (tonsils) 3 dorsal (bottom for inferior parathyroids) 3 ventral (to = thymus) 4 (top = superior parathyroids)
2nd pharyngeal pouch	Epithelial lining of palatine tonsil		
3rd pharyngeal pouch	Dorsal wings → inferior parathyroids Ventral wings → thymus	3rd pouch contributes to 3 structures (thymus, left and right inferior parathyroids) 3rd-pouch structures end up below 4th-pouch structures	
4th pharyngeal pouch	Dorsal wings → superior parathyroids Ventral wings → ultimopharyngeal body → parafollicular (C) cells of thyroid		

Cleft lip and cleft palate


Distinct, multifactorial etiologies, but often occur together.

Cleft lip

Due to failure of fusion of the maxillary and merged medial nasal processes (formation of 1° palate).

Cleft palate

Due to failure of fusion of the two lateral palatine shelves or failure of fusion of lateral palatine shelf with the nasal septum and/or 1° palate (formation of 2° palate).



Genital embryology

Female

Default development. Mesonephric duct degenerates and paramesonephric duct develops.

Male

SRY gene on Y chromosome—produces testis-determining factor → testes development. Sertoli cells secrete Müllerian inhibitory factor (MIF, also called antimüllerian hormone) that suppresses development of paramesonephric ducts.

Leydig cells secrete androgens that stimulate development of mesonephric ducts.


Paramesonephric (Müllerian) duct

Develops into female internal structures—fallopian tubes, uterus, upper portion of vagina (lower portion from urogenital sinus). Male remnant is appendix testis.


Müllerian agenesis (Mayer-Rokitansky-Küster-Hauser syndrome)—may present as 1° amenorrhea (due to a lack of uterine development) in females with fully developed 2° sexual characteristics (functional ovaries).

Mesonephric (Wolffian) duct

Develops into male internal structures (except prostate)—**S**eminal vesicles, **E**pididymis, **E**jaculatory duct, **D**uctus deferens (**SEED**). Female remnant is Gartner duct.



Sexual differentiation




- 1 Absence of Sertoli cells or lack of Müllerian inhibitory factor → develop both male and female internal genitalia and male external genitalia (streak gonads)
- 2 5α-reductase deficiency—inability to convert testosterone into DHT → male internal genitalia, ambiguous external genitalia until puberty (when ↑ testosterone levels cause masculinization)

In the testes:


- L**eydig **L**eads to male (internal and external) sexual differentiation.
- S**ertoli **S**huts down female (internal) sexual differentiation.


Uterine (Müllerian duct) anomalies

- Septate uterus** Common anomaly vs normal uterus **A**. Incomplete resorption of septum **B**. ↓ fertility and early miscarriage/pregnancy loss. Treat with septoplasty.
- Bicornuate uterus** Incomplete fusion of Müllerian ducts **C**. ↑ risk of complicated pregnancy, early pregnancy loss, malpresentation, prematurity.
- Uterus didelphys** Complete failure of fusion → double uterus, cervix, vagina **D**. Pregnancy possible.



Male/female genital homologs




Congenital penile abnormalities**Hypospadias**

Abnormal opening of penile urethra on ventral surface of penis due to failure of urethral folds to fuse.

Hypospadias is more common than epispadias. Associated with inguinal hernia, cryptorchidism, chordee (downward or upward bending of penis).

Hypo is **below**.

Can be seen in 5 α -reductase deficiency.

Epispadias

Abnormal opening of penile urethra on dorsal surface of penis due to faulty positioning of genital tubercle.

Exstrophy of the bladder is associated with Epispadias.

When you have Epispadias, you hit your Eye when you pEE.

Descent of testes and ovaries

	DESCRIPTION	MALE REMNANT	FEMALE REMNANT
Gubernaculum	Band of fibrous tissue	Anchors testes within scrotum	Ovarian ligament + round ligament of uterus
Processus vaginalis	Evagination of peritoneum	Forms tunica vaginalis Persistent patent processus vaginalis → hydrocele	Obliterated

▶ REPRODUCTIVE—ANATOMY**Gonadal drainage****Venous drainage**


Left ovary/testis → left gonadal vein → left renal vein → IVC.

Right ovary/testis → right gonadal vein → IVC.
Because the left spermatic vein enters the left renal vein at a 90° angle, flow is less laminar on left than on right → left venous pressure > right venous pressure → varicocele more common on the left.


“Left gonadal vein takes the **L**ongest way.”

Lymphatic drainage

Ovaries/testes → para-aortic lymph nodes.
Body of uterus/cervix/superior part of bladder → external iliac nodes.
Prostate/cervix/corpus cavernosum/proximal vagina → internal iliac nodes.
Distal vagina/vulva/scrotum/distal anus → superficial inguinal nodes.
Glans penis → deep inguinal nodes.



Female reproductive anatomy




LIGAMENT	CONNECTS	STRUCTURES CONTAINED	NOTES
Infundibulopelvic (suspensory) ligament	Ovaries to lateral pelvic wall	Ovarian vessels	Ligate vessels during oophorectomy to avoid bleeding Ureter courses retroperitoneally, close to gonadal vessels → at risk of injury during ligation of ovarian vessels
Cardinal (transverse cervical) ligament	Cervix to side wall of pelvis	Uterine vessels	Ureter at risk of injury during ligation of uterine vessels in hysterectomy
Round ligament of the uterus	Uterine horn to labia majora		Derivative of gubernaculum. Travels through round inguinal canal; above the artery of Sampson
Broad ligament	Uterus, fallopian tubes, and ovaries to pelvic side wall	Ovaries, fallopian tubes, round ligaments of uterus	Fold of peritoneum that comprises the mesosalpinx, mesometrium, and mesovarium
Ovarian ligament	Medial pole of ovary to uterine horn		Derivative of gubernaculum Ovarian ligament latches to lateral uterus

Adnexal torsion



Twisting of ovary and fallopian tube around infundibulopelvic ligament and ovarian ligament → compression of ovarian vessels in infundibulopelvic ligament → blockage of lymphatic and venous outflow. Continued arterial perfusion → ovarian edema → complete blockage of arterial inflow → necrosis, local hemorrhage.
Associated with ovarian masses. Presents with acute pelvic pain, adnexal mass, nausea/vomiting.

Female reproductive epithelial histology



TISSUE	HISTOLOGY/NOTES
Vulva	Stratified squamous epithelium
Vagina	Stratified squamous epithelium, nonkeratinized
Ectocervix	Stratified squamous epithelium, nonkeratinized
Transformation zone	Squamocolumnar junction A (most common area for cervical cancer)
Endocervix	Simple columnar epithelium
Uterus	Simple columnar epithelium with long tubular glands in proliferative phase; coiled glands in secretory phase
Fallopian tube	Simple columnar epithelium, ciliated
Ovary, outer surface	Simple cuboidal epithelium (germinal epithelium covering surface of ovary)

Male reproductive anatomy



Pathway of sperm during ejaculation—



SEVEN UP:

- Seminiferous tubules
- Epididymis
- Vas deferens
- Ejaculatory ducts
- (Nothing)
- Urethra
- Penis

Urethral injury

Occurs almost exclusively in men. Suspect if blood seen at urethral meatus. Urethral catheterization is relatively contraindicated.

	Anterior urethral injury	Posterior urethral injury
PART OF URETHRA	Bulbar (spongy) urethra	Membranous urethra
MECHANISM	Perineal straddle injury	Pelvic fracture
LOCATION OF URINE LEAK/BLOOD ACCUMULATION	Blood accumulates in scrotum If Buck fascia is torn, urine escapes into perineal space	Urine leaks into retropubic space
PRESENTATION	Blood at urethral meatus and scrotal hematoma	Blood at urethral meatus and high-riding prostate

**Autonomic innervation of male sexual response**

Erection—**P**arasympathetic nervous system (pelvic splanchnic nerves, S2-S4):

- NO → ↑ cGMP → smooth muscle relaxation → vasodilation → proerectile.
- Norepinephrine → ↑ $[Ca^{2+}]_{in}$ → smooth muscle contraction → vasoconstriction → antierectile.

Emission—**S**ympathetic nervous system (hypogastric nerve, T11-L2).

Expulsion—visceral and **S**omatic nerves (pudendal nerve).



Point, **S**queeze, and **S**hoot.

S2, **3**, **4** keep the penis off the **f**loor.

PDE-5 inhibitors (eg, sildenafil) → ↓ cGMP breakdown.

Seminiferous tubules

CELL	FUNCTION	LOCATION/NOTES
Spermatogonia	Maintain germ cell pool and produce 1° spermatocytes	Line seminiferous tubules A Germ cells
Sertoli cells	Secrete inhibin B → inhibit FSH Secrete androgen-binding protein → maintain local levels of testosterone Produce MIF Tight junctions between adjacent Sertoli cells form blood-testis barrier → isolate gametes from autoimmune attack Support and nourish developing spermatozoa Regulate spermatogenesis Temperature sensitive; ↓ sperm production and ↓ inhibin B with ↑ temperature	Line seminiferous tubules Non-germ cells Convert testosterone and androstenedione to estrogens via aromatase Sertoli cells are inSide Seminiferous tubules, Support Sperm Synthesis, and inhibit FSH Homolog of female granulosa cells
Leydig cells	Secrete testosterone in the presence of LH ; testosterone production unaffected by temperature	Interstitialium Endocrine cells Homolog of female theca interna cells Leydies (ladies) dig testosterone



▶ REPRODUCTIVE—PHYSIOLOGY


Spermatogenesis

Begins at puberty with spermatogonia. Full development takes 2 months. Occurs in seminiferous tubules. Produces spermatids that undergo spermiogenesis (loss of cytoplasmic contents, gain of acrosomal cap) to form mature spermatozoa.

“Gonium” is going to be a sperm; “Zoon” is “Zooming” to egg.


Tail mobility impaired in ciliary dyskinesia/ Kartagener syndrome → infertility.

Tail mobility normal in cystic fibrosis (in CF, absent vas deferens → infertility).



Estrogen

SOURCE	Ovary (17 β -estradiol), placenta (estriol), adipose tissue (estrone via aromatization).	Potency: estradiol > estrone > estriol.
FUNCTION	<p>Development of genitalia and breast, female fat distribution.</p> <p>Growth of follicle, endometrial proliferation, \uparrow myometrial excitability.</p> <p>Upregulation of estrogen, LH, and progesterone receptors; feedback inhibition of FSH and LH, then LH surge; stimulation of prolactin secretion.</p> <p>\uparrow transport proteins, SHBG; \uparrow HDL; \downarrow LDL.</p>	<p>Pregnancy:</p> <ul style="list-style-type: none"> 50-fold \uparrow in estradiol and estrone 1000-fold \uparrow in estriol (indicator of fetal well-being) <p>Estrogen receptors expressed in cytoplasm; translocate to nucleus when bound by estrogen.</p>

**Progesterone**

SOURCE	Corpus luteum, placenta, adrenal cortex, testes.	Fall in progesterone after delivery disinhibits prolactin \rightarrow lactation. \uparrow progesterone is indicative of ovulation.
FUNCTION	<p>During luteal phase, prepares uterus for implantation of fertilized egg:</p> <ul style="list-style-type: none"> Stimulation of endometrial glandular secretions and spiral artery development Production of thick cervical mucus \rightarrow inhibits sperm entry into uterus Prevention of endometrial hyperplasia \uparrow body temperature \downarrow estrogen receptor expression \downarrow gonadotropin (LH, FSH) secretion <p>During pregnancy:</p> <ul style="list-style-type: none"> Maintenance of pregnancy \downarrow myometrial excitability \rightarrow \downarrow contraction frequency and intensity \downarrow prolactin action on breasts 	<p>Progesterone is pro-gestation.</p> <p>Prolactin is pro-lactation.</p>


Oogenesis

1° oocytes begin meiosis I during fetal life and complete meiosis I just prior to ovulation.

Meiosis I is arrested in **pr**Ophase I for years until **O**vulation (1° oocytes).

Meiosis II is arrested in **met**aphase II until fertilization (2° oocytes). “An egg **met** a sperm.”

If fertilization does not occur within 1 day, the 2° oocyte degenerates.

**Ovulation**

↑ estrogen, ↑ GnRH receptors on anterior pituitary. Estrogen surge then stimulates LH release → ovulation (rupture of follicle).

↑ temperature (progesterone induced).

Mittelschmerz—transient mid-cycle ovulatory pain (“Middle hurts”); classically associated with peritoneal irritation (eg, follicular swelling/rupture, fallopian tube contraction). Can mimic appendicitis.

Menstrual cycle


Follicular phase can vary in length. Luteal phase is 14 days. Ovulation day + 14 days = menstruation.

Follicular growth is fastest during 2nd week of the follicular phase.

Estrogen stimulates endometrial proliferation.

Progesterone maintains endometrium to support implantation.

↓ progesterone → ↓ fertility.



Abnormal uterine bleeding

Characterized as either heavy menstrual bleeding (AUB/HMB) or intermenstrual bleeding (AUB/IMB).

These are further subcategorized by **PALM-COEIN**:

- Structural causes (**PALM**): Polyp, Adenomyosis, Leiomyoma, or Malignancy/hyperplasia
- Non-structural causes (**COEIN**): Coagulopathy, Ovulatory, Endometrial, Iatrogenic, Not yet classified

Terms such as dysfunctional uterine bleeding, menorrhagia, oligomenorrhea are no longer recommended.

Pregnancy


Fertilization most commonly occurs in upper end of fallopian tube (the ampulla). Occurs within 1 day of ovulation.

Implantation within the wall of the uterus occurs 6 days after fertilization. Syncytiotrophoblasts secrete hCG, which is detectable in blood 1 week after conception and on home test in urine 2 weeks after conception.

Gestational age—calculated from date of last menstrual period.

Physiologic adaptations in pregnancy:

- ↑ GFR → ↓ BUN and creatinine, ↓ glucosuria threshold
- ↑ cardiac output (↑ preload, ↓ afterload, ↑ HR → ↑ placental and uterus perfusion)
- Anemia (↑↑ plasma, ↑ RBCs)
- Hypercoagulability (to ↓ blood loss at delivery)
- Hyperventilation (eliminate fetal CO₂)
- ↑ lipolysis and fat utilization (due to maternal hypoglycemia and insulin resistance) → preserves glucose and amino acids for utilization by the fetus



Placental hormone secretion generally increases over the course of pregnancy, but hCG peaks at 8–10 weeks.

Human chorionic gonadotropin**SOURCE**

Syncytiotrophoblast of placenta.

FUNCTION

Maintains corpus luteum (and thus progesterone) for first 8–10 weeks of pregnancy by acting like LH (otherwise no luteal cell stimulation → abortion). After 8–10 weeks, placenta synthesizes its own estriol and progesterone and corpus luteum degenerates.

Used to detect pregnancy because it appears early in urine (see above).

Has identical α subunit as LH, FSH, TSH (states of ↑ hCG can cause hyperthyroidism). β subunit is unique (pregnancy tests detect β subunit). hCG is ↑ in multiple gestations, hydatidiform moles, choriocarcinomas, and Down syndrome; hCG is ↓ in ectopic/failing pregnancy, Edwards syndrome, and Patau syndrome.

Human placental lactogen

Also called chorionic somatomammotropin.







SOURCE

Syncytiotrophoblast of placenta.

FUNCTION

Stimulates insulin production; overall ↑ insulin resistance. Gestational diabetes can occur if maternal pancreatic function cannot overcome the insulin resistance.

Apgar score

	Score 2	Score 1	Score 0
A ppearance	 Pink	 Extremities blue	 Pale or blue
P ulse	≥ 100 bpm	< 100 bpm	No pulse
G rimace	Cries and pulls away	Grimaces or weak cry	No response to stimulation
A ctivity	 Active movement	 Arms, legs flexed	 No movement
R espiration	Strong cry	Slow, irregular	No breathing

Assessment of newborn vital signs following delivery via a 10-point scale evaluated at 1 minute and 5 minutes. **A**pgar score is based on **A**ppearance, **P**ulse, **G**rimace, **A**ctivity, and **R**espiration. Apgar scores < 7 may require further evaluation. If Apgar score remains low at later time points, there is ↑ risk the child will develop long-term neurologic damage.



Infant and child development

Milestone dates are ranges that have been approximated and vary by source. Children not meeting milestones may need assessment for potential developmental delay.

AGE	MOTOR	SOCIAL	VERBAL/COGNITIVE
Infant	Parents	Start	Observing,
0–12 mo	<p>Primitive reflexes disappear—Moro (by 3 mo), rooting (by 4 mo), palmar (by 6 mo), Babinski (by 12 mo)</p> <p>Posture—lifts head up prone (by 1 mo), rolls and sits (by 6 mo), crawls (by 8 mo), stands (by 10 mo), walks (by 12–18 mo)</p> <p>Picks—passes toys hand to hand (by 6 mo), Pincer grasp (by 10 mo)</p> <p>Points to objects (by 12 mo)</p>	<p>Social smile (by 2 mo)</p> <p>Stranger anxiety (by 6 mo)</p> <p>Separation anxiety (by 9 mo)</p>	<p>Orients—first to voice (by 4 mo), then to name and gestures (by 9 mo)</p> <p>Object permanence (by 9 mo)</p> <p>Oratory—says “mama” and “dada” (by 10 mo)</p>
Toddler	Child	Rearing	Working,
12–36 mo	<p>Cruises, takes first steps (by 12 mo)</p> <p>Climbs stairs (by 18 mo)</p> <p>Cubes stacked—number = age (yr) × 3</p> <p>Cutlery—feeds self with fork and spoon (by 20 mo)</p> <p>Kicks ball (by 24 mo)</p>	<p>Recreation—parallel play (by 24–36 mo)</p> <p>Rapprochement—moves away from and returns to mother (by 24 mo)</p> <p>Realization—core gender identity formed (by 36 mo)</p>	<p>Words—uses 50–200 words by 2 yr, uses 300+ words by 3 yr.</p>
Preschool	Don't	Forget, they're still	Learning!
3–5 yr	<p>Drive—tricycle (3 wheels at 3 yr)</p> <p>Drawings—copies line or circle, stick figure (by 4 yr)</p> <p>Dexterity—hops on one foot (by 4 yr), uses buttons or zippers, grooms self (by 5 yr)</p>	<p>Freedom—comfortably spends part of day away from mother (by 3 yr)</p> <p>Friends—cooperative play, has imaginary friends (by 4 yr)</p>	<p>Language—understands 1000 words by 3 yr (3 zeros), uses complete sentences and prepositions (by 4 yr)</p> <p>Legends—can tell detailed stories (by 4 yr)</p>

Low birth weight

Defined as < 2500 g. Caused by prematurity or intrauterine growth restriction (IUGR). Associated with ↑ risk of sudden infant death syndrome (SIDS) and with ↑ overall mortality.

Lactation

After parturition and delivery of placenta, rapid ↓ in progesterone disinhibits prolactin → initiation of lactation. Suckling is required to maintain milk production and ejection, since ↑ nerve stimulation → ↑ oxytocin and prolactin.

Prolactin—induces and maintains lactation and ↓ reproductive function.

Oxytocin—assists in milk letdown; also promotes uterine contractions.

Breast milk is the ideal nutrition for infants < 6 months old. Contains maternal immunoglobulins (conferring passive immunity; mostly IgA), macrophages, lymphocytes. Breast milk reduces infant infections and is associated with ↓ risk for child to develop asthma, allergies, diabetes mellitus, and obesity. Guidelines recommend exclusively breastfed infants get vitamin D and possibly iron supplementation.

Breastfeeding ↓ maternal risk of breast and ovarian cancer and facilitates mother-child bonding.

Menopause

Diagnosed by amenorrhea for 12 months. ↓ estrogen production due to age-linked decline in number of ovarian follicles. Average age at onset is 51 years (earlier in smokers).

Usually preceded by 4–5 years of abnormal menstrual cycles. Source of estrogen (estrone) after menopause becomes peripheral conversion of androgens, ↑ androgens → hirsutism.

↑↑ FSH is specific for menopause (loss of negative feedback on FSH due to ↓ estrogen).

Hormonal changes: ↓ estrogen, ↑↑ FSH, ↑ LH (no surge), ↑ GnRH.

Causes **HAVOCS**: **H**ot flashes, **A**trophy of the **V**agina, **O**steoporosis, **C**oronary artery disease, **S**leep disturbances.

Menopause before age 40 suggests 1° ovarian insufficiency (premature ovarian failure); may occur in women who have received chemotherapy and/or radiation therapy.

Androgens

Testosterone, dihydrotestosterone (DHT), androstenedione.

SOURCE

DHT and testosterone (testis), **AnD**rostenedione (**AD**renal)

Potency: DHT > testosterone > androstenedione.

FUNCTION

Testosterone:

- Differentiation of epididymis, vas deferens, seminal vesicles (internal genitalia, except prostate)
- Growth spurt: penis, seminal vesicles, sperm, muscle, RBCs
- Deepening of voice
- Closing of epiphyseal plates (via estrogen converted from testosterone)
- Libido

DHT:

- Early—differentiation of penis, scrotum, prostate
- Late—prostate growth, balding, sebaceous gland activity

Testosterone is converted to DHT by


5α-reductase, which is inhibited by finasteride. In the male, androgens are converted to estrogen by cytochrome P-450 aromatase (primarily in adipose tissue and testis).

Aromatase is the key enzyme in conversion of androgens to estrogen.

Androgenic steroid abuse—abuse of anabolic steroids to ↑ fat-free mass, muscle strength, and performance. Suspect in men who present with changes in behavior (eg, aggression), acne, gynecomastia, ↑ Hb and Hct, small testes (exogenous testosterone → hypothalamic-pituitary-gonadal axis inhibition → ↓ intratesticular testosterone → ↓ testicular size, ↓ sperm count, azoospermia). Women may present with virilization (eg, hirsutism, acne, breast atrophy, male pattern baldness).

Tanner stages of sexual development

Tanner stage is assigned independently to genitalia, pubic hair, and breast (eg, a person can have Tanner stage 2 genitalia, Tanner stage 3 pubic hair). Earliest detectable secondary sexual characteristic is breast bud development in girls, testicular enlargement in boys.



Precocious puberty

Appearance of 2° sexual characteristics (eg, adrenarche, thelarche, menarche) before age 8 years in girls and 9 years in boys. ↑ sex hormone exposure or production → ↑ linear growth, somatic and skeletal maturation (eg, premature closure of epiphyseal plates → short stature). Types include:

- Central precocious puberty (↑ GnRH secretion): idiopathic (most common; early activation of hypothalamic-pituitary gonadal axis), CNS tumors.
- Peripheral precocious puberty (GnRH-independent; ↑ sex hormone production or exposure to exogenous sex steroids): congenital adrenal hyperplasia, estrogen-secreting ovarian tumor (eg, granulosa cell tumor), Leydig cell tumor, McCune-Albright syndrome.

▶ REPRODUCTIVE—PATHOLOGY

Sex chromosome disorders

Aneuploidy most commonly due to meiotic nondisjunction.

Klinefelter syndrome

Male, 47,XXY.

Testicular atrophy, eunuchoid body shape, tall, long extremities, gynecomastia, female hair distribution **A**. May present with developmental delay. Presence of inactivated X chromosome (Barr body). Common cause of hypogonadism seen in infertility work-up.

Dysgenesis of seminiferous tubules

→ ↓ inhibin B → ↑ FSH.

Abnormal Leydig cell function → ↓ testosterone
→ ↑ LH → ↑ estrogen.

Turner syndrome

Female, 45,XO.

Short stature (associated with *SHOX* gene, preventable with growth hormone therapy), ovarian dysgenesis (streak ovary), shield chest **B**, bicuspid aortic valve, coarctation of the aorta (femoral < brachial pulse), lymphatic defects (result in webbed neck or cystic hygroma; lymphedema in feet, hands), horseshoe kidney, high-arched palate, shortened 4th metacarpals.

Most common cause of 1° amenorrhea. No Barr body.

Menopause before menarche.

↓ estrogen leads to ↑ LH, FSH.

Sex chromosome (X, or rarely Y) loss often due to nondisjunction during meiosis or mitosis.

Meiosis errors usually occur in paternal gametes
→ sperm missing the sex chromosome.

Mitosis errors occur after zygote formation → loss of sex chromosome in some but not all cells
→ mosaic karyotype (eg. 45,X/46XX).

(45,X/46,XY) mosaicism associated with increased risk for gonadoblastoma.

Pregnancy is possible in some cases (IVF, exogenous estradiol-17β and progesterone).

Double Y males

47, XYY.

Phenotypically normal (usually undiagnosed), very tall. Normal fertility. May be associated with severe acne, learning disability, autism spectrum disorders.

Ovotesticular disorder of sex development

46,XX > 46,XY.

Both ovarian and testicular tissue present (ovotestis); ambiguous genitalia. Previously called true hermaphroditism.

Diagnosing disorders of sex hormones	Testosterone	LH	Diagnosis
	↑	↑	Defective androgen receptor
	↑	↓	Testosterone-secreting tumor, exogenous steroids
	↓	↑	Hypergonadotropic hypogonadism (1°)
	↓	↓	Hypogonadotropic hypogonadism (2°)

Other disorders of sex development	Disagreement between the phenotypic sex (external genitalia, influenced by hormonal levels) and the gonadal sex (testes vs ovaries, corresponds with Y chromosome). Includes the terms pseudohermaphrodite, hermaphrodite, and intersex.		
46,XX DSD	Ovaries present, but external genitalia are virilized or ambiguous. Due to excessive and inappropriate exposure to androgenic steroids during early gestation (eg, congenital adrenal hyperplasia or exogenous administration of androgens during pregnancy).		
46,XY DSD	Testes present, but external genitalia are female or ambiguous. Most common form is androgen insensitivity syndrome (testicular feminization).		
Disorders by physical characteristics	UTERUS	BREASTS	DISORDERS
	⊕	⊖	Hypergonadotropic hypogonadism (eg, Turner syndrome, genetic mosaicism, pure gonadal dysgenesis) Hypogonadotropic hypogonadism (eg, CNS lesions, Kallmann syndrome)
	⊖	⊕	Uterovaginal agenesis in genotypic female or androgen insensitivity in genotypic male
	⊖	⊖	Male genotype with insufficient production of testosterone


Placental aromatase deficiency	Inability to synthesize estrogens from androgens. Masculinization of female (46,XX DSD) infants (ambiguous genitalia), ↑ serum testosterone and androstenedione. Can present with maternal virilization during pregnancy (fetal androgens cross the placenta).
Androgen insensitivity syndrome	Defect in androgen receptor resulting in normal-appearing female (46,XY DSD); female external genitalia with scant axillary and pubic hair, rudimentary vagina; uterus and fallopian tubes absent due to persistence of anti-Müllerian hormone from testes. Patients develop normal functioning testes (often found in labia majora; surgically removed to prevent malignancy). ↑ testosterone, estrogen, LH (vs sex chromosome disorders).
5α-reductase deficiency	Autosomal recessive; sex limited to genetic males (46,XY DSD). Inability to convert testosterone to DHT. Ambiguous genitalia until puberty, when ↑ testosterone causes masculinization/↑ growth of external genitalia. Testosterone/estrogen levels are normal; LH is normal or ↑. Internal genitalia are normal.
Kallmann syndrome	Failure to complete puberty; a form of hypogonadotropic hypogonadism. Defective migration of neurons and subsequent failure of olfactory bulbs to develop → ↓ synthesis of GnRH in the hypothalamus; hyposmia/anosmia; ↓ GnRH, FSH, LH, testosterone. Infertility (low sperm count in males; amenorrhea in females).

Pregnancy complications

Abruptio placentae

Premature separation (partial or complete) of placenta from uterine wall before delivery of infant. Risk factors: trauma (eg, motor vehicle accident), smoking, hypertension, preeclampsia, cocaine abuse.

Presentation: **abrupt**, painful bleeding (concealed or apparent) in third trimester; possible DIC (mediated by tissue factor activation), maternal shock, fetal distress. May be life threatening for mother and fetus.

**Morbidly adherent placenta**


Defective decidual layer → abnormal attachment and separation after delivery. Risk factors: prior C-section or uterine surgery involving myometrium, inflammation, placenta previa, advanced maternal age, multiparity. Three types distinguishable by the depth of penetration:

Placenta accreta—placenta **attaches** to myometrium without penetrating it; most common type.

Placenta increta—placenta penetrates **into** myometrium.


Placenta percreta—placenta penetrates (“**perforates**”) through myometrium and into uterine serosa (invades entire uterine wall); can result in placental attachment to rectum or bladder (can result in hematuria).

Presentation: often detected on ultrasound prior to delivery. No separation of placenta after delivery → postpartum bleeding (can cause Sheehan syndrome).

**Placenta previa**


Attachment of placenta over internal cervical os. Risk factors: multiparity, prior C-section. Associated with painless third-trimester bleeding. A “**preview**” of the **placenta** is visible through cervix.

Low-lying placenta (< 2 cm from internal cervical os, but not over it) is managed differently from placenta previa.



Pregnancy complications (continued)**Vasa previa**

Fetal vessels run over, or in close proximity to, cervical os. May result in vessel rupture, exsanguination, fetal death. Presents with triad of membrane rupture, painless vaginal bleeding, fetal bradycardia (< 110 beats/min). Emergency C-section usually indicated. Frequently associated with velamentous umbilical cord insertion (cord inserts in chorioamniotic membrane rather than placenta → fetal vessels travel to placenta unprotected by Wharton jelly).

**Postpartum hemorrhage**


Due to **4 T's**: **T**one (uterine atony; most common), **T**rauma (lacerations, incisions, uterine rupture), **T**hrombin (coagulopathy), **T**issue (retained products of conception). Treatment: uterine massage, oxytocin. If refractory, surgical ligation of uterine or internal iliac artery (will preserve fertility since ovarian arteries provide collateral circulation).

Ectopic pregnancy

Implantation of fertilized ovum in a site other than the uterus, most often in ampulla of fallopian tube **A**. Suspect with history of amenorrhea, lower-than-expected rise in hCG based on dates, and sudden lower abdominal pain; confirm with ultrasound, which may show extraovarian adnexal mass. Often clinically mistaken for appendicitis. Pain +/- bleeding.

Risk factors:

- Prior ectopic pregnancy
- History of infertility
- Salpingitis (PID)
- Ruptured appendix
- Prior tubal surgery
- Smoking
- Advanced maternal age



**Amniotic fluid abnormalities****Polyhydramnios**

Too much amniotic fluid. Often idiopathic, but associated with fetal malformations (eg, esophageal/duodenal atresia, anencephaly; both result in inability to swallow amniotic fluid), maternal diabetes, fetal anemia, multiple gestations.

Oligohydramnios

Too little amniotic fluid. Associated with placental insufficiency, bilateral renal agenesis, posterior urethral valves (in males) and resultant inability to excrete urine. Any profound oligohydramnios can cause Potter sequence.

Hydatidiform mole




Cystic swelling of chorionic villi and proliferation of chorionic epithelium (only trophoblast). Presents with vaginal bleeding, emesis, uterine enlargement more than expected, pelvic pressure/pain. Associated with hCG-mediated sequelae: early preeclampsia (before 20 weeks), theca-lutein cysts, hyperemesis gravidarum, hyperthyroidism.

Treatment: dilation and curettage and methotrexate. Monitor hCG.


	Complete mole	Partial mole
KARYOTYPE	46,XX; 46,XY	69,XXX; 69,XXY; 69,XYY
COMPONENTS	Most commonly enucleated egg + single sperm (subsequently duplicates paternal DNA)	2 sperm + 1 egg
HISTOLOGY	Hydropic villi, circumferential and diffuse trophoblastic proliferation	Only some villi are hydropic, focal/minimal trophoblastic proliferation
FETAL PARTS	No	Yes (partial = fetal parts)
STAINING FOR P57 PROTEIN	⊖ (paternally imprinted)	⊕ (maternally expressed)
UTERINE SIZE	↑	—
hCG	↑↑↑↑	↑
IMAGING	“Honeycombed” uterus or “clusters of grapes” A , “snowstorm” B on ultrasound	Fetal parts
RISK OF INVASIVE MOLE	15–20%	< 5%
RISK OF CHORIOCARCINOMA	2%	Rare

Choriocarcinoma



Rare; can develop during or after pregnancy in mother or baby. Malignancy of trophoblastic tissue **A** (cytotrophoblasts, syncytiotrophoblasts); **no** chorionic villi present. ↑ frequency of bilateral/multiple theca-lutein cysts. Presents with abnormal ↑ hCG, shortness of breath, hemoptysis. Hematogenous spread to lungs → “cannonball” metastases **B**.

Treatment: methotrexate.



Hypertension in pregnancy


Gestational hypertension	BP > 140/90 mm Hg after 20th week of gestation. No pre-existing hypertension. No proteinuria or end-organ damage.	Treatment: antihypertensives (H ydralazine, α-M ethyldopa, L abetalol, N ifedipine), deliver at 37–39 weeks. H ypertensive M oms L ove N ifedipine.
Preeclampsia	New-onset hypertension with either proteinuria or end-organ dysfunction after 20th week of gestation (< 20 weeks suggests molar pregnancy). Caused by abnormal placental spiral arteries → endothelial dysfunction, vasoconstriction, ischemia. Incidence ↑ in patients with pre-existing hypertension, diabetes, chronic kidney disease, autoimmune disorders (eg, antiphospholipid antibody syndrome), age > 40 years. Complications: placental abruption, coagulopathy, renal failure, pulmonary edema, uteroplacental insufficiency; may lead to eclampsia (+ seizures) and/or HELLP syndrome.	Treatment: antihypertensives, IV magnesium sulfate (to prevent seizure); definitive is delivery of fetus. P roteinuria, R ising BP (new-onset HTN), E nd-organ dysfunction (eg, pulmonary edema).
Eclampsia	Preeclampsia + maternal seizures. Maternal death due to stroke, intracranial hemorrhage, or ARDS.	Treatment: IV magnesium sulfate, antihypertensives, immediate delivery.
HELLP syndrome	H emolysis, E levated L iver enzymes, L ow P latelets. A manifestation of severe preeclampsia. Blood smear shows schistocytes. Can lead to DIC (due to release of tissue factor from injured placenta) and hepatic subcapsular hematomas → rupture → severe hypotension.	Treatment: immediate delivery.
Gynecologic tumor epidemiology	Incidence (US)—endometrial > ovarian > cervical; cervical cancer is more common worldwide due to lack of screening or HPV vaccination. Prognosis: C ervical (best prognosis, diagnosed < 45 years old) > E ndometrial (middle-aged, about 55 years old) > O varian (worst prognosis, > 65 years).	CEO s often go from best to worst as they get older .

Vulvar pathology**Non-neoplastic**

Bartholin cyst and abscess	Due to blockage of Bartholin gland duct causing accumulation of gland fluid. May lead to abscess 2° to obstruction and inflammation A . Usually in reproductive-age females.
Lichen sclerosus	Thinning of epidermis with fibrosis/sclerosis of dermis. Presents with porcelain-white plaques with a red or violet border. Skin fragility with erosions can be observed B . Most common in postmenopausal women. Benign, but slightly increased risk for SCC.
Lichen simplex chronicus	Hyperplasia of vulvar squamous epithelium. Presents with leathery, thick vulvar skin with enhanced skin markings due to chronic rubbing or scratching. Benign, no risk of SCC.

Neoplastic


Vulvar carcinoma	Carcinoma from squamous epithelial lining of vulva C . Rare. Presents with leukoplakia, biopsy often required to distinguish carcinoma from other causes. HPV-related vulvar carcinoma—associated with high-risk HPV types 16, 18. Risk factors: multiple partners, early coitarche. Usually in reproductive-age females. Non-HPV vulvar carcinoma—usually from long-standing lichen sclerosus. Females > 70 years old.
Extramammary Paget disease	Intraepithelial adenocarcinoma. Carcinoma in situ, low risk of underlying carcinoma (vs Paget disease of the breast, which is always associated with underlying carcinoma). Presents with pruritus, erythema, crusting, ulcers D .



Imperforate hymen	Incomplete degeneration of the central portion of the hymen. Accumulation of vaginal mucus at birth → self-resolving bulge in introitus. If untreated, leads to 1° amenorrhea, cyclic abdominal pain, hematocolpos (accumulation of menstrual blood in vagina → bulging and bluish hymenal membrane).
--------------------------	---

Vaginal tumors

Vaginal squamous cell carcinoma	Usually 2° to cervical SCC; 1° vaginal carcinoma rare.
Clear cell adenocarcinoma	Affects women who had exposure to DES in utero.
Sarcoma botryoides	Embryonal rhabdomyosarcoma variant. Affects girls < 4 years old; spindle-shaped cells; desmin ⊕. Presents with clear, grape-like, polypoid mass emerging from vagina.

Cervical pathology**Dysplasia and carcinoma in situ**

Disordered epithelial growth; begins at basal layer of squamocolumnar junction (transformation zone) and extends outward. Classified as CIN 1, CIN 2, or CIN 3 (severe, irreversible dysplasia or carcinoma in situ), depending on extent of dysplasia. Associated with HPV-16 and HPV-18, which produce both the E6 gene product (inhibits *TP53*) and E7 gene product (inhibits *pRb*) (6 before 7; P before R). Koilocytes **A** are pathognomonic of HPV infection. May progress slowly to invasive carcinoma if left untreated. Typically asymptomatic (detected with Pap smear) or presents as abnormal vaginal bleeding (often postcoital).

Risk factors: multiple sexual partners, HPV, smoking, early coitarche, DES exposure, immunocompromise (eg, HIV, transplant).

Invasive carcinoma

Often squamous cell carcinoma. Pap smear can detect cervical dysplasia before it progresses to invasive carcinoma. Diagnose via colposcopy and biopsy. Lateral invasion can block ureters → hydronephrosis → renal failure.

Primary ovarian insufficiency

Also called premature ovarian failure.

Premature atresia of ovarian follicles in women of reproductive age. Most often idiopathic; associated with chromosomal abnormalities (especially in females < 30 years), autoimmunity. Need karyotype screening. Patients present with signs of menopause after puberty but before age 40. ↓ estrogen, ↑ LH, ↑ FSH.


Most common causes of anovulation

Pregnancy, polycystic ovarian syndrome, obesity, HPO axis abnormalities/immaturity, premature ovarian failure, hyperprolactinemia, thyroid disorders, eating disorders, competitive athletics, Cushing syndrome, adrenal insufficiency, chromosomal abnormalities (eg, Turner syndrome).

Functional hypothalamic amenorrhea

Also called exercise-induced amenorrhea. Severe caloric restriction, ↑ energy expenditure, and/or stress → functional disruption of pulsatile GnRH secretion → ↓ LH, FSH, estrogen. Pathogenesis includes ↓ leptin (due to ↓ fat) and ↑ cortisol (stress, excessive exercise).

Associated with eating disorders and “female athlete triad” (↓ calorie availability/excessive exercise, ↓ bone mineral density, menstrual dysfunction).

Polycystic ovarian syndrome

Hyperinsulinemia and/or insulin resistance hypothesized to alter hypothalamic hormonal feedback response → ↑ LH:FSH, ↑ androgens (eg, testosterone) from theca interna cells, ↓ rate of follicular maturation → unruptured follicles (cysts) + anovulation. Common cause of ↓ fertility in women.

Enlarged, bilateral cystic ovaries **A**; presents with amenorrhea/oligomenorrhea, hirsutism, acne, ↓ fertility. Associated with obesity, acanthosis nigricans. ↑ risk of endometrial cancer 2° to unopposed estrogen from repeated anovulatory cycles.

Treatment: cycle regulation via weight reduction (↓ peripheral estrone formation), OCPs (prevent endometrial hyperplasia due to unopposed estrogen); clomiphene (ovulation induction); spironolactone, finasteride, flutamide to treat hirsutism.

Primary dysmenorrhea


Painful menses, caused by uterine contractions to ↓ blood loss → ischemic pain. Mediated by prostaglandins. Treatment: NSAIDs.

Ovarian cysts

Follicular cyst	Distention of unruptured Graafian follicle. May be associated with hyperestrogenism, endometrial hyperplasia. Most common ovarian mass in young women.
Theca-lutein cyst	Often bilateral/multiple. Due to gonadotropin stimulation. Associated with choriocarcinoma and hydatidiform moles.

Ovarian neoplasms


Most common adnexal mass in women >55 years old. Present with abdominal distention, bowel obstruction, pleural effusion.
 Risk ↑ with advanced age, infertility, endometriosis, PCOS, genetic predisposition (eg, *BRCA1* or *BRCA2* mutations, Lynch syndrome, strong family history).
 Risk ↓ with previous pregnancy, history of breastfeeding, OCPs, tubal ligation.
 Epithelial tumors are typically serous (lined by serous epithelium natively found in fallopian tubes, and often bilateral) or mucinous (lined by mucinous epithelium natively found in cervix). Monitor response to therapy/relapse by measuring CA 125 levels (not good for screening).
 Germ cell tumors can differentiate into somatic structures (eg, teratomas), or extra-embryonic structures (eg, yolk sac tumors), or can remain undifferentiated (eg, dysgerminoma).
 Sex cord stromal tumors develop from embryonic sex cord (develops into theca and granulosa cells of follicle, Sertoli and Leydig cells of seminiferous tubules) and stromal (ovarian cortex) derivatives.



TYPE	MALIGNANT?	CHARACTERISTICS
Epithelial tumors		
Serous cystadenoma	Benign	Most common ovarian neoplasm.
Serous cystadenocarcinoma	Malignant	Most common malignant ovarian neoplasm. Psammoma bodies.
Mucinous cystadenoma	Benign	Multiloculated, large. Lined by mucus-secreting epithelium A .
Mucinous cystadenocarcinoma	Malignant	Rare. May be metastatic from appendiceal or GI tumors. Can result in pseudomyxoma peritonei (intraperitoneal accumulation of mucinous material).
Brenner tumor	Usually benign	Solid, pale yellow-tan tumor that appears encapsulated. “Coffee bean” nuclei on H&E stain.

Ovarian neoplasms (continued)

Germ cell tumors		
Dysgerminoma	Malignant	Most common in adolescents. Equivalent to male seminoma but rarer. Sheets of uniform “fried egg” cells B . Tumor markers: ↑ hCG, LDH.
Mature cystic teratoma	Benign	Also called dermoid cyst. Most common ovarian tumor in young females. Cystic mass with elements from all 3 germ layers (eg, teeth, hair, sebum) C . May be painful 2° to ovarian enlargement or torsion. Monodermal form with thyroid tissue (struma ovarii D) may present with hyperthyroidism. Malignant transformation rare (usually to squamous cell carcinoma).
Immature teratoma	Malignant, aggressive	Contains fetal tissue, neuroectoderm. Commonly diagnosed before age 20. Typically represented by immature/embryonic-like neural tissue.
Yolk sac (endodermal sinus) tumor	Malignant, aggressive	Occur in ovaries and sacrococcygeal area in children. Yellow, friable (hemorrhagic) mass. 50% have Schiller-Duval bodies (resemble glomeruli, arrow in E). Tumor marker: ↑ AFP.
Sex cord stromal tumors		
Thecoma	Benign	May produce estrogen. Usually presents as abnormal uterine bleeding in a postmenopausal woman.
Granulosa cell tumor	Malignant	Most common malignant sex cord stromal tumor. Predominantly women in their 50s. Often produces estrogen and/or progesterone and presents with postmenopausal bleeding, endometrial hyperplasia, sexual precocity (in pre-adolescents), breast tenderness. Histology shows Call -Exner bodies (granulosa cells arranged haphazardly around collections of eosinophilic fluid, resembling primordial follicles; arrow in F). “Give Granny a Call !”
Sertoli-Leydig cell tumor	Benign	Small, grey to yellow-brown mass. Resembles testicular histology with tubules/cords lined by pink Sertoli cells. May produce androgens → virilization (eg, hirsutism, male pattern baldness, clitoral enlargement).
Fibromas	Benign	Bundles of spindle-shaped fibroblasts. Meigs syndrome—triad of ovarian fibroma, ascites, pleural effusion. “Pulling” sensation in groin.




Uterine conditions**Non-neoplastic uterine conditions**


Adenomyosis	Extension of endometrial tissue (glandular) into uterine myometrium. Caused by hyperplasia of basal layer of endometrium. Presents with dysmenorrhea, AUB/HMB, and uniformly enlarged, soft, globular uterus. Treatment: GnRH agonists, hysterectomy, excision of an organized adenomyoma.
Asherman syndrome	Adhesions and/or fibrosis of the endometrium. Presents with ↓ fertility, recurrent pregnancy loss, AUB, pelvic pain. Often associated with dilation and curettage of intrauterine cavity.
Endometrial hyperplasia	Abnormal endometrial gland proliferation usually stimulated by excess estrogen. ↑ risk for endometrial carcinoma (especially with nuclear atypia). Presents as postmenopausal vaginal bleeding. ↑ risk with anovulatory cycles, hormone replacement therapy, PCOS, granulosa cell tumors.
Endometriosis	Endometrium-like glands/stroma outside endometrial cavity, most commonly in the ovary (frequently bilateral), pelvis, peritoneum (yellow-brown “powder burn” lesions). In ovary, appears as endometrioma (blood-filled “chocolate cysts” [oval structures above and below asterisks in A]). May be due to retrograde flow, metaplastic transformation of multipotent cells, transportation of endometrial tissue via lymphatic system. Characterized by cyclic pelvic pain, bleeding, dysmenorrhea, dyspareunia, dyschezia (pain with defecation), infertility; normal-sized uterus. Treatment: NSAIDs, OCPs, progestins, GnRH agonists, danazol, laparoscopic removal.
Endometritis	Inflammation of endometrium B associated with retained products of conception following delivery, miscarriage, abortion, or with foreign body (eg, IUD). Retained material is nidus for bacteria from vagina or GI tract. Chronic endometritis shows plasma cells on histology. Treatment: gentamicin + clindamycin +/- ampicillin.

Uterine neoplasms

Endometrial carcinoma	Most common gynecologic malignancy C . Presents with irregular vaginal bleeding. Two types: Endometrioid —most cases caused by unopposed estrogen exposure due to obesity, but also associated with early menarche, late menopause, nulliparity. Histology shows abnormally arranged endometrial glands. Early pathogenic events include loss of PTEN or mismatch repair proteins. Serous —associated with endometrial atrophy in postmenopausal women. Aggressive. Psammoma bodies often seen on histology. Characterized by formation of papillae and tufts.
Leiomyoma (fibroid)	Most common tumor in females. Often presents with multiple discrete tumors D . ↑ incidence in African Americans. Benign smooth muscle tumor; malignant transformation to leiomyosarcoma is rare. Estrogen sensitive; tumor size ↑ with pregnancy and ↓ with menopause. Peak occurrence at 20-40 years of age. May be asymptomatic, cause AUB, or result in miscarriage. Severe bleeding may lead to iron deficiency anemia. Whorled pattern of smooth muscle bundles with well-demarcated borders on histology E .
Leiomyosarcoma	Malignant proliferation of smooth muscle arising from myometrium; arises de novo (not from leiomyomas), usually in postmenopausal women. Exam shows single lesion with areas of necrosis.



Breast pathology



Benign breast diseases

Fibrocystic changes

Most common in premenopausal women 20-50 years old. Present with premenstrual breast pain or lumps; often bilateral and multifocal. Nonproliferative lesions include simple cysts (fluid-filled duct dilation, blue dome), papillary apocrine change/metaplasia, stromal fibrosis. Risk of cancer is usually not increased. Subtypes include:

- **Sclerosing adenosis**—acini and stromal fibrosis, associated with calcifications. Slight ↑ risk for cancer.
- **Epithelial hyperplasia**—cells in terminal ductal or lobular epithelium. ↑ risk of carcinoma with atypical cells.

Inflammatory processes


Fat necrosis—benign, usually painless, lump due to injury to breast tissue. Calcified oil cyst on mammography; necrotic fat and giant cells on biopsy. Up to 50% of patients may not report trauma.
Lactational mastitis—occurs during breastfeeding, ↑ risk of bacterial infection through cracks in nipple. *S aureus* is most common pathogen. Treat with antibiotics and continue breastfeeding.


Benign tumors

Fibroadenoma—most common in women < 35 years old. Small, well-defined, mobile mass **A**. Tumor composed of fibrous tissue and glands. ↑ size and tenderness with ↑ estrogen (eg, pregnancy, prior to menstruation). Risk of cancer is usually not increased.
Intraductal papilloma—small fibroepithelial tumor within lactiferous ducts, typically beneath areola. Most common cause of nipple discharge (serous or bloody). Slight ↑ risk for cancer.
Phyllodes tumor—large mass **B** of connective tissue and cysts with “leaf-like” lobulations **C**. Most common in 5th decade. Some may become malignant.

Gynecomastia

Breast enlargement in males due to ↑ estrogen compared with androgen activity. Physiologic in newborn, pubertal, and elderly males, but may persist after puberty. Other causes include cirrhosis, hypogonadism (eg, Klinefelter syndrome), testicular tumors, and drugs (**S**pirolactone, **H**ormones, **C**imetidine, **F**inasteride, **K**etoconazole: “**S**ome **H**ormones **C**reate **F**unny **K**nockers”).



Breast cancer

Commonly postmenopausal. Often presents as a palpable hard mass **A** most often in the upper outer quadrant. Invasive cancer can become fixed to pectoral muscles, deep fascia, Cooper ligaments, and overlying skin → nipple retraction/skin dimpling.

Usually arises from terminal duct lobular unit. Amplification/overexpression of estrogen/progesterone receptors or *c-erbB2* (HER2, an EGF receptor) is common; triple negative (ER ⊖, PR ⊖, and HER2/neu ⊖) form more aggressive.

Risk factors in women: ↑ age; history of atypical hyperplasia; family history of breast cancer; race (Caucasians at highest risk, African Americans at ↑ risk for triple ⊖ breast cancer); *BRCA1/BRCA2* mutations; ↑ estrogen exposure (eg, nulliparity); postmenopausal obesity (adipose tissue converts androstenedione to estrone); ↑ total number of menstrual cycles; absence of breastfeeding; later age of first pregnancy; alcohol intake. In men: *BRCA2* mutation, Klinefelter syndrome.

Axillary lymph node metastasis most important prognostic factor in early-stage disease.

TYPE	CHARACTERISTICS	NOTES
Noninvasive carcinomas		
Ductal carcinoma in situ	Fills ductal lumen (black arrow in B indicates neoplastic cells in duct; blue arrow shows engorged blood vessel). Arises from ductal atypia. Often seen early as microcalcifications on mammography.	Early malignancy without basement membrane penetration. Usually does not produce a mass. Comedocarcinoma —Subtype of DCIS. Cells have high-grade nuclei with extensive central necrosis C and dystrophic calcification.
Paget disease	Extension of underlying DCIS/invasive breast cancer up the lactiferous ducts and into the contiguous skin of nipple → eczematous patches over nipple and areolar skin D .	Paget cells = intraepithelial adenocarcinoma cells.
Lobular carcinoma in situ	↓ E-cadherin expression. No mass or calcifications → incidental biopsy finding.	↑ risk of cancer in either breast (vs DCIS, same breast and quadrant).
Invasive carcinomas^a		
Invasive ductal	Firm, fibrous, “rock-hard” mass with sharp margins and small, glandular, duct-like cells in desmoplastic stroma.	
Invasive lobular	↓ E-cadherin expression → orderly row of cells (“single file” E) and no duct formation. Often lacks desmoplastic response.	Often bilateral with multiple lesions in the same location. Lines of cells = Lobular.
Medullary	Large, anaplastic cells growing in sheets with associated lymphocytes and plasma cells.	Well-circumscribed tumor can mimic fibroadenoma.
Inflammatory	Dermal lymphatic space invasion → breast pain with warm, swollen, erythematous skin around exaggerated hair follicles, peau d’orange F .	Poor prognosis (50% survival at 5 years). Often mistaken for mastitis or Paget disease. Usually lacks a palpable mass.



^aAll types of invasive breast carcinoma can be either of tubular subtype (well-differentiated tubules that lack myoepithelium) or mucinous subtype (abundant extracellular mucin, seen in older women).

Penile pathology**Peyronie disease**

Abnormal curvature of penis **A** due to fibrous plaque within tunica albuginea. Associated with erectile dysfunction. Can cause pain, anxiety. Consider surgical repair or treatment with collagenase injections once curvature stabilizes. Distinct from penile fracture (rupture of corpora cavernosa due to forced bending).

Ischemic priapism

Painful sustained erection lasting > 4 hours. Associated with sickle cell disease (sickled RBCs block venous drainage of corpus cavernosum vascular channels), medications (eg, sildenafil, trazodone). Treat immediately with corporal aspiration, intracavernosal phenylephrine, or surgical decompression to prevent ischemia.

Squamous cell carcinoma

Seen in the US, but more common in Asia, Africa, South America. Precursor in situ lesions: Bowen disease (in penile shaft, presents as leukoplakia “white plaque”), erythroplasia of Queyrat (carcinoma in situ of the glans **B**, presents as erythroplakia “red plaque”). Bowenoid papulosis (carcinoma in situ of unclear malignant potential, presenting as reddish papules). Associated with uncircumcised males and HPV.


Cryptorchidism

Descent failure of one **A** or both testes; impaired spermatogenesis (since sperm develop best at temperatures < 37°C); can have normal testosterone levels (Leydig cells are mostly unaffected by temperature); associated with ↑ risk of germ cell tumors. Prematurity ↑ risk of cryptorchidism. ↓ inhibin B, ↑ FSH, ↑ LH; testosterone ↓ in bilateral cryptorchidism, normal in unilateral. Most cases resolve spontaneously; otherwise, orchiopexy performed before 2 years of age.

Testicular torsion

Rotation of testicle around spermatic cord and vascular pedicle. Commonly presents in males 12–18 years old. May occur after an inciting event (eg, trauma) or spontaneously. Characterized by acute, severe pain, high-riding testis, and absent cremasteric reflex.

Treatment: surgical correction (orchiopexy) within 6 hours, manual detorsion if surgical option unavailable in timeframe. If testis is not viable, orchiectomy. Orchiopexy, when performed, should be bilateral because the contralateral testis is at risk for subsequent torsion.

Varicocele

Dilated veins in pampiniform plexus due to ↑ venous pressure; most common cause of scrotal enlargement in adult males; most often on left side because of ↑ resistance to flow from left gonadal vein drainage into left renal vein; can cause infertility because of ↑ temperature; diagnosed by standing clinical exam/Valsalva maneuver (distension on inspection and “bag of worms” on palpation; augmented by Valsalva) or ultrasound **A**; does not transilluminate. Treatment: consider surgical ligation or embolization if associated with pain or infertility.

Extragenadal germ cell tumors Arise in midline locations. In adults, most commonly in retroperitoneum, mediastinum, pineal, and suprasellar regions. In infants and young children, sacrococcygeal teratomas are most common.


Scrotal masses Benign scrotal lesions present as testicular masses that can be transilluminated (vs solid testicular tumors).

Congenital hydrocele Common cause of scrotal swelling **A** in infants, due to incomplete obliteration of processus vaginalis. Most spontaneously resolve within 1 year. Transilluminating swelling.




Acquired hydrocele Scrotal fluid collection usually 2° to infection, trauma, tumor. If bloody → hematocele.

Spermatocele Cyst due to dilated epididymal duct or rete testis. Paratesticular fluctuant nodule.



Testicular tumors



Germ cell tumors account for ~95% of all testicular tumors. Arise from germ cells that produce sperm. Most often occur in young men. Risk factors: cryptorchidism, Klinefelter syndrome. Can present as a mixed germ cell tumor. Do not transilluminate. Usually not biopsied (risk of seeding scrotum), removed via radical orchiectomy.

Sex cord stromal tumors develop from embryonic sex cord (develops into Sertoli and Leydig cells of seminiferous tubules, theca and granulosa cells of follicle) derivatives. 5% of all testicular tumors. Mostly benign.

Testicular tumors (continued)

Germ cell tumors		
Seminoma	Malignant	Painless, homogenous testicular enlargement. Most common testicular tumor. Analogous to ovarian dysgerminoma. Does not occur in infancy. Large cells in lobules with watery cytoplasm and “fried egg” appearance on histology, ↑ placental ALP (PALP). Highly radiosensitive. Late metastasis, excellent prognosis.
Teratoma	May be malignant	Unlike in females, M ature teratoma in adult M ales may be M alignant. Benign in children.
Embryonal carcinoma	Malignant	Painful, hemorrhagic mass with necrosis. Often glandular/papillary morphology. “Pure” embryonal carcinoma is rare; most commonly mixed with other tumor types. May present with metastases. May be associated with ↑ hCG and normal AFP levels when pure (↑ AFP when mixed). Worse prognosis than seminoma.
Yolk sac (endodermal sinus) tumor	Malignant, aggressive	Yellow, mucinous. Analogous to ovarian yolk sac tumor. Schiller-Duval bodies resemble primitive glomeruli. ↑ AFP is highly characteristic. Most common testicular tumor in boys < 3 years old.
Choriocarcinoma	Malignant	Disordered syncytiotrophoblastic and cytotrophoblastic elements. Hematogenous metastases to lungs and brain. ↑ hCG, may produce gynecomastia, symptoms of hyperthyroidism (α-subunit of hCG is identical to LH, FSH, TSH).
Non-germ cell tumors		
Sertoli cell tumor	Mostly benign	Androblastoma from sex cord stroma.
Leydig cell tumor	Mostly benign	Golden brown color; contains Reinke crystals (eosinophilic cytoplasmic inclusions). Produces androgens or estrogens → gynecomastia in men, precocious puberty in boys.
Testicular lymphoma	Malignant, aggressive	Most common testicular cancer in older men. Not a 1° cancer; arises from metastatic lymphoma to testes.

Hormone levels in germ cell tumors

	SEMINOMA	YOLK SAC TUMOR	CHORIOCARCINOMA	TERATOMA	EMBRYONAL CARCINOMA
PALP	↑	–	–	–	–
AFP	–	↑↑	–	–	–/↑ (when mixed)
β-hCG	–/↑	–/↑	↑↑	–	↑

Epididymitis and orchitis

Most common causes:

- *C trachomatis* and *N gonorrhoeae* (young men)
- *E coli* and *Pseudomonas* (elderly, associated with UTI and BPH)
- Autoimmune (eg, granulomas involving seminiferous tubules)

Epididymitis

Inflammation of epididymis. Presents with localized pain and tenderness over posterior testis.
⊕ Prehn sign (pain relief with scrotal elevation). May progress to involve testis.


Orchitis

Inflammation of testis. Presents with testicular pain and swelling. Mumps orchitis ↑ infertility risk.
Rare in boys < 10 years old.

Benign prostatic hyperplasia

Common in men > 50 years old. Characterized by smooth, elastic, firm nodular enlargement (hyperplasia not hypertrophy) of periurethral (lateral and middle) lobes, which compress the urethra into a vertical slit. Not premalignant. Often presents with ↑ frequency of urination, nocturia, difficulty starting and stopping urine stream, dysuria. May lead to distention and hypertrophy of bladder, hydronephrosis, UTIs. ↑ free prostate-specific antigen (PSA).

Treatment: α_1 -antagonists (terazosin, tamsulosin), which cause relaxation of smooth muscle; 5α -reductase inhibitors (eg, finasteride); PDE-5 inhibitors (eg, tadalafil); surgical resection (eg, TURP, ablation).

**Prostatitis**


Characterized by dysuria, frequency, urgency, low back pain. Warm, tender, enlarged prostate. Acute bacterial prostatitis—in older men most common bacterium is *E coli*; in young men consider *C trachomatis*, *N gonorrhoeae*. Chronic prostatitis—either bacterial or nonbacterial (eg, 2° to previous infection, nerve problems, chemical irritation).

Prostatic adenocarcinoma

Common in men > 50 years old. Arises most often from posterior lobe (peripheral zone) of prostate gland and is most frequently diagnosed by ↑ PSA and subsequent needle core biopsies. Prostatic acid phosphatase (PAP) and PSA are useful tumor markers (↑ total PSA, with ↓ fraction of free PSA). Osteoblastic metastases in bone may develop in late stages, as indicated by lower back pain and ↑ serum ALP and PSA. Metastasis to the spine often occurs via Batson (vertebral) venous plexus.

► REPRODUCTIVE—PHARMACOLOGY

Control of reproductive hormones



8

Goserelin, leuprolide

MECHANISM	GnRH analogs. When used in pulsatile fashion act as GnRH agonists. When used in continuous fashion first transiently act as GnRH agonists (tumor flare), but subsequently act as GnRH antagonists (downregulate GnRH receptor in pituitary → ↓ FSH and ↓ LH).	Leuprolide can be used in lieu of GnRH.
CLINICAL USE	Uterine fibroids, endometriosis, precocious puberty, prostate cancer, infertility.	
ADVERSE EFFECTS	Hypogonadism, ↓ libido, erectile dysfunction, nausea, vomiting.	

Degarelix

MECHANISM	GnRH antagonist. No start-up flare.
CLINICAL USE	Prostate cancer.
ADVERSE EFFECTS	Hot flashes, liver toxicity.

Estrogens

	Ethinyl estradiol, DES, mestranol.
MECHANISM	Bind estrogen receptors.
CLINICAL USE	Hypogonadism or ovarian failure, menstrual abnormalities (combined OCPs), hormone replacement therapy in postmenopausal women.
ADVERSE EFFECTS	↑ risk of endometrial cancer (when given without progesterone), bleeding in postmenopausal women, clear cell adenocarcinoma of vagina in females exposed to DES in utero, ↑ risk of thrombi. Contraindications—ER ⊕ breast cancer, history of DVTs, tobacco use in women > 35 years old.

Selective estrogen receptor modulators

Clomiphene	Antagonist at estrogen receptors in hypothalamus. Prevents normal feedback inhibition and ↑ release of LH and FSH from pituitary, which stimulates ovulation. Used to treat infertility due to anovulation (eg, PCOS). May cause hot flashes, ovarian enlargement, multiple simultaneous pregnancies, visual disturbances.
Tamoxifen	Antagonist at breast; agonist at bone, uterus; ↑ risk of thromboembolic events (especially with smoking) and endometrial cancer. Used to treat and prevent recurrence of ER/PR ⊕ breast cancer.
Raloxifene	Antagonist at breast, uterus; agonist at bone; ↑ risk of thromboembolic events (especially with smoking) but no increased risk of endometrial cancer (vs tamoxifen); used primarily to treat osteoporosis.

Aromatase inhibitors

	Anastrozole, letrozole, exemestane.
MECHANISM	Inhibit peripheral conversion of androgens to estrogen.
CLINICAL USE	ER ⊕ breast cancer in postmenopausal women.

Hormone replacement therapy	Used for relief or prevention of menopausal symptoms (eg, hot flashes, vaginal atrophy), osteoporosis (↑ estrogen, ↓ osteoclast activity). Unopposed estrogen replacement therapy ↑ risk of endometrial cancer, progesterone/progestin is added. Possible increased cardiovascular risk.
Progestins	Levonorgestrel, medroxyprogesterone, etonogestrel, norethindrone, megestrol.
MECHANISM	Bind progesterone receptors, ↓ growth and ↑ vascularization of endometrium, thicken cervical mucus.
CLINICAL USE	Contraception (forms include pill, intrauterine device, implant, depot injection), endometrial cancer, abnormal uterine bleeding. Progestin challenge: presence of withdrawal bleeding excludes anatomic defects (eg, Asherman syndrome) and chronic anovulation without estrogen.
Antiprogestins	Mifepristone, ulipristal.
MECHANISM	Competitive inhibitors of progestins at progesterone receptors.
CLINICAL USE	Termination of pregnancy (mifepristone with misoprostol); emergency contraception (ulipristal).
Combined contraception	Progestins and ethinyl estradiol; forms include pill, patch, vaginal ring. Estrogen and progestins inhibit LH/FSH and thus prevent estrogen surge. No estrogen surge → no LH surge → no ovulation. Progestins cause thickening of cervical mucus, thereby limiting access of sperm to uterus. Progestins also inhibit endometrial proliferation → endometrium is less suitable to the implantation of an embryo. Adverse effects: breakthrough menstrual bleeding, breast tenderness, VTE, hepatic adenomas. Contraindications: smokers > 35 years old (↑ risk of cardiovascular events), patients with ↑ risk of cardiovascular disease (including history of venous thromboembolism, coronary artery disease, stroke), migraine (especially with aura), breast cancer, liver disease.
Copper intrauterine device	
MECHANISM	Produces local inflammatory reaction toxic to sperm and ova, preventing fertilization and implantation; hormone free.
CLINICAL USE	Long-acting reversible contraception. Most effective emergency contraception.
ADVERSE EFFECTS	Heavier or longer menses, dysmenorrhea. Risk of PID with insertion (contraindicated in active pelvic infection).
Tocolytics	Medications that relax the uterus; include terbutaline (β ₂ -agonist action), nifedipine (Ca ²⁺ channel blocker), indomethacin (NSAID). Used to ↓ contraction frequency in preterm labor and allow time for administration of steroids (to promote fetal lung maturity) or transfer to appropriate medical center with obstetrical care.

Danazol

MECHANISM	Synthetic androgen that acts as partial agonist at androgen receptors.
CLINICAL USE	Endometriosis, hereditary angioedema.
ADVERSE EFFECTS	Weight gain, edema, acne, hirsutism, masculinization, ↓ HDL levels, hepatotoxicity, idiopathic intracranial hypertension.

Testosterone, methyltestosterone

MECHANISM	Agonists at androgen receptors.
CLINICAL USE	Treat hypogonadism and promote development of 2° sex characteristics; stimulate anabolism to promote recovery after burn or injury.
ADVERSE EFFECTS	Masculinization in females; ↓ intratesticular testosterone in males by inhibiting release of LH (via negative feedback) → gonadal atrophy. Premature closure of epiphyseal plates. ↑ LDL, ↓ HDL.

Antiandrogens

Finasteride	5 α -reductase inhibitor (↓ conversion of testosterone to DHT). Used for BPH and male-pattern baldness. Adverse effects: gynecomastia and sexual dysfunction.	Testosterone $\xrightarrow{5\alpha\text{-reductase}}$ DHT (more potent).
Flutamide, bicalutamide, apalutamide, enzalutamide	Nonsteroidal competitive inhibitors at androgen receptors. Used for prostate carcinoma.	
Ketoconazole	Inhibits steroid synthesis (inhibits 17,20 desmolase/17 α -hydroxylase).	Used in PCOS to reduce androgenic symptoms.
Spirolactone	Inhibits steroid binding, 17,20 desmolase/17 α -hydroxylase.	Both can cause gynecomastia and amenorrhea.

Tamsulosin

α_1 -antagonist used to treat BPH by inhibiting smooth muscle contraction. Selective for $\alpha_{1A/D}$ receptors (found on prostate) vs vascular α_{1B} receptors.

Minoxidil

MECHANISM	Direct arteriolar vasodilator.
CLINICAL USE	Androgenetic alopecia (pattern baldness), severe refractory hypertension.

HIGH-YIELD SYSTEMS

Respiratory

“There’s so much pollution in the air now that if it weren’t for our lungs, there’d be no place to put it all.”

—Robert Orben

“Freedom is the oxygen of the soul.”

—Moshe Dayan

“Whenever I feel blue, I start breathing again.”

—L. Frank Baum

“Life is not the amount of breaths you take; it’s the moments that take your breath away.”

—Will Smith, *Hitch*


Group key respiratory, cardiovascular, and renal concepts together for study whenever possible. Know obstructive vs restrictive lung disorders, \dot{V}/\dot{Q} mismatch, lung volumes, mechanics of respiration, and hemoglobin physiology. Lung cancers and other causes of lung masses are high yield. Be comfortable reading basic chest x-rays, CT scans, and PFTs.

▶ Embryology	660
▶ Anatomy	662
▶ Physiology	664
▶ Pathology	671
▶ Pharmacology	686

▶ RESPIRATORY—EMBRYOLOGY

Lung development Occurs in five stages. Initial development includes development of lung bud from distal end of respiratory diverticulum during week 4. **Every Pulmonologist Can See Alveoli.**

STAGE	STRUCTURAL DEVELOPMENT	NOTES
Embryonic (weeks 4–7)	Lung bud → trachea → bronchial buds → mainstem bronchi → secondary (lobar) bronchi → tertiary (segmental) bronchi.	Errors at this stage can lead to tracheoesophageal fistula.
Pseudoglandular (weeks 5–17)	Endodermal tubules → terminal bronchioles. Surrounded by modest capillary network.	Respiration impossible, incompatible with life.
Canalicular (weeks 16–25)	Terminal bronchioles → respiratory bronchioles → alveolar ducts. Surrounded by prominent capillary network.	Airways increase in diameter. Respiration capable at 25 weeks. Pneumocytes develop starting at 20 weeks.
Saccular (week 26–birth)	Alveolar ducts → terminal sacs. Terminal sacs separated by 1° septae.	
Alveolar (week 36–8 years)	Terminal sacs → adult alveoli (due to 2° septation). In utero, “breathing” occurs via aspiration and expulsion of amniotic fluid → ↑ vascular resistance through gestation. At birth, fluid gets replaced with air → ↓ in pulmonary vascular resistance.	At birth: 20–70 million alveoli. By 8 years: 300–400 million alveoli.

**Congenital lung malformations**

Pulmonary hypoplasia Poorly developed bronchial tree with abnormal histology. Associated with congenital diaphragmatic hernia (usually left-sided), bilateral renal agenesis (Potter sequence).

Bronchogenic cysts Caused by abnormal budding of the foregut and dilation of terminal or large bronchi. Discrete, round, sharply defined, fluid-filled densities on CXR (air-filled if infected). Generally asymptomatic but can drain poorly, causing airway compression and/or recurrent respiratory infections.

Club cells

Nonciliated; low columnar/cuboidal with secretory granules. Located in bronchioles. Degrade toxins; secrete component of surfactant; act as reserve cells.

Alveolar cell types**Type I pneumocytes**



Squamous. 97% of alveolar surfaces. Thinly line the alveoli (two black arrows in **A**) for optimal gas exchange.

Type II pneumocytes

Cuboidal and clustered **A**.

2 functions:

1. Serve as stem cell precursors for 2 cell types (type I and type II cells); proliferate during lung damage.
2. Secrete surfactant from lamellar bodies (arrowheads in **B**)



Surfactant— ↓ alveolar surface tension, ↓ alveolar collapse, ↓ lung recoil, and ↑ compliance.

Composed of multiple lecithins, mainly dipalmitoylphosphatidylcholine (DPPC).

Synthesis begins ~week 20 of gestation and achieves mature levels ~week 35.


Corticosteroids important for fetal surfactant synthesis and lung development.

$$\text{Collapsing pressure (P)} = \frac{2 (\text{surface tension})}{\text{radius}}$$

Law of Laplace—Alveoli have ↑ tendency to collapse on expiration as radius ↓.

Alveolar macrophages

Phagocytose foreign materials; release cytokines and alveolar proteases. Hemosiderin-laden macrophages (“HF cells”) may be found in the setting of pulmonary edema or alveolar hemorrhage.

Neonatal respiratory distress syndrome


Surfactant deficiency → ↑ surface tension → alveolar collapse (“ground-glass” appearance of lung fields) **A**.

Risk factors: prematurity, maternal diabetes (due to ↑ fetal insulin), C-section delivery (↓ release of fetal glucocorticoids; less stressful than vaginal delivery).

Treatment: maternal steroids before birth; exogenous surfactant for infant.

Therapeutic supplemental O₂ can result in **R**etinopathy of prematurity, **I**ntraventricular hemorrhage, **B**ronchopulmonary dysplasia (**RIB**).

Screening tests for fetal lung maturity: lecithin-sphingomyelin (L/S) ratio in amniotic fluid (≥ 2 is healthy; < 1.5 predictive of NRDS), foam stability index, surfactant-albumin ratio. Persistently low O₂ tension → risk of PDA.



► RESPIRATORY—ANATOMY


Respiratory tree

Conducting zone


Large airways consist of nose, pharynx, larynx, trachea, and bronchi. Airway resistance highest in the large- to medium-sized bronchi. Small airways consist of bronchioles that further divide into terminal bronchioles (large numbers in parallel → least airway resistance).
 Warms, humidifies, and filters air but does not participate in gas exchange → “anatomic dead space.”
 Cartilage and goblet cells extend to the end of bronchi.
 Pseudostratified ciliated columnar cells primarily make up epithelium of bronchus and extend to beginning of terminal bronchioles, then transition to cuboidal cells. Clear mucus and debris from lungs (mucociliary escalator).
 Airway smooth muscle cells extend to end of terminal bronchioles (sparse beyond this point).

Respiratory zone

Lung parenchyma; consists of respiratory bronchioles, alveolar ducts, and alveoli. Participates in gas exchange.
 Mostly cuboidal cells in respiratory bronchioles, then simple squamous cells up to alveoli. Cilia terminate in respiratory bronchioles. Alveolar macrophages clear debris and participate in immune response.



Lung anatomy







Right lung has 3 lobes; Left has **Less Lobes** (2) and **Lingula** (homolog of right middle lobe). Instead of a middle lobe, left lung has a space occupied by the heart **A**.


Relation of the pulmonary artery to the bronchus at each lung hilum is described by **RALS**—**R**ight **A**nterior; **L**eft **S**uperior. Carina is posterior to ascending aorta and anteromedial to descending aorta **B**.

Right lung is a more common site for inhaled foreign bodies because right main stem bronchus is wider, more vertical, and shorter than the left. If you aspirate a peanut:

- While supine—usually enters superior segment of right lower lobe.
- While lying on right side—usually enters right upper lobe.
- While upright—usually enters right lower lobe.



Diaphragm structures



Structures perforating diaphragm:

- At T8: IVC, right phrenic nerve
- At T10: esophagus, vagus (CN 10; 2 trunks)
- At T12: aorta (red), thoracic duct (white), azygos vein (blue) (“At **T-1-2** it’s the **red, white, and blue**”)

Diaphragm is innervated by C3, 4, and 5 (phrenic nerve). Pain from diaphragm irritation (eg, air, blood, or pus in peritoneal cavity) can be referred to shoulder (C5) and trapezius ridge (C3, 4).

Number of letters = T level:

- T8: vena cava (**IVC**)
- T10: (**O**)esophagus
- T12: **aortic** hiatus

I ate (**8**) **ten** eggs at **twelve**.


C3, 4, 5 keeps the diaphragm **alive**.

Other bifurcations:

- The common carotid **bifourcates** at **C4**.
- The trachea **bifourcates** at **T4**.
- The abdominal aorta **bifourcates** at **L4**.

► RESPIRATORY—PHYSIOLOGY

Lung volumes	Note: a capacity is a sum of ≥ 2 physiologic volumes.
Inspiratory reserve volume	Air that can still be breathed in after normal inspiration
Tidal volume	Air that moves into lung with each quiet inspiration, typically 500 mL
Expiratory reserve volume	Air that can still be breathed out after normal expiration
Residual volume	Air in lung after maximal expiration; RV and any lung capacity that includes RV cannot be measured by spirometry
Inspiratory capacity	IRV + TV Air that can be breathed in after normal exhalation
Functional residual capacity	RV + ERV Volume of gas in lungs after normal expiration
Vital capacity	TV + IRV + ERV Maximum volume of gas that can be expired after a maximal inspiration
Total lung capacity	IRV + TV + ERV + RV Volume of gas present in lungs after a maximal inspiration

**Determination of physiologic dead space**

$$V_D = V_T \times \frac{P_{aCO_2} - P_{ECo_2}}{P_{aCO_2}}$$

V_D = physiologic dead space = anatomic dead space of conducting airways plus alveolar dead space; apex of healthy lung is largest contributor of alveolar dead space. Volume of inspired air that does not take part in gas exchange.

V_T = tidal volume.

P_{aCO_2} = arterial P_{CO_2} .

P_{ECo_2} = expired air P_{CO_2} .

T_{aco} , P_{aco} , P_{Eco} , P_{aco} (refers to order of variables in equation)

Physiologic dead space—approximately equivalent to anatomic dead space in normal lungs. May be greater than anatomic dead space in lung diseases with \dot{V}/\dot{Q} defects.


Ventilation

Minute ventilation	Total volume of gas entering lungs per minute $V_E = V_T \times RR$	Normal values: Respiratory rate (RR) = 12–20 breaths/min
Alveolar ventilation	Volume of gas that reaches alveoli each minute $V_A = (V_T - V_D) \times RR$	$V_T = 500$ mL/breath $V_D = 150$ mL/breath

Lung and chest wall**Elastic recoil**

Tendency for lungs to collapse inward and chest wall to spring outward.

At FRC, airway and alveolar pressures equal atmospheric pressure (called zero), and intrapleural pressure is negative (preventing atelectasis). The inward pull of the lung is balanced by the outward pull of the chest wall. System pressure is atmospheric. Pulmonary vascular resistance (PVR) is at a minimum.


**Compliance**

Change in lung volume for a change in pressure ($\Delta V/\Delta P$). Inversely proportional to wall stiffness and increased by surfactant.

- \uparrow compliance = lung easier to fill (eg, emphysema, aging)
- \downarrow compliance = lung harder to fill (eg, pulmonary fibrosis, pneumonia, ARDS, pulmonary edema)


Hysteresis

Lung inflation follows a different pressure-volume curve than lung deflation due to need to overcome surface tension forces in inflation.

**Respiratory system changes in the elderly**

Aging is associated with progressive \downarrow in lung function. TLC remains the same.

INCREASED	DECREASED
Lung compliance (loss of elastic recoil)	Chest wall compliance (\uparrow chest wall stiffness)
RV	FVC and FEV ₁
\dot{V}/\dot{Q} mismatch	Respiratory muscle strength (can impair cough)
A-a gradient	Ventilatory response to hypoxia/hypercapnia

Hemoglobin

Normal adult hemoglobin (Hb) is composed of 4 polypeptide subunits (2 α and 2 β) and exists in 2 forms:

- Deoxygenated form has low affinity for O₂, thus promoting release/unloading of O₂.
- Oxygenated form has high affinity for O₂ (300 \times). Hb exhibits positive cooperativity and positive allostery.

Hemoglobin acts as buffer for H⁺ ions.

Myoglobin is composed of a single polypeptide chain associated with one heme moiety.

Higher affinity for oxygen than Hb.

Oxygen content of blood

$$O_2 \text{ content} = (1.34 \times \text{Hb} \times \text{Sao}_2) + (0.003 \times \text{Pao}_2)$$

Hb = hemoglobin concentration; Sao₂ = arterial O₂ saturation

Pao₂ = partial pressure of O₂ in arterial blood

Normally 1 g Hb can bind 1.34 mL O₂; normal Hb amount in blood is 15 g/dL.

O₂ binding capacity ≈ 20 mL O₂/dL of blood.

With ↓ Hb there is ↓ O₂ content of arterial blood, but no change in O₂ saturation and Pao₂.

O₂ delivery to tissues = cardiac output × O₂ content of blood.

	Hb CONCENTRATION	% O ₂ SAT OF Hb	DISSOLVED O ₂ (Pao ₂)	TOTAL O ₂ CONTENT
CO poisoning	Normal	↓ (CO competes with O ₂)	Normal	↓
Anemia	↓	Normal	Normal	↓
Polycythemia	↑	Normal	Normal	↑

Methemoglobin

Iron in Hb is normally in a reduced state (ferrous Fe²⁺; “just the **2** of us”).

Oxidized form of Hb (ferric, Fe³⁺) does not bind O₂ as readily as Fe²⁺, but has ↑ affinity for cyanide → tissue hypoxia from ↓ O₂ saturation and ↓ O₂ content.

Methemoglobinemia may present with cyanosis and chocolate-colored blood.

Nitrites (eg, from dietary intake or polluted/high-altitude water sources) and benzocaine cause poisoning by oxidizing Fe²⁺ to Fe³⁺.

Methemoglobinemia can be treated with **methylene blue** and vitamin C.

Oxygen-hemoglobin dissociation curve


ODC has sigmoidal shape due to positive cooperativity (ie, tetrameric Hb molecule can bind 4 O₂ molecules and has higher affinity for each subsequent O₂ molecule bound).

Myoglobin is monomeric and thus does not show positive cooperativity; curve lacks sigmoidal appearance.

Shifting ODC to the right → ↓ Hb affinity for O₂ (facilitates unloading of O₂ to tissue) → ↑ P₅₀ (higher Po₂ required to maintain 50% saturation).

Shifting ODC to the left → ↓ O₂ unloading → renal hypoxia → ↑ EPO synthesis → compensatory erythrocytosis.

Fetal Hb (2 α and 2 γ subunits) has higher affinity for O₂ than adult Hb (due to ↓ affinity for 2,3-BPG) → dissociation curve is shifted left, driving diffusion of O₂ across the placenta from mother to fetus.




Left shift (↓ O ₂ unloading to tissue) Left = Lower	Right shift (↑ O ₂ unloading to tissues) ACE BATs right handed
↓ H ⁺ (↑ pH, base) ↓ Pco ₂ ↓ 2,3-BPG ↓ Temperature ↑ CO ↑ MetHb ↑ HbF	↑ H ⁺ (↓ pH, Acid) ↑ Pco ₂ Exercise ↑ 2,3-BPG High Altitude ↑ Temperature

Cyanide vs carbon monoxide poisoning

Both inhibit aerobic metabolism via inhibition of complex IV (cytochrome c oxidase) → hypoxia that does not fully correct with supplemental O₂ and ↑ anaerobic metabolism. Both can lead to pink or cherry red skin (usually postmortem finding), seizures, and coma.

	Cyanide	Carbon monoxide
SOURCE	Byproduct of synthetic product combustion, ingestion of amygdalin (cyanogenic glucoside found in apricot seeds) or cyanide.	Odorless gas from fires, car exhaust, or gas heaters.
TREATMENT	Hydroxocobalamin (binds cyanide → cyanocobalamin → renal excretion). Nitrites (oxidize Hb → methemoglobin → binds cyanide → cyanomethemoglobin → less toxicity). Sodium thiosulfate (↑ cyanide conversion to thiocyanate → renal excretion).	100% O ₂ , hyperbaric O ₂ .

SIGNS/SYMPTOMS




Breath has bitter almond odor; cardiovascular collapse.

Headache, dizziness. Multiple individuals may be involved (eg, family with similar symptoms in winter). Classically associated with bilateral globus pallidus lesions on MRI **A**, although rarely seen with cyanide toxicity as well.

EFFECT ON OXYGEN-HEMOGLOBIN DISSOCIATION CURVE

Curve normal; oxygen saturation may appear normal initially.

Left shift in curve → ↑ affinity for O₂ → ↓ O₂ unloading in tissues. Binds competitively to Hb with 200× greater affinity than O₂ to form carboxyhemoglobin → ↓ %O₂ saturation of Hb.



Pulmonary circulation

Normally a low-resistance, high-compliance system. A ↓ in PAO_2 causes a hypoxic vasoconstriction that shifts blood away from poorly ventilated regions of lung to well-ventilated regions of lung.

Perfusion limited— O_2 (normal health), CO_2 , N_2O . Gas equilibrates early along the length of the capillary. Exchange can be ↑ only if blood flow ↑.

Diffusion limited— O_2 (emphysema, fibrosis, exercise), CO . Gas does not equilibrate by the time blood reaches the end of the capillary.


A consequence of pulmonary hypertension is cor pulmonale and subsequent right ventricular failure.

Diffusion: $\dot{V}_{gas} = A \times D_k \times \frac{P_1 - P_2}{\Delta_x}$ where

A = area, Δ_x = alveolar wall thickness, D_k = diffusion coefficient of gas, $P_1 - P_2$ = difference in partial pressures.

- A ↓ in emphysema.
- Δ_x ↑ in pulmonary fibrosis.

DLCO is the extent to which CO passes from air sacs of lungs into blood.



P_a = partial pressure of gas in pulmonary capillary blood
 P_A = partial pressure of gas in alveolar air

**Pulmonary vascular resistance**

$$PVR = \frac{P_{\text{pulm artery}} - P_{L\text{atrium}}}{Q}$$

Remember: $\Delta P = Q \times R$, so $R = \Delta P / Q$

$$R = \frac{8\eta l}{\pi r^4}$$

$P_{\text{pulm artery}}$ = pressure in pulmonary artery
 $P_{L\text{atrium}} \approx$ pulmonary capillary wedge pressure
 Q = cardiac output (flow)
 R = resistance
 η = viscosity of blood
 l = vessel length
 r = vessel radius

Alveolar gas equation

$$PAO_2 = PIO_2 - \frac{PaCO_2}{R}$$

$$\approx 150 \text{ mm Hg}^a - \frac{PaCO_2}{0.8}$$

^aAt sea level breathing room air

PAO_2 = alveolar PO_2 (mm Hg)
 PIO_2 = PO_2 in inspired air (mm Hg)
 $PaCO_2$ = arterial PCO_2 (mm Hg)
 R = respiratory quotient = CO_2 produced/
 O_2 consumed

A-a gradient = $PAO_2 - PaO_2$. Normal A-a gradient estimated as $(\text{age}/4) + 4$ (eg, for a person <40 years old, gradient should be <14).

Oxygen deprivation

Hypoxia (↓ O ₂ delivery to tissue)	Hypoxemia (↓ Pao ₂)	Ischemia (loss of blood flow)
<ul style="list-style-type: none"> ↓ cardiac output Hypoxemia Ischemia Anemia CO poisoning 	<ul style="list-style-type: none"> Normal A-a gradient <ul style="list-style-type: none"> ▪ High altitude ▪ Hypoventilation (eg, opioid use, obesity hypoventilation syndrome) ↑ A-a gradient <ul style="list-style-type: none"> ▪ \dot{V}/\dot{Q} mismatch ▪ Diffusion limitation (eg, fibrosis) ▪ Right-to-left shunt 	<ul style="list-style-type: none"> Impeded arterial flow ↓ venous drainage

Ventilation/perfusion mismatch

Ideally, ventilation is matched to perfusion (ie, $\dot{V}/\dot{Q} = 1$) for adequate gas exchange.

Lung zones:


- \dot{V}/\dot{Q} at apex of lung = 3 (wasted ventilation)
- \dot{V}/\dot{Q} at base of lung = 0.6 (wasted perfusion)

Both ventilation and perfusion are greater at the base of the lung than at the apex of the lung. With exercise (↑ cardiac output), there is vasodilation of apical capillaries → \dot{V}/\dot{Q} ratio approaches 1.

Certain organisms that thrive in high O₂ (eg, TB) flourish in the apex.

$\dot{V}/\dot{Q} = 0$ = “airway” obstruction (shunt). In shunt, 100% O₂ does not improve Pao₂ (eg, foreign body aspiration).

$\dot{V}/\dot{Q} = \infty$ = blood flow obstruction (physiologic dead space). Assuming < 100% dead space, 100% O₂ improves Pao₂ (eg, pulmonary embolus).



Carbon dioxide transport


CO₂ is transported from tissues to lungs in 3 forms:

- ① HCO₃⁻ (70%).
- ② Carbaminohemoglobin or HbCO₂ (21–25%). CO₂ bound to Hb at N-terminus of globin (not heme). CO₂ favors deoxygenated form (O₂ unloaded).
- ③ Dissolved CO₂ (5–9%).

In lungs, oxygenation of Hb promotes dissociation of H⁺ from Hb. This shifts equilibrium toward CO₂ formation; therefore, CO₂ is released from RBCs (Haldane effect).

In peripheral tissue, ↑ H⁺ from tissue metabolism shifts curve to right, unloading O₂ (Bohr effect).

Majority of blood CO₂ is carried as HCO₃⁻ in the plasma.

**Response to high altitude**

↓ atmospheric oxygen (P_iO₂) → ↓ P_aO₂ → ↑ ventilation → ↓ P_aCO₂ → respiratory alkalosis → altitude sickness.

Chronic ↑ in ventilation.

↑ erythropoietin → ↑ Hct and Hb (due to chronic hypoxia).

↑ 2,3-BPG (binds to Hb causing rightward shift of the ODC so that Hb releases more O₂).

Cellular changes (↑ mitochondria).

↑ renal excretion of HCO₃⁻ to compensate for respiratory alkalosis (can augment with acetazolamide).

Chronic hypoxic pulmonary vasoconstriction results in pulmonary hypertension and RVH.

Response to exercise

↑ CO₂ production.

↑ O₂ consumption.

Right shift of ODC.

↑ ventilation rate to meet O₂ demand.


V̇/Q̇ ratio from apex to base becomes more uniform.

↑ pulmonary blood flow due to ↑ cardiac output.

↓ pH during strenuous exercise (2° to lactic acidosis).

No change in P_aO₂ and P_aCO₂, but ↑ in venous CO₂ content and ↓ in venous O₂ content.

▶ RESPIRATORY—PATHOLOGY

Rhinosinusitis

Obstruction of sinus drainage into nasal cavity → inflammation and pain over affected area.

Typically affects maxillary sinuses, which drain against gravity due to ostia located superomedially (red arrow points to fluid-filled right maxillary sinus in **A**).

Superior meatus—drains sphenoid, posterior ethmoid; middle meatus—drains frontal, maxillary, and anterior ethmoid; inferior meatus—drains nasolacrimal duct.

Most common acute cause is viral URI; may lead to superimposed bacterial infection, most commonly *H influenzae*, *S pneumoniae*, *M catarrhalis*.

Paranasal sinus infections may extend to the orbits, cavernous sinus, and brain, causing complications (eg, orbital cellulitis, cavernous sinus syndrome, meningitis).

Epistaxis

Nose bleed. Most commonly occurs in anterior segment of nostril (**Kiesselbach plexus**). Life-threatening hemorrhages occur in posterior segment (sphenopalatine artery, a branch of maxillary artery). Common causes include foreign body, trauma, allergic rhinitis, and nasal angiofibromas (common in adolescent males).

Kiesselbach drives his **Lexus** with his **LEGS**: superior **L**abial artery, anterior and posterior **E**thmoidal arteries, **G**reater palatine artery, **S**phenopalatine artery.

Head and neck cancer

Mostly squamous cell carcinoma. Risk factors include tobacco, alcohol, HPV-16 (oropharyngeal), EBV (nasopharyngeal). Field cancerization: carcinogen damages wide mucosal area → multiple tumors that develop independently after exposure.

Deep venous thrombosis

Blood clot within a deep vein → swelling, redness **A**, warmth, pain. Predisposed by Virchow triad (**SHE**):

- **S**tasis (eg, post-op, long drive/flight)
- **H**ypercoagulability (eg, defect in coagulation cascade proteins, such as factor V Leiden; oral contraceptive use; pregnancy)
- **E**ndothelial damage (exposed collagen triggers clotting cascade)

Most pulmonary emboli arise from proximal deep veins of lower extremity.

D-dimer lab test used clinically to rule out DVT in low-to-moderate risk patients (high sensitivity, low specificity).

Imaging test of choice is compression ultrasound with Doppler.

Use unfractionated heparin or low-molecular weight heparins (eg, enoxaparin) for prophylaxis and acute management.

Use oral anticoagulants (eg, rivaroxaban, apixaban) for treatment and long-term prevention.

Pulmonary emboli


\dot{V}/\dot{Q} mismatch, hypoxemia, respiratory alkalosis. Sudden-onset dyspnea, pleuritic chest pain, tachypnea, tachycardia. Large emboli or saddle embolus **A** may cause sudden death due to electromechanical dissociation (pulseless electrical activity). CT pulmonary angiography is imaging test of choice for PE (look for filling defects) **B**. May have S1Q3T3 abnormality on ECG. Lines of Zahn **C** are interdigitating areas of pink (platelets, fibrin) and red (RBCs) found only in thrombi formed before death; help distinguish pre- and postmortem thrombi.

Types: **F**at, **A**ir, **T**hrombus, **B**acteria, **A**mniotic fluid, **T**umor. An embolus moves like a **FAT BAT**.

Fat emboli—associated with long bone fractures and liposuction; classic triad of hypoxemia, neurologic abnormalities, petechial rash.

Air emboli—nitrogen bubbles precipitate in ascending divers (caisson disease/decompression sickness); treat with hyperbaric O₂; or, can be iatrogenic 2° to invasive procedures (eg, central line placement).

Amniotic fluid emboli—typically occurs during labor or postpartum, but can be due to uterine trauma. Can lead to DIC. Rare, but high mortality.

**Mediastinal pathology**

Normal mediastinum contains heart, thymus, lymph nodes, esophagus, and aorta.

Mediastinal masses

Some pathologies (eg, lymphoma, lung cancer, abscess) can occur in any compartment, but there are common associations:

- Anterior—**4T**'s: **T**hyroid (substernal goiter), **T**hymic neoplasm, **T**eratoma, "**T**errible" lymphoma.
- Middle—esophageal carcinoma, metastases, hiatal hernia, bronchogenic cysts.
- Posterior—neurogenic tumor (eg, neurofibroma), multiple myeloma.

Mediastinitis

Inflammation of mediastinal tissues. Commonly due to postoperative complications of cardiothoracic procedures (≤ 14 days), esophageal perforation, or contiguous spread of odontogenic/retropharyngeal infection.

Chronic mediastinitis—also known as fibrosing mediastinitis; due to \uparrow proliferation of connective tissue in mediastinum. *Histoplasma capsulatum* is common cause.


Clinical features: fever, tachycardia, leukocytosis, chest pain, and sternal wound drainage.

Pneumomediastinum

Presence of gas (usually air) in the mediastinum (black arrows show air around the aorta, red arrow shows air dissecting into the neck **A**). Can either be spontaneous (due to rupture of pulmonary bleb) or 2° (eg, trauma, iatrogenic, Boerhaave syndrome).


Ruptured alveoli allow tracking of air into the mediastinum via peribronchial and perivascular sheaths.

Clinical features: chest pain, dyspnea, voice change, subcutaneous emphysema, \oplus Hamman sign (crepitus on cardiac auscultation).




Flow-volume loops

FLOW-VOLUME PARAMETER	Obstructive lung disease	Restrictive lung disease
RV	↑	↓
FRC	↑	↓
TLC	↑	↓
FEV ₁	↓↓	↓
FVC	↓	↓
FEV ₁ /FVC	↓ FEV ₁ decreased more than FVC	Normal or ↑ FEV ₁ decreased proportionately to FVC





Obstructive lung diseases

Obstruction of air flow → air trapping in lungs. Airways close prematurely at high lung volumes → ↑ **FRC**, ↑ **RV**, ↑ **TLC**. PFTs: ↓↓ **FEV₁**, ↓ **FVC** → ↓ **FEV₁/FVC** ratio (hallmark), \dot{V}/\dot{Q} mismatch. Chronic hypoxic pulmonary vasoconstriction can lead to cor pulmonale. Chronic obstructive pulmonary disease (**COPD**) includes chronic bronchitis and emphysema. “**FRiCKin’ RV** needs some increased **TLC**, but it’s hard with **COPD!**”

TYPE	PRESENTATION	PATHOLOGY	OTHER
Chronic bronchitis (“blue bloater”)	Findings: wheezing, crackles, cyanosis (hypoxemia due to shunting), dyspnea, CO ₂ retention, 2° polycythemia.	Hypertrophy and hyperplasia of mucus-secreting glands in bronchi → Reid index (thickness of mucosal gland layer to thickness of wall between epithelium and cartilage) > 50%. DLCO usually normal.	Diagnostic criteria: productive cough for ≥ 3 months in a year for > 2 consecutive years.
Emphysema (“pink puffer”) 	Findings: barrel-shaped chest D , exhalation through pursed lips (increases airway pressure and prevents airway collapse).	Centriacinar—affects respiratory bronchioles while sparing distal alveoli, associated with smoking A B . Frequently in upper lobes (smoke rises up) . Panacinar—affects respiratory bronchioles and alveoli, associated with α ₁ -antitrypsin deficiency. Frequently in lower lobes. Enlargement of air spaces ↓ recoil, ↑ compliance, ↓ DLCO from destruction of alveolar walls (arrow in C) and ↓ blood volume in pulmonary capillaries. Imbalance of proteases and antiproteases → ↑ elastase activity → ↑ loss of elastic fibers → ↑ lung compliance.	CXR: ↑ AP diameter, flattened diaphragm, ↑ lung field lucency.
Asthma	Findings: cough, wheezing, tachypnea, dyspnea, hypoxemia, ↓ inspiratory/expiratory ratio, pulsus paradoxus, mucus plugging E . Triggers: viral URIs, allergens, stress.	Hyperresponsive bronchi → reversible bronchoconstriction. Smooth muscle hypertrophy and hyperplasia, Curschmann spirals F (shed epithelium forms whorled mucous plugs), and Charcot-Leyden crystals G (eosinophilic, hexagonal, double-pointed crystals formed from breakdown of eosinophils in sputum). DLCO normal or ↑.	Type I hypersensitivity reaction. Diagnosis supported by spirometry and methacholine challenge. NSAID-exacerbated respiratory disease is a combination of COX inhibition (leukotriene overproduction → airway constriction), chronic sinusitis with nasal polyps, and asthma symptoms.

Obstructive lung diseases (continued)

TYPE	PRESENTATION	PATHOLOGY	OTHER
Bronchiectasis	Findings: purulent sputum, recurrent infections (most often <i>P aeruginosa</i>), hemoptysis, digital clubbing.	Chronic necrotizing infection of bronchi or obstruction → permanently dilated airways.	Associated with bronchial obstruction, poor ciliary motility (eg, smoking, Kartagener syndrome), cystic fibrosis [H], allergic bronchopulmonary aspergillosis.

**Restrictive lung diseases**

Restricted lung expansion causes ↓ lung volumes (↓ FVC and TLC). PFTs: ↑ FEV₁/FVC ratio. Patient presents with short, shallow breaths.

Types:

- Poor breathing mechanics (extrapulmonary, normal D_{LCO}, normal A-a gradient):
 - Poor muscular effort—polio, myasthenia gravis, Guillain-Barré syndrome
 - Poor structural apparatus—scoliosis, morbid obesity
- Interstitial lung diseases (pulmonary, ↓ D_{LCO}, ↑ A-a gradient):
 - Pneumoconioses (eg, coal workers' pneumoconiosis, silicosis, asbestosis)
 - Sarcoidosis: bilateral hilar lymphadenopathy, noncaseating granulomas; ↑ ACE and Ca²⁺
 - Idiopathic pulmonary fibrosis (repeated cycles of lung injury and wound healing with ↑ collagen deposition, “honeycomb” lung appearance [red arrows in A], traction bronchiectasis [blue arrow in A] and digital clubbing).
 - Granulomatosis with polyangiitis (Wegener)
 - Pulmonary Langerhans cell histiocytosis (eosinophilic granuloma)
 - Hypersensitivity pneumonitis
 - Drug toxicity (eg, bleomycin, busulfan, amiodarone, methotrexate)


Hypersensitivity pneumonitis—mixed type III/IV hypersensitivity reaction to environmental antigen. Causes dyspnea, cough, chest tightness, fever, headache. Often seen in farmers and those exposed to birds. Reversible in early stages if stimulus is avoided.

Sarcoidosis

Characterized by immune-mediated, widespread noncaseating granulomas **A**, elevated serum ACE levels, and elevated CD4/CD8 ratio in bronchoalveolar lavage fluid. More common in African-American females. Often asymptomatic except for enlarged lymph nodes. CXR shows bilateral adenopathy and coarse reticular opacities **B**; CT of the chest better demonstrates the extensive hilar and mediastinal adenopathy **C**.


Associated with **Bell palsy**, **Uveitis**, **Granulomas** (noncaseating epithelioid, containing microscopic Schaumann and asteroid bodies), **Lupus pernio** (skin lesions on face resembling lupus), **Interstitial fibrosis** (restrictive lung disease), **Erythema nodosum**, **Rheumatoid arthritis-like arthropathy**, hypercalcemia (due to ↑ 1 α -hydroxylase-mediated vitamin D activation in macrophages). A **facial droop** is **UGLIER**.

Treatment: steroids (if symptomatic).

**Inhalation injury and sequelae**

Complication of inhalation of noxious stimuli (eg, smoke). Caused by heat, particulates (< 1 μ m diameter), or irritants (eg, NH_3) → chemical tracheobronchitis, edema, pneumonia, ARDS. Many patients present 2° to burns, CO inhalation, cyanide poisoning, or arsenic poisoning. Singed nasal hairs or soot in oropharynx common on exam.

Bronchoscopy shows severe edema, congestion of bronchus, and soot deposition (**A**, 18 hours after inhalation injury; **B**, resolution at 11 days after injury).




Pneumoconioses

Asbestos is from the **roof** (was common in insulation), but affects the **base** (lower lobes).

Silica and **coal** are from the **base** (earth), but affect the **roof** (upper lobes).

Asbestosis	Associated with shipbuilding, roofing, plumbing. “Ivory white,” calcified, supradiaphragmatic A and pleural B plaques are pathognomonic of asbestosis. Risk of bronchogenic carcinoma > risk of mesothelioma. ↑ risk of Caplan syndrome (rheumatoid arthritis and pneumoconioses with intrapulmonary nodules).	Affects lower lobes. Asbestos (ferruginous) bodies are golden-brown fusiform rods resembling dumbbells C , found in alveolar sputum sample, visualized using Prussian blue stain, often obtained by bronchoalveolar lavage. ↑ risk of pleural effusions.
Berylliosis	Associated with exposure to beryllium in aerospace and manufacturing industries. Granulomatous (noncaseating) D on histology and therefore occasionally responsive to steroids. ↑ risk of cancer and cor pulmonale.	Affects upper lobes.
Coal workers’ pneumoconiosis	Prolonged coal dust exposure → macrophages laden with carbon → inflammation and fibrosis. Also known as black lung disease. ↑ risk of Caplan syndrome.	Affects upper lobes. Small, rounded nodular opacities seen on imaging. Anthracosis —asymptomatic condition found in many urban dwellers exposed to sooty air.
Silicosis	Associated with sandblasting, foundries, mines. Macrophages respond to silica and release fibrogenic factors, leading to fibrosis. It is thought that silica may disrupt phagolysosomes and impair macrophages, increasing susceptibility to TB. ↑ risk of cancer, cor pulmonale, and Caplan syndrome.	Affects upper lobes. “Eggshell” calcification of hilar lymph nodes on CXR. The silly egg sandwich I found is mine!



Mesothelioma



Malignancy of the pleura associated with asbestosis. May result in hemorrhagic pleural effusion (exudative), pleural thickening **A**.

Psammoma bodies seen on histology. Calretinin and cytokeratin 5/6 ⊕ in almost all mesotheliomas, ⊖ in most carcinomas. Smoking not a risk factor.

Acute respiratory distress syndrome

PATHOPHYSIOLOGY

Alveolar insult → release of pro-inflammatory cytokines → neutrophil recruitment, activation, and release of toxic mediators (eg, reactive oxygen species, proteases, etc) → capillary endothelial damage and ↑ vessel permeability → leakage of protein-rich fluid into alveoli → formation of intra-alveolar hyaline membranes (arrows in **A**) and noncardiogenic pulmonary edema (normal PCWP).

Loss of surfactant also contributes to alveolar collapse.

CAUSES

Sepsis (most common), aspiration, pneumonia, trauma, pancreatitis.

DIAGNOSIS

Diagnosis of exclusion with the following criteria (**ARDS**):

- **A** Abnormal chest X-ray (bilateral lung opacities) **B**
- **R**espiratory failure within 1 week of alveolar insult
- **D**ecreased Pao₂/FiO₂ (ratio < 300, hypoxemia due to ↑ intrapulmonary shunting and diffusion abnormalities)
- **S**ymptoms of respiratory failure are not due to HF/fluid overload


CONSEQUENCES

Impaired gas exchange, ↓ lung compliance; pulmonary hypertension.

MANAGEMENT

Treat the underlying cause.


Mechanical ventilation: ↓ tidal volume, ↑ PEEP.



Sleep apnea	Repeated cessation of breathing > 10 seconds during sleep → disrupted sleep → daytime somnolence. Diagnosis confirmed by sleep study. Nocturnal hypoxia → systemic/pulmonary hypertension, arrhythmias (atrial fibrillation/flutter), sudden death. Hypoxia → ↑ EPO release → ↑ erythropoiesis.
Obstructive sleep apnea	Respiratory effort against airway obstruction. Normal PaO ₂ during the day. Associated with obesity, loud snoring, daytime sleepiness. Caused by excess parapharyngeal tissue in adults, adenotonsillar hypertrophy in children. Treatment: weight loss, CPAP, dental devices.
Central sleep apnea	Impaired respiratory effort due to CNS injury/toxicity, HF, opioids. May be associated with Cheyne-Stokes respirations (oscillations between apnea and hyperpnea). Think 3 C's: C ongestive HF, C NS toxicity, C heyne-Stokes respirations. Treat with positive airway pressure.
Obesity hypoventilation syndrome	Obesity (BMI ≥ 30 kg/m ²) → hypoventilation → ↑ PaCO ₂ during waking hours (retention); ↓ PaO ₂ and ↑ PaCO ₂ during sleep. Also known as Pickwickian syndrome.
Pulmonary hypertension	Normal mean pulmonary artery pressure = 10–14 mm Hg; pulmonary hypertension ≥ 25 mm Hg at rest. Results in arteriosclerosis, medial hypertrophy, intimal fibrosis of pulmonary arteries, plexiform lesions. Course: severe respiratory distress → cyanosis and RVH → death from decompensated cor pulmonale.
ETIOLOGIES	
Pulmonary arterial hypertension	Often idiopathic. Heritable PAH can be due to an inactivating mutation in <i>BMPR2</i> gene (normally inhibits vascular smooth muscle proliferation); poor prognosis. Pulmonary vasculature endothelial dysfunction results in ↑ vasoconstrictors (eg, endothelin) and ↓ vasodilators (eg, NO and prostacyclins). Other causes include drugs (eg, amphetamines, cocaine), connective tissue disease, HIV infection, portal hypertension, congenital heart disease, schistosomiasis.
Left heart disease	Causes include systolic/diastolic dysfunction and valvular disease.
Lung diseases or hypoxia	Destruction of lung parenchyma (eg, COPD), lung inflammation/fibrosis (eg, interstitial lung diseases), hypoxemic vasoconstriction (eg, obstructive sleep apnea, living in high altitude).
Chronic thromboembolic	Recurrent microthrombi → ↓ cross-sectional area of pulmonary vascular bed.
Multifactorial	Causes include hematologic, systemic, and metabolic disorders, along with compression of the pulmonary vasculature by a tumor.


Physical findings in select lung diseases

ABNORMALITY	BREATH SOUNDS	PERCUSSION	FREMITUS	TRACHEAL DEVIATION
Pleural effusion	↓	Dull	↓	None if small Away from side of lesion if large
Atelectasis	↓	Dull	↓	Toward side of lesion
Simple pneumothorax	↓	Hyperresonant	↓	None
Tension pneumothorax	↓	Hyperresonant	↓	Away from side of lesion
Consolidation (lobar pneumonia, pulmonary edema)	Bronchial breath sounds; late inspiratory crackles, egophony, whispered pectoriloquy	Dull	↑	None

Atelectasis

Alveolar collapse (right upper lobe collapse against mediastinum in **A**). Multiple causes:

- Obstructive—airway obstruction prevents new air from reaching distal airways, old air is resorbed (eg, foreign body, mucous plug, tumor)
- Compressive—external compression on lung decreases lung volumes (eg, space-occupying lesion, pleural effusion)
- Contraction (cicatrization)—scarring of lung parenchyma that distorts alveoli (eg, sarcoidosis)
- Adhesive—due to lack of surfactant (eg, NRDS in premature babies)



Pleural effusions

Excess accumulation of fluid **A** between pleural layers → restricted lung expansion during inspiration. Can be treated with thoracentesis to remove/reduce fluid **B**.

Lymphatic



Also known as chylothorax. Due to thoracic duct injury from trauma or malignancy. Milky-appearing fluid; ↑ triglycerides.

Exudate


↑ protein content (> 2.9 g/dL), cloudy (cellular). Due to malignancy, inflammation/infection (eg, pneumonia, collagen vascular disease), trauma (occurs in states of ↑ vascular permeability). Must be drained due to risk of infection.

Transudate

↓ protein content (< 2.5 g/dL), clear (hypoalbuminemic). Due to ↑ hydrostatic pressure (eg, HF, Na⁺ retention) or ↓ oncotic pressure (eg, nephrotic syndrome, cirrhosis).




Pneumothorax	Accumulation of air in pleural space A . Dyspnea, uneven chest expansion. Chest pain, ↓ tactile fremitus, hyperresonance, and diminished breath sounds, all on the affected side.
Primary spontaneous pneumothorax	Due to rupture of apical subpleural bleb or cysts. Occurs most frequently in tall, thin, young males and smokers.
Secondary spontaneous pneumothorax	Due to diseased lung (eg, bullae in emphysema, infections), mechanical ventilation with use of high pressures → barotrauma.
Traumatic pneumothorax	Caused by blunt (eg, rib fracture), penetrating (eg, gunshot), or iatrogenic (eg, central line placement, lung biopsy, barotrauma due to mechanical ventilation) trauma.
Tension pneumothorax	Can be from any of the above. Air enters pleural space but cannot exit. Increasing trapped air → tension pneumothorax. Trachea deviates away from affected lung B . May lead to increased intrathoracic pressure → mediastinal displacement → kinking of IVC → ↓ venous return → ↓ cardiac output. Needs immediate needle decompression and chest tube placement.



Pneumonia

TYPE	TYPICAL ORGANISMS	CHARACTERISTICS
Lobar pneumonia	<i>S pneumoniae</i> most frequently, also <i>Legionella</i> , <i>Klebsiella</i>	Intra-alveolar exudate → consolidation A ; may involve entire lobe B or the whole lung.
Bronchopneumonia	<i>S pneumoniae</i> , <i>S aureus</i> , <i>H influenzae</i> , <i>Klebsiella</i>	Acute inflammatory infiltrates C from bronchioles into adjacent alveoli; patchy distribution involving ≥ 1 lobe D .
Interstitial (atypical) pneumonia	<i>Mycoplasma</i> , <i>Chlamydophila pneumoniae</i> , <i>Chlamydophila psittaci</i> , <i>Legionella</i> , viruses (RSV, CMV, influenza, adenovirus)	Diffuse patchy inflammation localized to interstitial areas at alveolar walls; CXR shows bilateral multifocal opacities E . Generally follows a more indolent course (“walking” pneumonia).
Cryptogenic organizing pneumonia	Etiology unknown. Secondary organizing pneumonia is caused by chronic inflammatory diseases (eg, rheumatoid arthritis) or medication side effects (eg, amiodarone). ⊖ sputum and blood cultures, often responds to steroids but not to antibiotics.	Formerly known as bronchiolitis obliterans organizing pneumonia (BOOP). Noninfectious pneumonia characterized by inflammation of bronchioles and surrounding structure.

**Natural history of lobar pneumonia**

	Congestion	Red hepatization	Gray hepatization	Resolution
DAYS	1–2	3–4	5–7	8+
FINDINGS	Red-purple, partial consolidation of parenchyma Exudate with mostly bacteria	Red-brown consolidation Exudate with fibrin, bacteria, RBCs, WBCs Reversible	Uniformly gray Exudate full of WBCs, lysed RBCs, and fibrin	Enzymatic digestion of exudate by macrophages


Lung cancer


Leading cause of cancer death.
 Presentation: cough, hemoptysis, bronchial obstruction, wheezing, pneumonic “coin” lesion on CXR or noncalcified nodule on CT.
 Sites of metastases from lung cancer: **L**iver (jaundice, hepatomegaly), **A**drenals, **B**one (pathologic fracture), **B**rain; “Lung ‘mets’ **L**ove **A**ffective **B**oneheads and **B**rainiacs.”
 In the lung, metastases (usually multiple lesions) are more common than 1° neoplasms. Most often from breast, colon, prostate, and bladder cancer.

SPHERE of complications:

- S**uperior vena cava/thoracic outlet syndromes
 - P**ancoast tumor
 - H**orner syndrome
 - E**ndocrine (paraneoplastic)
 - R**ecurrent laryngeal nerve compression (hoarseness)
 - E**ffusions (pleural or pericardial)
- Risk factors include smoking, secondhand smoke, radon, asbestos, family history.
Squamous and **S**mall cell carcinomas are **S**entral (central) and often caused by **S**moking.

TYPE	LOCATION	CHARACTERISTICS	HISTOLOGY
Small cell			
Small cell (oat cell) carcinoma	Central	Undifferentiated → very aggressive. May produce A CTH (Cushing syndrome), A DH (SIADH), or A ntibodies against presynaptic Ca ²⁺ channels (Lambert-Eaton myasthenic syndrome) or neurons (paraneoplastic myelitis, encephalitis, subacute cerebellar degeneration). A mplification of <i>myc</i> oncogenes common. Managed with chemotherapy +/- radiation.	Neoplasm of neuroendocrine Kulchitsky cells → small dark blue cells A . Chromogranin A ⊕, neuron-specific enolase ⊕, synaptophysin ⊕.
Non-small cell			
Adenocarcinoma	Peripheral	Most common 1° lung cancer. More common in women than men, most likely to arise in nonsmokers. Activating mutations include <i>KRAS</i> , <i>EGFR</i> , and <i>ALK</i> . Associated with hypertrophic osteoarthropathy (clubbing). Bronchioloalveolar subtype (adenocarcinoma in situ): CXR often shows hazy infiltrates similar to pneumonia; better prognosis.	Glandular pattern on histology, often stains mucin ⊕ B . Bronchioloalveolar subtype: grows along alveolar septa → apparent “thickening” of alveolar walls. Tall, columnar cells containing mucus.
Squamous cell carcinoma	Central	Hilar mass C arising from bronchus; C avitation; C igarettes; hyper C alcemia (produces PTHrP).	Keratin pearls D and intercellular bridges.
Large cell carcinoma	Peripheral	Highly anaplastic undifferentiated tumor; poor prognosis. Less responsive to chemotherapy; removed surgically. Strong association with smoking.	Pleomorphic giant cells E .
Bronchial carcinoid tumor	Central or peripheral	Excellent prognosis; metastasis rare. Symptoms due to mass effect or carcinoid syndrome (flushing, diarrhea, wheezing).	Nests of neuroendocrine cells; chromogranin A ⊕.




Lung abscess

Localized collection of pus within parenchyma **A**. Caused by aspiration of oropharyngeal contents (especially in patients predisposed to loss of consciousness [eg, alcoholics, epileptics]) or bronchial obstruction (eg, cancer).

Air-fluid levels **B** often seen on CXR; presence suggests cavitation. Due to anaerobes (eg, *Bacteroides*, *Fusobacterium*, *Peptostreptococcus*) or *S aureus*.

Treatment: antibiotics, drainage, or surgery.


Lung abscess 2° to aspiration is most often found in right lung. Location depends on patient's position during aspiration: RLL if upright, RUL or RML if recumbent.

Pancoast tumor


Also known as superior sulcus tumor. Carcinoma that occurs in the apex of lung **A** may cause Pancoast syndrome by invading/compressing local structures.

Compression of locoregional structures may cause array of findings:

- Recurrent laryngeal nerve → hoarseness
- Stellate ganglion → Horner syndrome (ipsilateral ptosis, miosis, anhidrosis)
- Superior vena cava → SVC syndrome
- Brachiocephalic vein → brachiocephalic syndrome (unilateral symptoms)
- Brachial plexus → sensorimotor deficits
- Phrenic nerve → hemidiaphragm paralysis (hemidiaphragm elevation on CXR)

Superior vena cava syndrome



An obstruction of the SVC that impairs blood drainage from the head ("facial plethora"; note blanching after fingertip pressure in **A**), neck (jugular venous distention), and upper extremities (edema). Commonly caused by malignancy (eg, mediastinal mass, Pancoast tumor) and thrombosis from indwelling catheters **B**. Medical emergency. Can raise intracranial pressure (if obstruction is severe) → headaches, dizziness, ↑ risk of aneurysm/rupture of intracranial arteries.



▶ RESPIRATORY—PHARMACOLOGY

Histamine-1 blockers	Reversible inhibitors of H ₁ histamine receptors.	
First generation	Diph en hydram ine , dimen en hydrin ate , chlorph en iramin ine , doxylamin ine .	Names usually contain “-en/-ine” or “-en/-ate.”
CLINICAL USE	Allergy, motion sickness, sleep aid.	
ADVERSE EFFECTS	Sedation, antimuscarinic, anti- α -adrenergic.	
Second generation	Lorat ad ine, fexofen ad ine, deslorat ad ine, cetirizine.	Names usually end in “-adine.”
CLINICAL USE	Allergy.	
ADVERSE EFFECTS	Far less sedating than 1st generation because of ↓ entry into CNS.	
Guafenesin	Expectorant—thins respiratory secretions; does not suppress cough reflex.	
N-acetylcysteine	Mucolytic—liquifies mucus in chronic bronchopulmonary diseases (eg, COPD, CF) by disrupting disulfide bonds. Also used as an antidote for acetaminophen overdose.	
Dextromethorphan	Antitussive (antagonizes NMDA glutamate receptors). Synthetic codeine analog. Has mild opioid effect when used in excess. Naloxone can be given for overdose. Mild abuse potential. May cause serotonin syndrome if combined with other serotonergic agents.	
Pseudoephedrine, phenylephrine	α -adrenergic agonists.	
MECHANISM	α -adrenergic agonists.	
CLINICAL USE	Reduce hyperemia, edema (used as nasal decongestants); open obstructed eustachian tubes.	
ADVERSE EFFECTS	Hypertension. Rebound congestion if used more than 4–6 days. Can also cause CNS stimulation/anxiety (pseudoephedrine).	
Pulmonary hypertension drugs		
DRUG	MECHANISM	CLINICAL NOTES
Endothelin receptor antagonists	Competitively antagonizes endothelin-1 receptors → ↓ pulmonary vascular resistance.	Hepatotoxic (monitor LFTs). Example: bosentan.
PDE-5 inhibitors	Inhibits PDE-5 → ↑ cGMP → prolonged vasodilatory effect of NO.	Also used to treat erectile dysfunction. Contraindicated when taking nitroglycerin or other nitrates (due to risk of severe hypotension). Example: sildenafil.
Prostacyclin analogs	PGI ₂ (prostacyclin) with direct vasodilatory effects on pulmonary and systemic arterial vascular beds. Inhibits platelet aggregation.	Side effects: flushing, jaw pain. Examples: epoprostenol, iloprost.

Asthma drugs	Bronchoconstriction is mediated by (1) inflammatory processes and (2) parasympathetic tone; therapy is directed at these 2 pathways.
β₂-agonists	Albuterol —relaxes bronchial smooth muscle (short acting β ₂ -agonist). For acute exacerbations. Can cause tremor, arrhythmia. Salmeterol, formoterol —long-acting agents for prophylaxis. Can cause tremor, arrhythmia.
Inhaled corticosteroids	Fluticasone, budesonide —inhibit the synthesis of virtually all cytokines. Inactivate NF-κB, the transcription factor that induces production of TNF-α and other inflammatory agents. 1st-line therapy for chronic asthma. Use a spacer or rinse mouth after use to prevent oral thrush.
Muscarinic antagonists	Tiotropium, ipratropium —competitively block muscarinic receptors, preventing bronchoconstriction. Also used for COPD. Tiotropium is long acting.
Antileukotrienes	Montelukast, zafirlukast —block leukotriene receptors (CysLT1). Especially good for aspirin-induced and exercise-induced asthma. Zileuton —5-lipoxygenase pathway inhibitor. Blocks conversion of arachidonic acid to leukotrienes. Hepatotoxic.
Anti-IgE monoclonal therapy	Omalizumab —binds mostly unbound serum IgE and blocks binding to FcεRI. Used in allergic asthma with ↑ IgE levels resistant to inhaled steroids and long-acting β ₂ -agonists.
Methylxanthines	Theophylline —likely causes bronchodilation by inhibiting phosphodiesterase → ↑ cAMP levels due to ↓ cAMP hydrolysis. Limited use due to narrow therapeutic index (cardiotoxicity, neurotoxicity); metabolized by cytochrome P-450. Blocks actions of adenosine.
Chronones	Cromolyn —prevents mast cell degranulation. Prevents acute asthma symptoms. Rarely used.
Anti-IL-5 monoclonal therapy	Prevents eosinophil differentiation, maturation, activation, and survival mediated by IL-5 stimulation. For maintenance therapy in severe eosinophilic asthma. Mepolizumab, reslizumab —against IL-5. Benralizumab —against IL-5 receptor α.



HIGH-YIELD SYSTEMS

Rapid Review

“Study without thought is vain: thought without study is dangerous.”

—Confucius

“It is better, of course, to know useless things than to know nothing.”

—Lucius Annaeus Seneca

“For every complex problem there is an answer that is clear, simple, and wrong.”

—H. L. Mencken

The following tables represent a collection of high-yield associations between diseases and their clinical findings, treatments, and key associations. They can be quickly reviewed in the days before the exam.

▶ Classic Presentations	690
▶ Classic Labs/ Findings	695
▶ Classic/Relevant Treatments	699
▶ Key Associations	702
▶ Equation Review	707
▶ Easily Confused Medications	709

► CLASSIC PRESENTATIONS

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Gout, intellectual disability, self-mutilating behavior in a boy	Lesch-Nyhan syndrome (HGPRT deficiency, X-linked recessive)	37
Situs inversus, chronic sinusitis, bronchiectasis, infertility	Kartagener syndrome (dynein arm defect affecting cilia)	49
Blue sclera	Osteogenesis imperfecta (type I collagen defect)	51
Elastic skin, hypermobility of joints, ↑ bleeding tendency	Ehlers-Danlos syndrome (type V collagen defect, type III collagen defect seen in vascular subtype of ED)	51
Arachnodactyly, lens dislocation (upward and temporal), aortic dissection, hyperflexible joints	Marfan syndrome (fibrillin defect)	52
Café-au-lait spots (unilateral), polyostotic fibrous dysplasia, precocious puberty, multiple endocrine abnormalities	McCune-Albright syndrome (G_s -protein activating mutation)	57
Calf pseudohypertrophy	Muscular dystrophy (most commonly Duchenne, due to X-linked recessive frameshift mutation of dystrophin gene)	61
Child uses arms to stand up from squat	Duchenne muscular dystrophy (Gowers sign)	61
Slow, progressive muscle weakness in boys	Becker muscular dystrophy (X-linked non-frameshift deletions in dystrophin; less severe than Duchenne)	61
Infant with cleft lip/palate, microcephaly or holoprosencephaly, polydactyly, cutis aplasia	Patau syndrome (trisomy 13)	63
Infant with microcephaly, rocker-bottom feet, clenched hands, and structural heart defect	Edwards syndrome (trisomy 18)	63
Single palmar crease	Down syndrome	63
Dilated cardiomyopathy, edema, alcoholism or malnutrition	Wet beriberi (thiamine [vitamin B ₁] deficiency)	66
Dermatitis, dementia, diarrhea	Pellagra (niacin [vitamin B ₃] deficiency)	67
Swollen gums, mucosal bleeding, poor wound healing, petechiae	Scurvy (vitamin C deficiency: can't hydroxylate proline/lysine for collagen synthesis)	69
Chronic exercise intolerance with myalgia, fatigue, painful cramps, myoglobinuria	McArdle disease (skeletal muscle glycogen phosphorylase deficiency)	87
Infant with hypoglycemia, hepatomegaly	Cori disease (debranching enzyme deficiency) or Von Gierke disease (glucose-6-phosphatase deficiency, more severe)	87
Myopathy (infantile hypertrophic cardiomyopathy), exercise intolerance	Pompe disease (lysosomal α -1,4-glucosidase deficiency)	87
“Cherry-red spots” on macula	Tay-Sachs (ganglioside accumulation) or Niemann-Pick (sphingomyelin accumulation), central retinal artery occlusion	88
Hepatosplenomegaly, pancytopenia, osteoporosis, avascular necrosis of femoral head, bone crises	Gaucher disease (glucocerebrosidase [β -glucosidase] deficiency)	88
Achilles tendon xanthoma	Familial hypercholesterolemia (\downarrow LDL receptor signaling)	94
Anaphylaxis following blood transfusion	IgA deficiency	116
Male child, recurrent infections, no mature B cells	Bruton disease (X-linked agammaglobulinemia)	116

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Recurrent cold (noninflamed) abscesses, eczema, high serum IgE, ↑ eosinophils	Hyper-IgE syndrome (Job syndrome: neutrophil chemotaxis abnormality)	116
“Strawberry tongue”	Scarlet fever Kawasaki disease	136, 314
Abdominal pain, diarrhea, leukocytosis, recent antibiotic use	<i>Clostridium difficile</i> infection	138
Back pain, fever, night sweats	Pott disease (vertebral TB)	140
Adrenal hemorrhage, hypotension, DIC	Waterhouse-Friderichsen syndrome (meningococemia)	142, 349
Red “currant jelly” sputum in alcoholic or diabetic patients	<i>Klebsiella pneumoniae</i> pneumonia	145
Large rash with bull’s-eye appearance	Erythema migrans from <i>Ixodes</i> tick bite (Lyme disease: <i>Borrelia</i>)	146
Ulcerated genital lesion	Nonpainful, indurated: chancre (1° syphilis, <i>Treponema pallidum</i>) Painful, with exudate: chancroid (<i>Haemophilus ducreyi</i>)	147, 184
Pupil accommodates but doesn’t react	Neurosyphilis (Argyll Robertson pupil)	147
Smooth, moist, painless, wart-like white lesions on genitals	Condylomata lata (2° syphilis)	147
Fever, chills, headache, myalgia following antibiotic treatment for syphilis	Jarisch-Herxheimer reaction (rapid lysis of spirochetes results in endotoxin-like release)	148
Dog or cat bite resulting in infection	<i>Pasteurella multocida</i> (cellulitis at inoculation site)	149
Rash on palms and soles	Coxsackie A, 2° syphilis, Rocky Mountain spotted fever	150
Black eschar on face of patient with diabetic ketoacidosis	<i>Mucor</i> or <i>Rhizopus</i> fungal infection	153
Chorioretinitis, hydrocephalus, intracranial calcifications	Congenital toxoplasmosis	156
Child with fever later develops red rash on face that spreads to body	Erythema infectiosum/fifth disease (“slapped cheeks” appearance, caused by parvovirus B19)	164
Fever, cough, conjunctivitis, coryza, diffuse rash	Measles	170
Small, irregular red spots on buccal/lingual mucosa with blue-white centers	Koplik spots (measles [rubeola] virus)	170
Bounding pulses, wide pulse pressure, diastolic heart murmur, head bobbing	Aortic regurgitation	291
Systolic ejection murmur (crescendo-decrescendo)	Aortic stenosis	291
Continuous “machine-like” heart murmur	PDA (close with indomethacin; keep open with PGE analogs)	291
Chest pain on exertion	Angina (stable: with moderate exertion; unstable: with minimal exertion or at rest)	304
Chest pain with ST depressions on ECG	Angina (⊖ troponins) or NSTEMI (⊕ troponins)	304
Chest pain, pericardial effusion/friction rub, persistent fever following MI	Dressler syndrome (autoimmune-mediated post-MI fibrinous pericarditis, 2 weeks to several months after acute episode)	307
Painful, raised red lesions on pads of fingers/toes	Osler nodes (infective endocarditis, immune complex deposition)	311

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Painless erythematous lesions on palms and soles	Janeway lesions (infective endocarditis, septic emboli/microabscesses)	311
Splinter hemorrhages in fingernails	Bacterial endocarditis	311
Retinal hemorrhages with pale centers	Roth spots (bacterial endocarditis)	311
Distant heart sounds, distended neck veins, hypotension	Beck triad of cardiac tamponade	310
Cervical lymphadenopathy, desquamating rash, coronary aneurysms, red conjunctivae and tongue, hand-foot changes	Kawasaki disease (mucocutaneous lymph node syndrome, treat with IVIG and aspirin)	314
Palpable purpura on buttocks/legs, joint pain, abdominal pain (child), hematuria	Immunoglobulin A vasculitis (Henoch-Schönlein purpura, affects skin and kidneys)	315
Telangiectasias, recurrent epistaxis, skin discoloration, arteriovenous malformations, GI bleeding, hematuria	Hereditary hemorrhagic telangiectasia (Osler-Weber-Rendu syndrome)	316
Skin hyperpigmentation, hypotension, fatigue	1° adrenocortical insufficiency → ↑ ACTH, ↑ α-MSH (eg, Addison disease)	349
Cutaneous flushing, diarrhea, bronchospasm	Carcinoid syndrome (right-sided cardiac valvular lesions, ↑ 5-HIAA)	352
Cold intolerance, weight gain, brittle hair	Hypothyroidism	341
Cutaneous/dermal edema due to deposition of mucopolysaccharides in connective tissue	Myxedema (caused by hypothyroidism, Graves disease [pretibial])	340
Facial muscle spasm upon tapping	Chvostek sign (hypocalcemia)	344
No lactation postpartum, absent menstruation, cold intolerance	Sheehan syndrome (postpartum hemorrhage leading to pituitary infarction)	339
Deep, labored breathing/hyperventilation	Diabetic ketoacidosis (Kussmaul respirations)	347
Pancreatic, pituitary, parathyroid tumors	MEN 1 (autosomal dominant)	351
Thyroid tumors, pheochromocytoma, ganglioneuromatosis, Marfanoid habitus	MEN 2B (autosomal dominant <i>RET</i> mutation)	351
Thyroid and parathyroid tumors, pheochromocytoma	MEN 2A (autosomal dominant <i>RET</i> mutation)	351
Jaundice, palpable distended non-tender gallbladder	Courvoisier sign (distal malignant obstruction of biliary tree)	398
Vomiting blood following gastroesophageal lacerations	Mallory-Weiss syndrome (alcoholic and bulimic patients)	377
Dysphagia (esophageal webs), glossitis, iron deficiency anemia	Plummer-Vinson syndrome (may progress to esophageal squamous cell carcinoma)	377
Enlarged, hard left supraclavicular node	Virchow node (abdominal metastasis)	379
Arthralgias, adenopathy, cardiac and neurological symptoms, diarrhea	Whipple disease (<i>Tropheryma whipplei</i>)	381
Severe RLQ pain with palpation of LLQ	Rovsing sign (acute appendicitis)	383
Severe RLQ pain with deep tenderness	McBurney sign (acute appendicitis)	383
Hamartomatous GI polyps, hyperpigmented macules on mouth, feet, hands, genitalia	Peutz-Jeghers syndrome (inherited, benign polyposis can cause bowel obstruction; ↑ cancer risk, mainly GI)	387
Multiple colon polyps, osteomas/soft tissue tumors, impacted/supernumerary teeth	Gardner syndrome (subtype of FAP)	387
Abdominal pain, ascites, hepatomegaly	Budd-Chiari syndrome (posthepatic venous thrombosis)	392

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Severe jaundice in neonate	Crigler-Najjar syndrome (congenital unconjugated hyperbilirubinemia)	394
Golden brown rings around peripheral cornea	Wilson disease (Kayser-Fleischer rings due to copper accumulation)	395
Fat, female, forty, fertile	Cholelithiasis (gallstones)	396
Painless jaundice	Cancer of the pancreatic head obstructing bile duct	398
Bluish line on gingiva	Burton line (lead poisoning)	419
Short stature, café-au-lait spots, thumb/radial defects, ↑ incidence of tumors/leukemia, aplastic anemia	Fanconi anemia (genetic loss of DNA crosslink repair; often progresses to AML)	421
Red/pink urine, fragile RBCs	Paroxysmal nocturnal hemoglobinuria	422
Painful blue fingers/toes, hemolytic anemia	Cold agglutinin disease (autoimmune hemolytic anemia caused by <i>Mycoplasma pneumoniae</i> , infectious mononucleosis, CLL)	423
Petechiae, mucosal bleeding, prolonged bleeding time	Platelet disorders (eg, Glanzmann thrombasthenia, Bernard Soulier, HUS, TTP, ITP)	427
Fever, night sweats, weight loss	B symptoms of malignancy	429
Skin patches/plaques, Pautrier microabscesses, atypical T cells	Mycosis fungoides (cutaneous T-cell lymphoma) or Sézary syndrome (mycosis fungoides + malignant T cells in blood)	430
WBCs that look “smudged”	CLL	432
Neonate with arm paralysis following difficult birth, arm in “waiter’s tip” position	Erb-Duchenne palsy (superior trunk [C5–C6] brachial plexus injury)	448
Anterior drawer sign ⊕	Anterior cruciate ligament injury	454
Bone pain, bone enlargement, arthritis	Osteitis deformans (Paget disease of bone, ↑ osteoblastic and osteoclastic activity)	463
Swollen, hard, painful finger joints in an elderly individual, pain worse with activity	Osteoarthritis (osteophytes on PIP [Bouchard nodes], DIP [Heberden nodes])	466
Sudden swollen/painful big toe joint, tophi	Gout/podagra (hyperuricemia)	467
Dry eyes, dry mouth, arthritis	Sjögren syndrome (autoimmune destruction of exocrine glands)	468
Urethritis, conjunctivitis, arthritis in a male	Reactive arthritis associated with HLA-B27	469
“Butterfly” facial rash and Raynaud phenomenon in a young female	Systemic lupus erythematosus	470
Painful fingers/toes changing color from white to blue to red with cold or stress	Raynaud phenomenon (vasospasm in extremities)	472
Anticentromere antibodies	Scleroderma (CREST)	473
Dark purple skin/mouth nodules in a patient with AIDS	Kaposi sarcoma, associated with HHV-8	478
Anti-desmoglein (anti-desmosome) antibodies	Pemphigus vulgaris (blistering)	480
Pruritic, purple, polygonal planar papules and plaques (6 P’s)	Lichen planus	482
↑ AFP in amniotic fluid/maternal serum	Dating error, anencephaly, spina bifida (open neural tube defects)	491
Ataxia, nystagmus, vertigo, dysarthria	Cerebellar lesion	499

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Toe extension/fanning upon plantar scrape	Babinski sign (UMN lesion)	510
Hyperphagia, hypersexuality, hyperorality	Klüver-Bucy syndrome (bilateral amygdala lesion)	511
Resting tremor, athetosis, chorea	Basal ganglia lesion	511
Lucid interval after traumatic brain injury	Epidural hematoma (middle meningeal artery rupture)	513
“Worst headache of my life”	Subarachnoid hemorrhage	513
Resting tremor, rigidity, akinesia, postural instability, shuffling gait	Parkinson disease (loss of dopaminergic neurons in substantia nigra pars compacta)	520
Chorea, dementia, caudate degeneration	Huntington disease (autosomal dominant CAG repeat expansion)	520
Nystagmus, intention tremor, scanning speech, bilateral internuclear ophthalmoplegia	Multiple sclerosis	523
Rapidly progressive limb weakness that ascends following GI/upper respiratory infection	Guillain-Barré syndrome (acute inflammatory demyelinating polyradiculopathy subtype)	524
Café-au-lait spots, Lisch nodules (iris hamartoma), cutaneous neurofibromas, pheochromocytomas, optic gliomas	Neurofibromatosis type I	525
Vascular birthmark (port-wine stain) of the face	Nevus flammeus (benign, but associated with Sturge-Weber syndrome)	525
Renal cell carcinoma (bilateral), hemangioblastomas, angiomas, pheochromocytoma	von Hippel-Lindau disease (dominant tumor suppressor gene mutation)	525
Bilateral vestibular schwannomas	Neurofibromatosis type 2	525
Hyperreflexia, hypertonia, Babinski sign present	UMN damage	529
Hyporeflexia, hypotonia, atrophy, fasciculations	LMN damage	529
Spastic weakness, sensory loss, bowel/bladder dysfunction	Spinal cord lesion	530
Unilateral facial drooping involving forehead	LMN facial nerve (CN VII) palsy; UMN lesions spare the forehead	532
Episodic vertigo, tinnitus, hearing loss	Ménière disease	534
Ptosis, miosis, anhidrosis	Horner syndrome (sympathetic chain lesion)	540
Conjugate horizontal gaze palsy, horizontal diplopia	Internuclear ophthalmoplegia (damage to MLF; may be unilateral or bilateral)	543
Polyuria, renal tubular acidosis type II, growth failure, electrolyte imbalances, hypophosphatemic rickets	Fanconi syndrome (multiple combined dysfunction of the proximal convoluted tubule)	586
Athlete with polycythemia	2° to erythropoietin injection	589
Periorbital and/or peripheral edema, proteinuria (> 3.5g/day), hypoalbuminemia, hypercholesterolemia	Nephrotic syndrome	597
Hereditary nephritis, sensorineural hearing loss, retinopathy, lens dislocation	Alport syndrome (mutation in collagen IV)	596
Streak ovaries, congenital heart disease, horseshoe kidney, cystic hygroma at birth, short stature, webbed neck, lymphedema	Turner syndrome (45,XO)	638
Red, itchy, swollen rash of nipple/areola	Paget disease of the breast (sign of underlying neoplasm)	650

CLINICAL PRESENTATION	DIAGNOSIS/DISEASE	PAGE
Fibrous plaques in tunica albuginea of penis with abnormal curvature	Peyronie disease (connective tissue disorder)	651
Hypoxemia, polycythemia, hypercapnia	Chronic bronchitis (hyperplasia of mucous cells, “blue bloater”)	674
Pink complexion, dyspnea, hyperventilation	Emphysema (“pink puffer,” centriacinar [smoking] or panacinar [α_1 -antitrypsin deficiency])	674
Bilateral hilar adenopathy, uveitis	Sarcoidosis (noncaseating granulomas)	676

▶ CLASSIC LABS/FINDINGS

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
↓ AFP in amniotic fluid/maternal serum	Down syndrome, Edwards syndrome	63
Large granules in phagocytes, immunodeficiency	Chédiak-Higashi disease (congenital failure of phagolysosome formation)	117
Recurrent infections, eczema, thrombocytopenia	Wiskott-Aldrich syndrome	117
Optochin sensitivity	Sensitive: <i>S pneumoniae</i> ; resistant: viridans streptococci (<i>S mutans</i> , <i>S sanguis</i>)	134
Novobiocin response	Sensitive: <i>S epidermidis</i> ; resistant: <i>S saprophyticus</i>	134
Bacitracin response	Sensitive: <i>S pyogenes</i> (group A); resistant: <i>S agalactiae</i> (group B)	134
<i>Streptococcus bovis</i> bacteremia	Colon cancer	137
Branching gram ⊕ rods with sulfur granules	<i>Actinomyces israelii</i>	139
Hilar lymphadenopathy, peripheral granulomatous lesion in middle or lower lung lobes (can calcify)	Ghon complex (1° TB: <i>Mycobacterium bacilli</i>)	140
“Thumb sign” on lateral neck x-ray	Epiglottitis (<i>Haemophilus influenzae</i>)	142
Bacteria-covered vaginal epithelial cells	“Clue cells” (<i>Gardnerella vaginalis</i>)	148
Cardiomegaly with apical atrophy	Chagas disease (<i>Trypanosoma cruzi</i>)	158
Atypical lymphocytes	EBV	165
Enlarged cells with intranuclear inclusion bodies	“Owl eye” appearance of CMV	165
Heterophile antibodies	Infectious mononucleosis (EBV)	165
Intranuclear eosinophilic droplet-like bodies	Cowdry type A bodies (HSV or VZV)	166
Eosinophilic globule in liver	Councilman body (viral hepatitis, yellow fever), represents hepatocyte undergoing apoptosis	168
“Steeple” sign on frontal CXR	Croup (parainfluenza virus)	170
Eosinophilic inclusion bodies in cytoplasm of hippocampal and cerebellar neurons	Negri bodies of rabies	171
Ring-enhancing brain lesion on CT/MRI in AIDS	<i>Toxoplasma gondii</i> , CNS lymphoma	177
Psammoma bodies	Meningiomas, papillary thyroid carcinoma, mesothelioma, papillary serous carcinoma of the endometrium and ovary	211

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
“Delta wave” on ECG, short PR interval, supraventricular tachycardia	Wolff-Parkinson-White syndrome (Bundle of Kent bypasses AV node)	294
“Boot-shaped” heart on x-ray	Tetralogy of Fallot (due to RVH)	298
Rib notching (inferior surface, on x-ray)	Coarctation of the aorta	299
Heart nodules (granulomatous)	Aschoff bodies (rheumatic fever)	312
Electrical alternans (alternating amplitude on ECG)	Cardiac tamponade	310
Antineutrophil cytoplasmic antibodies (ANCA)	Microscopic polyangiitis and eosinophilic granulomatosis with polyangiitis (MPO-ANCA/p-ANCA); granulomatosis with polyangiitis (Wegener; PR3-ANCA/c-ANCA); primary sclerosing cholangitis (MPO-ANCA/p-ANCA)	315
Hypertension, hypokalemia, metabolic alkalosis	1° hyperaldosteronism (Conn syndrome)	349
Enlarged thyroid cells with ground-glass nuclei with central clearing	“Orphan Annie” eyes nuclei (papillary carcinoma of the thyroid)	343
Mucin-filled cell with peripheral nucleus	“Signet ring” (gastric carcinoma)	379
Anti-transglutaminase/anti-gliadin/anti-endomysial antibodies	Celiac disease (diarrhea, weight loss)	381
Narrowing of bowel lumen on barium x-ray	“String sign” (Crohn disease)	382
“Lead pipe” appearance of colon on abdominal imaging	Ulcerative colitis (loss of haustra)	382
Thousands of polyps on colonoscopy	Familial adenomatous polyposis (autosomal dominant, mutation of APC gene)	387
“Apple core” lesion on barium enema x-ray	Colorectal cancer (usually left-sided)	388
Eosinophilic cytoplasmic inclusion in liver cell	Mallory body (alcoholic liver disease)	391
Triglyceride accumulation in liver cell vacuoles	Fatty liver disease (alcoholic or metabolic syndrome)	391
“Nutmeg” appearance of liver	Chronic passive congestion of liver due to right heart failure or Budd-Chiari syndrome	392
Antimitochondrial antibodies (AMAs)	1° biliary cholangitis (female, cholestasis, portal hypertension)	395
Low serum ceruloplasmin	Wilson disease (hepatolenticular degeneration; Kayser-Fleischer rings due to copper accumulation)	395
Migratory thrombophlebitis (leading to migrating DVTs and vasculitis)	Trousseau syndrome (adenocarcinoma of pancreas or lung)	398
Basophilic nuclear remnants in RBCs	Howell-Jolly bodies (due to splenectomy or nonfunctional spleen)	416
Basophilic stippling of RBCs	Lead poisoning or sideroblastic anemia	416
Hypochromic, microcytic anemia	Iron deficiency anemia, lead poisoning, thalassemia (fetal hemoglobin sometimes present)	418, 419
“Hair on end” (“Crew-cut”) appearance on x-ray	β-thalassemia, sickle cell disease (marrow expansion)	422
Hypersegmented neutrophils	Megaloblastic anemia (B ₁₂ deficiency: neurologic symptoms; folate deficiency: no neurologic symptoms)	420
Antiplatelet antibodies	Idiopathic thrombocytopenic purpura	427
High level of D-dimers	DVT, PE, DIC	428
Giant B cells with bilobed nuclei with prominent inclusions (“owl’s eye”)	Reed-Sternberg cells (Hodgkin lymphoma)	429

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Sheets of medium-sized lymphoid cells with scattered pale, tingible body-laden macrophages (“starry sky” histology)	Burkitt lymphoma (t[8:14] c-myc activation, associated with EBV; “starry sky” made up of malignant cells)	430
Lytic (“punched-out”) bone lesions on x-ray	Multiple myeloma	431
Monoclonal antibody spike	<ul style="list-style-type: none"> ▪ Multiple myeloma (usually IgG or IgA) ▪ Monoclonal gammopathy of undetermined significance (MGUS consequence of aging) ▪ Waldenström (M protein = IgM) macroglobulinemia ▪ Primary amyloidosis 	431
Stacks of RBCs	Rouleaux formation (high ESR, multiple myeloma)	423
Azurophilic peroxidase ⊕ granular inclusions in granulocytes and myeloblasts	Auer rods (AML, especially the promyelocytic [M3] type)	432
WBCs that look “smudged”	CLL (almost always B cell)	432
“Tennis racket”-shaped cytoplasmic organelles (EM) in Langerhans cells	Birbeck granules (Langerhans cell histiocytosis)	434
“Brown” tumor of bone	Hyperparathyroidism or osteitis fibrosa cystica (deposited hemosiderin from hemorrhage gives brown color)	464
“Soap bubble” in femur or tibia on x-ray	Giant cell tumor of bone (generally benign)	464
Raised periosteum (creating a “Codman triangle”)	Aggressive bone lesion (eg, osteosarcoma, Ewing sarcoma, osteomyelitis)	465
“Onion skin” periosteal reaction	Ewing sarcoma (malignant small blue cell tumor)	465
Anti-IgG antibodies	Rheumatoid arthritis (systemic inflammation, joint pannus, boutonniere and swan neck deformities)	466
Rhomboid crystals, ⊕ birefringent	Pseudogout (calcium pyrophosphate dihydrate crystals)	467
Needle-shaped, ⊖ birefringent crystals	Gout (monosodium urate crystals)	467
↑ uric acid levels	Gout, Lesch-Nyhan syndrome, tumor lysis syndrome, loop and thiazide diuretics	467
“Bamboo spine” on x-ray	Ankylosing spondylitis (chronic inflammatory arthritis: HLA-B27)	469
Antinuclear antibodies (ANAs: anti-Smith and anti-dsDNA)	SLE (type III hypersensitivity)	470
Anti-histone antibodies	Drug-induced SLE (eg, hydralazine, isoniazid, phenytoin, procainamide)	250
Anti-topoisomerase antibodies	Diffuse scleroderma	473
Keratin pearls on a skin biopsy	Squamous cell carcinoma	484
Bloody or yellow tap on lumbar puncture	Xanthochromia (due to subarachnoid hemorrhage)	513
Eosinophilic cytoplasmic inclusion in neuron	Lewy body (Parkinson disease and Lewy body dementia)	520
Extracellular amyloid deposition in gray matter of brain	Senile plaques (Alzheimer disease)	520
Depigmentation of neurons in substantia nigra	Parkinson disease (basal ganglia disorder: rigidity, resting tremor, bradykinesia)	520
Protein aggregates in neurons from hyperphosphorylation of tau protein	Neurofibrillary tangles (Alzheimer disease) and Pick bodies (Pick disease)	520
Silver-staining spherical aggregation of tau proteins in neurons	Pick bodies (Pick disease: progressive dementia, changes in personality)	520

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Pseudopalisading tumor cells on brain biopsy	Glioblastoma multiforme	526
Circular grouping of dark tumor cells surrounding pale neurofibrils	Homer-Wright rosettes (neuroblastoma, medulloblastoma)	528
“Waxy” casts with very low urine flow	Chronic end-stage renal disease	594
Nodular hyaline deposits in glomeruli	Kimmelstiel-Wilson nodules (diabetic nephropathy)	597
Podocyte fusion or “effacement” on electron microscopy	Minimal change disease (child with nephrotic syndrome)	597
“Spikes” on basement membrane, “dome-like” subepithelial deposits	Membranous nephropathy (nephrotic syndrome)	597
RBC casts in urine	Glomerulonephritis	594
“Tram-track” appearance of capillary loops of glomerular basement membranes on light microscopy	Membranoproliferative glomerulonephritis	596
Anti-glomerular basement membrane antibodies	Goodpasture syndrome (glomerulonephritis and hemoptysis)	596
Cellular crescents in Bowman capsule	Rapidly progressive (crescentic) glomerulonephritis	596
“Wire loop” glomerular capillary appearance on light microscopy	Diffuse proliferative glomerulonephritis (usually seen with lupus)	596
Linear appearance of IgG deposition on glomerular and alveolar basement membranes	Goodpasture syndrome	596
“Lumpy bumpy” appearance of glomeruli on immunofluorescence	Poststreptococcal glomerulonephritis (due to deposition of IgG, IgM, and C3)	596
Necrotizing vasculitis (lungs) and necrotizing glomerulonephritis	Granulomatosis with polyangiitis (Wegener; PR3-ANCA/c-ANCA) and Goodpasture syndrome (anti-basement membrane antibodies)	596
Thyroid-like appearance of kidney	Chronic pyelonephritis (usually due to recurrent infections)	600
WBC casts in urine	Acute pyelonephritis	600
Renal epithelial casts in urine	Intrinsic renal failure (eg, ischemia or toxic injury)	601
hCG elevated	Choriocarcinoma, hydatidiform mole (occurs with and without embryo, and multiple pregnancy)	633
Dysplastic squamous cervical cells with “raisinoid” nuclei and hyperchromasia	Koilocytes (HPV: predisposes to cervical cancer)	645
Disarrayed granulosa cells arranged around collections of eosinophilic fluid	Call-Exner bodies (granulosa cell tumor of the ovary)	647
“Chocolate cyst” of ovary	Endometriosis (frequently involves both ovaries)	648
Mammary gland (“blue domed”) cyst	Fibrocystic change of the breast	649
Glomerulus-like structure surrounding vessel in germ cells	Schiller-Duval bodies (yolk sac tumor)	647
Rectangular, crystal-like, cytoplasmic inclusions in Leydig cells	Reinke crystals (Leydig cell tumor)	653
Thrombi made of white/red layers	Lines of Zahn (arterial thrombus, layers of platelets/RBCs)	672
Hexagonal, double-pointed, needle-like crystals in bronchial secretions	Bronchial asthma (Charcot-Leyden crystals: eosinophilic granules)	674

LAB/DIAGNOSTIC FINDING	DIAGNOSIS/DISEASE	PAGE
Desquamated epithelium casts in sputum	Curschmann spirals (bronchial asthma; can result in whorled mucous plugs)	674
“Honeycomb lung” on x-ray or CT	Idiopathic pulmonary fibrosis	675
Colonies of mucoid <i>Pseudomonas</i> in lungs	Cystic fibrosis (autosomal recessive mutation in <i>CFTR</i> gene → fat-soluble vitamin deficiency and mucous plugs)	675
Iron-containing nodules in alveolar septum	Ferruginous bodies (asbestosis: ↑ chance of lung cancer)	677
Bronchogenic apical lung tumor on imaging	Pancoast tumor (can compress cervical sympathetic chain and cause Horner syndrome)	685

▶ CLASSIC/RELEVANT TREATMENTS

CONDITION	COMMON TREATMENT(S)	PAGE
Ethylene glycol/methanol intoxication	Fomepizole (alcohol dehydrogenase inhibitor)	72
Chronic hepatitis B or C	IFN- α (HBV and HCV); ribavirin, simeprevir, sofosbuvir (HCV)	121
<i>Streptococcus bovis</i>	Penicillin prophylaxis; evaluation for colon cancer if linked to endocarditis	137
<i>Clostridium botulinum</i>	Antitoxin	138
<i>Clostridium tetani</i>	Antitoxin	138
<i>Haemophilus influenzae</i> (B)	Amoxicillin \pm clavulanate (mucosal infections), ceftriaxone (meningitis), rifampin (prophylaxis)	142
<i>Neisseria gonorrhoeae</i>	Ceftriaxone (add doxycycline to cover likely concurrent <i>C trachomatis</i>)	142
<i>Neisseria meningitidis</i>	Penicillin/ceftriaxone, rifampin (prophylaxis)	142
<i>Legionella pneumophila</i>	Macrolides (eg, azithromycin)	143
<i>Pseudomonas aeruginosa</i>	Piperacillin/tazobactam, aminoglycosides, carbapenems	143
<i>Treponema pallidum</i>	Penicillin G	147
<i>Chlamydia trachomatis</i>	Doxycycline (+ ceftriaxone for gonorrhea coinfection), oral erythromycin to treat chlamydial conjunctivitis in infants	148
<i>Candida albicans</i>	Topical azoles (vaginitis); nystatin, fluconazole, caspofungin (oral/esophageal); fluconazole, caspofungin, amphotericin B (systemic)	153
<i>Cryptococcus neoformans</i>	Induction with amphotericin B and flucytosine, maintenance with fluconazole (in AIDS patients)	153
<i>Sporothrix schenckii</i>	Itraconazole, oral potassium iodide	154
<i>Pneumocystis jirovecii</i>	TMP-SMX (prophylaxis and treatment in immunosuppressed patients, CD4 < 200/mm ³)	154
<i>Toxoplasma gondii</i>	Sulfadiazine + pyrimethamine	156
Malaria	Chloroquine, mefloquine, atovaquone/proguanil (for blood schizont), primaquine (for liver hypnozoite)	157

CONDITION	COMMON TREATMENT(S)	PAGE
<i>Trichomonas vaginalis</i>	Metronidazole (patient and partner)	158
<i>Streptococcus pyogenes</i>	Penicillin prophylaxis	187
<i>Streptococcus pneumoniae</i>	Penicillin/cephalosporin (systemic infection, pneumonia), vancomycin (meningitis)	187, 190
<i>Staphylococcus aureus</i>	MSSA: nafcillin, oxacillin, dicloxacillin (antistaphylococcal penicillins); MRSA: vancomycin, daptomycin, linezolid, ceftaroline	188, 190, 195
Enterococci	Vancomycin, aminopenicillins/cephalosporins	189, 190
<i>Rickettsia rickettsii</i>	Doxycycline, chloramphenicol	192
<i>Clostridium difficile</i>	Oral metronidazole; if refractory, oral vancomycin	190, 195
<i>Mycobacterium tuberculosis</i>	RIPE (rifampin, isoniazid, pyrazinamide, ethambutol)	196
UTI prophylaxis	TMP-SMX	198
Influenza	Oseltamivir, zanamivir	201
CMV	Ganciclovir, foscarnet, cidofovir	202
Patent ductus arteriosus	Close with indomethacin; keep open with PGE analogs	282
Stable angina	Sublingual nitroglycerin	304
Buerger disease	Smoking cessation	314
Kawasaki disease	IVIg, aspirin	314
Temporal arteritis	High-dose steroids	314
Granulomatosis with polyangiitis (Wegener)	Cyclophosphamide, corticosteroids	315
Hypercholesterolemia	Statin (first-line)	320
Hypertriglyceridemia	Fibrate	320
Arrhythmia in damaged cardiac tissue	Class IB antiarrhythmic (lidocaine, mexiletine)	322
Prolactinoma	Cabergoline/bromocriptine (dopamine agonists)	330
Diabetes insipidus	Desmopressin (central); hydrochlorothiazide, indomethacin, amiloride (nephrogenic)	338
SIADH	Fluid restriction, IV hypertonic saline, conivaptan/tolvaptan, demeclocycline	338
Diabetic ketoacidosis	Fluids, insulin, K ⁺	347
Diabetes mellitus type 1	Dietary intervention (low carbohydrate) + insulin replacement	347
Diabetes mellitus type 2	Dietary intervention, oral hypoglycemics, and insulin (if refractory)	347
Pheochromocytoma	α-antagonists (eg, phenoxybenzamine)	350
Carcinoid syndrome	Octreotide	352
Crohn disease	Corticosteroids, infliximab, azathioprine	382
Ulcerative colitis	5-ASA preparations (eg, mesalamine), 6-mercaptopurine, infliximab, colectomy	382
Sickle cell disease	Hydroxyurea (↑ fetal hemoglobin)	422

CONDITION	COMMON TREATMENT(S)	PAGE
Chronic myelogenous leukemia	Imatinib	433
Acute promyelocytic leukemia (M3)	All- <i>trans</i> retinoic acid, arsenic trioxide	432
Drug of choice for anticoagulation in pregnancy or renal failure	Low-molecular-weight heparin	436
Heparin reversal	Protamine sulfate	436
Immediate anticoagulation	Heparin	436
Long-term anticoagulation	Warfarin, dabigatran, rivaroxaban and apixaban	436, 437
Warfarin reversal	Fresh frozen plasma (acute), vitamin K (non-acute)	436
Cyclophosphamide-induced hemorrhagic cystitis	Mesna	441
HER2/neu ⊕ breast cancer	Trastuzumab	443
Osteoporosis	Calcium/vitamin D supplementation (prophylaxis); bisphosphonates, PTH analogs, SERMs, calcitonin, denosumab (treatment)	462
Osteomalacia/rickets	Vitamin D supplementation	463
Chronic gout	Xanthine oxidase inhibitors (eg, allopurinol, febuxostat); pegloticase; probenecid	467
Acute gout attack	NSAIDs, colchicine, glucocorticoids	467
Neural tube defect prevention	Prenatal folic acid	491
Migraine	Abortive therapies (eg, sumatriptan, NSAIDs); prophylaxis (eg, propranolol, topiramate, CCBs, amitriptyline)	518
Multiple sclerosis	Disease-modifying therapies (eg, β-interferon, natalizumab); for acute flares, use IV steroids	523
Tonic-clonic seizures	Levetiracetam, phenytoin, valproate, carbamazepine	544
Absence seizures	Ethosuximide	544
Trigeminal neuralgia (tic douloureux)	Carbamazepine	544
Malignant hyperthermia	Dantrolene	551
Anorexia	Nutrition, psychotherapy, SSRIs	567
Bulimia nervosa	SSRIs	567
Alcoholism	Disulfiram, acamprosate, naltrexone, supportive care	571
ADHD	Methylphenidate, amphetamines, CBT, atomoxetine, guanfacine, clonidine	572
Alcohol withdrawal	Long-acting benzodiazepines	572
Bipolar disorder	Mood stabilizers (eg, lithium, valproic acid, carbamazepine), atypical antipsychotics	572
Depression	SSRIs (first-line)	572
Generalized anxiety disorder	SSRIs, SNRIs (first line); buspirone (second line)	572
Schizophrenia (positive symptoms)	Typical and atypical antipsychotics	573
Schizophrenia (negative symptoms)	Atypical antipsychotics	573

CONDITION	COMMON TREATMENT(S)	PAGE
Hyperaldosteronism	Spironolactone	609
Benign prostatic hyperplasia	α_1 -antagonists, 5 α -reductase inhibitors, PDE-5 inhibitors	654
Infertility	Leuprolide, GnRH (pulsatile), clomiphene	656
Breast cancer in postmenopausal woman	Aromatase inhibitor (anastrozole)	656
ER \oplus breast cancer	Tamoxifen	656
Prostate adenocarcinoma/uterine fibroids	Leuprolide, GnRH (continuous)	656
Medical abortion	Mifepristone	657
Prostate adenocarcinoma	Flutamide	658
Erectile dysfunction	Sildenafil, tadalafil, vardenafil	686
Pulmonary arterial hypertension (idiopathic)	Sildenafil, bosentan, epoprostenol	686

▶ KEY ASSOCIATIONS

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Mitochondrial inheritance	Disease occurs in both males and females, inherited through females only	59
Intellectual disability	Down syndrome, fragile X syndrome	62, 63
Vitamin deficiency (USA)	Folate (pregnant women are at high risk; body stores only 3- to 4-month supply; prevents neural tube defects)	68
Lysosomal storage disease	Gaucher disease	88
Bacterial meningitis (adults and elderly)	<i>S pneumoniae</i>	180
Bacterial meningitis (newborns and kids)	Group B streptococcus/ <i>E coli</i> / <i>Listeria monocytogenes</i> (newborns), <i>S pneumoniae</i> / <i>N meningitidis</i> (kids/teens)	180
HLA-DR3	Diabetes mellitus type 1, SLE, Graves disease, Hashimoto thyroiditis (also associated with HLA-DR5), Addison disease	100
HLA-DR4	Diabetes mellitus type 1, rheumatoid arthritis, Addison disease	100
Bacteria associated with gastritis, peptic ulcer disease, and gastric malignancies (eg, adenocarcinoma, MALToma)	<i>H pylori</i>	146
Opportunistic infection in AIDS	<i>Pneumocystis jirovecii</i> pneumonia	154
Helminth infection (US)	<i>Enterobius vermicularis</i>	159
Viral encephalitis affecting temporal lobe	HSV-1	164
Infection 2° to blood transfusion	Hepatitis C	172
Food poisoning (exotoxin mediated)	<i>S aureus</i> , <i>B cereus</i>	178
Osteomyelitis	<i>S aureus</i> (most common overall)	180
Osteomyelitis in sickle cell disease	<i>Salmonella</i>	180
Osteomyelitis with IV drug use	<i>Pseudomonas</i> , <i>Candida</i> , <i>S aureus</i>	180

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
UTI	<i>E coli</i> , <i>Staphylococcus saprophyticus</i> (young women)	181
Sexually transmitted disease	<i>C trachomatis</i> (usually coinfects with <i>N gonorrhoeae</i>)	184
Nosocomial pneumonia	<i>S aureus</i> , <i>Pseudomonas</i> , other enteric gram \ominus rods	185
Pelvic inflammatory disease	<i>C trachomatis</i> , <i>N gonorrhoeae</i>	185
Infections in chronic granulomatous disease	<i>S aureus</i> , <i>E coli</i> , <i>Aspergillus</i> (catalase \oplus)	186
Metastases to bone	Prostate, breast > kidney, thyroid, lung	223
Metastases to brain	Lung > breast > melanoma, colon, kidney	223
Metastases to liver	Colon >> stomach > pancreas	223
S3 heart sound	\uparrow ventricular filling pressure (eg, mitral regurgitation, HF), common in dilated ventricles	287
S4 heart sound	Stiff/hypertrophic ventricle (aortic stenosis, restrictive cardiomyopathy)	287
Constrictive pericarditis	TB (developing world); idiopathic, viral illness (developed world)	287
Holosystolic murmur	VSD, tricuspid regurgitation, mitral regurgitation	291
Ejection click	Aortic stenosis	291
Mitral valve stenosis	Rheumatic heart disease	291
Opening snap	Mitral stenosis	291
Heart murmur, congenital	Mitral valve prolapse	291
Chronic arrhythmia	Atrial fibrillation (associated with high risk of emboli)	295
Cyanosis (early; less common)	Tetralogy of Fallot, transposition of great vessels, truncus arteriosus, total anomalous pulmonary venous return, tricuspid atresia	298
Late cyanotic shunt (uncorrected left to right becomes right to left)	Eisenmenger syndrome (caused by ASD, VSD, PDA; results in pulmonary hypertension/polycythemia)	299
Congenital cardiac anomaly	VSD	299
Hypertension, 2 $^{\circ}$	Renal artery stenosis, chronic kidney disease (eg, polycystic kidney disease, diabetic nephropathy), hyperaldosteronism	300
Aortic aneurysm, thoracic	Marfan syndrome (idiopathic cystic medial degeneration)	302
Aortic aneurysm, abdominal	Atherosclerosis, smoking is major risk factor	302
Aortic aneurysm, ascending or arch	3 $^{\circ}$ syphilis (syphilitic aortitis), vasa vasorum destruction	303
Sites of atherosclerosis	Abdominal aorta > coronary artery > popliteal artery > carotid artery	302
Aortic dissection	Hypertension	303
Right heart failure due to a pulmonary cause	Cor pulmonale	309
Heart valve in bacterial endocarditis	Mitral > aortic (rheumatic fever), tricuspid (IV drug abuse)	310
Endocarditis presentation associated with bacterium	<i>S aureus</i> (acute, IVDA, tricuspid valve), viridans streptococci (subacute, dental procedure), <i>S bovis</i> (colon cancer), culture negative (<i>Coxiella</i> , <i>Bartonella</i> , HACEK)	310
Temporal arteritis	Risk of ipsilateral blindness due to occlusion of ophthalmic artery; polymyalgia rheumatica	314

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Recurrent inflammation/thrombosis of small/medium vessels in extremities	Buerger disease (strongly associated with tobacco)	314
Cardiac 1° tumor (kids)	Rhabdomyoma, often seen in tuberous sclerosis	316
Cardiac tumor (adults)	Metastasis, myxoma (90% in left atrium; “ball valve”)	316
Congenital adrenal hyperplasia, hypotension	21-hydroxylase deficiency	335
Hypopituitarism	Pituitary adenoma (usually benign tumor)	339
Cretinism	Iodine deficit/congenital hypothyroidism	341
Thyroid cancer	Papillary carcinoma (childhood irradiation)	343
Hypoparathyroidism	Accidental excision during thyroidectomy	344
1° hyperparathyroidism	Adenomas, hyperplasia, carcinoma	345
2° hyperparathyroidism	Hypocalcemia of chronic kidney disease	345
Cushing syndrome	<ul style="list-style-type: none"> ▪ Iatrogenic (from corticosteroid therapy) ▪ Adrenocortical adenoma (secretes excess cortisol) ▪ ACTH-secreting pituitary adenoma (Cushing disease) ▪ Paraneoplastic (due to ACTH secretion by tumors) 	348
1° hyperaldosteronism	Adrenal hyperplasia or adenoma	349
Tumor of the adrenal medulla (kids)	Neuroblastoma (malignant)	350
Tumor of the adrenal medulla (adults)	Pheochromocytoma (usually benign)	350
Refractory peptic ulcers and high gastrin levels	Zollinger-Ellison syndrome (gastrinoma of duodenum or pancreas), associated with MEN1	351, 352
Esophageal cancer	Squamous cell carcinoma (worldwide); adenocarcinoma (US)	378
Acute gastric ulcer associated with CNS injury	Cushing ulcer (↑ intracranial pressure stimulates vagal gastric H ⁺ secretion)	379
Acute gastric ulcer associated with severe burns	Curling ulcer (greatly reduced plasma volume results in sloughing of gastric mucosa)	379
Bilateral ovarian metastases from gastric carcinoma	Krukenberg tumor (mucin-secreting signet ring cells)	379
Chronic atrophic gastritis (autoimmune)	Predisposition to gastric carcinoma (can also cause pernicious anemia)	379
Gastric cancer	Adenocarcinoma	379
Alternating areas of transmural inflammation and normal colon	Skip lesions (Crohn disease)	382
Site of diverticula	Sigmoid colon	383
Diverticulum in pharynx	Zenker diverticulum (diagnosed by barium swallow)	384
Hepatocellular carcinoma	Cirrhotic liver (associated with hepatitis B and C, alcoholism, and hemochromatosis)	392
Liver disease	Alcoholic cirrhosis	391
1° liver cancer	Hepatocellular carcinoma (chronic hepatitis, cirrhosis, hemochromatosis, α_1 -antitrypsin deficiency, Wilson disease)	392
Congenital conjugated hyperbilirubinemia (black liver)	Dubin-Johnson syndrome (inability of hepatocytes to secrete conjugated bilirubin into bile)	394

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Hereditary harmless jaundice	Gilbert syndrome (benign congenital unconjugated hyperbilirubinemia)	394
Hemochromatosis	Multiple blood transfusions or hereditary <i>HFE</i> mutation (can result in heart failure, “bronze diabetes,” and ↑ risk of hepatocellular carcinoma)	395
Pancreatitis (acute)	Gallstones, alcohol	397
Pancreatitis (chronic)	Alcohol (adults), cystic fibrosis (kids)	397
Microcytic anemia	Iron deficiency	418
Autosplenectomy (fibrosis and shrinkage)	Sickle cell disease (hemoglobin S)	422
Bleeding disorder with GpIb deficiency	Bernard-Soulier syndrome (defect in platelet adhesion to von Willebrand factor)	427
Hereditary bleeding disorder	von Willebrand disease	428
DIC	Severe sepsis, obstetric complications, cancer, burns, trauma, major surgery, acute pancreatitis, APL	428
Malignancy associated with noninfectious fever	Hodgkin lymphoma	429
Type of Hodgkin lymphoma	Nodular sclerosis (vs mixed cellularity, lymphocytic predominance, lymphocytic depletion)	429
t(14;18)	Follicular lymphomas (<i>BCL-2</i> activation, anti-apoptotic oncogene)	430
t(8;14)	Burkitt lymphoma (<i>c-myc</i> fusion, transcription factor oncogene)	430
Type of non-Hodgkin lymphoma	Diffuse large B-cell lymphoma	430
1° bone tumor (adults)	Multiple myeloma	431
Age ranges for patient with ALL/CLL/AML/CML	ALL: child, CLL: adult > 60, AML: adult ~ 65, CML: adult 45–85	432, 433
Malignancy (kids)	Leukemia, brain tumors	432, 526
Death in CML	Blast crisis	433
t(9;22)	Philadelphia chromosome, CML (<i>BCR-ABL</i> oncogene, tyrosine kinase activation), more rarely associated with ALL	434
Vertebral compression fracture	Osteoporosis (type I: postmenopausal woman; type II: elderly man or woman)	462
HLA-B27	Psoriatic arthritis, ankylosing spondylitis, IBD-associated arthritis, reactive arthritis (formerly Reiter syndrome)	469
Death in SLE	Lupus nephropathy	470
Tumor of infancy	Strawberry hemangioma (grows rapidly and regresses spontaneously by childhood)	478
Actinic (solar) keratosis	Precursor to squamous cell carcinoma	482
Cerebellar tonsillar herniation	Chiari I malformation	492
Atrophy of the mammillary bodies	Wernicke encephalopathy (thiamine deficiency causing ataxia, ophthalmoplegia, and confusion)	511

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Epidural hematoma	Rupture of middle meningeal artery (trauma; lentiform shaped)	513
Subdural hematoma	Rupture of bridging veins (crescent shaped)	513
Dementia	Alzheimer disease, multiple infarcts (vascular dementia)	520, 521
Demyelinating disease in young women	Multiple sclerosis	523
Brain tumor (adults)	Supratentorial: metastasis, astrocytoma (including glioblastoma multiforme), meningioma, schwannoma	526
Pituitary tumor	Prolactinoma, somatotrophic adenoma	527
Brain tumor (kids)	Infratentorial: medulloblastoma (cerebellum) or supratentorial: craniopharyngioma	528
Mixed (UMN and LMN) motor neuron disease	Amyotrophic lateral sclerosis	530
Degeneration of dorsal column fibers	Tabes dorsalis (3° syphilis), subacute combined degeneration (dorsal columns, lateral corticospinal, spinocerebellar tracts affected)	530
Nephrotic syndrome (adults)	Membranous nephropathy	597
Nephrotic syndrome (kids)	Minimal change disease	597
Glomerulonephritis (adults)	Berger disease (IgA nephropathy)	596
Kidney stones	<ul style="list-style-type: none"> ▪ Calcium = radiopaque ▪ Struvite (ammonium) = radiopaque (formed by urease ⊕ organisms such as <i>Klebsiella</i>, <i>Proteus</i> species, and <i>S saprophyticus</i>) ▪ Uric acid = radiolucent ▪ Cystine = faintly radiopaque 	598
Renal tumor	Renal cell carcinoma: associated with von Hippel-Lindau and cigarette smoking; paraneoplastic syndromes (EPO, renin, PTHrP, ACTH)	605
Obstruction of male urinary tract	BPH	654
1° amenorrhea	Turner syndrome (45,XO or 45,XO/46,XX mosaic)	638
Neuron migration failure	Kallmann syndrome (hypogonadotropic hypogonadism and anosmia)	639
Clear cell adenocarcinoma of the vagina	DES exposure in utero	644
Ovarian tumor (benign, bilateral)	Serous cystadenoma	646
Ovarian tumor (malignant)	Serous cystadenocarcinoma	646
Tumor in women	Leiomyoma (estrogen dependent, not precancerous)	648
Gynecologic malignancy	Endometrial carcinoma (most common in US); cervical carcinoma (most common worldwide)	648
Breast mass	Fibrocystic change, carcinoma (in postmenopausal women)	649
Breast tumor (benign, young woman)	Fibroadenoma	649
Breast cancer	Invasive ductal carcinoma	650
Testicular tumor	Seminoma (malignant, radiosensitive), ↑ placental ALP	652, 653

DISEASE/FINDING	MOST COMMON/IMPORTANT ASSOCIATIONS	PAGE
Pulmonary hypertension	Idiopathic, heritable, left heart disease (eg, HF), lung disease (eg, COPD), hypoxemic vasoconstriction (eg, OSA), thromboembolic (eg, PE)	679
Hypercoagulability, endothelial damage, blood stasis	Virchow triad (↑ risk of thrombosis)	671
SIADH	Small cell carcinoma of the lung	684

► EQUATION REVIEW

TOPIC	EQUATION	PAGE
Volume of distribution	$V_d = \frac{\text{amount of drug in the body}}{\text{plasma drug concentration}}$	231
Half-life	$t_{1/2} = \frac{0.7 \times V_d}{CL}$	231
Drug clearance	$CL = \frac{\text{rate of elimination of drug}}{\text{plasma drug concentration}} = V_d \times K_e$ (elimination constant)	231
Loading dose	$LD = \frac{C_p \times V_d}{F}$	231
Maintenance dose	$D = \frac{C_p \times CL \times \tau}{F}$	231
Sensitivity	$\text{Sensitivity} = TP / (TP + FN)$	257
Specificity	$\text{Specificity} = TN / (TN + FP)$	257
Positive predictive value	$PPV = TP / (TP + FP)$	257
Negative predictive value	$NPV = TN / (FN + TN)$	257
Odds ratio (for case-control studies)	$OR = \frac{a/c}{b/d} = \frac{ad}{bc}$	258
Relative risk	$RR = \frac{a/(a+b)}{c/(c+d)}$	258
Attributable risk	$AR = \frac{a}{a+b} - \frac{c}{c+d}$	258
Relative risk reduction	$RRR = 1 - RR$	258
Absolute risk reduction	$ARR = \frac{c}{c+d} - \frac{a}{a+b}$	258
Number needed to treat	$NNT = 1/ARR$	258
Number needed to harm	$NNH = 1/AR$	258
Cardiac output	$CO = \frac{\text{rate of } O_2 \text{ consumption}}{(\text{arterial } O_2 \text{ content} - \text{venous } O_2 \text{ content})}$	285
	$CO = \text{stroke volume} \times \text{heart rate}$	285

TOPIC	EQUATION	PAGE
Mean arterial pressure	MAP = cardiac output × total peripheral resistance	285
	MAP = $\frac{2}{3}$ diastolic + $\frac{1}{3}$ systolic	285
Stroke volume	SV = EDV – ESV	285
Ejection fraction	EF = $\frac{SV}{EDV} = \frac{EDV - ESV}{EDV}$	285
Resistance	Resistance = $\frac{\text{driving pressure } (\Delta P)}{\text{flow } (Q)} = \frac{8\eta \text{ (viscosity)} \times \text{length}}{\pi r^4}$	286
Capillary fluid exchange	$J_v = \text{net fluid flow} = K_f[(P_c - P_i) - \sigma(\pi_c - \pi_i)]$	297
Renal clearance	$C_x = (U_x V)/P_x$	582
Glomerular filtration rate	$C_{\text{inulin}} = \text{GFR} = U_{\text{inulin}} \times V/P_{\text{inulin}}$ $= K_f [(P_{\text{GC}} - P_{\text{BS}}) - (\pi_{\text{GC}} - \pi_{\text{BS}})]$	582
Effective renal plasma flow	$\text{eRPF} = U_{\text{PAH}} \times \frac{V}{P_{\text{PAH}}} = C_{\text{PAH}}$	582
Renal blood flow	$\text{RBF} = \frac{\text{RPF}}{1 - \text{Hct}}$	582
Filtration fraction	$\text{FF} = \frac{\text{GFR}}{\text{RPF}}$	583
Henderson-Hasselbalch equation (for extracellular pH)	$\text{pH} = 6.1 + \log \frac{[\text{HCO}_3^-]}{0.03 P_{\text{CO}_2}}$	592
Winters formula	$P_{\text{CO}_2} = 1.5 [\text{HCO}_3^-] + 8 \pm 2$	592
Anion gap	$\text{Na}^+ - (\text{Cl}^- + \text{HCO}_3^-)$	592
Physiologic dead space	$V_D = V_T \times \frac{P_{\text{aCO}_2} - P_{\text{ECO}_2}}{P_{\text{aCO}_2}}$	664
Pulmonary vascular resistance	$\text{PVR} = \frac{P_{\text{pulm artery}} - P_{\text{L atrium}}}{\text{cardiac output}}$	668
Alveolar gas equation	$P_{\text{AO}_2} = P_{\text{IO}_2} - \frac{P_{\text{aCO}_2}}{R}$	668

▶ EASILY CONFUSED MEDICATIONS

DRUG	CLINICAL USE/MECHANISM OF ACTION
Amiloride	K ⁺ -sparing diuretic
Amiodarone	Class III antiarrhythmic
Amlodipine	Dihydropyridine Ca ²⁺ channel blocker
Benztropine	Cholinergic antagonist
Bromocriptine	Dopamine agonist
Bupirone	Generalized anxiety disorder (5-HT _{1A} -receptor agonist)
Bupropion	Depression, smoking cessation (NE-DA reuptake inhibitor)
Cimetidine	H ₂ -receptor antagonist
Cetirizine	2nd-generation antihistamine
Chloramphenicol	Antibiotic (blocks 50S subunit)
Chlordiazepoxide	Long-acting benzodiazepine
Chlorpromazine	Typical antipsychotic
Chlorpropamide	1st-generation sulfonylurea
Chlorpheniramine	1st-generation antihistamine
Chlorthalidone	Thiazide diuretic
Clozapine	5-HT _{2A} -agonist
Clomipramine	Tricyclic antidepressant
Clomiphene	Selective estrogen receptor modulator
Clonidine	α ₂ -agonist
Doxepin	Tricyclic antidepressant
Doxazosin	α ₁ -antagonist
Eplerenone	K ⁺ -sparing diuretic
Propafenone	Class IC antiarrhythmic
Fluoxetine	Selective serotonin reuptake inhibitor
Fluphenazine	Typical antipsychotic
Duloxetine	Serotonin-norepinephrine reuptake inhibitor
Guaifenesin	Expectorant (thins respiratory secretions)
Guanfacine	α ₂ -agonist
Mifepristone	Progesterone receptor antagonist
Misoprostol	PGE ₁ synthetic analog
Naloxone	Opioid receptor antagonist (treats toxicity)
Naltrexone	Opioid receptor antagonist (prevents relapse)
Nitroprusside	Hypertensive emergency (↑ cGMP/NO)
Nitroglycerin	Antianginal (↑ cGMP/NO)
Omeprazole	Proton pump inhibitor
Ketoconazole	Antifungal (inhibits fungal sterol synthesis)

DRUG	CLINICAL USE/MECHANISM OF ACTION
Aripiprazole	Atypical antipsychotic
Anastrozole	Aromatase inhibitor
Rifaximin	Hepatic encephalopathy (↓ ammoniagenic bacteria)
Rifampin	Antimicrobial (inhibits DNA-dependent RNA polymerase)
Sertraline	Selective serotonin reuptake inhibitor
Selegiline	MAO-B inhibitor
Trazodone	Insomnia (blocks 5-HT ₂ , α ₁ -adrenergic, and H ₁ receptors)
Tramadol	Chronic pain (weak opioid agonist)
Varenicline	Smoking cessation (nicotinic ACh receptor partial agonist)
Venlafaxine	Serotonin-norepinephrine reuptake inhibitor

SECTION IV

Top-Rated Review Resources

“Some books are to be tasted, others to be swallowed, and some few to be chewed and digested.”

—Sir Francis Bacon

“Always read something that will make you look good if you die in the middle of it.”

—P.J. O’Rourke

“So many books, so little time.”

—Frank Zappa

“If one cannot enjoy reading a book over and over again, there is no use in reading it at all.”

—Oscar Wilde

▶ How to Use the Database	712
▶ Question Banks and Books	714
▶ Web and Mobile Apps	714
▶ Comprehensive	715
▶ Anatomy, Embryology, and Neuroscience	715
▶ Behavioral Science	716
▶ Biochemistry	716
▶ Cell Biology and Histology	716
▶ Microbiology and Immunology	717
▶ Pathology	717
▶ Pharmacology	718
▶ Physiology	718

▶ HOW TO USE THE DATABASE

This section is a database of top-rated basic science review books, sample examination books, software, websites, and apps that have been marketed to medical students studying for the USMLE Step 1. For each recommended resource, we list (where applicable) the **Title**, the **First Author** (or editor), the **Current Publisher**, the **Copyright Year**, the **Number of Pages**, the **Approximate List Price**, the **Format** of the resource, and the **Number of Test Questions**. Finally, each recommended resource receives a **Rating**. Within each section, resources are arranged first by Rating and then alphabetically by the first author within each Rating group.

For a complete list of resources, including summaries that describe their overall style and utility, go to www.firstaidteam.com/bonus.

A letter rating scale with six different grades reflects the detailed student evaluations for **Rated Resources**. Each rated resource receives a rating as follows:

A+	Excellent for boards review.
A A-	Very good for boards review; choose among the group.
B+ B	Good, but use only after exhausting better resources.
B-	Fair, but there are many better resources in the discipline; or low-yield subject material.

The Rating is meant to reflect the overall usefulness of the resource in helping medical students prepare for the USMLE Step 1. This is based on a number of factors, including:

- The cost
- The readability of the text or usability of the app
- The appropriateness and accuracy of the material
- The quality and number of sample questions
- The quality of written answers to sample questions
- The quality and appropriateness of the illustrations (eg, graphs, diagrams, photographs)
- The length of the text (longer is not necessarily better)
- The quality and number of other resources available in the same discipline
- The importance of the discipline for the USMLE Step 1

Please note that ratings do not reflect the quality of the resources for purposes other than reviewing for the USMLE Step 1. Many books with lower ratings are well written and informative but are not ideal for boards

preparation. We have not listed or commented on general textbooks available in the basic sciences.

Evaluations are based on the cumulative results of formal and informal surveys of thousands of medical students at many medical schools across the country. The ratings represent a consensus opinion, but there may have been a broad range of opinion or limited student feedback on any particular resource.

Please note that the data listed are subject to change in that:

- Publishers' prices change frequently.
- Bookstores often charge an additional markup.
- New editions come out frequently, and the quality of updating varies.
- The same book may be reissued through another publisher.

We actively encourage medical students and faculty to submit their opinions and ratings of these basic science review materials so that we may update our database. (See p. xvii, How to Contribute.) In addition, we ask that publishers and authors submit for evaluation review copies of basic science review books, including new editions and books not included in our database. We also solicit reviews of new books or suggestions for alternate modes of study that may be useful in preparing for the examination, such as flash cards, computer software, commercial review courses, apps, and websites.

Disclaimer/Conflict of Interest Statement

No material in this book, including the ratings, reflects the opinion or influence of the publisher. All errors and omissions will gladly be corrected if brought to the attention of the authors through our blog at www.firstaidteam.com. Please note that USMLE-Rx and the entire *First Aid for the USMLE* series are publications by certain authors of this book; the following ratings are based solely on recommendations from the student authors of this book as well as data from the student survey and feedback forms.

▶ TOP-RATED REVIEW RESOURCES

Question Banks and Books

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁺	<i>UWorld Qbank</i>	UWorld	www.uworld.com	Test/2400 q	\$249–\$749
A	<i>NBME Practice Exams</i>	National Board of Medical Examiners	www.nbme.org/students/sas/Comprehensive.html	Test/200 q	\$60
A⁻	<i>AMBOSS</i>	Amboss	www.amboss.com	Test/3500 q	\$9–\$365
A⁻	<i>USMLE-Rx Qmax</i>	USMLE-Rx	www.usmle-rx.com	Test/2300 q	\$89–\$339
B⁺	<i>Kaplan Qbank</i>	Kaplan	www.kaptest.com	Test/2100 q	\$99–\$349
B	<i>BoardVitals</i>		www.boardvitals.com	Test/1750 q	\$59–\$179
B	<i>Kaplan USMLE Step 1 Qbook</i>	Kaplan	Kaplan, 2017, 468 pages	Test/850 q	\$50
B	<i>Pastest</i>		www.pastest.com	Test/2100 q	\$79–\$249
B	<i>TrueLearn Review</i>		www.truelearn.com	Test/2200 q	\$159–\$399

Web and Mobile Apps

		AUTHOR	PUBLISHER	TYPE	PRICE
A	<i>Anki</i>		www.ankisrs.net	Flash cards	Free
A	<i>Boards and Beyond</i>		www.boardsbeyond.com	Review/ Test/1300 q	\$19–\$249
A	<i>Physeio</i>		www.physeio.com	Review	\$30–\$150
A	<i>SketchyMedical</i>		www.sketchymedical.com	Review	\$99–\$369
A⁻	<i>Cram Fighter</i>		www.cramfighter.com	Study plan	\$29–\$159
A⁻	<i>First Aid Step 1 Express</i>		www.usmle-rx.com	Review/Test	\$69–\$299
B⁺	<i>First Aid Step 1 Flash Facts</i>		www.usmle-rx.com	Flash cards	\$29–\$149
B⁺	<i>Medbullets</i>		www.medbullets.com	Review/ Test/1000 q	Free
B⁺	<i>Medical School Pathology</i>		www.medicalschoolpathology.com	Review	Free
B⁺	<i>OnlineMedEd</i>		www.onlinemeded.org	Review	Free
B⁺	<i>Osmosis</i>		www.osmosis.org	Test	\$179–\$279
B⁺	<i>USMLE Step 1 Mastery</i>		builtbyhlt.com/medical/usmle-step-1-mastery	Test/1400 q	\$2–\$10
B⁺	<i>WebPath: The Internet Pathology Laboratory</i>		webpath.med.utah.edu	Review/ Test/1300 q	Free
B	<i>Blue Histology</i>		www.lab.anhb.uwa.edu.au/mb140	Review/Test	Free
B	<i>Digital Anatomist Project: Interactive Atlases</i>	University of Washington	da.si.washington.edu/da.html	Review	Free
B	<i>Dr. Najeeb Lectures</i>		www.drnajeeblectures.com	Review	\$99

B	<i>Firecracker</i>	Firecracker Inc.	firecracker.lww.com	Review/ Test/2800 q	\$39–\$660
B	<i>KISSPrep</i>		www.kissprep.com	Review	\$99–\$135
B	<i>Lecturio</i>		www.lecturio.com	Review/ Test/2150 q	\$50–\$300
B	<i>Memorang</i>	Memorang Inc.	www.memorangapp.com	Flash cards	\$19–\$239
B	<i>Picmonic</i>		www.picmonic.com	Review	\$25–\$480
B⁻	<i>Radiopaedia.org</i>		www.radiopaedia.org	Cases/Test	Free
B⁻	<i>The Pathology Guy</i>	Friedlander	www.pathguy.com	Review	Free

Comprehensive

		AUTHOR	PUBLISHER	TYPE	PRICE
A	<i>First Aid for the Basic Sciences: General Principles</i>	Le	McGraw-Hill, 2017, 528 pages	Review	\$55
A	<i>First Aid Cases for the USMLE Step 1</i>	Le	McGraw-Hill, 2018, 496 pages	Cases	\$50
A⁻	<i>First Aid for the Basic Sciences: Organ Systems</i>	Le	McGraw-Hill, 2017, 912 pages	Review	\$72
A⁻	<i>Crush Step 1: The Ultimate USMLE Step 1 Review</i>	O'Connell	Elsevier, 2017, 704 pages	Review	\$45
A⁻	<i>Cracking the USMLE Step 1</i>	Princeton Review	Princeton Review, 2013, 832 pages	Review	\$45
B⁺	<i>USMLE Step 1 Secrets in Color</i>	Brown	Elsevier, 2016, 800 pages, ISBN 9780323396790	Review	\$43
B⁺	<i>Step-Up to USMLE Step 1 2015</i>	Jenkins	Lippincott Williams & Wilkins, 2014, 528 pages	Review	\$50
B⁺	<i>USMLE Step 1 Lecture Notes 2018</i>	Kaplan	Kaplan Medical, 2018, ~2700 pages	Review	\$330
B⁺	<i>USMLE Images for the Boards: A Comprehensive Image-Based Review</i>	Tully	Elsevier, 2012, 296 pages	Review	\$42
B	<i>USMLE Step 1 Made Ridiculously Simple</i>	Carl	MedMaster, 2017, 416 pages,	Review/Test 1000 q	\$30
B	<i>medEssentials for the USMLE Step 1</i>	Manley	Kaplan, 2012, 588 pages	Review	\$55

Anatomy, Embryology, and Neuroscience

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	<i>High-Yield Gross Anatomy</i>	Dudek	Lippincott Williams & Wilkins, 2014, 320 pages	Review	\$43
A⁻	<i>Clinical Anatomy Made Ridiculously Simple</i>	Goldberg	MedMaster, 2016, 175 pages	Review	\$30
B⁺	<i>High-Yield Embryology</i>	Dudek	Lippincott Williams & Wilkins, 2013, 176 pages	Review	\$56
B⁺	<i>High-Yield Neuroanatomy</i>	Fix	Lippincott Williams & Wilkins, 2015, 208 pages	Review/ Test/50 q	\$40

Anatomy, Embryology, and Neuroscience (continued)

		AUTHOR	PUBLISHER	TYPE	PRICE
B⁺	<i>Anatomy—An Essential Textbook</i>	Gilroy	Thieme, 2017, 528 pages	Text/ Test/400 q	\$48
B⁺	<i>Netter's Anatomy Flash Cards</i>	Hansen	Saunders, 2018, 688 flash cards	Flash cards	\$40
B⁺	<i>Crash Course: Anatomy</i>	Stenhouse	Elsevier, 2015, 288 pages	Review	\$45
B	<i>BRS Embryology</i>	Dudek	Lippincott Williams & Wilkins, 2014, 336 pages	Review/ Test/220 q	\$56
B	<i>Anatomy Flash Cards: Anatomy on the Go</i>	Gilroy	Thieme, 2013, 752 flash cards	Flash cards	\$60
B	<i>Clinical Neuroanatomy Made Ridiculously Simple</i>	Goldberg	MedMaster, 2014, 90 pages + CD-ROM	Review/Test/ Few q	\$26
B	<i>Netter's Anatomy Coloring Book</i>	Hansen	Elsevier, 2018, 392 pages	Review	\$20
B	<i>Case Files: Anatomy</i>	Toy	McGraw-Hill, 2014, 416 pages	Cases	\$35
B⁻	<i>Case Files: Neuroscience</i>	Toy	McGraw-Hill, 2014, 432 pages	Cases	\$35

Behavioral Science

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	<i>BRS Behavioral Science</i>	Fadem	Lippincott Williams & Wilkins, 2016, 384 pages	Review/ Test/700 q	\$52
B⁺	<i>High-Yield Biostatistics, Epidemiology, and Public Health</i>	Glaser	Lippincott Williams & Wilkins, 2013, 168 pages	Review	\$43

Biochemistry

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	<i>Pixorize</i>		www.pixorize.com	Review	\$100–\$130
B⁺	<i>Medical Biochemistry—An Illustrated Review</i>	Panini	Thieme, 2013, 441 pages	Review/ Test/400 q	\$40
B	<i>Lange Flash Cards Biochemistry and Genetics</i>	Baron	McGraw-Hill, 2017, 196 flash cards	Flash cards	\$40
B	<i>Lippincott Illustrated Reviews: Biochemistry</i>	Ferrier	Lippincott Williams & Wilkins, 2017, 560 pages	Review/ Test/200 q	\$78
B	<i>BRS Biochemistry, Molecular Biology, and Genetics</i>	Lieberman	Lippincott Williams & Wilkins, 2013, 432 pages	Review/Test	\$54
B	<i>Case Files: Biochemistry</i>	Toy	McGraw-Hill, 2014, 480 pages	Cases	\$35
B	<i>PreTest Biochemistry and Genetics</i>	Wilson	McGraw-Hill, 2017, 592 pages	Test/500 q	\$38

Cell Biology and Histology

		AUTHOR	PUBLISHER	TYPE	PRICE
B⁺	<i>BRS Cell Biology and Histology</i>	Gartner	Lippincott Williams & Wilkins, 2018, 448 pages	Review/ Test/320 q	\$54
B⁺	<i>Crash Course: Cell Biology and Genetics</i>	Stubbs	Elsevier, 2015, 216 pages	Review/Print + online	\$47
B	<i>Wheater's Functional Histology</i>	Young	Elsevier, 2013, 464 pages	Text	\$83

Microbiology and Immunology

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	<i>Basic Immunology</i>	Abbas	Elsevier, 2019, 336 pages	Review	\$70
A⁻	<i>Clinical Microbiology Made Ridiculously Simple</i>	Gladwin	MedMaster, 2019, 418 pages	Review	\$38
A⁻	<i>Medical Microbiology and Immunology Flash Cards</i>	Rosenthal	Elsevier, 2016, 192 flash cards	Flash cards	\$40
B⁺	<i>Lippincott Illustrated Reviews: Immunology</i>	Doan	Lippincott Williams & Wilkins, 2012, 384 pages	Reference/ Test/Few q	\$75
B⁺	<i>Microcards: Microbiology Flash Cards</i>	Harpavat	Lippincott Williams & Wilkins, 2015, 312 flash cards	Flash cards	\$53
B⁺	<i>Review of Medical Microbiology and Immunology</i>	Levinson	McGraw-Hill, 2018, 832 pages	Review/ Test/654 q	\$63
B⁺	<i>How the Immune System Works</i>	Sompayrac	Wiley-Blackwell, 2019, 168 pages	Review	\$50
B	<i>Case Studies in Immunology: Clinical Companion</i>	Geha	W. W. Norton & Company, 2016, 384 pages	Cases	\$62
B	<i>Pretest: Microbiology</i>	Kettering	McGraw-Hill, 2013, 480 pages	Test/500 q	\$38
B	<i>Case Files: Microbiology</i>	Toy	McGraw-Hill, 2014, 416 pages	Cases	\$36
B	<i>Lange Microbiology and Infectious Diseases Flash Cards, 3e</i>	Somers	McGraw-Hill Education, 2017, 358 pages	Flash cards	\$46
B⁻	<i>Lippincott Illustrated Reviews: Microbiology</i>	Cornelissen	Lippincott Williams & Wilkins, 2019, 448 pages	Review/Test/ Few q	\$73

Pathology

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁺	<i>Pathoma: Fundamentals of Pathology</i>	Sattar	Pathoma, 2019, 218 pages	Review/ Lecture	\$85–\$120
A⁻	<i>Rapid Review: Pathology</i>	Goljan	Elsevier, 2018, 864 pages	Review/ Test/500 q	\$65
A⁻	<i>Robbins and Cotran Review of Pathology</i>	Klatt	Elsevier, 2014, 504 pages	Test/1100 q	\$55
A⁻	<i>Crash Course: Pathology</i>	Xiu	Elsevier, 2019, 438 pages	Review	\$40
B	<i>High-Yield Histopathology</i>	Dudek	Lippincott Williams & Wilkins, 2017, 320 pages	Review	\$36
B	<i>Pathophysiology of Disease: Introduction to Clinical Medicine</i>	Hammer	McGraw-Hill, 2018, 832 pages	Text	\$90
B	<i>Haematology at a Glance</i>	Mehta	Blackwell Science, 2014, 136 pages	Review	\$49
B	<i>Pocket Companion to Robbins and Cotran Pathologic Basis of Disease</i>	Mitchell	Elsevier, 2016, 896 pages	Review	\$40
B	<i>BRS Pathology</i>	Schneider	Lippincott Williams & Wilkins, 2013, 480 pages	Review/ Test/450 q	\$54

Pharmacology

		AUTHOR	PUBLISHER	TYPE	PRICE
B⁺	<i>Crash Course: Pharmacology</i>	Battista	Elsevier, 2019, 336 pages	Review	\$40
B⁺	<i>Master the Boards USMLE Step 1 Pharmacology Flashcards</i>	Fischer	Kaplan, 2015, 200 flash cards	Flash cards	\$55
B⁺	<i>BRS Pharmacology</i>	Rosenfeld	Lippincott Williams & Wilkins, 2019, 384 pages	Review/ Test/200 q	\$55
B	<i>Lange Pharmacology Flash Cards</i>	Baron	McGraw-Hill, 2017, 266 flash cards	Flash cards	\$39
B	<i>Pharmacology Flash Cards</i>	Brenner	Elsevier, 2017, 230 flash cards	Flash cards	\$45
B	<i>Katzung & Trevor's Pharmacology: Examination and Board Review</i>	Trevor	McGraw-Hill, 2018, 592 pages	Review/ Test/800 q	\$54
B	<i>Lippincott Illustrated Reviews: Pharmacology</i>	Whalen	Lippincott Williams & Wilkins, 2018, 576 pages	Review/ Test/380 q	\$75

Physiology

		AUTHOR	PUBLISHER	TYPE	PRICE
A⁻	<i>BRS Physiology</i>	Costanzo	Lippincott Williams & Wilkins, 2018, 304 pages	Review/ Test/350 q	\$54
A⁻	<i>Pathophysiology of Heart Disease</i>	Lilly	Lippincott Williams & Williams, 2015, 480 pages	Review	\$57
A⁻	<i>PreTest Physiology</i>	Metting	McGraw-Hill, 2013, 528 pages	Test/500 q	\$38
A⁻	<i>Color Atlas of Physiology</i>	Silbernagl	Thieme, 2015, 472 pages	Review	\$50
B⁺	<i>BRS Physiology Cases and Problems</i>	Costanzo	Lippincott Williams & Wilkins, 2012, 368 pages	Cases	\$58
B⁺	<i>Physiology</i>	Costanzo	Saunders, 2017, 528 pages	Text	\$60
B⁺	<i>Vander's Renal Physiology</i>	Eaton	McGraw-Hill, 2018, 224 pages	Text	\$49
B⁺	<i>Acid-Base, Fluids, and Electrolytes Made Ridiculously Simple</i>	Preston	MedMaster, 2017, 166 pages	Review	\$24
B⁺	<i>Pulmonary Pathophysiology: The Essentials</i>	West	Lippincott Williams & Wilkins, 2017, 264 pages	Review/ Test/75 q	\$57
B	<i>Rapid Review: Physiology</i>	Brown	Elsevier, 2011, 384 pages	Test/350 q	\$39
B	<i>Endocrine Physiology</i>	Molina	McGraw-Hill, 2018, 320 pages	Review	\$59
B⁻	<i>Netter's Physiology Flash Cards</i>	Mulroney	Saunders, 2015, 450 flash cards	Flash cards	\$40

SECTION IV

Abbreviations and Symbols

ABBREVIATION	MEANING
1st MC*	1st metacarpal
A-a	alveolar-arterial [gradient]
AA	Alcoholics Anonymous, amyloid A
AAMC	Association of American Medical Colleges
AAo*	ascending aorta
Ab	antibody
ABPA	allergic bronchopulmonary aspergillosis
AC	adenyl cyclase
ACA	anterior cerebral artery
Acetyl-CoA	acetyl coenzyme A
ACD	anemia of chronic disease
ACE	angiotensin-converting enzyme
ACh	acetylcholine
AChE	acetylcholinesterase
ACL	anterior cruciate ligament
ACom	anterior communicating [artery]
ACTH	adrenocorticotrophic hormone
AD	Alzheimer disease, autosomal dominant
ADA	adenosine deaminase, Americans with Disabilities Act
ADH	antidiuretic hormone
ADHD	attention-deficit hyperactivity disorder
ADP	adenosine diphosphate
ADPKD	autosomal-dominant polycystic kidney disease
AFP	α -fetoprotein
Ag	antigen, silver
AICA	anterior inferior cerebellar artery
AIDS	acquired immunodeficiency syndrome
AIHA	autoimmune hemolytic anemia
AKI	acute kidney injury
AKT	protein kinase B
AL	amyloid light [chain]
ALA	aminolevulinate
ALI	acute lung injury
ALL	acute lymphoblastic (lymphocytic) leukemia
ALP	alkaline phosphatase
ALS	amyotrophic lateral sclerosis
ALT	alanine transaminase
AMA	American Medical Association, antimitochondrial antibody
AML	acute myelogenous (myeloid) leukemia
AMP	adenosine monophosphate
ANA	antinuclear antibody
ANCA	antineutrophil cytoplasmic antibody
ANOVA	analysis of variance

ABBREVIATION	MEANING
ANP	atrial natriuretic peptide
ANS	autonomic nervous system
Ant*	anterior
anti-CCP	anti-cyclic citrullinated peptide
Ao*	aorta
AOA	American Osteopathic Association
AP	action potential, A & P [ribosomal binding sites]
APC	antigen-presenting cell, activated protein C
Apo	apolipoprotein
APP	amyloid precursor protein
APRT	adenine phosphoribosyltransferase
aPTT	activated partial thromboplastin time
APUD	amine precursor uptake decarboxylase
AR	attributable risk, autosomal recessive, aortic regurgitation
ARB	angiotensin receptor blocker
ARDS	acute respiratory distress syndrome
Arg	arginine
ARPKD	autosomal-recessive polycystic kidney disease
ART	antiretroviral therapy
AS	aortic stenosis
ASA	anterior spinal artery
ASD	atrial septal defect
ASO	anti-streptolysin O
AST	aspartate transaminase
AT	angiotensin, antithrombin
ATN	acute tubular necrosis
ATP	adenosine triphosphate
ATPase	adenosine triphosphatase
ATTR	transthyretin-mediated amyloidosis
AUB	abnormal uterine bleeding
AV	atrioventricular
AZT	azidothymidine
BAL	British anti-Lewisite [dimercaprol]
BBB	blood-brain barrier
BCCG	bacille Calmette-Guérin
BH ₄	tetrahydrobiopterin
BM	basement membrane
BOOP	bronchiolitis obliterans organizing pneumonia
BP	bisphosphate, blood pressure
BPG	bisphosphoglycerate
BPH	benign prostatic hyperplasia
BT	bleeding time
BUN	blood urea nitrogen
Ca*	capillary

*Image abbreviation only

ABBREVIATION	MEANING
Ca ²⁺	calcium ion
CAD	coronary artery disease
CAF	common application form
cAMP	cyclic adenosine monophosphate
CBG	corticosteroid-binding globulin
Cbm*	cerebellum
CBSE	Comprehensive Basic Science Examination
CBSSA	Comprehensive Basic Science Self-Assessment
CBT	computer-based test, cognitive behavioral therapy
CC*	corpus callosum
CCA*	common carotid artery
CCK	cholecystokinin
CCS	computer-based case simulation
CD	cluster of differentiation
CDK	cyclin-dependent kinase
cDNA	complementary deoxyribonucleic acid
CEA	carcinoembryonic antigen
CETP	cholesterol-ester transfer protein
CF	cystic fibrosis
CFTR	cystic fibrosis transmembrane conductance regulator
CGD	chronic granulomatous disease
cGMP	cyclic guanosine monophosphate
CGRP	calcitonin gene-related peptide
C _H 1–C _H 3	constant regions, heavy chain [antibody]
ChAT	choline acetyltransferase
CHD*	common hepatic duct
χ ²	chi-squared
CI	confidence interval
CIN	candidate identification number, carcinoma in situ, cervical intraepithelial neoplasia
CIS	Communication and Interpersonal Skills
CK	clinical knowledge, creatine kinase
CKD	chronic kidney disease
CK-MB	creatine kinase, MB fraction
C _L	constant region, light chain [antibody]
CL	clearance
Cl ⁻	chloride ion
CLL	chronic lymphocytic leukemia
CMC	carpometacarpal (joint)
CML	chronic myelogenous (myeloid) leukemia
CMV	cytomegalovirus
CN	cranial nerve
CN ⁻	cyanide ion
CNS	central nervous system
CNV	copy number variation
CO	carbon monoxide, cardiac output
CO ₂	carbon dioxide
CoA	coenzyme A
COL1A1	collagen, type I, alpha 1
COL1A2	collagen, type I, alpha 2
COMT	catechol-O-methyltransferase
COP	coat protein
COPD	chronic obstructive pulmonary disease
CoQ	coenzyme Q

ABBREVIATION	MEANING
COX	cyclooxygenase
C _p	plasma concentration
CPAP	continuous positive airway pressure
CPR	cardiopulmonary resuscitation
Cr	creatinine
CRC	colorectal cancer
CREST	calcinosis, Raynaud phenomenon, esophageal dysfunction, sclerosis, and telangiectasias [syndrome]
CRH	corticotropin-releasing hormone
CRP	C-reactive protein
CS	clinical skills
C-section	cesarean section
CSF	cerebrospinal fluid
CT	computed tomography
CTP	cytidine triphosphate
CXR	chest x-ray
DA	dopamine
DAF	decay-accelerating factor
DAG	diacylglycerol
dATP	deoxyadenosine triphosphate
DCIS	ductal carcinoma in situ
DCT	distal convoluted tubule
ddI	didanosine
DES	diethylstilbestrol
DH	dehydrogenase
DHAP	dihydroxyacetone phosphate
DHEA	dehydroepiandrosterone
DHF	dihydrofolic acid
DHT	dihydrotestosterone
DI	diabetes insipidus
DIC	disseminated intravascular coagulation
DIP	distal interphalangeal [joint]
DKA	diabetic ketoacidosis
DLCO	diffusing capacity for carbon monoxide
DM	diabetes mellitus
DNA	deoxyribonucleic acid
DNR	do not resuscitate
dNTP	deoxynucleotide triphosphate
DO	doctor of osteopathy
DPGN	diffuse proliferative glomerulonephritis
DPM	doctor of podiatric medicine
DPP-4	dipeptidyl peptidase-4
DPPC	dipalmitoylphosphatidylcholine
DS	double stranded
dsDNA	double-stranded deoxyribonucleic acid
dsRNA	double-stranded ribonucleic acid
DRG	dorsal root ganglion
d4T	didehydrodeoxythymidine [stavudine]
dTMP	deoxythymidine monophosphate
DTR	deep tendon reflex
DTs	delirium tremens
dUDP	deoxyuridine diphosphate
dUMP	deoxyuridine monophosphate
DVT	deep venous thrombosis
E*	euthromatin, esophagus

*Image abbreviation only

ABBREVIATION	MEANING
EBV	Epstein-Barr virus
ECA*	external carotid artery
ECF	extracellular fluid
ECFMG	Educational Commission for Foreign Medical Graduates
ECCG	electrocardiogram
ECL	enterochromaffin-like [cell]
ECM	extracellular matrix
ECT	electroconvulsive therapy
ED ₅₀	median effective dose
EDRF	endothelium-derived relaxing factor
EDTA	ethylenediamine tetra-acetic acid
EDV	end-diastolic volume
EEG	electroencephalogram
EF	ejection fraction
EGF	epidermal growth factor
EHEC	enterohemorrhagic <i>E coli</i>
EIEC	enteroinvasive <i>E coli</i>
ELISA	enzyme-linked immunosorbent assay
EM	electron micrograph/microscopy
EMB	eosin–methylene blue
EPEC	eneteropathogenic <i>E coli</i>
Epi	epinephrine
EPO	erythropoietin
EPS	extrapyramidal system
ER	endoplasmic reticulum, estrogen receptor
ERAS	Electronic Residency Application Service
ERCP	endoscopic retrograde cholangiopancreatography
ERP	effective refractory period
eRPF	effective renal plasma flow
ERT	estrogen replacement therapy
ERV	expiratory reserve volume
ESR	erythrocyte sedimentation rate
ESRD	end-stage renal disease
ESV	end-systolic volume
ETEC	enterotoxigenic <i>E coli</i>
EtOH	ethyl alcohol
EV	esophageal vein
F	bioavailability
FA	fatty acid
Fab	fragment, antigen-binding
FAD	flavin adenine dinucleotide
FADH ₂	reduced flavin adenine dinucleotide
FAP	familial adenomatous polyposis
F1,6BP	fructose-1,6-bisphosphate
F2,6BP	fructose-2,6-bisphosphate
FBPase	fructose bisphosphatase
FBPase-2	fructose bisphosphatase-2
Fc	fragment, crystallizable
FcR	Fc receptor
5f-dUMP	5-fluorodeoxyuridine monophosphate
Fe ²⁺	ferrous ion
Fe ³⁺	ferric ion
Fem*	femur
FENa	excreted fraction of filtered sodium

ABBREVIATION	MEANING
FEV ₁	forced expiratory volume in 1 second
FF	filtration fraction
FFA	free fatty acid
FGF	fibroblast growth factor
FGFR	fibroblast growth factor receptor
FISH	fluorescence in situ hybridization
FIT	fecal immunochemical testing
FKBP	FK506 binding protein
fMet	formylmethionine
FMG	foreign medical graduate
FMN	flavin mononucleotide
FN	false negative
FP, FP*	false positive, foot process
FRC	functional residual capacity
FSH	follicle-stimulating hormone
FSMB	Federation of State Medical Boards
FTA-ABS	fluorescent treponemal antibody—absorbed
FTD*	frontotemporal dementia
5-FU	5-fluorouracil
FVC	forced vital capacity
GABA	γ-aminobutyric acid
GAG	glycosaminoglycan
Gal	galactose
GBM	glomerular basement membrane
GC	glomerular capillary
G-CSF	granulocyte colony-stimulating factor
GERD	gastroesophageal reflux disease
GFAP	glial fibrillary acid protein
GFR	glomerular filtration rate
GGT	γ-glutamyl transpeptidase
GH	growth hormone
GHB	γ-hydroxybutyrate
GHRH	growth hormone–releasing hormone
G ₁	G protein, I polypeptide
GI	gastrointestinal
GIP	gastric inhibitory peptide
GIST	gastrointestinal stromal tumor
GLUT	glucose transporter
GM	granulocyte macrophage
GM-CSF	granulocyte-macrophage colony stimulating factor
GMP	guanosine monophosphate
GnRH	gonadotropin-releasing hormone
GP	glycoprotein
G6P	glucose-6-phosphate
G6PD	glucose-6-phosphate dehydrogenase
GPe	globus pallidus externa
GPi	globus pallidus interna
GPI	glycosyl phosphatidylinositol
GRP	gastrin-releasing peptide
G _s	G protein, S polypeptide
GSH	reduced glutathione
GSSG	oxidized glutathione
GTP	guanosine triphosphate
GTPase	guanosine triphosphatase

*Image abbreviation only

ABBREVIATION	MEANING
GU	genitourinary
H*	heterochromatin
H ⁺	hydrogen ion
H ₁ , H ₂	histamine receptors
H ₂ S	hydrogen sulfide
HAV	hepatitis A virus
HAVAb	hepatitis A antibody
Hb	hemoglobin
HBcAb/HBcAg	hepatitis B core antibody/antigen
HBsAb/HBsAg	hepatitis B early antibody/antigen
HBsAb/HBsAg	hepatitis B surface antibody/antigen
HbCO ₂	carbaminohemoglobin
HBV	hepatitis B virus
HCC	hepatocellular carcinoma
hCG	human chorionic gonadotropin
HCO ₃ ⁻	bicarbonate
Hct	hematocrit
HCTZ	hydrochlorothiazide
HCV	hepatitis C virus
HDL	high-density lipoprotein
HDN	hemolytic disease of the newborn
HDV	hepatitis D virus
H&E	hematoxylin and eosin
HEV	hepatitis E virus
HF	heart failure
Hfr	high-frequency recombination [cell]
HFpEF	heart failure with preserved ejection fraction
HFrEF	heart failure with reduced ejection fraction
HGPRT	hypoxanthine-guanine phosphoribosyltransferase
HHb	deoxygenated hemoglobin
HHS	hyperosmolar hyperglycemic state
HHV	human herpesvirus
5-HIAA	5-hydroxyindoleacetic acid
HIT	heparin-induced thrombocytopenia
HIV	human immunodeficiency virus
HL	hepatic lipase
HLA	human leukocyte antigen
HMG-CoA	hydroxymethylglutaryl-coenzyme A
HMP	hexose monophosphate
HMWK	high-molecular-weight kininogen
HNPCC	hereditary nonpolyposis colorectal cancer
hnRNA	heterogeneous nuclear ribonucleic acid
H ₂ O ₂	hydrogen peroxide
HOCM	hypertrophic obstructive cardiomyopathy
HPA	hypothalamic-pituitary-adrenal [axis]
HPL	human placental lactogen
HPO	hypothalamic-pituitary-ovarian [axis]
HPV	human papillomavirus
HR	heart rate
HSP	Henoch-Schönlein purpura
HSV	herpes simplex virus
5-HT	5-hydroxytryptamine (serotonin)
HTLV	human T-cell leukemia virus
HTN	hypertension

ABBREVIATION	MEANING
HUS	hemolytic-uremic syndrome
HVA	homovanillic acid
IBD	inflammatory bowel disease
IBS	irritable bowel syndrome
IC	inspiratory capacity, immune complex
I _{Ca}	calcium current [heart]
I _f	funny current [heart]
ICA	internal carotid artery
ICAM	intercellular adhesion molecule
ICD	implantable cardioverter defibrillator
ICE	Integrated Clinical Encounter
ICF	intracellular fluid
ICP	intracranial pressure
ID	identification
ID ₅₀	median infective dose
IDL	intermediate-density lipoprotein
IF	immunofluorescence, initiation factor
IFN	interferon
Ig	immunoglobulin
IGF	insulin-like growth factor
I _K	potassium current [heart]
IL	interleukin
IM	intramuscular
IMA	inferior mesenteric artery
IMG	international medical graduate
IMP	inosine monophosphate
IMV	inferior mesenteric vein
I _{Na}	sodium current [heart]
INH	isoniazid
INO	internuclear ophthalmoplegia
INR	International Normalized Ratio
IO	inferior oblique [muscle]
IOP	intraocular pressure
IP ₃	inositol triphosphate
IPV	inactivated polio vaccine
IR	current × resistance [Ohm's law], inferior rectus [muscle]
IRV	inspiratory reserve volume
ITP	idiopathic thrombocytopenic purpura
IUD	intrauterine device
IUGR	intrauterine growth restriction
IV	intravenous
IVC	inferior vena cava
IVDU	intravenous drug use
IVIG	intravenous immunoglobulin
JAK/STAT	Janus kinase/signal transducer and activator of transcription [pathway]
JGA	juxtaglomerular apparatus
JVD	jugular venous distention
JVP	jugular venous pulse
K ⁺	potassium ion
KatG	catalase-peroxidase produced by <i>M tuberculosis</i>
K _e	elimination constant
K _f	filtration constant
KG	ketoglutarate

*Image abbreviation only

ABBREVIATION	MEANING
K_m	Michaelis-Menten constant
KOH	potassium hydroxide
L	left, liver
LA	left atrial, left atrium
LAD	left anterior descending coronary artery
LAP	leukocyte alkaline phosphatase
Lat cond*	lateral condyle
Lb*	lamellar body
LCA	left coronary artery
LCAT	lecithin-cholesterol acyltransferase
LCC*	left common carotid artery
LCFA	long-chain fatty acid
LCL	lateral collateral ligament
LCME	Liaison Committee on Medical Education
LCMV	lymphocytic choriomeningitis virus
LCX	left circumflex coronary artery
LD	loading dose
LD_{50}	median lethal dose
LDH	lactate dehydrogenase
LDL	low-density lipoprotein
LES	lower esophageal sphincter
LFA	leukocyte function-associated antigen
LFT	liver function test
LH	luteinizing hormone
LLL*	left lower lobe (of lung)
LLQ	left lower quadrant
LM	lateral meniscus, left main coronary artery, light microscopy
LMN	lower motor neuron
LOS	lipooligosaccharide
LPA*	left pulmonary artery
LPL	lipoprotein lipase
LPS	lipopolysaccharide
LR	lateral rectus [muscle]
LT	labile toxin, leukotriene
LUL*	left upper lobe (of lung)
LV	left ventricle, left ventricular
M_1-M_3	muscarinic (parasympathetic) ACh receptors
MAC	membrane attack complex, minimum alveolar concentration
MALT	mucosa-associated lymphoid tissue
MAO	monoamine oxidase
MAOI	monoamine oxidase inhibitor
MAP	mean arterial pressure, mitogen-activated protein
Max*	maxillary sinus
MC	midsystolic click
MCA	middle cerebral artery
MCAT	Medical College Admissions Test
MCHC	mean corpuscular hemoglobin concentration
MCL	medial collateral ligament
MCP	metacarpophalangeal [joint]
MCV	mean corpuscular volume
MD	maintenance dose
MDD	major depressive disorder
Med cond*	medial condyle

ABBREVIATION	MEANING
MELAS syndrome	mitochondrial encephalopathy, lactic acidosis, and stroke-like episodes
MEN	multiple endocrine neoplasia
Mg^{2+}	magnesium ion
$MgSO_4$	magnesium sulfate
MGUS	monoclonal gammopathy of undetermined significance
MHC	major histocompatibility complex
MI	myocardial infarction
MIF	müllerian inhibiting factor
MIRL	membrane inhibitor of reactive lysis
MLCK	myosin light-chain kinase
MLF	medial longitudinal fasciculus
MMC	migrating motor complex
MMR	measles, mumps, rubella [vaccine]
6-MP	6-mercaptopurine
MPGN	membranoproliferative glomerulonephritis
MPO	myeloperoxidase
MPO-ANCA/p-ANCA	myeloperoxidase/perinuclear antineutrophil cytoplasmic antibody
MR	medial rectus [muscle], mitral regurgitation
MRI	magnetic resonance imaging
miRNA	microribonucleic acid
mRNA	messenger ribonucleic acid
MRSA	methicillin-resistant <i>S aureus</i>
MS	mitral stenosis, multiple sclerosis
MSH	melanocyte-stimulating hormone
mtDNA	mitochondrial DNA
mTOR	mammalian target of rapamycin
MTP	metatarsophalangeal [joint]
MTX	methotrexate
MVO_2	myocardial oxygen consumption
MVP	mitral valve prolapse
N*	nucleus
Na^+	sodium ion
NAT	nucleic acid testing
NAD	nicotinamide adenine dinucleotide
NAD^+	oxidized nicotinamide adenine dinucleotide
NADH	reduced nicotinamide adenine dinucleotide
$NADP^+$	oxidized nicotinamide adenine dinucleotide phosphate
NADPH	reduced nicotinamide adenine dinucleotide phosphate
NBME	National Board of Medical Examiners
NBOME	National Board of Osteopathic Medical Examiners
NBPME	National Board of Podiatric Medical Examiners
NE	norepinephrine
NF	neurofibromatosis
NFAT	nuclear factor of activated T-cell
NH_3	ammonia
NH_4^+	ammonium
NK	natural killer [cells]
N_M	muscarinic ACh receptor in neuromuscular junction
NMDA	N-methyl-d-aspartate
NMJ	neuromuscular junction
NMS	neuroleptic malignant syndrome
N_N	nicotinic ACh receptor in autonomic ganglia

*Image abbreviation only

ABBREVIATION	MEANING
NRMP	National Residency Matching Program
NNRTI	non-nucleoside reverse transcriptase inhibitor
NO	nitric oxide
N ₂ O	nitrous oxide
NPH	neutral protamine Hagedorn, normal pressure hydrocephalus
NPV	negative predictive value
NRTI	nucleoside reverse transcriptase inhibitor
NSAID	nonsteroidal anti-inflammatory drug
NSE	neuron-specific enolase
NSTEMI	non-ST-segment elevation myocardial infarction
Nu*	nucleolus
OAA	oxaloacetic acid
OCD	obsessive-compulsive disorder
OCP	oral contraceptive pill
ODC	oxygen-hemoglobin dissociation curve
OH	hydroxy
1,25-OH D ₃	calcitriol (active form of vitamin D)
25-OH D ₃	storage form of vitamin D
OPV	oral polio vaccine
OR	odds ratio
OS	opening snap
OSA	obstructive sleep apnea
OVLIT	organum vasculosum of the lamina terminalis
P-body	processing body (cytoplasmic)
P-450	cytochrome P-450 family of enzymes
PA	posteroanterior, pulmonary artery
PABA	<i>para</i> -aminobenzoic acid
Paco ₂	arterial PCO ₂
PACO ₂	alveolar PCO ₂
PAH	<i>para</i> -aminohippuric acid
PAN	polyarteritis nodosa
Pao ₂	partial pressure of oxygen in arterial blood
PAO ₂	partial pressure of oxygen in alveolar blood
PAP	Papanicolaou [smear], prostatic acid phosphatase
PAPPA	pregnancy-associated plasma protein A
PAS	periodic acid-Schiff
Pat*	patella
PBP	penicillin-binding protein
PC	platelet count, pyruvate carboxylase
PCA	posterior cerebral artery
PCC	prothrombin complex concentrate
PCL	posterior cruciate ligament
Pco ₂	partial pressure of carbon dioxide
PCom	posterior communicating [artery]
PCOS	polycystic ovarian syndrome
PCP	phencyclidine hydrochloride, <i>Pneumocystis jirovecii</i> pneumonia
PCR	polymerase chain reaction
PCT	proximal convoluted tubule
PCV13	pneumococcal conjugate vaccine
PCWP	pulmonary capillary wedge pressure
PDA	patent ductus arteriosus, posterior descending artery
PDE	phosphodiesterase

ABBREVIATION	MEANING
PDGF	platelet-derived growth factor
PDH	pyruvate dehydrogenase
PE	pulmonary embolism
PECAM	platelet-endothelial cell adhesion molecule
P _E CO ₂	expired air PCO ₂
PEP	phosphoenolpyruvate
PF	platelet factor
PFK	phosphofructokinase
PFK-2	phosphofructokinase-2
PFT	pulmonary function test
PG	phosphoglycerate
P _i	plasma interstitial osmotic pressure, inorganic phosphate
PICA	posterior inferior cerebellar artery
PID	pelvic inflammatory disease
P _I O ₂	PO ₂ in inspired air
PIP	proximal interphalangeal [joint]
PIP ₂	phosphatidylinositol 4,5-bisphosphate
PIP ₃	phosphatidylinositol 3,4,5-bisphosphate
PKD	polycystic kidney disease
PKR	interferon- α -induced protein kinase
PKU	phenylketonuria
PLP	pyridoxal phosphate
PML	progressive multifocal leukoencephalopathy
PMN	polymorphonuclear [leukocyte]
P _{net}	net filtration pressure
PNET	primitive neuroectodermal tumor
PNS	peripheral nervous system
Po ₂	partial pressure of oxygen
PO ₄ ³⁻	phosphate
Pop*	popliteal artery
Pop a*	popliteal artery
Post*	posterior
PPAR	peroxisome proliferator-activated receptor
PPD	purified protein derivative
PPI	proton pump inhibitor
PPM	parts per million
PPSV23	pneumococcal polysaccharide vaccine
PPV	positive predictive value
PR3-ANCA/ c-ANCA	cytoplasmic antineutrophil cytoplasmic antibody
PrP	prion protein
PRPP	phosphoribosylpyrophosphate
PSA	prostate-specific antigen
PSS	progressive systemic sclerosis
PT	prothrombin time
<i>PTEN</i>	phosphatase and tensin homolog
PTH	parathyroid hormone
PTHrP	parathyroid hormone-related protein
PTSD	post-traumatic stress disorder
PTT	partial thromboplastin time
PV	plasma volume, venous pressure
P _v *	pulmonary vein
PVC	polyvinyl chloride
PVR	pulmonary vascular resistance

*Image abbreviation only

ABBREVIATION	MEANING
R	correlation coefficient, right, R variable [group]
R ₃	Registration, Ranking, & Results [system]
RA	right atrium
RAAS	renin-angiotensin-aldosterone system
RANK-L	receptor activator of nuclear factor- κ B ligand
RAS	reticular activating system
RBF	renal blood flow
RCA	right coronary artery
REM	rapid eye movement
RER	rough endoplasmic reticulum
Rh	<i>rhesus</i> antigen
RLL*	right lower lobe (of lungs)
RLQ	right lower quadrant
RML*	right middle lobe (of lung)
RNA	ribonucleic acid
RNP	ribonucleoprotein
ROS	reactive oxygen species
RPF	renal plasma flow
RPGN	rapidly progressive glomerulonephritis
RPR	rapid plasma reagin
RR	relative risk, respiratory rate
rRNA	ribosomal ribonucleic acid
RS	Reed-Sternberg [cells]
RSC*	right subclavian artery
RSV	respiratory syncytial virus
RTA	renal tubular acidosis
RUL*	right upper lobe (of lung)
RUQ	right upper quadrant
RV	residual volume, right ventricle, right ventricular
RVH	right ventricular hypertrophy
[S]	substrate concentration
SA	sinoatrial
SAA	serum amyloid-associated [protein]
SAM	S-adenosylmethionine
SARS	severe acute respiratory syndrome
SCC	squamous cell carcinoma
SCD	sudden cardiac death
SCID	severe combined immunodeficiency disease
SCJ	squamocolumnar junction
SCM	sternocleidomastoid muscle
SCN	suprachiasmatic nucleus
SD	standard deviation
SE	standard error [of the mean]
SEP	Spoken English Proficiency
SER	smooth endoplasmic reticulum
SERM	selective estrogen receptor modulator
SGLT	sodium-glucose transporter
SHBG	sex hormone-binding globulin
SIADH	syndrome of inappropriate [secretion of] antidiuretic hormone
SIDS	sudden infant death syndrome
SJS	Stevens-Johnson syndrome
SLE	systemic lupus erythematosus
SLL	small lymphocytic lymphoma

ABBREVIATION	MEANING
SLT	Shiga-like toxin
SMA	superior mesenteric artery
SMX	sulfamethoxazole
SNARE	soluble NSF attachment protein receptor
SNe	substantia nigra pars compacta
SNP	single nucleotide polymorphism
SNr	substantia nigra pars reticulata
SNRI	serotonin and norepinephrine receptor inhibitor
snRNA	small nuclear RNA
snRNP	small nuclear ribonucleoprotein
SO	superior oblique [muscle]
SOAP	Supplemental Offer and Acceptance Program
Sp*	spleen
spp	species
SR	superior rectus [muscle]
SS	single stranded
ssDNA	single-stranded deoxyribonucleic acid
SSPE	subacute sclerosing panencephalitis
SSRI	selective serotonin reuptake inhibitor
ssRNA	single-stranded ribonucleic acid
St*	stomach
ST	Shiga toxin
StAR	steroidogenic acute regulatory protein
STEMI	ST-segment elevation myocardial infarction
STI	sexually transmitted infection
STN	subthalamic nucleus
SV	splenic vein, stroke volume
SVC	superior vena cava
SVR	systemic vascular resistance
SVT	supraventricular tachycardia
T*	trachea
t _{1/2}	half-life
T ₃	triiodothyronine
T ₄	thyroxine
TAPVR	total anomalous pulmonary venous return
TB	tuberculosis
TBG	thyroxine-binding globulin
TBV	total blood volume
3TC	dideoxythiacytidine [lamivudine]
TCA	tricarboxylic acid [cycle], tricyclic antidepressant
Tc cell	cytotoxic T cell
TCR	T-cell receptor
TDF	tenofovir disoproxil fumarate
TdT	terminal deoxynucleotidyl transferase
TE	tracheoesophageal
TFT	thyroid function test
TG	triglyceride
TGF	transforming growth factor
Th cell	helper T cell
THF	tetrahydrofolic acid
TI	therapeutic index
TIA	transient ischemic attack
Tib*	tibia
TIBC	total iron-binding capacity
TIPS	transjugular intrahepatic portosystemic shunt

*Image abbreviation only

ABBREVIATION	MEANING
TLC	total lung capacity
T_m	maximum rate of transport
TMP	trimethoprim
TN	true negative
TNF	tumor necrosis factor
TNM	tumor, node, metastases [staging]
TOP	topoisomerase
ToRCHeS	<i>Toxoplasma gondii</i> , rubella, CMV, HIV, HSV-2, syphilis
TP	true positive
tPA	tissue plasminogen activator
TPO	thyroid peroxidase, thrombopoietin
TPP	thiamine pyrophosphate
TPPA	<i>Treponema pallidum</i> particle agglutination assay
TPR	total peripheral resistance
TR	tricuspid regurgitation
TRAP	tartrate-resistant acid phosphatase
TRECs	T-cell receptor excision circles
TRH	thyrotropin-releasing hormone
tRNA	transfer ribonucleic acid
TSH	thyroid-stimulating hormone
TSI	triple sugar iron
TSS	toxic shock syndrome
TSST	toxic shock syndrome toxin
TTP	thrombotic thrombocytopenic purpura
TTR	transthyretin
TV	tidal volume
TXA_2	thromboxane A_2
UDP	uridine diphosphate
UMN	upper motor neuron
UMP	uridine monophosphate
UPD	uniparental disomy
URI	upper respiratory infection
USMLE	United States Medical Licensing Examination
UTI	urinary tract infection
UTP	uridine triphosphate


*Image abbreviation only

ABBREVIATION	MEANING
UV	ultraviolet
V_1, V_2	vasopressin receptors
VC	vital capacity
V_d	volume of distribution
VD	physiologic dead space
V(D)J	variable, (diversity), joining gene segments rearranged to form Ig genes
VDRL	Venereal Disease Research Laboratory
VEGF	vascular endothelial growth factor
V_H	variable region, heavy chain [antibody]
VHL	von Hippel-Lindau [disease]
VIP	vasoactive intestinal peptide
VIPoma	vasoactive intestinal polypeptide-secreting tumor
VJ	light-chain hypervariable region [antibody]
V_L	variable region, light chain [antibody]
VLCFA	very-long-chain fatty acids
VLDL	very low density lipoprotein
VMA	vanillylmandelic acid
VMAT	vesicular monoamine transporter
V_{max}	maximum velocity
VPL	ventral posterior nucleus, lateral
VPM	ventral posterior nucleus, medial
VPN	vancomycin, polymyxin, nystatin [media]
\dot{V}/\dot{Q}	ventilation/perfusion [ratio]
VRE	vancomycin-resistant enterococcus
VSD	ventricular septal defect
V_T	tidal volume
VTE	venous thromboembolism
vWF	von Willebrand factor
VZV	varicella-zoster virus
VMAT	vesicular monoamine transporter
XR	X-linked recessive
XXXY	normal complement of sex chromosomes for female/male
ZDV	zidovudine [formerly AZT]


SECTION IV


Image Acknowledgments

In this edition, in collaboration with MediQ Learning, LLC, and a variety of other partners, we are pleased to include the following clinical images and diagrams for the benefit of integrative student learning.

Portions of this book identified with the symbol  are copyright © USMLE-Rx.com (MediQ Learning, LLC).

Portions of this book identified with the symbol  are copyright © Dr. Richard Usatine and are provided under license through MediQ Learning, LLC.















Portions of this book identified with the symbol  are listed below by page number.












This symbol  refers to material that is available in the public domain.

This symbol  refers to the Creative Commons Attribution license, full text at <http://creativecommons.org/licenses/by/4.0/legalcode>.



This symbol  refers to the Creative Commons Attribution-Share Alike license, full text at: <http://creativecommons.org/licenses/by-sa/4.0/legalcode>.






Biochemistry

- 34 Chromatin structure.** Electron micrograph showing heterochromatin, euchromatin, and nucleolus. This image is a derivative work, adapted from the following source, available under : Røller RA, Rickett JD, Stickle WB. The hypobranchial gland of the estuarine snail *Stramonita haemastoma canaliculata* (Gray) (Prosobranchia: Muricidae): a light and electron microscopical study. *Am Malac Bull.* 1995;11(2):177-190. Available at <https://archive.org/details/americanm101119931994amer>.
- 49 Cilia structure: Image A.** Nine doublet + 2 singlet arrangement of microtubule.  Courtesy of Louisa Howard and Michael Binder. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 49 Cilia structure: Image B.** Cilia structure of basal body. This image is a derivative work, adapted from the following source, available under : Riparbelli MG, Cabrera OA, Callaini G, et al. Unique properties of *Drosophila* spermatocyte primary cilia. *Biol Open.* 2013 Nov 15; 2(11): 1137–1147. DOI: 10.1242/bio.20135355.
- 49 Cilia structure: Image C.** Dextrocardia. This image is a derivative work, adapted from the following source, available under : Oluwadare O, Ayoka AO, Akomolafe RO, et al. The role of electrocardiogram in the diagnosis of dextrocardia with mirror image atrial arrangement and ventricular position in a young adult Nigerian in Ile-Ife: a case report. *J Med Case Rep.* 2015;9:222. DOI: 10.1186/s13256-015-0695-4.
- 51 Osteogenesis imperfecta: Image A.** Skeletal deformities in upper extremity of child. This image is a derivative work, adapted from the following source, available under : Vanakker OM, Hemelsoet D, De Paepe. Hereditary connective tissue diseases in young adult stroke: a comprehensive synthesis. *Stroke Res Treat.* 2011;712903. DOI: 10.4061/2011/712903. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 51 Osteogenesis imperfecta: Image B.** Blue sclera. This image is a derivative work, adapted from the following source, available under : Wheatley K, Heng EL, Sheppard M, et al. A case of spontaneous intestinal perforation in osteogenesis imperfects. *J Clin Med Res.* 2010;2(4):198–200. DOI: 10.4021/jocmr369w.
- 51 Ehlers-Danlos syndrome: Images A and B.** Hyperextensibility of skin and DIP joint. This image is a derivative work, adapted from the following source, available under : Whitaker JK, Alexander, P, Chau DYS, et al. Severe conjunctivochalasis in association with classic type Ehlers-Danlos syndrome. *BMC Ophthalmol.* 2012;2:47. DOI: 10.1186/1471-2415-12-47.
- 52 Elastin: Image A.** Pes escavatium. This image is a derivative work, adapted from the following source, available under : De Maio F, Fichera A, De Luna V, et al. Orthopaedic aspects of Marfan syndrome: the experience of a referral center for diagnosis of rare diseases. *Adv Orthop.* 2016; 2016: 8275391. DOI 10.1155/2016/8275391.
- 55 Karyotyping.** Paar C, Herber G, Voskova, et al. This image is a derivative work, adapted from the following source, available under : A case of acute myeloid leukemia (AML) with an unreported combination of chromosomal abnormalities: gain of isochromosome 5p, tetrasomy 8 and unbalanced translocation der(19)t(17;19)(q23;p13). *Mol Cytogenet.* 2013;6:40. DOI: 10.1186/1755-8166-6-40.
- 55 Fluorescence in situ hybridization.** This image is a derivative work, adapted from the following source, available under : Paar C, Herber G, Voskova, et al. A case of acute myeloid leukemia (AML) with an unreported combination of chromosomal abnormalities: gain of isochromosome 5p, tetrasomy 8 and unbalanced translocation der(19)t(17;19)(q23;p13). *Mol Cytogenet.* 2013;6:40. DOI: 10.1186/1755-8166-6-40.
- 57 Genetic terms.** Café-au-lait spots. This image is a derivative work, adapted from the following source, available under : Dumitrescu CE and Collins MT. *Orphanet J Rare Dis.* 2008;3:12. DOI: 10.1186/1750-1172-3-12.
- 61 Muscular dystrophies: Image A.** Fibrofatty replacement of muscle.  Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 64 Williams syndrome.** This image is a derivative work, adapted from the following source, available under : Mazumdar J, Sarkar R, Badveli A, et al. Double chamber right ventricle in Williams syndrome: a rare cardiac anomaly reported. *Springerplus.* 2016; 5: 275. DOI: 10.1186/s40064-016-1897-y.
- 66 Vitamin A.** Bitot spots on conjunctiva. This image is a derivative work, adapted from the following source, available under : Baiyerolu A, Bowman R, Gilbert C, et al. Managing eye health in young children. *Comm Eye Health.* 2010;23(72):4-11. Available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2873666>.












- 67 **Vitamin B₃. Pellagra.** This image is a derivative work, adapted from the following source, available under : van Dijk HA, Fred H. Images of memorable cases: case 2. Connexions Web site. Dec 4, 2008. Available at: <http://cnx.org/contents/3d3db2e-8e98-496f-91c2-fe94e93428a1@3@3/>.
- 70 **Vitamin D.** X-ray of lower extremity in child with rickets. This image is a derivative work, adapted from the following source, available under : Dr. Michael L. Richardson. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 71 **Protein-energy malnutrition: Image A.** Child with kwashiorkor.  Courtesy of the Department of Health and Human Services and Dr. Lyle Conrad.
- 71 **Protein-energy malnutrition: Image B.** Child with marasmus.  Courtesy of the Department of Health and Human Services.
- 84 **Alkaptonuria.** Pigment granules on dorsum of hand. This image is a derivative work, adapted from the following source, available under : Vasudevan B, Sawhney MPS, Radhakrishnan S. Alkaptonuria associated with degenerative collagenous palmar plaques. *Indian J Dermatol.* 2009;54:299-301. DOI: 10.4103/0019-5154.55650.
- 85 **Cystinuria.** Hexagonal cystine stones in urine. This image is a derivative work, adapted from the following source, available under : Courtesy of Cayla Devine.
- 88 **Lysosomal storage diseases: Image A.** “Cherry-red” spot on macula in Tay-Sachs disease. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Jonathan Trobe.
- 88 **Lysosomal storage diseases: Image B.** Angiokeratomas. This image is a derivative work, adapted from the following source, available under : Burlina AP, Sims KB, Politei JM, et al. Early diagnosis of peripheral nervous system involvement in Fabry disease and treatment of neuropathic pain: the report of an expert panel. *BMC Neurol.* 2011;11:61. DOI: 10.1186/1471-2377-11-61. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 88 **Lysosomal storage diseases: Image C.** Gaucher cells in Gaucher disease. This image is a derivative work, adapted from the following source, available under : Sokołowska B, Skomra D, Czartoryska B, et al. Gaucher disease diagnosed after bone marrow trephine biopsy—a report of two cases. *Folia Histochem Cytobiol.* 2011;49:352-356. DOI: 10.5603/FHC.2011.0048. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 88 **Lysosomal storage diseases: Image D.** Foam cells in Niemann-Pick disease. This image is a derivative work, adapted from the following source, available under : Prieto-Potin I, Roman-Blas JA, Martinez-Calatrava MJ, et al. Hypercholesterolemia boosts joint destruction in chronic arthritis. An experimental model aggravated by foam macrophage infiltration. *Arthritis Res Ther.* 2013;15:R81. DOI: 10.1186/ar4261.

Immunology
























- 96 **Lymph node: Images A and B.** Lymph node histology. This image is a derivative work, adapted from the following source, available under : Navid Golpur.
- 98 **Spleen.** Red and white pulp. This image is a derivative work, adapted from the following source, available under : Heinrichs S, Conover LF, Bueso-Ramos CE, et al. MYBL2 is a sub-haploinsufficient tumor suppressor gene in myeloid malignancy. *eLife.* 2013;2:e00825. DOI: 10.7554/eLife.00825. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
























- 98 **Thymus: Image A.** Hassall corpuscles. This image is a derivative work, adapted from the following source, available under : Minato H, Kinoshita E, Nakada S, et al. Thymic lymphoid hyperplasia with multilocular thymic cysts diagnosed before the Sjögren syndrome diagnosis. *Diagn Pathol.* 2015;10:103. DOI: 10.1186/s13000-015-0332-y.
- 98 **Thymus: Image B.** “Sail sign” on x-ray of normal thymus in neonate. This image is a derivative work, adapted from the following source, available under : Di Serafino M, Esposito F, Severino R, et al. Think thymus, think well: the chest x-ray thymic signs. *J Pediatr Moth Care.* 2016;1(2):108-109. DOI: 10.19104/japm.2016.108.
- 107 **Complement disorders.** Paroxysmal nocturnal hemoglobinuria. This image is a derivative work, adapted from the following source, available under : Nakamura N, Sugawara T, Shirato K, et al. *J Med Case Reports.* 2011;5:550. doi: 10.1186/1752-1947-5-550
- 117 **Immunodeficiencies: Image A.** Spider angioma (telangiectasia). This image is a derivative work, adapted from the following source, available under : Liapakis IE, Englander M, Sinani R, et al. Management of facial telangiectasias with hand cautery. *World J Plast Surg.* 2015 Jul;4(2):127-133.
- 117 **Immunodeficiencies: Image B.** Giant granules in granulocytes in Chédiak-Higashi syndrome. This image is a derivative work, adapted from the following source, available under : Bharti S, Bhatia P, Bansal D, et al. The accelerated phase of Chediak-Higashi syndrome: the importance of hematological evaluation. *Turk J Haematol.* 2013;30:85-87. DOI: 10.4274/tjh.2012.0027. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.




















Microbiology

- 125 **Stains: Image A.** *Trypanosoma lewisi* on Giemsa stain.  Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.
- 125 **Stains: Image B.** Periodic acid–Schiff stain reveals *Tropheryma whipplei* infection. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman.
- 125 **Stains: Image C.** *Mycobacterium tuberculosis* on Ziehl-Neelsen stain.  Courtesy of the Department of Health and Human Services and Dr. George P. Kubica.
- 125 **Stains: Image D.** *Cryptococcus neoformans* on India ink stain.  Courtesy of the Department of Health and Human Services.
- 125 **Stains: Image E.** *Coccidioides immitis* on silver stain.  Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.
- 127 **Encapsulated bacteria.** Capsular swelling of *Streptococcus pneumoniae* using the Neufeld-Quellung test.  Courtesy of the Department of Health and Human Services.
- 128 **Catalase-positive organisms.** Oxygen bubbles released during catalase reaction. This image is a derivative work, adapted from the following source, available under : Stefano Nase. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 129 **Spore-forming bacteria.** This image is a derivative work, adapted from the following source, available under : Jones SW, Paredes CJ, Tracy B. The transcriptional program underlying the physiology of clostridial sporulation. *Genome Biol.* 2008;9:R114. DOI: 10.1186/gb-2008-9-7-r114.
- 135 **α-hemolytic bacteria.** α-hemolysis. This image is a derivative work, adapted from the following source, available under : Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .




















- 135 **β -hemolytic bacteria.** β -hemolysis. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Wikimedia Commons.
- 135 ***Staphylococcus aureus*.** [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. Richard Facklam.
- 136 ***Streptococcus pneumoniae*.** [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. Mike Miller.
- 136 ***Streptococcus pyogenes*: (group A streptococci).** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 137 ***Bacillus anthracis*.** Ulcer with black eschar. [CC-BY](#) Courtesy of the Department of Health and Human Services and James H. Steele.
- 138 **Clostridia: Image A.** Gas gangrene due to *Clostridium perfringens*. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Schröpfer E, Rauthe S, Meyer T. Diagnosis and misdiagnosis of necrotizing soft tissue infections: three case reports. *Cases J.* 2008;1:252. DOI: 10.1186/1757-1626-1-252.
- 138 **Clostridia: Image B.** Pseudomembranous enterocolitis on colonoscopy. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Klinikum Dritter Orden für die Überlassung des Bildes zur Veröffentlichung. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 139 ***Corynebacterium diphtheriae*.** Pseudomembranous pharyngitis. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 139 ***Listeria monocytogenes*.** Actin rockets. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Schuppler M, Loessner MJ. The opportunistic pathogen *Listeria monocytogenes*: pathogenicity and interaction with the mucosal immune system. *Int J Inflam.* 2010;2010:704321. DOI: 10.4061/2010/704321. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 139 ***Nocardia* vs *Actinomyces*: Image A.** *Nocardia* on acid-fast stain. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Venkataramana K. Human *Nocardia* infections: a review of pulmonary nocardiosis. *Cereus.* 2015;7(8):e304. DOI: 10.7759/cureus.304.
- 139 ***Nocardia* vs *Actinomyces*: Image B.** *Actinomyces israelii* on Gram stain. [CC-BY](#) Courtesy of the Department of Health and Human Services.
- 140 **Mycobacteria.** Acid-fast stain. [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. George P. Kubica
- 140 **Tuberculosis.** Langhans giant cell in caseating granuloma. [CC-BY](#) Courtesy of J. Hayman.
- 141 **Leprosy: Image A.** “Glove and stocking” distribution. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Courtesy of Bruno Jehle.
- 142 ***Neisseria*: Image A.** Intracellular *N gonorrhoeae*. [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. Mike Miller.
- 142 ***Haemophilus influenzae*: Image A.** Epiglottitis. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 143 ***Legionella pneumophila*.** Lung findings of unilateral and lobar infiltrate. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Robbins NM, Kumar A, Blair BM. *Legionella pneumophila* infection presenting as headache, confusion and dysarthria in a human immunodeficiency virus-1 (HIV-1) positive patient: case report. *BMC Infect Dis.* 2012;12:225. DOI: 10.1186/1471-2334-12-225.
- 143 ***Pseudomonas aeruginosa*: Image A.** Blue-green pigment on centrimide agar. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Hansen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 143 ***Pseudomonas aeruginosa*: Image B.** Ecthyma gangrenosum. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Uludokumaci S, Balkan II, Mete B, et al. Ecthyma gangrenosum-like lesions in a febrile neutropenic patient with simultaneous *Pseudomonas* sepsis and disseminated fusariosis. *Turk J Haematol.* 2013 Sep;30(3):321-4. DOI: 10.4274/Tjh.2012.0030.
- 145 ***Klebsiella*.** [CC-BY](#) Courtesy of the Department of Health and Human Services.
- 145 ***Campylobacter jejuni*.** [CC-BY](#) Courtesy of the Department of Health and Human Services.
- 146 ***Vibrio cholerae*.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Phetsouvanh R, Nakatsu M, Arakawa E, et al. Fatal bacteremia due to immotile *Vibrio cholerae* serogroup O21 in Vientiane, Laos—a case report. *Ann Clin Microbiol Antimicrob.* 2008;7:10. DOI: 10.1186/1476-0711-7-10.
- 146 ***Helicobacter pylori*.** [CC-BY](#) Courtesy of the Department of Health and Human Services, Dr. Patricia Fields, and Dr. Collette Fitzgerald.
- 146 **Spirochetes.** Appearance on darkfield microscopy. [CC-BY](#) Courtesy of the Department of Health and Human Services.
- 146 **Lyme disease: Image A.** *Ixodes* tick. [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. Michael L. Levin.
- 146 **Lyme disease: Image B.** Erythema migrans. [CC-BY](#) Courtesy of the Department of Health and Human Services and James Gathany.
- 147 **Syphilis: Image A.** Painless chancre in primary syphilis. [CC-BY](#) Courtesy of the Department of Health and Human Services and M. Rein.
- 147 **Syphilis: Image B.** Treponeme on darkfield microscopy. [CC-BY](#) Courtesy of the Department of Health and Human Services and Renelle Woodall.
- 147 **Syphilis: Image D.** Rash on palms. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Drahansky M, Dolezel M, Urbanek J, et al. Influence of skin diseases on fingerprint recognition. *J Biomed Biotechnol.* 2012;626148. DOI: 10.1155/2012/626148.
- 147 **Syphilis: Image E.** Condyloma lata. [CC-BY](#) Courtesy of the Department of Health and Human Services and Susan Lindsley.
- 147 **Syphilis: Image F.** Gumma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Chakir K, Benchikhi H. Granulome centro-facial révélant une syphilis tertiaire. *Pan Afr Med J.* 2013;15:82. DOI: 10.11604/pamj.2013.15.82.3011.
- 147 **Syphilis: Image G.** Congenital syphilis. [CC-BY](#) Courtesy of the Department of Health and Human Services and Dr. Norman Cole.
- 147 **Syphilis: Image H.** Hutchinson teeth. [CC-BY](#) Courtesy of the Department of Health and Human Services and Susan Lindsley.
- 148 ***Gardnerella vaginalis*.** [CC-BY](#) Courtesy of the Department of Health and Human Services and M. Rein.
- 150 **Rickettsial diseases and vector-borne illnesses: Image A.** Rash of Rocky Mountain spotted fever. [CC-BY](#) Courtesy of the Department of Health and Human Services.

- 150 **Rickettsial diseases and vector-borne illnesses: Image B.** *Ehrlichia morulae*. This image is a derivative work, adapted from the following source, available under : Dantas-Torres F. Canine vector-borne diseases in Brazil. *Parasit Vectors*. 2008;1:25. DOI: 10.1186/1756-3305-1-25. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 150 **Rickettsial diseases and vector-borne illnesses: Image C.** *Anaplasma phagocytophilum* in neutrophil.  Courtesy of the Department of Health and Human Services and Dumler JS, Choi K, Garcia-Garcia JC, et al. Human granulocytic anaplasmosis. *Emerg Infect Dis*. 2005. DOI 10.3201/eid1112.050898.
- 150 ***Mycoplasma pneumoniae*.** This image is a derivative work, adapted from the following source, available under : Rottem S, Kosower ND, Kornspan JD. Contamination of tissue cultures by *Mycoplasma*. In: Ceccherini-Nelli L, ed: *Biomedical tissue culture*. 2016. DOI: 10.5772/51518.
- 151 **Systemic mycoses: Image A.** *Histoplasma*.  Courtesy of the Department of Health and Human Services and Dr. D.T. McClean.
- 151 **Systemic mycoses: Image B.** *Blastomyces dermatitidis* undergoing broad-base budding.  Courtesy of the Department of Health and Human Services and Dr. Libero Ajello.
- 151 **Systemic mycoses: Image C.** Coccidiomycosis with endospores.  Courtesy of the Department of Health and Human Services.
- 151 **Systemic mycoses: Image D.** “Captain’s wheel” shape of *Paracoccidioides*.  Courtesy of the Department of Health and Human Services and Dr. Lucille K. Georg.
- 152 **Cutaneous mycoses: Image G.** Tinea versicolor. This image is a derivative work, adapted from the following source, available under : Sarah (Rosenau) Korf. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 153 **Opportunistic fungal infections: Image A.** Budding yeast of *Candida albicans*. This image is a derivative work, adapted from the following source, available under : Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 153 **Opportunistic fungal infections: Image B.** Germ tubes of *Candida albicans*. This image is a derivative work, adapted from the following source, available under : Y. Tambe. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 153 **Opportunistic fungal infections: Image C.** Oral thrush.  Courtesy of the Department of Health and Human Services and Dr. Sol Silverman, Jr.
- 153 **Opportunistic fungal infections: Image E.** Conidiophores of *Aspergillus fumigatus*.  Courtesy of the Department of Health and Human Services.
- 153 **Opportunistic fungal infections: Image F.** Aspergilloma in left lung. This image is a derivative work, adapted from the following source, available under : Souilamas R, Souilamas JI, Alkhamees K, et al. Extra corporal membrane oxygenation in general thoracic surgery: a new single veno-venous cannulation. *J Cardiothorac Surg*. 2011;6:52. DOI: 10.1186/1749-8090-6-52.
- 153 **Opportunistic fungal infections: Image G.** *Cryptococcus neoformans*.  Courtesy of the Department of Health and Human Services and Dr. Leanor Haley.
- 153 **Opportunistic fungal infections: Image H.** *Cryptococcus neoformans* on mucicarmine stain.  Courtesy of the Department of Health and Human Services and Dr. Leanor Haley.
- 153 **Opportunistic fungal infections: Image I.** *Mucor*.  Courtesy of the Department of Health and Human Services and Dr. Lucille K. Georg.
- 153 **Opportunistic fungal infections: Image J.** Mucormycosis. This image is a derivative work, adapted from the following source, available under : Jiang N, Zhao G, Yang S, et al. A retrospective analysis of eleven cases of invasive rhino-orbito-cerebral mucormycosis presented with orbital apex syndrome initially. *BMC Ophthalmol*. 2016; 16: 10. DOI: 10.1186/s12886-016-0189-1.
- 154 ***Pneumocystis jirovecii*: Image A.** Interstitial opacities in lung. This image is a derivative work, adapted from the following source, available under : Chuang C, Zhanhong X, Yinyin G, et al. Unsuspected *Pneumocystis pneumonia* in an HIV-seronegative patient with untreated lung cancer: circa case report. *J Med Case Rep*. 2007;1:15. DOI: 10.1186/1752-1947-1-115.
- 154 ***Pneumocystis jirovecii*: Image B.** CT of lung. This image is a derivative work, adapted from the following source, available under : Allen CM, Al-Jahdali HH, Irion KL, et al. Imaging lung manifestations of HIV/AIDS. *Ann Thorac Med*. 2010 Oct-Dec; 5(4): 201–216. DOI: 10.4103/1817-1737.69106.
- 154 ***Pneumocystis jirovecii*: Image C.** Disc-shaped yeast. This image is a derivative work, adapted from the following source, available under : Kirby S, Satoskar A, Brodsky S, et al. Histological spectrum of pulmonary manifestations in kidney transplant recipients on sirolimus inclusive immunosuppressive regimens. *Diagn Pathol*. 2012;7:25. DOI: 10.1186/1746-1596-7-25.
- 154 ***Sporothrix schenckii*.** Subcutaneous mycosis. This image is a derivative work, adapted from the following source, available under : Govender NP, Maphanga TG, Zulu TG, et al. An outbreak of lymphocutaneous sporotrichosis among mine-workers in South Africa. *PLoS Negl Trop Dis*. 2015 Sep; 9(9): e0004096. DOI: 10.1371/journal.pntd.0004096.
- 155 **Protozoa—GI infections: Image A.** *Giardia lamblia* trophozoite. This image is a derivative work, adapted from the following source, available under : Lipoldová M. *Giardia* and Vilém Dušan Lambl. *PLoS Negl Trop Dis*. 2014;8:e2686. DOI: 10.1371/journal.pntd.0002686.
- 155 **Protozoa—GI infections: Image B.** *Giardia lamblia* cyst. Courtesy of the Department of Health and Human Services.
- 155 **Protozoa—GI infections: Image C.** *Entamoeba histolytica* trophozoites. Courtesy of the Department of Health and Human Services.
- 155 **Protozoa—GI infections: Image D.** *Entamoeba histolytica* cyst. Courtesy of the Department of Health and Human Services.
- 155 **Protozoa—GI infections: Image E.** *Cryptosporidium* oocysts. Courtesy of the Department of Health and Human Services.
- 156 **Protozoa—CNS infections: Image A.** *Toxoplasma gondii*. This image is a derivative work, adapted from the following source, available under : Agrawal A, Bhake A, Sangole VM, et al. Multiple-ring enhancing lesions in an immunocompetent adult. *J Glob Infect Dis*. 2010 Sep-Dec;2(3):313-4. DOI: 10.4103/0974-777X.68545.
- 156 **Protozoa—CNS infections: Image B.** *Toxoplasma gondii* tachyzoite. Courtesy of the Department of Health and Human Services and Dr. L.L. Moore, Jr.
- 156 **Protozoa—CNS infections: Image C.** *Naegleria fowleri* amoebas. Courtesy of the Department of Health and Human Services.
- 156 **Protozoa—CNS infections: Image D.** *Trypanosoma brucei gambiense*. Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.























- 157 **Protozoa—hematologic infections: Image A.** *Plasmodium* trophozoite ring form.  Courtesy of the Department of Health and Human Services.
- 157 **Protozoa—hematologic infections: Image B.** *Plasmodium* schizont containing merozoites.  Courtesy of the Department of Health and Human Services and Steven Glenn.
- 157 **Protozoa—hematologic infections: Image C.** *Babesia* with ring form and with “Maltese cross” form.  Courtesy of the Department of Health and Human Services.
- 158 **Protozoa—others: Image A.** *Trypanosoma cruzi*.  Courtesy of the Department of Health and Human Services and Dr. Mae Melvin.
- 158 **Protozoa—others: Image B.** Cutaneous leishmaniasis. This image is a derivative work, adapted from the following source, available under : Sharara SL, Kanj SS. War and infectious diseases: challenges of the Syrian civil war. *PLoS Pathog*. 2014 Nov;10(11):e1004438. DOI: 10.1371/journal.ppat.1004438.
- 158 **Protozoa—others: Image C.** *Leishmania* spp.  Courtesy of the Department of Health and Human Services and Dr. Francis W. Chandler. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 158 **Protozoa—others: Image D.** *Trichomonas vaginalis*.  Courtesy of the Department of Health and Human Services.
- 159 **Nematodes (roundworms): Image A.** *Enterobius vermicularis* eggs.  Courtesy of the Department of Health and Human Services, BG Partin, and Dr. Moore.
- 159 **Nematodes (roundworms): Image B.** *Ascaris lumbricoides* egg.  Courtesy of the Department of Health and Human Services.
- 159 **Nematodes (roundworms): Image C.** *Ancylostoma* spp rash This image is a derivative work, adapted from the following source, available under : Archer M. Late presentation of cutaneous larva migrans: a case report. *Cases J*. 2009; 2: 7553. doi:10.4076/1757-1626-2-7553.
- 159 **Nematodes (roundworms): Image D.** *Trichinella spiralis* cysts in muscle. This image is a derivative work, adapted from the following source, available under : Franssen FFJ, Fonville M, Takumi K, et al. *Vet Res*. 2011; 42(1): 113. DOI: 10.1186/1297-9716-42-113.
- 159 **Nematodes (roundworms): Image E.** *Wuchereria bancrofti* Elephantiasis.  Courtesy of the Department of Health and Human Services.
- 160 **Cestodes (tapeworms): Image A.** *Taenia solium*.  Courtesy of the Department of Health and Human Services Robert J. Galindo. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 160 **Cestodes (tapeworms): Image B.** Neurocysticercosis. This image is a derivative work, adapted from the following source, available under : Coyle CM, Tanowitz HB. Diagnosis and treatment of neurocysticercosis. *Interdiscip Perspect Infect Dis*. 2009;2009:180742. DOI: 10.1155/2009/180742. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 160 **Cestodes (tapeworms): Image C.** *Echinococcus granulosus*.  Courtesy of the Department of Health and Human Services.
- 160 **Cestodes (tapeworms): Image D.** Hyatid cyst of *Echinococcus granulosus*.  Courtesy of the Department of Health and Human Services and Dr. I. Kagan.
- 160 **Cestodes (tapeworms): Image E.** *Echinococcus granulosus* cyst in liver. This image is a derivative work, adapted from the following source, available under : Ma Z, Yang W, Yao Y, et al. The adventitia resection in treatment of liver hydatid cyst: a case report of a 15-year-old boy. *Case Rep Surg*. 2014;2014:123149. DOI: 10.1155/2014/123149.
- 160 **Trematodes (flukes): Image A.** *Schistosoma mansoni* egg with lateral spine.  Courtesy of the Department of Health and Human Services.
- 160 **Trematodes (flukes): Image B.** *Schistosoma haematobium* egg with terminal spine.  Courtesy of the Department of Health and Human Services.
- 161 **Ectoparasites: Image A.** Scabies. This image is a derivative work, adapted from the following source, available under : Siegfried EC, Hebert AA. Diagnosis of atopic dermatitis: mimics, overlaps, and complications. *Clin Med*. 2015 May; 4(5): 884–917. DOI: 10.3390/jcm4050884.
- 161 **Ectoparasites: Image B.** Nit of a louse.  Courtesy of the Department of Health and Human Services and Joe Miller.
- 164 **DNA viruses: Image A.** Febrile pharyngitis. Balfour HH Jr, Dunmire SK, Hogquist KA. *Clin Transl Immunology*. 2015 Feb 27. DOI: 10.1038/cti.2015.1.
- 165 **Herpesviruses: Image A.** Keratoconjunctivitis in HSV-1 infection. This image is a derivative work, adapted from the following source, available under : Yang HK, Han YK, Wee WR, et al. Bilateral herpetic keratitis presenting with unilateral neurotrophic keratitis in pemphigus foliaceus: a case report. *J Med Case Rep*. 2011;5:328. DOI: 10.1186/1752-1947-5-328.
- 165 **Herpesviruses: Image B.** Herpes labialis. Courtesy of the Department of Health and Human Services and Dr. Herrmann.
- 165 **Herpesviruses: Image E.** Shingles (varicella-zoster virus infection). This image is a derivative work, adapted from the following source, available under : Fisle. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 165 **Herpesviruses: Image F.** Hepatosplenomegaly due to EBV infection. This image is a derivative work, adapted from the following source, available under : Gow NJ, Davidson RN, Ticehurst R, et al. Case report: no response to liposomal daunorubicin in a patient with drug-resistant HIV-associated visceral leishmaniasis. *PLoS Negl Trop Dis*. 2015 Aug; 9(8):e0003983. DOI: 10.1371/journal.pntd.0003983.
- 165 **Herpesviruses: Image G.** Atypical lymphocytes in Epstein-Barr virus infection. This image is a derivative work, adapted from the following source, available under : Coutesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 165 **Herpesviruses: Image I.** Roseola. Courtesy of Emiliano Burzagli.
- 165 **Herpesviruses: Image J.** Kaposi sarcoma. Courtesy of the Department of Health and Human Services.
- 166 **HSV identification.** Positive Tzanck smear in HSV-2 infection. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 168 **Rotavirus.** Courtesy of the Department of Health and Human Services and Erskine Palmer.
- 169 **Rubella virus.** Rubella rash. Courtesy of the Department of Health and Human Services.
- 170 **Acute laryngotracheobronchitis.** Steeple sign. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 170 **Measles (rubeola) virus: Image A.** Koplik spots. Courtesy of the Department of Health and Human Services. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.

- 170 **Measles (rubeola) virus: Image B.** Rash of measles.  Courtesy of the Department of Health and Human Services.
- 170 **Mumps virus.** Swollen neck and parotid glands.  Courtesy of the Department of Health and Human Services.
- 171 **Rabies virus: Image A.** Transmission electron micrograph.  Courtesy of the Department of Health and Human Services Dr. Fred Murphy, and Sylvia Whitfield.
- 171 **Rabies virus: Image B.** Negri bodies.  Courtesy of the Department of Health and Human Services and Dr. Daniel P. Perl.
- 171 **Ebola virus.**  Courtesy of the Department of Health and Human Services and Cynthia Goldsmith.
- 180 **Osteomyelitis.** X-ray (left) and MRI (right) views. This image is a derivative work, adapted from the following source, available under : Huang P-Y, Wu P-K, Chen C-F, et al. Osteomyelitis of the femur mimicking bone tumors: a review of 10 cases. *World J Surg Oncol.* 2013;11:283. DOI: 10.1186/1477-7819-11-283.
- 181 **Common vaginal infections: Image B.** Motile trichomonads.  Courtesy of Joe Miller.
- 181 **Common vaginal infections: Image C.** *Candida* vulvovaginitis.  Courtesy of Mikael Häggström.
- 182 **TORCH infections: Image A.** “Blueberry muffin” rash. This image is a derivative work, adapted from the following source, available under : Benmiloud S, Elhaddou G, Belghiti ZA, et al. Blueberry muffin syndrome. *Pan Afr Med J.* 2012;13:23.
- 182 **TORCH infections: Image B.** Cataract in infant with congenital rubella.  Courtesy of the Department of Health and Human Services .
- 182 **TORCH infections: Image C.** Periventricular calcifications in congenital cytomegalovirus infection. This image is a derivative work, adapted from the following source, available under : Bonthius D, Perlman S. Congenital viral infections of the brain: lessons learned from lymphocytic choriomeningitis virus in the neonatal rat. *PLoS Pathog.* 2007;3:e149. DOI: 10.1371/journal.ppat.0030149. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 183 **Red rashes of childhood: Image C.** Child with scarlet fever. This image is a derivative work, adapted from the following source, available under : www.badobadop.co.uk.
- 183 **Red rashes of childhood: Image D.** Chicken pox.  Courtesy of the Department of Health and Human Services.
- 184 **Sexually transmitted infections: Image A.** Chancroid.  Courtesy of the Department of Health and Human Services and Susan Lindsley.
- 184 **Sexually transmitted infections: Image B.** Donovanosis.  Courtesy of the Department of Health and Human Services and Dr. Pinozzi.
- 185 **Pelvic inflammatory disease: Image A.** Purulent cervical discharge. This image is a derivative work, adapted from the following source, available under : SOS-AIDS Amsterdam The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 185 **Pelvic inflammatory disease: Image B.** Adhesions in Fitz-Hugh–Curtis syndrome.  Courtesy of Hic et nunc.
- 190 **Vancomycin.** Red man syndrome. This image is a derivative work, adapted from the following source, available under : O’Meara P, Borici-Mazi R, Morton R, et al. DRESS with delayed onset acute interstitial nephritis and profound refractory eosinophilia secondary to vancomycin. *Allergy Asthma Clin Immunol.* 2011;7:16. DOI: 10.1186/1710-1492-7-16.









Pathology

- 209 **Necrosis: Image A.** Coagulative necrosis.  Courtesy of the Department of Health and Human Services and Dr. Steven Rosenberg.
- 209 **Necrosis: Image B.** Liquefactive necrosis.  Courtesy of Daftblogger.
- 209 **Necrosis: Image C.** Caseous necrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 209 **Necrosis: Image D.** Fat necrosis. This image is a derivative work, adapted from the following source, available under : Patho. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 209 **Necrosis: Image E.** Fibrinoid necrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 209 **Necrosis: Image F.** Acral gangrene.  Courtesy of the Department of Health and Human Services and William Archibald.
- 210 **Ischemia.** This image is a derivative work, adapted from the following source, available under : Van Assche LM, Kim HW, Jensen CJ, et al. A new CMR protocol for non-destructive, high resolution, ex-vivo assessment of the area at risk simultaneous with infarction: validation with histopathology. *J Cardiovasc Magn Reson.* 2012; 14(Suppl 1): O7. DOI: 10.1186/1532-429X-14-S1-O7.
- 210 **Types of infarcts: Image B.** Pale infarct.  Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 211 **Types of calcification: Image A.** Dystrophic calcification. This image is a derivative work, adapted from the following source, available under : Chun J-S, Hong R, Kim J-A. Osseous metaplasia with mature bone formation of the thyroid gland: three case reports. *Oncol Lett.* 2013;6:977-979. DOI: 10.3892/ol.2013.1475. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 211 **Lipofuscin.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 212 **Amyloidosis: Image A.** Amyloid deposits on Congo red stain. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.
- 212 **Amyloidosis: Image B.** Apple green birefringence under polarized light. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.
- 214 **Acute inflammation: Image A.** Pericardium with severe inflammation, neutrophilic infiltration and fibrin with entrapped clusters of bacteria. This image is a derivative work, adapted from the following source, available under : Faida Ajili, et al. Coexistence of pyoderma gangrenosum and sweet’s syndrome in a patient with ulcerative colitis. *Pan Afr Med J.* 2015 Jun 24. DOI: 10.11604/pamj.2015.21.151.6364.
- 217 **Granulomatous diseases.** Granuloma.  Courtesy of Sanjay Mukhopadhyay.
- 218 **Scar formation: Image A.** Hypertrophic scar. This image is a derivative work, adapted from the following source, available under : Baker R, Urso-Baiarda F, Linge C, et al. Cutaneous scarring: a clinical review. *Dermatol Res Pract.* 2009;2009:625376. DOI: 10.1155/2009/625376.




- 218 Scar formation: Image B.** Keloid scar. This image is a derivative work, adapted from the following source, available under : Dr. Andreas Settje. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 219 Neoplasia and neoplastic progression: Image A.** Cervical tissue. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 223 Common metastases: Image A.** Brain metastases from breast cancer. This image is a derivative work, adapted from the following source, available under : Jmarchn. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 223 Common metastases: Image B.** Brain metastasis. Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 223 Common metastases: Image C.** Liver metastasis. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 223 Common metastases: Image D.** Liver metastasis. Courtesy of J. Hayman.
- 223 Common metastases: Image E.** Bone metastasis. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff.
- 223 Common metastases: Image F.** Bone metastasis. This image is a derivative work, adapted from the following source, available under : Courtesy of M Emmanuel.
- 227 Psammoma bodies.** Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- Cardiovascular**
- 283 Anatomy of the heart: Image A.** MRI showing normal cardiac anatomy. This image is a derivative work, adapted from the following source, available under : Zhang J, Chen L, Wang X, et al. Compounding local invariant features and global deformable geometry for medical image registration. *PLoS One*. 2014;9(8):e105815. DOI: 10.1371/journal.pone.0105815.
- 298 Congenital heart diseases: Image A.** Tetralogy of Fallot. This image is a derivative work, adapted from the following source, available under : Rashid AKM: Heart diseases in Down syndrome. In: Dey S, ed: Down syndrome. DOI: 10.5772/46009. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 299 Congenital heart diseases: Image B.** Ventricular septal defect. This image is a derivative work, adapted from the following source, available under : Bardo DME, Brown P. Cardiac multidetector computed tomography: basic physics of image acquisition and clinical applications. *Curr Cardiol Rev*. 2008 Aug;4(3):231–243. DOI: 10.2174/157340308785160615.
- 299 Congenital heart diseases: Image C.** Atrial septal defect. This image is a derivative work, adapted from the following source, available under : Teo KSL, Dundon BK, Molae P, et al. Percutaneous closure of atrial septal defects leads to normalisation of atrial and ventricular volumes. *J Cardiovasc Magn Reson*. 2008;10(1):55. DOI: 10.1186/1532-429X-10-55.
- 299 Congenital heart diseases: Image D.** Patent ductus arteriosus. This image is a derivative work, adapted from the following source, available under : Henjes CR, Nolte I, Wesfaedt P. Multidetector-row computed tomography of thoracic aortic anomalies in dogs and cats: patent ductus arteriosus and vascular rings. *BMC Vet Res*. 2011;7:57. DOI: 10.1186/1746-6148-7-57.
- 299 Congenital heart diseases: Image E.** Clubbing of fingers. Courtesy of Ann McGrath.
- 299 Congenital heart diseases: Image F.** MRI showing coarctation of the aorta. This image is a derivative work, adapted from the following source, available under : Vergales JE, Gangemi JJ, Rhueban KS, Lim DS. Coarctation of the aorta — the current state of surgical and transcatheter therapies. *Curr Cardiol Rev*. 2013 Aug; 9(3): 211–219. DOI: 10.2174/1573403X113099990032
- 300 Hypertension: Image A.** “String of beads” appearance in fibromuscular dysplasia. This image is a derivative work, adapted from the following source, available under : Plouin PF, Perdu J, LaBatide-Alanore A, et al. Fibromuscular dysplasia. *Orphanet J Rare Dis*. 2007;7:28. DOI: 10.1186/1750-1172-2-28. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 301 Hyperlipidemia signs: Image C.** Tendinous xanthoma. This image is a derivative work, adapted from the following source, available under : Raffa W, Hassam B. Xanthomes tendineux et tubéreux révélant une hypercholestérolémie familiale. *Pan Afr Med J*. 2013; 15: 49. DOI: 10.11604/pamj.2013.15.49.2636.
- 301 Arteriosclerosis: Image A.** Hyaline type. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 301 Arteriosclerosis: Image B.** Hyperplastic type. This image is a derivative work, adapted from the following source, available under : Paco Larosa. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 301 Arteriosclerosis: Image C.** Monckeberg sclerosis (medial calcific sclerosis). This image is a derivative work, adapted from the following source, available under : Couri CE, da Silva GA, Martinez JA, et al. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 303 Aortic dissection.** This image is a derivative work, adapted from the following source, available under : Qi Y, Ma X, Li G, et al. Three-dimensional visualization and imaging of the entry tear and intimal flap of aortic dissection using CT virtual intravascular endoscopy. *PLoS One*. 2016; 11(10): e0164750. DOI: 10.1371/journal.pone.0164750.
- 307 Myocardial infarction complications: Image A.** Papillary muscle rupture. This image is a derivative work, adapted from the following source, available under : Routy B, Huynh T, Fraser R, et al. Vascular endothelial cell function in catastrophic antiphospholipid syndrome: a case report and review of the literature. *Case Rep Hematol*. 2013;2013:710365. DOI: 10.1155/2013/710365.
- 307 Myocardial infarction complications: Image B.** Drawing of pseudoaneurysm. This image is a derivative work, adapted from the following source, available under : Patrick J. Lynch and Dr. C. Carl Jaffe.
- 307 Myocardial infarction complications: Image C.** Free wall rupture of left ventricle. This image is a derivative work, adapted from the following source, available under : Zacarias ML, da Trindade H, Tsutsu J, et al. Left ventricular free wall impeding rupture in post-myocardial infarction period diagnosed by myocardial contrast echocardiography: case report. *Cardiovasc Ultrasound*. 2006;4:7. DOI: 10.1186/1476-7120-4-7.
- 308 Cardiomyopathies: Image A.** Dilated cardiomyopathy. This image is a derivative work, adapted from the following source, available under

- : Cho JMIH, van Es R, Stathonikos N, et al. High resolution systematic digital histological quantification of cardiac fibrosis and adipose tissue in phospholamban p.Arg14del mutation associated cardiomyopathy. *PLoS One*. 2014;9:e94820. DOI: 10.1371/journal.pone.0094820.
- 308 Cardiomyopathies: Image B.** Hypertrophic obstructive cardiomyopathy. This image is a derivative work, adapted from the following source, available under : Benetti MA, Belo Nunes RA, Benvenuti LA. Case 2/2016 - 76-year-old male with hypertensive heart disease, renal tumor and shock. *Arq Bras Cardiol*. 2016 May; 106(5): 439–446. DOI: 10.5935/abc.20160067.
- 309 Heart failure.** Pedal edema. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 310 Cardiac tamponade: Image A.** This image is a derivative work, adapted from the following source, available under : Yousuf T, Kramer J, Kopiec A, et al. A rare case of cardiac tamponade induced by chronic rheumatoid arthritis. *J Clin Med Res*. 2015 Sep;7(9):720–723. DOI: 10.14740/jocmr.2226w.
- 310 Cardiac tamponade: Image B.** This image is a derivative work, adapted from the following source, available under : Maharaj SS, Chang SM. Cardiac tamponade as the initial presentation of systemic lupus erythematosus: a case report and review of the literature. *Pediatr Rheumatol Online J*. 2015; 13: 9. DOI: 10.1186/s12969-015-0005-0.
- 311 Bacterial endocarditis: Image A.**  Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.
- 311 Bacterial endocarditis: Image C.** Osler nodes. This image is a derivative work, adapted from the following source, available under : Yang ML, Chen YH, Lin WR, et al. Case report: infective endocarditis caused by *Brevundimonas vesicularis*. *BMC Infect Dis*. 2006;6:179. DOI: 10.1186/1471-2334-6-179.
- 311 Bacterial endocarditis: Image D.** Janeway lesions on sole. This image is a derivative work, adapted from the following source, available under : Courtesy of DeNanneke.
- 312 Rheumatic fever.** Aschoff body and Anitschkow cells. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 313 Acute pericarditis.** This image is a derivative work, adapted from the following source, available under : Bogaert J, Francone M. Cardiovascular magnetic resonance in pericardial diseases. *J Cardiovasc Magn Reson*. 2009;11:14. DOI: 10.1186/1532-429X-11-14. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 315 Vasculitides: Image A.** Temporal arteritis histology. This image is a derivative work, adapted from the following source, available under : Marvin. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 315 Vasculitides: Image B.** Angiogram in patient with Takayasu arteritis.  Courtesy of the Department of Health and Human Services and Justin Ly.
- 315 Vasculitides: Image C.** Gangrene as a consequence of Buerger disease. This image is a derivative work, adapted from the following source, available under : Afsjarfard A, Mozaffar M, Malekpour F, et al. The wound healing effects of iloprost in patients with Buerger's disease: claudication and prevention of major amputations. *Iran Red Crescent Med J*. 2011;13:420-423.
- 315 Vasculitides: Image D.** Strawberry tongue in patient with Kawasaki disease. This image is a derivative work, adapted from the following source, available under : Courtesy of Natr.
- 315 Vasculitides: Image E.** Coronary artery aneurysm in Kawasaki disease. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 315 Vasculitides.** Polyarteritis nodosa. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 315 Vasculitides: Image G.** Churg-Strauss syndrome histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 315 Vasculitides: Image H.** Granulomatosis with polyangiitis (formerly Wegener) and PR3-ANCA/c-ANCA.  Courtesy of M.A. Little.
- 315 Vasculitides: Image I.** Henoch-Schönlein purpura.  Courtesy of Okwikikim.
- 315 Vasculitides: Image J.** MPO-ANCA/p-ANCA in microscopic polyangiitis.  Courtesy of and M.A. Little.

































Endocrine



























- 326 Thyroid development.** Thyroglossal duct cyst. This image is a derivative work, adapted from the following source, available under : Adelchi C, Mara P, Melissa L, et al. Ectopic thyroid tissue in the head and neck: a case series. *BMC Res Notes*. 2014;7:790. DOI: 10.1186/1756-0500-7-790.
- 340 Hypothyroidism vs hyperthyroidism.** Onycholysis. This image is a derivative work, adapted from the following source, available under : Rajebi MR, Shahrokni A, Chaisson M. Uncommon osseous involvement in multisystemic sarcoidosis. *Ann Saudi Med*. 2009 Nov-Dec;29(6):485–486.
- 341 Hypothyroidism: Image B.** Before and after treatment of congenital hypothyroidism.  Courtesy of the Department of Health and Human Services.
- 341 Hypothyroidism: Image C.** Congenital hypothyroidism. This image is a derivative work, adapted from the following source, available under : Sadasiv Swain. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 342 Thyroid adenoma: Image A.** This image is a derivative work, adapted from the following source, available under : Terada T. Brain metastasis from thyroid adenomatous nodules or an encapsulated thyroid follicular tumor without capsular and vascular invasion: a case report. *Cases J*. 2009; 2: 7180. DOI: 10.4076/1757-1626-2-7180.
- 344 Hypoparathyroidism.** Shortened 4th and 5th digits. This image is a derivative work, adapted from the following source, available under : Ferrario C, Gastaldi G, Portmann L, et al. Bariatric surgery in an obese patient with Albright hereditary osteodystrophy: a case report. *J Med Case Rep*. 2013; 7: 111. DOI: 10.1186/1752-1947-7-111.
- 345 Hyperparathyroidism.** Multiple lytic lesions. This image is a derivative work, adapted from the following source, available under : Khaoula BA, Kaouther BA, Ines C, et al. An unusual presentation of primary hyperparathyroidism: pathological fracture. *Case Rep Orthop*. 2011;2011:521578. DOI: 10.1155/2011/521578. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 349 Adrenal insufficiency: Image A.** Mucosal hyperpigmentation in primary adrenal insufficiency.  Courtesy of FlatOut. The image

may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.

- 350 **Pheochromocytoma.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Feldman.
- 351 **Multiple endocrine neoplasias.** Mucosal neuroma. This image is a derivative work, adapted from the following source, available under : Martucciello G, Lerone M, Bricco L, et al. Multiple endocrine neoplasias type 2B and RET proto-oncogene. *Ital J Pediatr.* 2012;38:9. DOI: 10.1186/1824-7288-38-9.
- 352 **Carcinoid syndrome.**  Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.

Gastrointestinal






























- 358 **Ventral wall defects.** Gastroschisis. This image is a derivative work, adapted from the following source, available under : Zvadic Z. Gastroschisis with concomitant jejuno-ileal atresia complicated by jejunal perforation. *J Neonatal Surg.* 2016 Apr-Jun; 5(2): 25.
- 358 **Ventral wall defects.** Omphalocele. This image is a derivative work, adapted from the following source, available under : Khan YA, Qureshi MA, Akhtar J. Omphalomesenteric duct cyst in an omphalocele: a rare association. *Pak J Med Sci.* 2013 May-Jun; 29(3): 866–868.
- 358 **Ventral wall defects.** Drawings of gastroschisis (left) and omphalocele (right).  Courtesy of the Department of Health and Human Services.
- 359 **Intestinal atresia.** This image is a derivative work, adapted from the following source, available under : Saha M. Alimentary tract atresias associated with anorectal malformations: 10 years' experience. *J Neonatal Surg.* 2016 Oct-Dec; 5(4): 43. DOI: 10.21699/jns.v5i4.449.
- 359 **Hypertrophic pyloric stenosis.** This image is a derivative work, adapted from the following source, available under : Hassan RAA, Choo YU, Noraida R, et al. Infantile hypertrophic pyloric stenosis in postoperative esophageal atresia with tracheoesophageal fistula. *J Neonatal Surg.* 2015 Jul-Sep;4(3):32.
- 360 **Pancreas and spleen embryology.** Annular pancreas. This image is a derivative work, adapted from the following source, available under : Mahdi B, Selim S, Hassen T, et al. A rare cause of proximal intestinal obstruction in adults—annular pancreas: a case report. *Pan Afr Med J.* 2011;10:56. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 360 **Retroperitoneal structures.** This image is a derivative work, adapted from the following source, available under : Sammut J, Ahiaku E, Williams DT. Complete regression of renal tumour following ligation of an accessory renal artery during repair of an abdominal aortic aneurysm. *Ann R Coll Surg Engl.* 2012 Sep; 94(6): e198–e200. DOI: 10.1308/003588412X13373405384972.
- 362 **Digestive tract anatomy.** Histology of stomach wall. This image is a derivative work, adapted from the following source, available under : Alexander Klepnev.
- 362 **Digestive tract histology: Image A.**  Courtesy of Dr. Michale Bonert.
- 362 **Digestive tract histology: Image B.**  Courtesy of W. Ben Smith.
- 362 **Digestive tract histology: Images C, D, E.** This image is a derivative work, adapted from the following source, available under : Wikimedia Commons.
- 367 **Liver tissue architecture: Image A.** Portal triad. This image is a derivative work, adapted from the following source, available under : Liver development. In: Zorn AM. Stem book. Cambridge: Harvard Stem Cell Institute, 2008.
- 367 **Liver tissue architecture: Image B.** Kupffer cells. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 368 **Biliary structures.** Gallstones. This image is a derivative work, adapted from the following source, available under : J. Guntau. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 370 **Hernias: Image A.** Congenital diaphragmatic hernia. This image is a derivative work, adapted from the following source, available under : Tovar J. Congenital diaphragmatic hernia. *Orphanet J Rare Dis.* 2012;7:1. DOI: 10.1186/1750-1172-7-1.
- 372 **Gastrointestinal secretory products.** Histology of gastric pit. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 374 **Peyer patches.** This image is a derivative work, adapted from the following source, available under : Plainpaper. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 376 **Sialolithiasis.** This image is a derivative work, adapted from the following source, available under : Pastor-Ramos V, Cuervo-Diaz A, Aracil-Kessler L. Sialolithiasis. Proposal for a new minimally invasive procedure: piezoelectric surgery. *J Clin Exp Dent.* 2014 Jul;6(3):e295–e298. DOI: 10.4317/jced.51253.
- 376 **Salivary gland tumors.** Pleomorphic adenoma histology. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 376 **Achalasia.** This image is a derivative work, adapted from the following source, available under : Farnooosh Farrokhi and Michael F. Vaezi. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 377 **Esophageal pathologies: Image A.** Pneumomediastinum in Boerhaave syndrome. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 377 **Esophageal pathologies: Image B.** Esophageal varices on endoscopy. This image is a derivative work, adapted from the following source, available under : Costaguta A, Alvarez F. Etiology and management of hemorrhagic complications of portal hypertension in children. *Int J Hepatol.* 2012;2012:879163. DOI: 10.1155/2012/879163.
- 377 **Esophageal pathologies: Image C.** Esophageal varices on CT. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 377 **Esophageal pathologies: Image D.** Esophagitis. This image is a derivative work, adapted from the following source, available under : Takahashi Y, Nagata N, Shimbo T. Long-term trends in esophageal candidiasis prevalence and associated risk factors with or without HIV infection: lessons from an endoscopic study of 80,219 patients. *PLoS One.* 2015; 10(7): e0133589. DOI: 10.1371/journal.pone.0133589.
- 378 **Barrett esophagus: Image A.** Endoscopy. This image is a derivative work, adapted from the following source, available under : Coda S, Thillainayagam AV. State of the art in advanced endoscopic


- imaging for the detection and evaluation of dysplasia and early cancer of the gastrointestinal tract. *Clin Exp Gastroenterol*. 2014;7:133-150. DOI: 10.2147/CEG.S58157. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 378 Barrett esophagus: Image B.** Goblet cells. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 379 Ménériere disease.** This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 379 Gastric cancer Tan Y, Fu J, Li X.** This image is a derivative work, adapted from the following source, available under : A minor (<50%) signet-ring cell component associated with poor prognosis in colorectal cancer patients: a 26-year retrospective study in China. *PLoS One*. 2015; 10(3): e0121944. DOI: 10.1371/journal.pone.0121944.
- 380 Ulcer complications.** Free air under diaphragm in perforated ulcer. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 381 Malabsorption syndromes: Image A.** This image is a derivative work, adapted from the following source, available under : Celiac disease. Sedda S, Caruso R, Marafini I, et al. Pyoderma gangrenosum in refractory celiac disease: a case report. *BMC Gastroenterol*. 2013; 13: 162. DOI: 10.1186/1471-230X-13-162.
- 381 Malabsorption syndromes: Image B.** *Tropheryma whippeli* on PAS stain. This image is a derivative work, adapted from the following source, available under : Tran HA. Reversible hypothyroidism and Whipple's disease. *BMC Endocr Disord*. 2006;6:3. DOI: 10.1186/1472-6823-6-3.
- 382 Inflammatory bowel diseases: Image A.** "String sign" on barium swallow in Crohn disease. This image is a derivative work, adapted from the following source, available under : Al-Mofarreh MA, Al Mofleh IA, Al-Teimi IN, et al. Crohn's disease in a Saudi outpatient population: is it still rare? *Saudi J Gastroenterol*. 2009;15:111-116. DOI: 10.4103/1319-3767.45357. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 382 Inflammatory bowel diseases: Images B (normal mucosa) and C (punched-out ulcers) in ulcerative colitis.** This image is a derivative work, adapted from the following source, available under : Ishikawa D, Ando T, Watanabe O, et al. Images of colonic real-time tissue sonoelastography correlate with those of colonoscopy and may predict response to therapy in patients with ulcerative colitis. *BMC Gastroenterol*. 2011;11:29. DOI: 10.1186/1471-230X-11-29.
- 383 Appendicitis.** Fecalith. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 383 Diverticula of the GI tract: Image B.** Diverticulosis. This image is a derivative work, adapted from the following source, available under : Sartelli M, Moore FA, Ansaloni L, et al. A proposal for a CT driven classification of left colon acute diverticulitis. *World J Emerg Surg*. 2015;10:3. DOI: 10.1186/1749-7922-10-3.
- 383 Diverticula of the GI tract: Image C.** This image is a derivative work, adapted from the following source, available under : Hupfeld L, Burcharth J, Pommergaard HC, Rosenberg J. The best choice of treatment for acute colonic diverticulitis with purulent peritonitis is uncertain. *Biomed Res Int*. 2014; 2014: 380607. DOI: 10.1155/2014/380607.
- 384 Zenker diverticulum.** This image is a derivative work, adapted from the following source, available under : Courtesy of Bernd Brägelmann.
- 385 Maltotation.** This image is a derivative work, adapted from the following source, available under : Mathews R, Thenabadu S, Jaiganesh T. Abdominal pain with a twist. *Int J Emerg Med*. 2011;4:21. DOI: 10.1186/1865-1380-4-21.
- 385 Intussusception: Image A.** Intraoperative image of intussusception. This image is a derivative work, adapted from the following source, available under : Vasiliadis K, Kogopoulos E, Katsamakas M, et al. Ileocecal intussusception induced by a gastrointestinal stromal tumor. *World J Surg Oncol*. 2008;6:133. DOI: 10.1186/1477-7819-6-133.
- 385 Intussusception: Image B.** Ultrasound showing target sign. This image is a derivative work, adapted from the following source, available under : Abbo O, Pinnagoda K, Micol LA. Osteosarcoma metastasis causing ileo-ileal intussusception. *World J Surg Oncol*. 2013 Aug 12;11(1):188. DOI: 10.1186/1477-7819-11-188.
- 386 Volvulus.** Coffee bean sign. This image is a derivative work, adapted from the following source, available under : Yigit M, Turkdogan KA. Coffee bean sign, whirl sign and bird's beak sign in the diagnosis of sigmoid volvulus. *Pan Afr Med J*. 2014;19:56. DOI: 10.11604/pamj.2014.19.56.5142.
- 386 Other intestinal disorders: Image A.** Necrosis due to occlusion of SMA. This image is a derivative work, adapted from the following source, available under : Van De Winkel N, Cheragwandi A, Nieboer K, et al. Superior mesenteric arterial branch occlusion causing partial jejunal ischemia: a case report. *J Med Case Rep*. 2012;6:48. DOI: 10.1186/1752-1947-6-48.
- 386 Other intestinal disorders: Image B.** Loops of dilated bowel suggestive of small bowel obstruction. This image is a derivative work, adapted from the following source, available under : Welte FJ, Crosso M. Left-sided appendicitis in a patient with congenital gastrointestinal malrotation: a case report. *J Med Case Rep*. 2007;1:92. DOI: 10.1186/1752-1947-1-92.
- 386 Other intestinal disorders: Image C.** Endoscopy showing dilated vessels. This image is a derivative work, adapted from the following source, available under : Gunjan D, Sharma V, Rana SS, et al. Small bowel bleeding: a comprehensive review. *Gastroenterol Rep*. 2014 Nov;2(4):262-75. DOI: 10.1093/gastro/gou025.
- 386 Other intestinal disorders: Image D.** Pneumatosis intestinalis. This image is a derivative work, adapted from the following source, available under : Pelizzo G, Nakib G, Goruppi I, et al. Isolated colon ischemia with norovirus infection in preterm babies: a case series. *J Med Case Rep*. 2013;7:108. DOI: 10.1186/1752-1947-7-108.
- 387 Colonic polyps: Image A.** This image is a derivative work, adapted from the following source, available under : M. Emmanuel.
- 387 Colonic polyps: Image B.** Adenomatous polyps. This image is a derivative work, adapted from the following source, available under : Shussman N, Wexner SD. Colorectal polyps and polyposis syndromes. *Gastroenterol Rep (Oxf)*. 2014 Feb;2(1):1-15. DOI: 10.1093/gastro/got041.
- 387 Colonic polyps: Image C.** This image is a derivative work, adapted from the following source, available under : Rehani B, Chasen RM, Dowdy Y, et al. Advanced adenoma diagnosis with FDG PET in a visibly normal mucosa: a case report. *J Med Case Reports*. 2007; 1: 99. DOI: 10.1186/1752-1947-1-99.
- 388 Colorectal cancer: Image A.** Polyp. This image is a derivative work, adapted from the following source, available under : Takiyama A, Nozawa H, Ishihara S, et al. Secondary metastasis in the lymph node of the bowel invaded by colon cancer: a report of three cases. *World J Surg Oncol*. 2016; 14: 273. DOI: 10.1186/s12957-016-1026-y.
- 389 Cirrhosis and portal hypertension: Image A.** Splenomegaly and liver nodularity in cirrhosis. This image is a derivative work, adapted from

- the following source, available under : Inversitus. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 389 Cirrhosis and portal hypertension: Image B.** This image is a derivative work, adapted from the following source, available under : Blackburn PR, Hickey RD, Nace RA, et al. Silent tyrosinemia type I without elevated tyrosine or succinylacetone associated with liver cirrhosis and hepatocellular carcinoma. *Hum Mutat.* 2016 Oct; 37(10): 1097–1105. DOI: 10.1002/humu.23047.
- 391 Alcoholic liver disease: Image B.** Mallory bodies. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 391 Alcoholic liver disease: Image C.** Sclerosis in alcoholic cirrhosis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 391 Non-alcoholic fatty liver disease.** This image is a derivative work, adapted from the following source, available under : El-Karakasy HM, El-Koofy NM, Anwar GM, et al. Predictors of non-alcoholic fatty liver disease in obese and overweight Egyptian children: single center study. *Saudi J Gastroenterol.* 2011;17:40-46. DOI: 10.4103/1319-3767.74476.
- 392 Hepatocellular carcinoma/hepatoma: Image A.** Gross specimen. Reproduced, with permission, from Jean-Christophe Fournet and Humpath.
- 392 Other liver tumors.** Cavernous liver hemangioma. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 392 α_1 -antitrypsin deficiency.** Liver histology. This image is a derivative work, adapted from the following source, available under : Dr. Jerad M. Gardner. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 393 Jaundice.** Yellow sclera. Courtesy of the Department of Health and Human Services and Dr. Thomas F. Sellers.
- 395 Wilson disease.** This image is a derivative work, adapted from the following source, available under : Kodama H, Fujisawa C, Bhadhrasit W. Inherited copper transport disorders: biochemical mechanisms, diagnosis, and treatment. *Curr Drug Metab.* 2012 Mar; 13(3): 237–250. DOI: 10.2174/138920012799320455.
- 395 Hemochromatosis.** Hemosiderin deposits. This image is a derivative work, adapted from the following source, available under : Mathew J, Leong MY, Morley N, et al. A liver fibrosis cocktail? Psoriasis, methotrexate and genetic hemochromatosis. *BMC Dermatol.* 2005;5:12. DOI: 10.1186/1471-5945-5-12.
- 396 Cholelithiasis and related pathologies: Image A.** Gross specimen of gallstones. This image is a derivative work, adapted from the following source, available under : Courtesy of M. Emmanuel.
- 396 Cholelithiasis and related pathologies: Image B.** Large gallstone. This image is a derivative work, adapted from the following source, available under : Spangler R, Van Pham T, Khoujah D, et al. Abdominal emergencies in the geriatric patient. *Int J Emerg Med.* 2014; 7: 43. DOI: 10.1186/s12245-014-0043-2.
- 397 Cholelithiasis and related pathologies: Image C.** Porcelain gallbladder. This image is a derivative work, adapted from the following source, available under : Fred H, van Dijk H. Images of memorable cases: case 19. Connexions Web site. December 4, 2008. Available at: <http://cnx.org/content/m14939/1.3/>. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 397 Acute pancreatitis: Image A.** Acute exudative pancreatitis. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 397 Acute pancreatitis: Image B.** Pancreatic pseudocyst. This image is a derivative work, adapted from the following source, available under : Thomas Zimmerman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 397 Chronic pancreatitis.** This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 398 Pancreatic adenocarcinoma: Image A.** Histology. This image is a derivative work, adapted from the following source, available under : KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 398 Pancreatic adenocarcinoma: Image B.** CT scan. Courtesy of MBq. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.










Hematology and Oncology




























- 406 Neutrophils.** Courtesy of B. Lennert.
- 407 Erythrocytes.** Courtesy of the Department of Health and Human Services and Drs. Noguchi, Rodgers, and Schechter.
- 407 Thrombocytes (platelets).** This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 407 Monocytes.** This image is a derivative work, adapted from the following source, available under : Dr. Graham Beards. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 407 Macrophages.** This image is a derivative work, adapted from the following source, available under : De Tommasi AS, Otranto D, Furlanello T, et al. Evaluation of blood and bone marrow in selected canine vector-borne diseases. *Parasit Vectors.* 2014;7:534. DOI: 10.1186/s13071-014-0534-2.
- 408 Eosinophils.** This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman.
- 408 Basophils.** This image is a derivative work, adapted from the following source, available under : Dr. Erhabor Osaro. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 408 Mast cells.** Courtesy of Wikimedia Commons.
- 408 Dendritic cells.** This image is a derivative work, adapted from the following source, available under : Cheng J-H, Lee S-Y, Lien Y-Y, et al. Immunomodulating activity of *Nymphaea rubra* roxb. extracts: activation of rat dendritic cells and improvement of the TH1 immune response. *Int J Mol Sci.* 2012;13:10722-10735. DOI: 10.3390/ijms130910722.
- 409 Lymphocytes.** This image is a derivative work, adapted from the following source, available under : Fickleandfreckled.
- 409 Plasma cells.** Courtesy of the Department of Health and Human Services and Dr. Francis W. Chandler. The image may have been

- modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 415 **RBC morphology: Image J.** Sickle cell.  Courtesy of the Department of Health and Human Services and the Sickle Cell Foundation of Georgia, Jackie George, and Beverly Sinclair.
- 416 **RBC inclusions: Image A.** Ringed sideroblast. This image is a derivative work, adapted from the following source, available under . Paulo Henrique Orlandi Mourao. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 416 **RBC inclusions: Image B.** Howell-Jolly bodies. This image is a derivative work, adapted from the following source, available under . Serio B, Pezzullo L, Giudice V, et al. OPSI threat in hematological patients. *Transl Med UniSa*. 2013 May-Aug;6:2-10.
- 416 **RBC inclusions: Image C.** Bsaophilic stippling. This image is a derivative work, adapted from the following source, available under . Dr. Erhabor Osaro.
- 416 **RBC inclusions: Image D.** Pappenheimer bodies. This image is a derivative work, adapted from the following source, available under . Paulo Henrique Orlandi Mourao. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 419 **Microcytic, hypochromic anemias: Image A.** This image is a derivative work, adapted from the following source, available under . Bock F, Borucki K, Vorwerk P, et al. A two-and-a-half-year-old breastfed toddler presenting with anemia: a case report. *BMC Res Notes*. 2014; 7: 917. DOI: 10.1186/1756-0500-7-917.
- 419 **Microcytic, hypochromic anemia: Image D.** Lead lines in lead poisoning. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 419 **Microcytic, hypochromic anemia: Image E.** Sideroblastic anemia. This image is a derivative work, adapted from the following source, available under . Paulo Henrique Orlandi Moura. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 420 **Macrocytic anemias.** Megaloblastic anemia. This image is a derivative work, adapted from the following source, available under . Courtesy of Dr. Ed Uthman.
- 422 **Intrinsic hemolytic anemias.** This image is a derivative work, adapted from the following source, available under . El Ariss AB, Younes M, Matar J. Prevalence of sickle cell trait in the southern suburb of Beirut, Lebanon. *Mediterr J Hematol Infect Dis*. 2016; 8(1): e2016015. DOI: 10.4084/MJHID.2016.015.
- 425 **Heme synthesis, porphyrias, and lead poisoning: Image A.** Basophilic stippling in lead poisoning. This image is a derivative work, adapted from the following source, available under . van Dijk HA, Fred HL. Images of memorable cases: case 81. Connexions Web site. December 3, 2008. Available at <http://cnx.org/contents/3196b3e-1e1e-4c4d-a1ac-d4fc9ab65443@4@4>.
- 425 **Heme synthesis, porphyrias, and lead poisoning: Image B.** Porphyria cutanea tarda. This image is a derivative work, adapted from the following source, available under . Bovenschen HJ, Vissers WHPM. Primary hemochromatosis presented by porphyria cutanea tarda: a case report. *Cases J*. 2009;2:7246. DOI: 10.4076/1757-1626-2-7246.
- 426 **Coagulation disorders.** This image is a derivative work, adapted from the following source, available under . Lakjiri S, Mernissi FZ. Tabetic arthropathy revealing neurosyphilis: a new observation. *Pan Afr Med J*. 2014; 18: 198. DOI: 10.11604/pamj.2014.18.198.4893.
- 429 **Hodgkin lymphoma.** This image is a derivative work, adapted from the following source, available under . Knecht H, Righolt C, Mai S. Genomic instability: the driving force behind refractory/relapsing Hodgkin's lymphoma. *Cancers (Basel)*. 2013 Jun; 5(2): 714–725. DOI: 10.3390/cancers5020714.
- 430 **Non-Hodgkin lymphoma: Image B.** This image is a derivative work, adapted from the following source, available under . Bi CF, Tang Y, Zhang WY, et al. Sporadic Burkitt lymphomas of children and adolescents in Chinese: a clinicopathological study of 43 cases. *Diagn Pathol*. 2012;7:72. DOI:10.1186/1746-1596-7-72.
- 430 **Non-Hodgkin lymphoma: Image C.** This image is a derivative work, adapted from the following source, available under . Mansour A, Qandeel M, Abdel-Razeq H, et al. MR imaging features of intracranial primary CNS lymphoma in immune competent patients. *Cancer Imaging*. 2014;14(1):22. DOI: 10.1186/1470-7330-14-22.
- 430 **Non-Hodgkin lymphoma: Image D.** This image is a derivative work, adapted from the following source, available under . Chaudhary S, Bansal C, Ranga U, et al. Erythrodermic mycosis fungoides with hypereosinophilic syndrome: a rare presentation. *Ecancermedicalscience*. 2013;7:337. DOI:10.3332/ecancer.2013.337
- 431 **Plasma cell dyscrasias: Image C.** This image is a derivative work, adapted from the following source, available under . Mehrotra R, Singh M, Singh PA, et al. Should fine needle aspiration biopsy be the first pathological investigation in the diagnosis of a bone lesion? An algorithmic approach with review of literature. *Cytojournal*. 2007; 4: 9. DOI: 10.1186/1742-6413-4-9.
- 432 **Myelodysplastic syndromes.** This image is a derivative work, adapted from the following source, available under . Lukaszewska J, Allison RW, Stepkowska J. Congenital Pelger-Huët anomaly in a Danish/Swedish farmdog: case report. *Acta Vet Scand*. 2011; 53(1): 14. DOI: 10.1186/1751-0147-53-14.
- 433 **Leukemias: Image A.** This image is a derivative work, adapted from the following source, available under . Chiaretti S, Zini G, Bassan R. Diagnosis and subclassification of acute lymphoblastic leukemia. *Mediterr J Hematol Infect Dis*. 2014; 6(1): e2014073. DOI: 10.4084/MJHID.2014.073.
- 433 **Leukemias: Image C.** Hairy cell leukemia. This image is a derivative work, adapted from the following source, available under . Chan SM, George T, Cherry AM, et al. Complete remission of primary plasma cell leukemia with bortezomib, doxorubicin, and dexamethasone: a case report. *Cases J*. 2009;2:121. DOI: 10.1186/1757-1626-2-121.
- 433 **Chronic myeloproliferative disorders: Image A.** Erythromelalgia in polycythemia vera. This image is a derivative work, adapted from the following source, available under . Fred H, van Dijk H. Images of memorable cases: case 151. Connexions Web site. December 4, 2008. Available at <http://cnx.org/content/m14932/1.3/>.
- 433 **Chronic myeloproliferative disorders: Image C.** Myelofibrosis. This image is a derivative work, adapted from the following source, available under . Courtesy of Dr. Ed Uthman.
- 434 **Langerhans cell histiocytosis: Image A.** Lytic bone lesion. This image is a derivative work, adapted from the following source, available under . Dehkordi NR, Rajabi P, Naimi A, et al. Langerhans cell histiocytosis following Hodgkin lymphoma: a case report from Iran. *J Res Med Sci*. 2010;15:58-61. PMCID PMC3082786.
- 434 **Langerhans cell histiocytosis: Image B.** Birbeck granules. This image is a derivative work, adapted from the following source, available under . Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 435 **Hemophagocytic lymphohistiocytosis.** This image is a derivative work, adapted from the following source, available under . Kashif M, Tariq H, Ijaz M. Disseminated histoplasmosis and secondary hemophagocytic syndrome in a non-HIV patient. *Case Rep Crit Care*. 2015; 2015: 295735. DOI: 10.1155/2015/295735.




- 436 Warfarin.** This image is a derivative work, adapted from the following source, available under : Bakoyiannis C, Karaolanis G, Patelis N. Dabigatran in the treatment of warfarin-induced skin necrosis: A new hope. *Case Rep Dermatol Med*. 2016; 2016: 3121469. DOI: 10.1155/2016/3121469.



















Musculoskeletal, Skin, and Connective Tissue
































- 446 Rotator cuff muscles.** Glenohumeral instability. This image is a derivative work, adapted from the following source, available under : Koike Y, Sano H, Imamura I, et al. Changes with time in skin temperature of the shoulders in healthy controls and a patient with shoulder-hand syndrome. *Ups J Med Sci* 2010;115:260-265. DOI: 10.3109/03009734.2010.503354. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 448 Brachial plexus lesions: Image A.** Cervical rib. This image is a derivative work, adapted from the following source, available under : Dahlin LB, Backman C, Duppe H, et al. Compression of the lower trunk of the brachial plexus by a cervical rib in two adolescent girls: case reports and surgical treatment. *J Brachial Plex Peripher Nerve Inj*. 2009;4:14. DOI: 10.1186/1749-7221-4-14.
- 448 Brachial plexus lesions: Image B.** Winged scapula. This image is a derivative work, adapted from the following source, available under : Boukhris J, Boussouga M, Jaafar A, et al. Stabilisation dynamique d'un winging scapula (à propos d'un cas avec revue de la littérature). *Pan Afr Med J*. 2014; 19: 331. DOI: 10.11604/pamj.2014.19.331.3429.
- 449 Wrist region: Image B.** Anatomic snuff box. This image is a derivative work, adapted from the following source, available under : Rhenrev SJ, Ootes D, Beeres FJP, et al. Current methods of diagnosis and treatment of scaphoid fractures. *Int J Emerg Med*. 2011;4:4. DOI: 10.1186/1865-1380-4-4.
- 456 Motoneuron action potential to muscle contraction: Image A.** This image is a derivative work, adapted from the following source, available under : Ottenheim CAC, Heunks LMA, Dekhuijzen RPN. Diaphragm adaptations in patients with COPD. *Respir Res*. 2008; 9(1): 12. DOI: 10.1186/1465-9921-9-12.
- 459 Wrist and hand injuries: Image A.** Metacarpal neck fracture. This image is a derivative work, adapted from the following source, available under : Bohr S, Pallua N. Early functional treatment and modern cast making for indications in hand surgery. *Adv Orthop*. 2016; 2016: 5726979. DOI: 10.1155/2016/5726979.
- 459 Wrist and hand injuries: Image B.** Thenar eminence atrophy in carpal tunnel syndrome.  Courtesy of Dr. Harry Gouvas.
- 460 Common hip and knee conditions: Image A.** ACL tear. This image is a derivative work, adapted from the following source, available under : Chang MJ, Chang CB, Choi J-Y, et al. Can magnetic resonance imaging findings predict the degree of knee joint laxity in patients undergoing anterior cruciate ligament reconstruction? *BMC Musculoskelet Disord*. 2014;15:214. DOI: 10.1186/1471-2474-15-214. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 460 Common hip and knee conditions: Images B (prepatellar bursitis) and C (Baker cyst).** This image is a derivative work, adapted from the following source, available under : Hirji Z, Hunhun JS, Choudur HN. Imaging of the bursae. *J Clin Imaging Sci*. 2011;1:22. DOI: 10.4103/2156-7514.80374. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 462 Common pediatric fractures: Image A.** Greenstick fracture. This image is a derivative work, adapted from the following source, available under : Randsborg PH, Sivertsen EA. Classification of distal radius fractures in children: good inter- and intraobserver reliability, which improves with clinical experience. *BMC Musculoskelet Disord*. 2013;13:6. DOI: 10.1186/1471-2474-13-6.
- 462 Common pediatric fractures: Image B.** Torus (buckle) fracture. This image is a derivative work, adapted from the following source, available under : Aksel Seyahi, et al. Tibial torus and toddler's fractures misdiagnosed as transient synovitis: a case series. *J Med Case Reports*. 2011; 5: 305. DOI: 10.1186/1752-1947-5-305.
- 462 Osteoporosis.** Vertebral compression fractures of spine. This image is a derivative work, adapted from the following source, available under : Imani F, Gharaei H, Rahimzadeh P, et al. Management of painful vertebral compression fracture with kyphoplasty in a severe cardio-respiratory compromised patient. *Anesth Pain Med*. 2012 summer;2(1):42-45. DOI: 10.5812/aapm.5030.
- 463 Osteopetrosis.** This image is a derivative work, adapted from the following source, available under : Kant P, Sharda N, Bhowate RR. Clinical and radiological findings of autosomal dominant osteopetrosis type II: a case report. *Case Rep Dent*. 2013;2013:707343. DOI: 10.1155/2013/707343.
- 463 Osteomalacia/rickets: Image A, left.** Clinical photo. This image is a derivative work, adapted from the following source, available under : Linglart A, Biosse-Duplan M, Briot K, et al. Therapeutic management of hypophosphatemic rickets from infancy to adulthood. *Endocr Connect*. 2014;3:R13-R30. DOI: 10.1530/EC-13-0103.
- 463 Osteomalacia/rickets: Image B.** Rachitic rosary on chest X-ray. This image is a derivative work, adapted from the following source, available under : Essabar L, Meskini T, Ettair S, et al. Malignant infantile osteopetrosis: case report with review of literature. *Pan Afr Med J*. 2014;17:63. DOI: 10.11604/pamj.2014.17.63.3759.
- 463 Osteitis deformans.** Thickened calvarium. This image is a derivative work, adapted from the following source, available under : Dawes L. Paget's disease. [Radiology Picture of the Day Website]. Published June 21, 2007. Available at <http://www.radpod.org/2007/06/21/pagets-disease/>.
- 463 Avascular necrosis of bone.** Bilateral necrosis of femoral head. This image is a derivative work, adapted from the following source, available under : Ding H, Chen S-B, Lin S, et al. The effect of postoperative corticosteroid administration on free vascularized fibular grafting for treating osteonecrosis of the femoral head. *Sci World J*. 2013;708014. DOI: 10.1155/2013/708014. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 465 Primary bone tumors: Image A.** Osteochondroma. This image is a derivative work, adapted from the following source, available under : Lucien Monfils. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 465 Primary bone tumors: Image B.** Osteoid osteoma. This image is a derivative work, adapted from the following source, available under : Jankharia B, Burute N. Percutaneous radiofrequency ablation for osteoid osteoma: how we do it. *Indian J Radiol Imaging*. 2009 Feb; 19(1): 36-42. DOI: 10.4103/0971-3026.44523.
- 465 Primary bone tumors: Image C.** Giant cell tumor. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 465 Primary bone tumors: Image D.** This image is a derivative work, adapted from the following source, available under : Xu SF, Yu XC, Zu M, et al. Limb function and quality of life after various reconstruction methods according to tumor location following resection of osteosarcoma in distal femur. *BMC Musculoskelet Disord*. 2014; 15: 453. DOI: 10.1186/1471-2474-15-453.
- 465 Primary bone tumors: Image E.** Starburst pattern in osteosarcoma. This image is a derivative work, adapted from the following source, available under : Ding H, Yu G, Tu Q, et al. Computer-aided resection and endoprosthesis design for the management of

- malignant bone tumors around the knee: outcomes of 12 cases. *BMC Musculoskelet Disord.* 2013; 14: 331. DOI: 10.1186/1471-2474-14-331.
- 466 **Osteoarthritis vs rheumatoid arthritis: Image A.** Histology of rheumatoid nodule. This image is a derivative work, adapted from the following source, available under : Gomez-Rivera F, El-Naggar AK, Guha-Thakurta N, et al. Rheumatoid arthritis mimicking metastatic squamous cell carcinoma. *Head Neck Oncol.* 2011;3:26. DOI: 10.1186/1758-3284-3-26.
- 467 **Gout: Image B.** Uric acid crystals under polarized light. This image is a derivative work, adapted from the following source, available under : Robert J. Galindo. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 467 **Gout: Image C.** Podagra. This image is a derivative work, adapted from the following source, available under : Roddy E. Revisiting the pathogenesis of podagra: why does gout target the foot? *J Foot Ankle Res.* 2011;4:13. DOI: 10.1186/1757-1146-4-13.
- 467 **Calcium pyrophosphate deposition disease.** Calcium phosphate crystals. This image is a derivative work, adapted from the following source, available under : Dieppe P, Swan A. Identification of crystals in synovial fluid. *Ann Rheum Dis.* 1999 May;58(5):261–263.
- 468 **Sjögren syndrome: Image A.** Lymphocytic infiltration.  Courtesy of the Department of Health and Human Services.
- 468 **Sjögren syndrome: Image B.** Dry tongue. This image is a derivative work, adapted from the following source, available under : Negrato CA, Tarzia O. Buccal alterations in diabetes mellitus. *Diabetol Metab Syndr.* 2010;2:3. DOI: 10.1186/1758-5996-2-3.
- 468 **Septic arthritis.** Joint effusion. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 469 **Seronegative spondyloarthropathies: Image C, left.** Bamboo spine. This image is a derivative work, adapted from the following source, available under : Stevenfruitsmaak. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 469 **Seronegative spondyloarthropathies: Image C, right.** Bamboo spine.  Courtesy of Heather Hawker.
- 471 **Polymyositis/dermatomyositis: Image A.** Groton papules of dermatomyositis. This image is a derivative work, adapted from the following source, available under : *Pan Afr Med J.* 2015; 21: 89. DOI: 10.11604/pamj.2015.21.89.6971.
- 472 **Raynaud phenomenon.** This image is a derivative work, adapted from the following source, available under : Jamclaassen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 474 **Epithelial cell junctions: Image A.** Intracellular membrane. This image is a derivative work, adapted from the following source, available under : Tang VW. Proteomic and bioinformatic analysis of epithelial tight junction reveals an unexpected cluster of synaptic molecules. *Biol Direct.* 2006; 1: 37. DOI: 10.1186/1745-6150-1-37.
- 474 **Epithelial cell junctions: Image B.** Large, electron-dense actin structures within adherens junction. This image is a derivative work, adapted from the following source, available under : Taylor RR, Jagger DJ, Saeed SR, et al. Characterizing human vestibular sensory epithelia for experimental studies: new hair bundles on old tissue and implications for therapeutic interventions in ageing. *Neurobiol Aging.* 2015 Jun;36(6):2068–2084. DOI: 10.1016/j.neurobiolaging.2015.02.013.
- 474 **Epithelial cell junctions: Image C.** Desmosome. This image is a derivative work, adapted from the following source, available under : Massa F, Devader C, Lacas-Gervais S, et al. Impairment of HT29 cancer cells cohesion by the soluble form of neurotensin receptor-3. *Genes Cancer.* 2014 Jul; 5(7-8):240–249. DOI: 10.18632/genesandcancer.22.
- 474 **Epithelial cell junctions: Image D.** Gap junction. This image is a derivative work, adapted from the following source, available under : Shu X, Lev-Ram V, Deerinck TJ. A Genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms. *PLoS Biol.* 2011 Apr; 9(4): e1001041. DOI: 10.1371/journal.pbio.1001041.
- 474 **Epithelial cell junctions: Image E.** Hemidesmosome. This image is a derivative work, adapted from the following source, available under : Nguyen NM, Pulkkinen L, Schlueter JA, et al. Lung development in laminin gamma2 deficiency: abnormal tracheal hemidesmosomes with normal branching morphogenesis and epithelial differentiation. *Respir Res.* 2006 Feb 16;7:28. DOI: 10.1186/1465-9921-7-28.
- 476 **Seborrheic dermatitis.** This image is a derivative work, adapted from the following source, available under : Roymishali.
- 477 **Common skin disorders: Image O.** Urticaria. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 478 **Vascular tumors of skin: Image C.** Glomus tumor under fingernail. This image is a derivative work, adapted from the following source, available under : Hazani R, Houle JM, Kasdan ML, et al. Glomus tumors of the hand. *Eplasty.* 2008;8:e48. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 479 **Skin infections: Image C.** Erysipelas. This image is a derivative work, adapted from the following source, available under : Courtesy of Klaus D. Peter.
- 480 **Autoimmune blistering skin disorders: Image D.** Bullous pemphigoid on immunofluorescence. This image is a derivative work, adapted from the following source, available under : Courtesy of M. Emmanuel.
- 484 **Skin cancer: Image D.** Basal cell palisading nuclei. This image is a derivative work, adapted from the following source, available under : Yuri T. Jadotte, MD, et al. Superficial spreading basal cell carcinoma of the face: a surgical challenge. *Eplasty.* 2010; 10: e46. Published online 2010 Jun 21.





































Neurology and Special Senses


































- 491 **Holoprosencephaly: Image A.** This image is a derivative work, adapted from the following source, available under : Alorainy IA, Barlas NB, Al-Boukai AA. Pictorial essay: infants of diabetic mothers. *Indian J Radiol Imaging.* 2010 Aug;20(3):174-81. DOI: 10.4103/0971-3026.69349.
- 492 **Posterior fossa malformations: Image A.** Chiari I malformation. This image is a derivative work, adapted from the following source, available under : Toldo I, De Carlo D, Mardari R, et al. Short lasting activity-related headaches with sudden onset in children: a case-based reasoning on classification and diagnosis. *J Headache Pain.* 2013;14(1):3. DOI: 10.1186/1129-2377-14-3.
- 492 **Posterior fossa malformations: Image B.** Dandy-Walker malformation. This image is a derivative work, adapted from the following source, available under : Krupa K, Bekiesinska-Figatowska M. Congenital and acquired abnormalities of the corpus callosum: a pictorial essay. *Biomed Res Int.* 2013;2013:265619. DOI: 10.1155/2013/265619.
















- 492 **Syringomyelia.** Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 494 **Myelin.** Myelinated neuron.  Courtesy of the Electron Microscopy Facility at Trinity College.
- 495 **Chromatolysis.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonnert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 499 **Limbic system: Image A.** This image is a derivative work, adapted from the following source, available under : Schopf V, Fischmeister FP, Windischberger C, et al. Effects of individual glucose levels on the neuronal correlates of emotions. *Front Hum Neurosci*. 2013 May 21;7:212. DOI: 10.3389/fnhum.2013.00212.
- 499 **Cerebellum.** This image is a derivative work, adapted from the following source, available under : Jarius S, Wandinger KP, Horn S, et al. A new Purkinje cell antibody (anti-Ca) associated with subacute cerebellar ataxia: immunological characterization. *J Neuroinflammation*. 2010;7: 21. DOI: 10.1186/1742-2094-7-21.
- 500 **Basal ganglia.** This image is a derivative work, adapted from the following source, available under : Rudger P, Jaunmuktane Z, Adlard P, et al. Iatrogenic CJD due to pituitary-derived growth hormone with genetically determined incubation times of up to 40 years. *Brain*. 2015 Nov; 138(11): 3386–3399. DOI: 10.1093/brain/awv235.
- 502 **Cerebral arteries—cortical distribution.** Cortical watershed areas. This image is a derivative work, adapted from the following source, available under : Isabel C, Lecler A, Turc G, et al. Relationship between watershed infarcts and recent intra plaque haemorrhage in carotid atherosclerotic plaque. *PLoS One*. 2014;9(10):e108712. DOI: 10.1371/journal.pone.0108712.
- 503 **Dural venous sinuses.** This image is a derivative work, adapted from the following source, available under : Cikla U, Aagaard-Kienitz B, Turski PA, et al. Familial perimesencephalic subarachnoid hemorrhage: two case reports. *J Med Case Rep*. 2014;8. DOI: 10.1186/1752-1947-8-380. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 508 **Spinal cord and associated tracts: Image A.** Spinal cord cross-section. This image is a derivative work, adapted from the following source, available under : Regents of University of Michigan Medical School.
- 512 **Neonatal interventricular hemorrhage.** This image is a derivative work, adapted from the following source, available under : Shooman D, Portess H, Sparrow O. A review of the current treatment methods for posthaemorrhagic hydrocephalus of infants. *Cerebrospinal Fluid Res*. 2009;6:1. DOI: 10.1186/1743-8454-6-1.
- 513 **Intracranial hemorrhage: Images A and B.** Axial CT of brain showing epidural blood. This image is a derivative work, adapted from the following source, available under : Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 513 **Intracranial hemorrhage: Image C.** Subdural hematoma. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 513 **Intracranial hemorrhage: Image E.** Subarachnoid hemorrhage. This image is a derivative work, adapted from the following source, available under : Hakan T, Turk CC, Celik H. Intra-operative real time intracranial subarachnoid haemorrhage during glial tumour resection: a case report. *Cases J*. 2008;1:306. DOI: 10.1186/1757-1626-1-306. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 515 **Effects of strokes: Image A.** Large abnormality of the left MCA territory. This image is a derivative work, adapted from the following source, available under : Hakimelahi R, Yoo AJ, He J, et al. Rapid identification of a major diffusion/perfusion mismatch in distal internal carotid artery or middle cerebral artery ischemic stroke. *BMC Neurol*. 2012 Nov 5;12:132. DOI: 10.1186/1471-2377-12-132. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 515 **Effects of strokes: Image B.** MRI diffusion weighted image shows a hypersensitive lesion on posterior limb of internal capsular. This image is a derivative work, adapted from the following source, available under : Zhou L, Ni J, Yao M, et al. High-resolution MRI findings in patients with capsular warning syndrome. *BMC Neurol*. 2014;14:16. DOI: 10.1186/1471-2377-14-16.
- 515 **Effects of strokes: Image C.** This image is a derivative work, adapted from the following source, available under : Nough A, Remke J, Ruland S. Ischemic posterior circulation stroke: a review of anatomy, clinical presentations, diagnosis, and current management. *Front Neurol*. 2014 Apr 7;5:30. DOI: 10.3389/fneur.2014.00030.
- 515 **Effects of strokes: Image D.** This image is a derivative work, adapted from the following source, available under : Mittal P, Kalia V, Dua S. Pictorial essay: Susceptibility-weighted imaging in cerebral ischemia. *Indian J Radiol Imaging*. 2010 Nov; 20(4): 250–253. DOI: 10.4103/0971-3026.73530.
- 515 **Diffuse axonal injury.** Moeninghoff C, Kraff O, Maderwald S, et al. Diffuse axonal injury at ultra-high field MRI. *PLoS One*. 2015;10(3):e0122329. DOI: 10.1371/journal.pone.0122329.
- 516 **Aneurysms.** This image is a derivative work, adapted from the following source, available under : Kayhan A, Koc O, Keskin S. The role of bone subtraction computed tomographic angiography in determining intracranial aneurysms in non-traumatic subarachnoid hemorrhage. *Iran J Radiol*. 2014 May; 11(2): e12670. DOI: 10.5812/iranradiol.12670.
- 521 **Neurodegenerative disorders: Image A.** Lewy body in substantia nigra. This image is a derivative work, adapted from the following source, available under : Werner CJ, Heyny-von Haussen R, Mall G, et al. Parkinson's disease. *Proteome Sci*. 2008;6:8. DOI: 10.1186/1477-5956-6-8. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MediQ Learning, LLC are reserved.
- 521 **Neurodegenerative disorders: Image B.** Gross specimen of normal brain. This image is a derivative work, adapted from the following source, available under : Niedowicz DM, Nelson PT, Murphy MP. Alzheimer's disease: pathological mechanisms and recent insights. *Curr Neuropharmacol*. 2011 Dec;9(4):674-84. DOI: 10.2174/157015911798376181.
- 521 **Neurodegenerative disorders: Images C (brain atrophy in Alzheimer disease) and F (atrophy in frontotemporal dementia).** This image is a derivative work, adapted from the following source, available under : Niedowicz DM, Nelson PT, Murphy MP. Alzheimer's disease: pathological mechanisms and recent insights. *Curr Neuropharmacol*. 2011 Dec;9(4):674-84. DOI: 10.2174/157015911798376181.
- 521 **Neurodegenerative disorders: Image G.** Frontotemporal dementia: Pick bodies in frontotemporal dementia (Pick disease). This image is a derivative work, adapted from the following source, available under : Neumann M. Molecular neuropathology of TDP-43 proteinopathies. *Int J Mol Sci*. 2009 Jan; 10(1): 232–246. DOI: 10.3390/ijms10010232.
- 521 **Neurodegenerative disorders: Image H.** Spongiform changes in brain in Creutzfeldt-Jacob disease. This image is a derivative work, adapted from the following source, available under : DRdoubleB. The

- image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 522 Hydrocephalus: Image B.** Communicating hydrocephalus. This image is a derivative work, adapted from the following source, available under : Torres-Martin M, Pena-Granero C, Carceller F, et al. Homozygous deletion of *TNFRSF4*, *TP73*, *PPAP2B* and *DPYD* at 1p and *PDCD5* at 19q identified by multiplex ligation-dependent probe amplification (MLPA) analysis in pediatric anaplastic glioma with questionable oligodendroglial component. *Mol Cytogenet.* 2014;7:1. DOI: 10.1186/1755-8166-7-1.
- 522 Hydrocephalus: Image C.** Ex vacuo ventriculomegaly. This image is a derivative work, adapted from the following source, available under : Ghetti B, Oblak AL, Boeve BF, et al. Frontotemporal dementia caused by microtubule-associated protein tau gene (*MAPT*) mutations: a chameleon for neuropathology and neuroimaging. *NeuroPathol Appl Neurobiol.* 2015 Feb;41(1):24-46. DOI: 10.1111/nan.12213.
- 523 Multiple sclerosis.** Periventricular plaques. This image is a derivative work, adapted from the following source, available under : Dooley MC, Foroozan R. Optic neuritis. *J Ophthalmic Vis Res.* 2010 Jul;5(3):182-187.
- 524 Other demyelinating and dysmyelinating diseases: Image B.** Progressive multifocal leukoencephalopathy. This image is a derivative work, adapted from the following source, available under : Garrote H, de la Fuente A, Ona R, et al. Long-term survival in a patient with progressive multifocal leukoencephalopathy after therapy with rituximab, fludarabine and cyclophosphamide for chronic lymphocytic leukemia. *Exp Hematol Oncol.* 2015;4:8. DOI: 10.1186/s40164-015-0003-4.
- 524 Other demyelinated and dysmyelinating disorders: Image A.** Central pontine myelinolysis. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 525 Neurocutaneous disorders: Image A.** Sturge-Weber syndrome and port wine stain. This image is a derivative work, adapted from the following source, available under : Babaji P, Bansal A, Krishna G, et al. Sturge-Weber syndrome with osteohypertrophy of maxilla. *Case Rep Pediatr.* 2013. DOI: 10.1155/2013/964596.
- 525 Neurocutaneous disorders: Image B.** Leptomeningeal angioma in Sturge-Weber syndrome. Reproduced, with permission, from Dr. Frank Gaillard and www.radiopaedia.org.
- 525 Neurocutaneous disorders: Image C.** Tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : Fred H, van Dijk H. Images of memorable cases: case 143. Connexions Web site. December 4, 2008. Available at: <http://cnx.org/content/m14923/1.3/>.
- 525 Neurocutaneous disorders: Image D.** Ash leaf spots in tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : Tonekaboni SH, Tousi P, Ebrahimi A, et al. Clinical and para clinical manifestations of tuberous sclerosis: a cross sectional study on 81 pediatric patients. *Iran J Child Neurol.* 2012;6:25-31. PMID: PMC3943027.
- 525 Neurocutaneous disorders: Image E.** Angiomyolipoma in tuberous sclerosis. This image is a derivative work, adapted from the following source, available under : KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 525 Neurocutaneous disorders: Image F.** Café-au-lait spots in neurofibromatosis. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 525 Neurocutaneous disorders: Image G.** Lisch nodules in neurofibromatosis.  Courtesy of the Department of Health and Human Services.
- 525 Neurocutaneous disorders: Image H.** Cutaneous neurofibromas. This image is a derivative work, adapted from the following source, available under : Kim BK, Choi YS, Gwoo S, et al. Neurofibromatosis type 1 associated with papillary thyroid carcinoma incidentally detected by thyroid ultrasonography: a case report. *J Med Case Rep.* 2012;6:179. DOI: 10.1186/1752-1947-6-179.
- 525 Neurocutaneous disorders: Image I.** Cerebellar hemangioblastoma histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 525 Neurocutaneous disorders: Image J.** Brainstem and spinal cord hemangioblastomas in von Hippel-Lindau disease. This image is a derivative work, adapted from the following source, available under : Park DM, Zhuang Z, Chen L, et al. von Hippel-Lindau disease-associated hemangioblastomas are derived from embryologic multipotent cells. *PLoS Med.* 2007 Feb;4(2):e60. DOI: 10.1371/journal.pmed.0040060.
- 526 Adult primary brain tumors: Image A.** This image is a derivative work, adapted from the following source, available under : Rossmesl JH, Clapp K, Pancotto TE. Canine butterfly glioblastomas: A neuroradiological review. *Front Vet Sci.* 2016; 3: 40. DOI: 10.3389/fvets.2016.00040.
- 526 Adult primary brain tumors: Image B.** Glioblastoma multiforme histology. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 526 Adult primary brain tumors: Image C.** Oligodendroglioma in frontal lobes. This image is a derivative work, adapted from the following source, available under : Celzo FG, Venstermans C, De Belder F, et al. Brain stones revisited—between a rock and a hard place. *Insights Imaging.* 2013 Oct;4(5):625-35. DOI: 10.1007/s13244-013-0279-z.
- 526 Adult primary brain tumors: Image D.** Oligodendroglioma, “fried egg” cells. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 526 Adult primary brain tumors: Image E.** Meningioma with dural tail. This image is a derivative work, adapted from the following source, available under : Smits A, Zetterling M, Lundin M, et al. Neurological impairment linked with cortico-subcortical infiltration of diffuse low-grade gliomas at initial diagnosis supports early brain plasticity. *Front Neurol.* 2015;6:137. DOI: 10.3389/fneur.2015.00137.
- 526 Adult primary brain tumors: Image F.** Meningioma, psammoma bodies. This image is a derivative work, adapted from the following source, available under : Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 526 Adult primary brain tumors: Image G.** Cerebellar hemangioblastoma. This image is a derivative work, adapted from the following source, available under : Park DM, Zhengping Z, Chen L, et al. von Hippel-Lindau disease-associated hemangioblastomas are derived from embryologic multipotent cells. *PLoS Med.* 2007 Feb;4(2):e60. DOI: 10.1371/journal.pmed.0040060.
- 526 Adult primary brain tumors: Image H.** Minimal parenchyma in hemangioblastoma. This image is a derivative work, adapted from the following source, available under : Marvin 101. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .








- 527 Adult primary brain tumors: Image I.** Field of vision in bitemporal hemianopia. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 527 Adult primary brain tumors: Image J.** Prolactinoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Wang CS, Yeh TC, Wu TC, et al. Pituitary macroadenoma co-existent with supraclinoid internal carotid artery cerebral aneurysm: a case report and review of the literature. *Cases J.* 2009;2:6459. DOI: 10.4076/1757-1626-2-6459.
- 527 Adult primary brain tumors: Image K.** Schwannoma at cerebellopontine angle. [CC-BY](#) Courtesy of MRT-Bild.
- 527 Adult primary brain tumors: Image L.** Schwann cell origin of schwannoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 528 Childhood primary brain tumors: Image A.** MRI of pilocytic astrocytoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Hafez RFA. Stereotaxic gamma knife surgery in treatment of critically located pilocytic astrocytoma: preliminary result. *World J Surg Oncol.* 2007;5:39. doi 10.1186/1477-7819-5-39.
- 528 Childhood primary brain tumors: Image C.** CT of medulloblastoma. [CC-BY](#) Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 528 Childhood primary brain tumors: Image D.** Medulloblastoma histology. This image is a derivative work, adapted from the following source, available under [CC-BY](#): KGH. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 528 Childhood primary brain tumors: Image E.** MRI of ependymoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Dr. Paul Hellerhoff. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 528 Childhood primary brain tumors: Image F.** Ependymoma histology. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 528 Childhood primary brain tumors: Image G.** CT of craniopharyngioma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Garnet MR, Puget S, Grill J, et al. Craniopharyngioma. *Orphanet J Rare Dis.* 2007;2:18. DOI: 10.1186/1750-1172-2-18.
- 528 Childhood primary brain tumors: Image H.** Craniopharyngioma histology. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Nephron. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 531 Friedreich ataxia: Image A.** Clinical kyphoscoliosis. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Axelrod FB, Gold-von Simson. Hereditary sensory and autonomic neuropathies: types II, III, and IV. *Orphanet J Rare Dis.* 2007;2:39. DOI: 10.1186/1750-1172-2-39.
- 531 Friedreich ataxia: Image B.** Radiograph showing kyphoscoliosis. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Bounakis N, Karampalis C, Tsirikos AI. Surgical treatment of scoliosis in Rubinstein-Taybi syndrome type 2: a case report. *J Med Case Rep.* 2015; 9: 10. doi 10.1186/1752-1947-9-10.
- 532 Facial nerve lesions.** Facial nerve palsy. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Socolovsky M, Paez MD, Di Masi G, et al. Bell's palsy and partial hypoglossal to facial nerve transfer: Case presentation and literature review. *Surg Neurol Int.* 2012;3:46. DOI: 10.4103/2152-7806.95391.
- 533 Cholesteatoma.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Welleschik. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 534 Normal eye anatomy.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Jan Kaláb. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under [CC-BY](#).
- 534 Conjunctivitis.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Baiyeroju A, Bowman R, Gilbert C, et al. Managing eye health in young children. *Community Eye Health.* 2010;23:4-11.
- 535 Cataract.** Juvenile cataract. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Roshan M, Vijaya PH, Lavanya GR, et al. A novel human CRYGD mutation in a juvenile autosomal dominant cataract. *Mol Vis.* 2010;16:887-896. PMID: PMC2875257.
- 536 Glaucoma: Image C.** Closed/narrow angle glaucoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Low S, Davidson AE, Holder GE, et al. Autosomal dominant Best disease with an unusual electrooculographic light rise and risk of angle-closure glaucoma: a clinical and molecular genetic study. *Mol Vis.* 2011;17:2272-2282. PMID: PMC3171497. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 536 Glaucoma: Image D.** Acute angle closure glaucoma. This image is a derivative work, adapted from the following source, available under [CC-BY](#): Courtesy of Dr. Jonathan Trobe.
- 536 Uveitis: Image A.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Weber AC, Levison AL, Srivastava, et al. A case of *Listeria monocytogenes* endophthalmitis with recurrent inflammation and novel management. *J Ophthalmic Inflamm Infect.* 2015;5(1):28. DOI: 10.1186/s12348-015-0058-8.
- 536 Age-related macular degeneration.** [CC-BY](#) Courtesy of the Department of Health and Human Services.
- 537 Diabetic retinopathy.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Sundling V, Gulbrandsen P, Straand J. Sensitivity and specificity of Norwegian optometrists' evaluation of diabetic retinopathy in single-field retinal images – a cross-sectional experimental study. *BMC Health Services Res.* 2013;13:17. DOI: 10.1186/1472-6963-13-17.
- 537 Hypertensive retinopathy.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Diallo JW, Méda N, Tougouma SJB, et al. Intérêts de l'examen du fond d'œil en pratique de ville: bilan de 438 cas. *Pan Afr Med J.* 2015;20:363. DOI: 10.11604/pamj.2015.20.363.6629.
- 537 Retinal vein occlusion.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Alasil T, Rauser ME. Intravitreal bevacizumab in the treatment of neovascular glaucoma secondary to central retinal vein occlusion: a case report. *Cases J.* 2009;2:176. DOI: 10.1186/1757-1626-2-176. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 537 Retinal detachment.** Courtesy of EyeRounds.
- 538 Retinitis pigmentosa.** Courtesy of EyeRounds.
- 538 Leukocoria.** This image is a derivative work, adapted from the following source, available under [CC-BY](#): Aerts I, Lumbroso-Le Rouic L,
































- Gauthier-Villars M, et al. Retinoblastoma. *Orphanet J Rare Dis*. 2006 Aug 25;1:31. DOI: 10.1186/1750-1172-1-31.
- 540 Ocular motility.** Testing ocular muscles. This image is a derivative work, adapted from the following source, available under : Au.yousef. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 541 CN III, IV, VI palsies: Image A.** Cranial nerve III damage. This image is a derivative work, adapted from the following source, available under : Hakim W, Sherman R, Rezk T, et al. An acute case of herpes zoster ophthalmicus with ophthalmoplegia. *Case Rep Ophthalmol Med*. 2012;2012:953910. DOI: 10.1155/2012/953910.
- 541 CN III, IV, VI palsies: Image B.** Cranial nerve IV damage. This image is a derivative work, adapted from the following source, available under : Mendez JA, Arias CR, Sanchez D, et al. Painful ophthalmoplegia of the left eye in a 19-year-old female, with an emphasis in Tolosa-Hunt syndrome: a case report. *Cases J*. 2009; 2: 8271. DOI: 10.4076/1757-1626-2-8271.
- 541 CN III, IV, VI palsies: Image C.** Cranial nerve VI damage. This image is a derivative work, adapted from the following source, available under : Jordi March i Nogué. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- Psychiatry**
- 563 Trichotillomania.**  Courtesy of Robodoc.
- Renal**
- 578 Potter sequence (syndrome).**  Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 580 Kidney anatomy and glomerular structure.** This image is a derivative work, adapted from the following source, available under : Ramidi GA, Kurukumbi MK, Sealy PL. Collapsing glomerulopathy in sickle cell disease: a case report. *J Med Case Reports*. 2011; 5: 71. DOI: 10.1186/1752-1947-5-71.
- 581 Course of ureters.** This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 581 Glomerular filtration barrier.** This image is a derivative work, adapted from the following source, available under : Feng J, Wei H, Sun Y, et al. Regulation of podocalyxin expression in the kidney of streptozotocin-induced diabetic rats with Chinese herbs (Yishen capsule). *BMC Complement Altern Med*. 2013;13:76. DOI: 10.1186/1472-6882-13-76.
- 594 Casts in urine: Image B.** WBC casts. This image is a derivative work, adapted from the following source, available under : Perazella MA. Diagnosing drug-induced AIN in the hospitalized patient: a challenge for the clinician. *Clin Nephrol*. 2014 Jun; 81(6): 381-8. DOI: 10.5414/CN108301.
- 594 Casts in urine: Image D.** Fatty casts. This image is a derivative work, adapted from the following source, available under : Li S, Wang ZJ, Chang TT. Temperature oscillation modulated self-assembly of periodic concentric layered magnesium carbonate microparticles. *PLoS One*. 2014;9(2):e88648. DOI:10.1371/journal.pone.0088648
- 596 Nephritic syndrome: Image A.** Histology of acute poststreptococcal glomerulonephritis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 596 Nephritic syndrome: Image B.** This image is a derivative work, adapted from the following source, available under : Immunofluorescence of acute poststreptococcal glomerulonephritis. Oda T, Yoshizawa N, Yamakami K, et al. The role of nephritis-associated plasmin receptor (naplr) in glomerulonephritis associated with streptococcal infection. *Biomed Biotechnol*. 2012;2012:417675. DOI 10.1155/2012/417675.
- 596 Nephritic syndrome: Image C.** Histology of rapidly progressive glomerulonephritis.  Courtesy of the Department of Health and Human Services and Uniformed Services University of the Health Sciences.
- 596 Nephritic syndrome: Image D.** This image is a derivative work, adapted from the following source, available under : Kiremitci S, Ensari A. Classifying lupus nephritis: an ongoing story. *Scientific World Journal*. 2014; 2014: 580620. DOI: 10.1155/2014/580620.
- 597 Nephrotic syndrome: Image A.** This image is a derivative work, adapted from the following source, available under : Teoh DCY, El-Modir A. Managing a locally advanced malignant thymoma complicated by nephrotic syndrome: a case report. *J Med Case Reports*. 2008; 2: 89. DOI: 10.1186/1752-1947-2-89.
- 597 Nephrotic syndrome: Image B.** Histology of focal segmental glomerulosclerosis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 597 Nephrotic syndrome: Image D.** Diabetic glomerulosclerosis with Kimmelstiel-Wilson lesions. This image is a derivative work, adapted from the following source, available under : Doc Mari. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 598 Kidney stones: Image A.** Nair S, George J, Kumar S, et al. Acute oxalate nephropathy following ingestion of *Averrhoa bilimbi* juice. *Case Rep Nephrol*. 2014; 2014: DOI: 10.1155/2014/240936.
- 598 Kidney stones: Image B.** This image is a derivative work, adapted from the following source, available under : Joel Mills. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 599 Hydronephrosis.** Ultrasound. This image is a derivative work, adapted from the following source, available under : Wikimedia Commons. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 600 Pyelonephritis: Image A.** This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 600 Pyelonephritis: Image B.** CT scan.  Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 602 Acute tubular necrosis: Image A.** Muddy brown casts. This image is a derivative work, adapted from the following source, available under : Dr. Serban Nicolescu.
- 602 Renal papillary necrosis.**  Courtesy of the Department of Health and Human Services and William D. Craig, Dr. Brent J. Wagner, and Mark D. Travis.
- 604 Renal cyst disorders: Image C.** Ultrasound of simple cyst. This image is a derivative work, adapted from the following source, available under : Nevit Dilmen. The image may have been modified by cropping, labeling, and/or captions. MediQ Learning, LLC makes this available under .
- 605 Renal cell carcinoma: Image A.** Histology. This image is a derivative work, adapted from the following source, available under .



















- Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 605 Renal cell carcinoma: Image C.** CT scan. This image is a derivative work, adapted from the following source, available under : Behnes CL, Schlegel C, Shoukier M, et al. Hereditary papillary renal cell carcinoma primarily diagnosed in a cervical lymph node: a case report of a 30-year-old woman with multiple metastases. *BMC Urol.* 2013;13:3. DOI: 10.1186/1471-2490-13-3.
- 605 Renal cell carcinoma: Image B.** Gross specimen.  Courtesy of Dr. Ed Uthman.
- 605 Renal oncocyoma: Image A.** Gross specimen. This image is a derivative work, adapted from the following source, available under : Courtesy of M. Emmanuel.
- 605 Renal oncocyoma: Image B.** Histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 606 Nephroblastoma.** This image is a derivative work, adapted from the following source, available under : Refaie H, Sarhan M, Hafez A. Role of CT in assessment of unresectable Wilms tumor response after preoperative chemotherapy in pediatrics. *Sci World J.* 2008;8:661-669. doi 10.1100/tsw.2008.96.
- 606 Urothelial carcinoma of the bladder: Image A.** This image is a derivative work, adapted from the following source, available under : Geavlete B, Stanescu F, Moldoveanu C, et al. NBI cystoscopy and bipolar electrosurgery in NMIBC management—an overview of daily practice. *J Med Life.* 2013;6:140-145. PMID: PMC3725437.
- Reproductive**
- 615 Fetal alcohol syndrome.** This image is a derivative work, adapted from the following source, available under : Courtesy of Teresa Kellerman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 618 Umbilical cord: Image A.** Cross-section of normal umbilical cord. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 618 Meckel diverticulum: Image B.** This image is a derivative work, adapted from the following source, available under : Mathur P, Gupta R, Simlot A, et al. Congenital pouch colon with double Meckel's diverticulae. *J Neonatal Surg.* 2013 Oct-Dec; 2(4): 48.
- 623 Uterine (Müllerian) duct anomalies: Images A-D.** This image is a derivative work, adapted from the following source, available under : Ahmadi F, Zafarani F, Haghighi H, et al. Application of 3D ultrasonography in detection of uterine abnormalities. *Int J Fertil Steril.* 2011; 4:144-147. PMID: PMC4023499.
- 626 Female reproductive epithelial histology.** Transformation zone. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Ed Uthman. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 628 Seminiferous tubules.** This image is a derivative work, adapted from the following source, available under : Dr. Anlt Rao. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 641 Pregnancy complications.** This image is a derivative work, adapted from the following source, available under : Li W, Wang G, Lin T, et al. Misdiagnosis of bilateral tubal pregnancy: a case report. *J Med Case Rep.* 2014;8:342. DOI: 10.1186/1752-1947-8-342.
- 642 Hydatidiform mole: Image A.** Cluster of cluster of grapes appearance in complete hydatidiform mole. This image is a derivative work, adapted from the following source, available under : Dr. Ed Uthman
- 642 Choriocarcinoma: Image B.** “Cannonball” metastases. This image is a derivative work, adapted from the following source, available under : Lekanidi K, Vlachou PA, Morgan B, et al. Spontaneous regression of metastatic renal cell carcinoma: case report. *J Med Case Rep.* 2007;1:89. DOI: 10.1186/1752-1947-1-89.
- 644 Vulvar pathology: Image A.** Bartholin cyst.  Courtesy of the Department of Health and Human Services and Susan Lindsley.
- 644 Vulvar pathology: Image B.** Lichen sclerosis. This image is a derivative work, adapted from the following source, available under : Lambert J. Pruritus in female patients. *Biomed Res Int.* 2014;2014:541867. DOI: 10.1155/2014/541867.
- 644 Vulvar pathology: Image C.** Vulvar carcinoma. This image is a derivative work, adapted from the following source, available under : Ramli I, Hassam B. Carcinome épidermoïde vulvaire: pourquoi surveiller un lichen scléro-atrophique. *Pan Afr Med J.* 2015;21:48. DOI: 10.11604/pamj.2015.21.48.6018.
- 644 Vulvar pathology: Image D.** Extramallary Paget disease. This image is a derivative work, adapted from the following source, available under : Wang X, Yang W, Yang J. Extramammary Paget's disease with the appearance of a nodule: a case report. *BMC Cancer.* 2010;10:405. DOI: 10.1186/1471-2407-10-405.
- 645 Polycystic ovarian syndrome.** This image is a derivative work, adapted from the following source, available under : Kopera D, Wehr E, Obermayer-Pietsch B. Endocrinology of hirsutism. *Int J Trichology.* 2010;2(1):30–35. doi:10.4103/0974-7753.66910
- 647 Dysgerminoma: Image B.** This image is a derivative work, adapted from the following source, available under : Montesinos L, Acien P, Martinez-Beltran M, et al. Ovarian dysgerminoma and synchronic contralateral tubal pregnancy followed by normal intra-uterine gestation: a case report. *J Med Rep.* 2012;6:399. DOI: 10.1186/1752-1947-6-399.
- 647 Ovarian neoplasms: Image D.** Mature cystic teratoma. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 647 Ovarian neoplasms: Image E.** Yolk sac tumor. This image is a derivative work, adapted from the following source, available under : Jensflorian. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 647 Ovarian neoplasms: Image F.** Call-Exner bodies. This image is a derivative work, adapted from the following source, available under : Katoh T, Yasuda M, Hasegawa K, et al. Estrogen-producing endometrioid adenocarcinoma resembling sex cord-stromal tumor of the ovary: a review of four postmenopausal cases. *Diagn Pathol.* 2012;7:164. DOI: 10.1186/1746-1596-7-164.
- 648 Uterine conditions: Image A.** Endometrial tissue found outside the uterus. This image is a derivative work, adapted from the following source, available under : Hastings JM, Fazleabas AT. A baboon model for endometriosis: implications for fertility. *Reprod Biol Endocrinol.* 2006;4(suppl 1):S7. DOI: 10.1186/1477-7827-4-S1-S7.
- 648 Uterine conditions: Image B.** Endometritis with inflammation of the endometrium. This image is a derivative work, adapted from the following source, available under : Montesinos L, Acien P, Martinez-Beltran M, et al. Ovarian dysgerminoma and synchronic contralateral tubal pregnancy followed by normal intra-uterine

- gestation: a case report. *J Med Rep.* 2012;6:399. DOI: 10.1186/1752-1947-6-399.
- 648 Uterine conditions: Image C.** Endometrial carcinoma. This image is a derivative work, adapted from the following source, available under : Izadi-Mood N, Yarmohammadi M, Ahmadi SA, et al. Reproducibility determination of WHO classification of endometrial hyperplasia/well differentiated adenocarcinoma and comparison with computerized morphometric data in curettage specimens in Iran. *Diagn Pathol.* 2009;4:10. DOI:10.1186/1746-1596-4-10.
- 648 Uterine conditions: Image D.** Leiomyoma (fibroid), gross specimen. This image is a derivative work, adapted from the following source, available under : Courtesy of Hic et nunc.
- 648 Uterine conditions: Image E.** Leiomyoma (fibroid) histology. This image is a derivative work, adapted from the following source, available under : Londero AP, Perego P, Mangioni C, et al. Locally relapsed and metastatic uterine leiomyoma: a case report. *J Med Case Rep.* 2008;2:308. DOI: 10.1186/1752-1947-2-308. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 649 Benign breast disease: Image A.** Fibroadenomas. This image is a derivative work, adapted from the following source, available under : Gokhale S. Ultrasound characterization of breast masses. *Indian J Radiol Imaging.* 2009 Aug;19(3):242-7. DOI: 10.4103/0971-3026.54878.
- 649 Benign breast disease: Images B (phyllodes tumor on ultrasound) and C (phyllodes cyst).** This image is a derivative work, adapted from the following source, available under : Muttarak MD, Lerttumnongtum P, Somwangjaroen A, et al. Phyllodes tumour of the breast. *Biomed Imaging Interv J.* 2006 Apr-Jun;2(2):e33. DOI: 10.2349/bij.2.2.e33.
- 650 Breast cancer: Image A.** Mammography of breast cancer. This image is a derivative work, adapted from the following source, available under : Molino C, Mocerino C, Braucci A, et al. Pancreatic solitary and synchronous metastasis from breast cancer: a case report and systematic review of controversies in diagnosis and treatment. *World J Surg Oncol.* 2014;12:2. DOI:10.1186/1477-7819-12-2
- 650 Breast cancer: Image C.** Comedocarcinoma. This image is a derivative work, adapted from the following source, available under : Costarelli L, Campagna D, Mauri M, et al. Intraductal proliferative lesions of the breast—terminology and biology matter: premalignant lesions or preinvasive cancer? *Int J Surg Oncol.* 2012;501904. DOI: 10.1155/2012/501904. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 650 Breast cancer: Image D.** Paget disease of breast. This image is a derivative work, adapted from the following source, available under : Muttarak M, Siriya B, Kongmebhoh P, et al. Paget's disease of the breast: clinical, imaging and pathologic findings: a review of 16 patients. *Biomed Imaging Interv J.* 2011;7:e16. DOI: 10.2349/bij.7.2.e16.
- 650 Breast cancer: Image E.** Invasive lobular carcinoma. This image is a derivative work, adapted from the following source, available under : Franceschini G, Manno A, Mule A, et al. Gastro-intestinal symptoms as clinical manifestation of peritoneal and retroperitoneal spread of an invasive lobular breast cancer: report of a case and review of the literature. *BMC Cancer.* 2006;6:193. DOI: 10.1186/1471-2407-6-193.
- 650 Breast cancer: Image E.** Peau d'orange of inflammatory breast cancer. This image is a derivative work, adapted from the following source, available under : Levine PH, Zolfaghari L, Young H, et al. What Is inflammatory breast cancer? Revisiting the case definition. *Cancers (Basel).* 2010 Mar;2(1):143–152. DOI: 10.3390/cancers2010143.
- 651 Penile pathology: Image A.** Peyronie disease. This image is a derivative work, adapted from the following source, available under : Tran VQ, Kim DH, Lesser TF, et al. Review of the surgical approaches for Peyronie's disease: corporeal plication and plaque incision with grafting. *Adv Urol.* 2008; 2008: 263450. DOI: 10.1155/2008/263450.
- 651 Penile pathology: Image B.** Squamous cell carcinoma. This image is a derivative work, adapted from the following source, available under : Antônio JR, Antônio CR, Trídico LA. Erythroplasia of queyrat treated with topical 5-fluorouracil. *An Bras Dermatol.* 2016 Sep-Oct; 91(5 Suppl 1): 42–44. DOI: 10.1590/abd1806-4841.20164595.
- 651 Cryptorchidism.** This image is a derivative work, adapted from the following source, available under : Pandey A, Gangopadhyay AN, Kumar V. High anorectal malformation in a five-month-old boy: a case report. *J Med Case Reports.* 2010; 4: 296. DOI: 10.1186/1752-1947-4-296.
- 651 Varicocele.** This image is a derivative work, adapted from the following source, available under : Mak CW, Tzeng WS. Sonography of the scrotum. DOI: 10.5772/27586.
- 652 Scrotal masses: Image A.** Congenital hydrocele. This image is a derivative work, adapted from the following source, available under : Leonardi S, Barone P, Gravina G, et al. Severe Kawasaki disease in a 3-month-old patient: a case report. *BMC Res Notes.* 2013;6:500. DOI: 10.1186/1756-0500-6-500.

Respiratory

- 661 Alveolar cell types: Image A.** Electron micrograph of type II pneumocyte. This image is a derivative work, adapted from the following source, available under : Fehrenbach H, Tews S, Fehrenbach A, et al. Improved lung preservation relates to an increase in tubular myelin-associated surfactant protein A. *Respir Res.* 2005 Jun 21;6:60. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 661 Alveolar cell types: Image B.** Micrograph of type II pneumocyte. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Thomas Cacceti.
- 661 Neonatal respiratory distress syndrome: Image A.** This image is a derivative work, adapted from the following source, available under : Alorainy IA, Balas NB, Al-Boukai AA. Pictorial essay: infants of diabetic mothers. *Indian J Radiol Imaging.* 2010;20:174-181. DOI: 10.4103/0971-3026.69349.
- 663 Lung anatomy: Image A.** X-ray of normal lung. This image is a derivative work, adapted from the following source, available under : Namkoong H, Fujiwara H, Ishii M, et al. Immune reconstitution inflammatory syndrome due to *Mycobacterium avium* complex successfully followed up using 18F-fluorodeoxyglucose positron emission tomography-computed tomography in a patient with human immunodeficiency virus infection: A case report. *BMC Med Imaging.* 2015;15:24. DOI 10.1186/s12880-015-0063-2.
- 663 Lung anatomy: Image B.** CT scan of the chest. This image is a derivative work, adapted from the following source, available under : Wang JF, Wang B, Jansen JA, et al. Primary squamous cell carcinoma of lung in a 13-year-old boy: a case report. *Cases J.* 2008 Aug 22;1(1):123. DOI: 10.1186/1757-1626-1-123. The image may have been modified by cropping, labeling, and/or captions. All rights to this adaptation by MedIQ Learning, LLC are reserved.
- 667 Cyanide vs carbon monoxide poisoning.** This image is a derivative work, adapted from the following source, available under : Subhaschandra S, Jatishwor W, Suraj Th. Isolated symmetrical bilateral basal ganglia T2 hyperintensity in carbon monoxide poisoning. *Ann Indian Acad Neurol.* 2008 Oct-Dec; 11(4): 251–253. DOI: 10.4103/0972-2327.44563.
- 671 Rhinosinusitis.** This image is a derivative work, adapted from the following source, available under : Strek P, Zagolski O, Sktdzien

- J. Fatty tissue within the maxillary sinus: a rare finding. *Head Face Med.* 2006;2:28. DOI: 10.1186/1746-160X-2-28.
- 671 **Deep venous thrombosis.** This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 672 **Pulmonary emboli: Image B.** CT scan. This image is a derivative work, adapted from the following source, available under : Dr. Carl Chartrand-Lefebvre. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 672 **Mediastinal pathology.** This image is a derivative work, adapted from the following source, available under : Aga Z, Avelino J, Darling GE. An unusual case of spontaneous esophageal rupture after swallowing a boneless chicken nugget. *Case Rep Emerg Med.* 2016; 2016: 5971656. DOI: 10.1155/2016/5971656.
- 675 **Obstructive lung diseases: Image A.** Lung tissue with enlarged alveoli in emphysema. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert.
- 675 **Obstructive lung diseases: Image B.** CT of centriacinar emphysema.  Courtesy of the Department of Health and Human Services and Dr. Edwin P. Ewing, Jr.
- 675 **Obstructive lung diseases: Image C.** Emphysema histology. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 675 **Obstructive lung diseases: Image D.** Barrel-shaped chest in emphysema. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 675 **Obstructive lung diseases: Image E.** Mucus plugs in asthma. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 675 **Obstructive lung disease: Image F.** Curschmann spirals. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under . Dr. James Heilman.
- 675 **Obstructive lung diseases: Image G.** Charcot-Leyden crystals on bronchial alveolar lavage. This image is a derivative work, adapted from the following source, available under : Gholamnejad M, Rezaie N. Unusual presentation of chronic eosinophilic pneumonia with “reversed halo sign”: a case report. *Iran J Radiol.* 2014 May;11(2):e7891. DOI: 10.5812/iranradiol.7891.
- 675 **Obstructive lung disease: Image H.** Bronchiectasis in cystic fibrosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 675 **Restrictive lung diseases: Image A.** Pulmonary fibrosis. This image is a derivative work, adapted from the following source, available under : Walsh SLF, Wells AU, Sverzellati N, et al. Relationship between fibroblastic foci profusion and high resolution CT morphology in fibrotic lung disease. *BMC Med.* 2015;13:241. DOI: 10.1186/s12916-015-0479-0.
- 676 **Sarcoidosis: Images A.** Kajal B, Harvey J, Alowami S. Melkerson-Rosenthal Syndrome, a rare case report of chronic eyelid swelling. *Diagn Pathol.* 2013; 8: 188. DOI: 10.1186/1746-1596-8-188.
- 676 **Sarcoidosis: Images B (X-ray of the chest) and C (CT of the chest).** X-ray of the chest) and C CT of the chest. This image is a derivative work, adapted from the following source, available under : Lønborg J, Ward M, Gill A, et al. Utility of cardiac magnetic resonance in assessing right-sided heart failure in sarcoidosis. *BMC Med Imaging.* 2013;13:2. DOI: 10.1186/1471-2342-13-2.
- 676 **Inhalational injury and sequelae: Images A (18 hours after inhalation injury) and B (11 days after injury).** This image is a derivative work, adapted from the following source, available under : Bai C, Huang H, Yao X, et al. Application of flexible bronchoscopy in inhalation lung injury. *Diagn Pathol.* 2013;8:174. DOI: 10.1186/1746-1596-8-174.
- 677 **Pneumoconioses: Image A.** Pleural plaques in asbestosis. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 677 **Pneumoconioses: Image B.** CT scan of asbestosis. This image is a derivative work, adapted from the following source, available under : Miles SE, Sandrini A, Johnson AR, et al. Clinical consequences of asbestos-related diffuse pleural thickening: a review. *J Occup Med Toxicol.* 2008;3:20. DOI: 10.1186/1745-6673-3-20.
- 677 **Pneumoconioses: Image C.** Ferruginous bodies in asbestosis. This image is a derivative work, adapted from the following source, available under : Dr. Michael Bonert. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 677 **Pneumoconioses: Image D.** Berylliosis: non-caseous granuloma. *Ann Saudi Med.* 2009 Nov-Dec; 29(6): 485–486. DOI: 10.4103/0256-4947.57175.
- 678 **Mesothelioma.** This image is a derivative work, adapted from the following source, available under : Weiner SJ, Neragi-Miandoab S. Pathogenesis of malignant pleural mesothelioma and the role of environmental and genetic factors. *J Carcinog.* 2008;7:3. DOI: 10.1186/1477-3163-7-3.
- 678 **Acute respiratory distress syndrome: Image A.** This image is a derivative work, adapted from the following source, available under : Pires-Neto RC, Del Carlo Bernardi F, de Araujo PA. The expression of water and ion channels in diffuse alveolar damage is not dependent on DAD etiology. *PLoS One.* 2016; 11(11): e0166184. DOI: 10.1371/journal.pone.0166184.
- 678 **Acute respiratory distress syndrome: Image B.** Bilateral lung opacities. This image is a derivative work, adapted from the following source, available under : Imanaka H, Takahara B, Yamaguchi H, et al. Chest computed tomography of a patient revealing severe hypoxia due to amniotic fluid embolism: a case report. *J Med Case Reports.* 2010;4:55. DOI: 10.1186/1752-1947-4-55.
- 680 **Atelectasis.** This image is a derivative work, adapted from the following source, available under : Khan AN, Al-Jahdali H, Al-Ghanem S, et al. Reading chest radiographs in the critically ill (Part II): Radiography of lung pathologies common in the ICU patient. *Ann Thorac Med.* 2009;4(3):149–157. DOI:10.4103/1817-1737.53349
- 681 **Pleural effusions: Images A and B.** This image is a derivative work, adapted from the following source, available under : Toshikazu A, Takeoka H, Nishioka K, et al. Successful management of refractory pleural effusion due to systemic immunoglobulin light chain amyloidosis by vincristine adriamycin dexamethasone chemotherapy: a case report. *Med Case Rep.* 2010;4:322. DOI: 10.1186/1752-1947-4-322.
- 682 **Pneumothorax: Image A.** This image is a derivative work, adapted from the following source, available under : Miura K, Kondo R, Kurai M, et al. Birt-Hogg-Dubé syndrome detected incidentally by asymptomatic bilateral pneumothorax in health screening: a

- case of a young Japanese woman. *Surg Case Rep.* 2015 Dec; 1: 17. DOI: 10.1186/s40792-015-0014-8.
- 682 Pneumothorax: Image B.** This image is a derivative work, adapted from the following source, available under : Rosat A, Díaz C. Reexpansion pulmonary edema after drainage of tension pneumothorax. *Pan Afr Med J.* 2015; 22: 143. DOI: 10.11604/pamj.2015.22.143.8097.
- 683 Pneumonia: Image A.** This image is a derivative work, adapted from the following source, available under : Yoon BW, Song YG, Lee SH. Severe community-acquired adenovirus pneumonia treated with oral ribavirin: a case report. *BMC Res Notes.* 2017; 10: 47. DOI: 10.1186/s13104-016-2370-2.
- 683 Pneumonia: Image B.** Lobar pneumonia, gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 683 Pneumonia: Image C.** Acute inflammatory infiltrates in bronchopneumonia. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 683 Pneumonia: Image D.** Bronchopneumonia, gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 683 Pneumonia: Image E.** This image is a derivative work, adapted from the following source, available under : Allen CM, AL-Jahdali HH, Irion KL, et al. Imaging lung manifestations of HIV/AIDS. *Ann Thorac Med.* 2010 Oct-Dec; 5(4): 201–216. DOI: 10.4103/1817-1737.69106.
- 684 Lung cancer: Image B.** Adenocarcinoma histology.  Courtesy of the Department of Health and Human Services and the Armed Forces Institute of Pathology.
- 684 Lung cancer: Image C.** Squamous cell carcinoma. This image is a derivative work, adapted from the following source, available under : Dr. James Heilman. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 684 Lung cancer: Image E.** Large cell lung cancer. This image is a derivative work, adapted from the following source, available under : Jala VR, Radde BN, Haribabu B, et al. Enhanced expression of G-protein coupled estrogen receptor (GPER/GPR30) in lung cancer. *BMC Cancer.* 2012;12:624. doi 10.1186/1471-2407-12-624.
- 685 Lung abscess: Image A.** Gross specimen. This image is a derivative work, adapted from the following source, available under : Dr. Yale Rosen. The image may have been modified by cropping, labeling, and/or captions. MedIQ Learning, LLC makes this available under .
- 685 Lung abscess: Image B.** X-ray. This image is a derivative work, adapted from the following source, available under : Courtesy of Dr. Yale Rosen.
- 685 Pancoast tumor.** This image is a derivative work, adapted from the following source, available under : Manenti G, Raguso M, D'Onofrio S, et al. Pancoast tumor: the role of magnetic resonance imaging. *Case Rep Radiol.* 2013; 2013:479120. DOI: 10.1155/2013/479120.
- 685 Superior vena cava syndrome: Images A (blanching of skin with pressure) and B (CT of chest).** This image is a derivative work, adapted from the following source, available under : Shaikh I, Berg K, Kman N. Thrombogenic catheter-associated superior vena cava syndrome. *Case Rep Emerg Med.* 2013; 2013: 793054. DOI 10.1155/2013/793054.

Index

A

- A-a gradient
 by age, 668
 with oxygen deprivation, 669
 restrictive lung disease, 675
- Abacavir, 203
- Abciximab
 Glycoprotein IIb/IIIa inhibitors, 438
 therapeutic antibodies, 122
 thrombogenesis and, 411
- Abdominal aorta
 atherosclerosis in, 302
 bifurcation of, 663
 branches, 363
- Abdominal aortic aneurysm, 302
- Abdominal pain
 bacterial peritonitis, 390
 Budd-Chiari syndrome, 392
 diabetic ketoacidosis and, 347
 ectopic pregnancy, 641
 Henoch-Schönlein purpura, 315
 hypercalcemia, 591
 hyperparathyroidism, 345
 intussusception, 385
 irritable bowel syndrome, 383
 Meckel diverticulum, 618
 pancreas divisum, 360
 pancreatic cancer, 398
 polyarteritis nodosa, 314
 porphyria, 425
 postprandial, 363
 RLQ pain, 384
 RUQ pain, 396
- Abdominal wall
 anatomy, 369
 inguinal hernias, **369**
 ventral defects, 358
- Abducens nerve (CN VI), 506
 ocular motility, 540
 palsy, 541
- Abduction
 arm, 446
 hand, 450
 hip, 451, 453
 passive, abnormal, 454
- Abductor digiti minimi muscle, 450
- Abductor pollicis brevis muscle, 450
- Abetalipoproteinemia, **94**, 414
- Abnormal uterine bleeding (AUB)
 non-structural causes (COEIN), **633**
 structural causes (PALM), 633
- ABO blood classification, 405
 hemolytic disease of the newborn, 405
- Abortion
 ethical situations, 268
 methotrexate for, 440
- Abruptio placentae, 640
 cocaine use, 614
 preeclampsia, 643
- Abscesses, 479
 acute inflammation and, 214
 brain, 156, 177, **180**
 calcification with, 211
 cold staphylococcal, 116
 frontal lobe, 153
Klebsiella spp, 145
 liver, 155, 179
 lung, 685
 necrosis with, 209
Staphylococcus aureus, 135
Toxoplasma gondii, 177
 treatment of lung, 192
 in unvaccinated children, 186
- Absence seizures
 characteristics of, 517
 drug therapy for, 544
- Absolute risk reduction (ARR), 258
- AB toxin, 132
- Abuse
 child, 269, **556**
 intimate partner violence, 269
- Acalculia, 511
- Acalculous cholecystitis, 396
- Acanthocytes, 414
- Acantholysis, 475, 480
- Acanthosis, 475
 psoriasis, 477
- Acanthosis nigricans, 228, 482
 stomach cancer, 379
- Acarbose, 353
- Accessory nerve (CN XI), 506
 arm abduction, 446
 lesion of, 532
- Accessory pancreatic duct, 360, 368
- Accommodation, eye, 506, 535
- Accuracy (validity), 259
- Acebutolol, 245, 319
- ACE inhibitors, **610**
 acute coronary syndromes, 307
 C1 esterase inhibitor deficiency, 107
 dilated cardiomyopathy, 308
 dry cough, 251
 heart failure, 309
 hypertension, 316
 naming convention for, 253
 preload/afterload effects, 284
 teratogenicity, 614
- Acetaldehyde, 72
- Acetaldehyde dehydrogenase, 72
- Acetaminophen, **485**
 vs aspirin for pediatric patients, 485
 free radical injury and, 210
 hepatic necrosis from, 249
 for osteoarthritis, 466
 toxicity effects, 485
 toxicity treatment for, 248
- Acetazolamide, 252, 552, **608**
 pseudotumor cerebri, 521
- Acetoacetate metabolism, 90
- Acetone breath, 347
- Acetylation
 chromatin, 34
 drug metabolism, 232
 histones, 34
 posttranslation, 45
- Acetylcholine (ACh)
 anticholinesterase effect on, 240
 change with disease, 495
Clostridium botulinum inhibition of release, 138
 opioid analgesics, 551
 pacemaker action potential and, 292
- Acetylcholine (ACh) receptor
 agonists, 551
- Acetylcholine receptors, 228, **236**
- Acetylcholinesterase (AChE)
 in amniotic fluid, 491
 malathion and, 200
 neural tube defects and, 491
- Acetylcholinesterase (AChE) inhibitors
 naming convention for, 253
 for neuromuscular junction disease, 472
 toxicity treatment for, 248
- Acetyl-CoA carboxylase
 fatty acid synthesis, 73
 vitamin B₇ and, 68
- Achalasia, **376**
 esophageal cancer and, 378
 LES tone in, 371
- Achilles reflex, 510
- Achilles tendon xanthomas, 301
- Achlorydria
 stomach cancer, 379
 VIPomas, 371
- Achondroplasia, **462**
 chromosome disorder, 64
 endochondral ossification in, 458
 inheritance, 60
- Acid-base physiology, **592**
- Acid-fast oocysts, 177
- Acid-fast organisms, 125, 155
- Acidic amino acids, 81
- Acid maltase, 86
- Acidosis, **592**
 acidemia diuretic effect on, 609
 cardiac contractility in, 284
 hyperkalemia with, 590
 metabolic, 85, 349
 renal tubular, 592
- Acid phosphatase in neutrophils, 406
- Acid reflux
 H₂ blockers for, 399
 proton pump inhibitors for, 399
- Acid suppression therapy, **398**
- Acinetobacter baumannii*
 highly resistant bacteria, 198
 nosocomial infections, **142**
- Acinetobacter* spp
 nosocomial infections, 185
- Acne, 475, 477
 danazol, 658
 tetracyclines for, 192
- Acquired hydrocele (scrotal), 652
- Acrodermatitis enteropathica, 71
- Acromegaly, **339**
 carpal tunnel syndrome, 459
 GH, 329
 octreotide for, 400
 somatostatin analogs for, 328
- Actin
 cytoskeleton, 48
 muscular dystrophies, 61
- Acting out, 554
- Actinic keratosis, 482
 squamous cell carcinoma, 484
- Actinomyces israelii*
 culture requirements of, 127
 oral infections, 186
 pigment production, 128
- Actinomyces* spp
 effects and treatment of, **139**
 penicillin G/V for, 187
- Action/willpower stage, substance addiction, 568
- Activated carrier molecules and form, **75**
- Active errors, 274
- Active immunity, 110
- Acute chest syndrome, 422
- Acute cholestatic hepatitis
 drug reactions and, 249
 macrolides, 193
- Acute coronary syndrome
 ADP receptor inhibitors for, 437
 heparin for, 436
 nitrates for, 318
 treatments for, **307**
- Acute cystitis, 594, **600**
- Acute disseminated (postinfectious) encephalomyelitis, 524
- Acute dystonia, 569
 treatment of, 241
- Acute gastritis, 379
- Acute hemolytic transfusion reactions, 114
- Acute hemorrhagic cystitis, 164
- Acute inflammation, **214**

- Acute inflammatory demyelinating polyradiculopathy, **524**
- Acute intermittent porphyria, 425
- Acute interstitial nephritis, **601**
- Acute kidney injury, **601**
- Acute laryngotracheobronchitis, 170
- Acute lymphoblastic leukemia (ALL), 432
- methotrexate for, 440
- oncogenes and, 224
- Acute mesenteric ischemia, 386
- Acute myelogenous leukemia (AML), 432
- chromosomal translocations, 434
- cytarabine for, 440
- myelodysplastic syndromes, 432
- Acute pancreatitis, **397**
- hyperparathyroidism, 345
- necrosis and, 209
- Acute pericarditis, **313**
- Acute-phase proteins, 108, 213
- Acute-phase reactants, **213**
- IL-6, 108
- Acute poststreptococcal glomerulonephritis, 596
- Acute promyelocytic leukemia
- vitamin A for, 66
- Acute pulmonary edema
- opioid analgesics, 551
- Acute pyelonephritis, 600
- WBC casts in, 594
- Acute respiratory distress syndrome (ARDS), **678**
- eclampsia and, 643
- Acute stress disorder, 564
- Acute transplant rejection, 119
- Acute tubular necrosis, **602**
- Acyclovir, **201**
- Adalimumab, 122, 487
- for Crohn disease, 382
- Adaptive immunity, 99
- Addiction, stages of change in
- overcoming, 568
- Addison disease, 349
- HLA subtype association, 100
- Additive effect
- of drugs, 235
- Adduction
- arm (rotator cuff), 446
- hand, 450
- hip, 451
- passive, abnormal, 454
- thigh, 452
- Adductor brevis, 452
- Adductor longus, 451, 452
- Adductor magnus, 452
- Adenine
- Shiga/Shiga-like toxins and, 132
- Adenocarcinomas
- carcinogens causing, 225
- esophagus, 378
- gastric, 216, 226
- lung, 224, 684
- nomenclature for, 220
- nonbacterial thrombotic endocarditis and, 228
- pancreas, 226, 368, 398
- paraneoplastic syndromes, 228
- pectinate line, 366
- prostatic, 654
- stomach, 379
- Adenohypophysis, 327
- embryologic derivatives, 613
- hypothalamus and, 498
- Adenomas
- bone, 464
- colorectal, 389
- nomenclature for, 220
- salivary gland, 376
- thyroid, 342
- Adenomatous colonic polyps, 387
- Adenomyosis (endometrial), 648
- uterine bleeding from, 633
- Adenopathy
- Kawasaki disease, 314
- Adenosine
- as antiarrhythmic drug, 324
- blood flow regulation, 297
- pacemaker action potential and, 292
- Adenosine deaminase deficiency, 37, 117
- Adenosine triphosphate (ATP)
- electron transport chain, 78
- production of, 74, 78
- in TCA cycle, 77
- in urea cycle, 82
- Adenosine triphosphate (ATP) synthase inhibitors, 78
- Adenoviruses
- characteristics of, 164
- conjunctivitis, 534
- pneumonia, 683
- Adherens junctions, 474
- Adhesions, 386
- Adipose lipolysis, 320
- Adipose tissue
- estrogen production, 630
- in starvation, 91
- Adjustment disorder, 564
- Adnexal (ovarian) torsion, **625**
- Adoption studies, 256
- ADPKD (Autosomal dominant polycystic kidney disease)
- saccular aneurysms and, 516
- ADP receptor inhibitors, **437**
- ADP ribosyltransferases, 132
- Adrenal adenomas
- Cushing syndrome, 348
- hyperaldosteronism, 349
- Adrenal carcinomas
- Li-Fraumeni syndrome, 224
- P-glycoprotein in, 227
- Adrenal cortex, **327**
- progesterone production, 630
- Adrenal hemorrhage
- Waterhouse-Friderichsen syndrome, 349
- Adrenal hyperplasia
- Cushing syndrome, 348
- hyperaldosteronism and, 349
- Adrenal insufficiency
- adrenoleukodystrophy, 47
- anovulation with, 645
- fludrocortisone for, 354
- mechanism and types of, **349**
- vitamin B₅ deficiency, 67
- Adrenal medulla, **327**
- neuroblastomas of, 350
- pheochromocytomas in, 350
- Adrenal steroids, **335**
- Adrenal zona fasciculata, 336
- Adrenocortical atrophy
- Addison disease, 349
- exogenous corticosteroids, 348
- Adrenocortical insufficiency
- drug reaction and, 249
- Adrenocorticotrophic hormone (ACTH)
- adrenal cortex regulation of, 327
- in Cushing syndrome, 228, 348
- secretion of, 327
- signaling pathways of, 337
- Adrenoleukodystrophy, 47, 524
- Adults
- common causes of death, 272
- primary brain tumors, **526**
- Adult T-cell leukemia, 226
- Adult T-cell lymphoma, 430
- Advance directives, **266**
- Aedes* mosquitoes
- yellow fever transmission, 168
- Aerobes, **126**
- Aerobic metabolism
- ATP production, 74
- fed state, 91
- vitamin B₁ (thiamine), 66
- Aerobic organisms
- culture requirements, 126
- Afferent arteriole, 580
- ANP/BNP effect on, 588
- constriction of, 583
- Afferent nerves, 296
- Aflatoxins, 153
- as carcinogen, 225
- AFP, 117, 653
- African sleeping sickness, 156
- Afterload
- cardiac output, 284
- hydralazine, 318
- in shock, 310
- Agammaglobulinemia
- chromosome affected, 64
- Agars (bacterial culture), 126
- Agensis, 613
- Müllerian, 622
- uterovaginal, 639
- Age-related amyloidosis, 212
- Age-related macular degeneration (ARMD), **536**
- Aging changes, 270
- Agnosia, 511
- Agonists
- indirect cholinomimetic, 234
- indirect general, 242
- indirect sympathomimetics, 242
- partial, 234
- potency and efficacy, 234
- Agoraphobia, 563
- Agranulocytosis, 573
- drug reaction and, 250
- sulfa drug allergies, 252
- thionamides, 354
- Agraphia, 511
- AIDS (acquired immunodeficiency syndrome)
- bacillary angiomatosis, 478
- brain abscess, 180
- Candida albicans*, 153
- cryptococcal meningitis, 199
- Cryptosporidium*, 155
- Cytomegalovirus (CMV), 165
- human herpesvirus 8, 165
- marijuana for, 571
- mycobacteria, 140
- Pneumocystis jirovecii*, 154
- primary central nervous system lymphoma, 430
- retinitis, 165
- retroviruses, 167
- time course (untreated), 176
- Air emboli, 672
- Airways (conducting zone), 662
- Akathisia, 499, 519
- antipsychotic drugs and, 573
- Akinesia, 520
- ALA dehydratase, 419, 425
- Alanine
- ammonia transport, 82
- gluconeogenesis in starvation, 91
- pyruvate dehydrogenase complex deficiency, 77
- Alanine aminotransferase (ALT), 77
- hepatitis viruses, 172
- in liver damage, 390
- toxic shock syndrome, 135
- Alar plate, 490
- Albendazole
- cestodes, 160
- Albinism, 476
- locus heterogeneity, 57
- ocular, 61
- Albumin, 213
- calcium and, 333
- as liver marker, 390
- Albuminocytologic dissociation (CSF), 524
- Albuterol, 242
- asthma, 687
- Alcohol dehydrogenase, 72
- Alcohol exposure
- in utero, 300
- Alcoholic cirrhosis, **391**
- cholelithiasis and, 396
- Alcoholic hepatitis, 391
- Alcoholic liver disease, **391**
- Alcoholism, 145
- anemia, 420
- cataracts and, 535
- cirrhosis and, 389
- common organisms affecting, 179
- esophageal cancer, 378
- ethanol metabolism and, 72
- folate deficiency, 420
- gastritis in, 379
- hepatitis, 367
- hypertension and, 300
- ketone bodies in, 90
- Korsakoff syndrome, 558
- liver serum markers in, 390
- magnesium levels in, 332
- Mallory-Weiss syndrome in, 377
- osteonecrosis in, 463
- pancreatitis, 249, 397
- porphyria, 425
- sideroblastic anemia, 419
- vitamin B₁ deficiency, 66
- vitamin B₉ deficiency, 68
- Alcohol-related disorders
- readmissions with, 272
- Alcohol use
- essential tremor, 519
- gout and, 467
- head and neck cancer, 671
- intoxication and withdrawal, 570
- sleep, 497
- teratogenic effects, 614
- Alcohol use disorder, **571**
- Alcohol withdrawal, 570
- drug therapy, 546, 572
- hallucinations in, 559, 570
- Aldesleukin, 121
- Aldolase B, 80
- Aldose reductase, 81
- Aldosterone, 588, 590
- adrenal cortex secretion of, 327
- SIADH, 338
- signaling pathways for, 337
- Aldosterone antagonists, 316
- Aldosterone resistance, 593
- Aldosterone synthase, 335

- Alemtuzumab, 122
 Alendronate, 486
 Alexia, 515
 Alirocumab, 320
 Aliskiren, **610**
 Alkaline phosphatase (ALP), 390, 463
 bone disorder lab values, 464
 hyperparathyroidism and, 345
 Paget disease of bone, 463
 in thyroid storm, 342
 as tumor marker, 226
 Alkalosis, **592**
 contraction, 60
 diuretic effects, 609
 hypokalemia with, 590
 metabolic, 349
 Alkaptonuria, **84**
 ALK gene, 224
 lung cancer, 684
 Alkylating agents, **441**
 as carcinogens, 225
 targets of, 438
 teratogenicity of, 614
 Allantois, 618
 Allelic heterogeneity, 57
 Allergic bronchopulmonary
 aspergillosis (ABPA), 153
 Allergic contact dermatitis, 477
 Allergic reactions, 112
 blood transfusion, 114
 All-trans retinoic acid
 for promyelocytic leukemia, 66
 Allopurinol
 for gout, 487
 kidney stones, 598
 Lesch-Nyhan syndrome, 37
 rash with, 250
 Alopecia
 doxorubicin, 439
 etoposide/teniposide, 442
 minoxidil for, 658
 syphilis, 147
 tinea capitis, 152
 vitamin A toxicity, 66
 vitamin B₅ deficiency, 67
 vitamin B₇ deficiency, 68
 α -1,4-glucosidase
 glycogen metabolism, 86
 α ₁-antagonists
 BPH treatment, 654
 α ₁-antitrypsin, 52
 α ₁-antitrypsin deficiency, 51, 392
 emphysema, 674
 α ₁-blockers
 tamsulosin, example of, 237
 α ₁ selective blockers, 244
 α ₂-agonists, 243
 α ₂-antagonists, 576
 α ₂ selective blockers, 244
 α -adrenergic agonists, 686
 α -agonists
 glaucoma treatment, 552
 muscle spasm treatment, 551
 α -amanitin
 RNA polymerase inhibition, 42
 α -amylase, 373
 α -antagonists
 for pheochromocytomas, 350
 α -blockers, 244
 Beers criteria, 247
 for cocaine overdose, 571
 nonspecific, 244
 α cells, 328
 glucagonomas in, 351
 glucagon production by, 333
 α -dystroglycan
 muscular dystrophy, 61
 α -fetoprotein
 as tumor marker, 226
 in hepatocellular carcinoma, 392
 neural tube defects, 491
 α -galactosidase A
 Fabry disease, 88
 α -glucosidase inhibitors, 353
 α -hemolytic cocci
 viridans group streptococci, 136
 α -hemolytic bacteria
 Streptococcus pneumoniae, 136
 α -intercalated cells, 593
 α -ketoglutarate
 hyperammonemia and, 82
 α -ketoglutarate dehydrogenase
 metabolic pathways, 74
 TCA cycle, 77
 vitamin B₁ and, 66
 α -methyl dopa, 243
 anemia and, 423
 α -oxidation, 47
 Alpha rhythm (EEG), 497
 α -synuclein, 520
 α -thalassemia, 418
 Alpha toxin, 133
 α -toxin
 Clostridium botulinum, 138
 α (type I) error, 263
 Alport syndrome, 596
 cataracts and, 535
 collagen deficiency in, 50
 Alprazolam, 546
 Alteplase (tPA), 413, 437
 Alternative hypothesis, 262
 Alternative splicing, 43
 Altitude sickness, 670
 Altruism, 555
 Aluminum hydroxide, 399
 Alveolar cell types, **661**
 Alveolar dead space, 664
 Alveolar gas equation, **668**
 Alveolar macrophages, 661, 662
 Alveolar PO₂, 668
 Alveolar sacs, 662
 Alveolar stage (development), 660
 Alveolar ventilation, 664
 Alveoli, 660
 pneumocytes, 661
 Alzheimer disease, 520
 amyloidosis in, 212
 drug therapy for, 240, 549
 neurotransmitters for, 495
Amanita phalloides
 necrosis caused by, 42, 249
 RNA polymerase inhibition, 42
 Amantadine, 548
 Ambiguous genitalia
 46,XY DSD, 639
 ovotesticular disorder of sex
 development, 638
 placental aromatase deficiency, 639
 Amebiasis
 Entamoeba histolytica
 amebiasis, 155
 Amenorrhea
 antiandrogens, 658
 cystic fibrosis, 60
 ectopic pregnancy and, 641
 functional hypothalamic, 645
 menopause diagnosis, 636
 Müllerian agenesis, 622
 pituitary prolactinomas, 328
 Amides (local anesthetics), 550
 Amikacin, 191
 Amiloride, 609
 for diabetes insipidus, 338
 Amines
 MAO inhibitors, 575
 Amine whiff test, 148
 Amino acids
 blood-brain barrier and, 496
 branched, 84
 classification of, **81**
 codons for, 37
 derivatives of, **83**
 genetic code for, 37
 in histones, 34
 metabolism, 90
 purine synthesis, 35
 tRNA, 44
 urea cycle, 82
 Aminoacyl-tRNA, 45
 Aminoglycosides, **191**
 magnesium levels and, 332
 pregnancy use, 204
 Pseudomonas aeruginosa, 143
 teratogenicity, 614
 toxicity of, 251
 Aminopenicillins
 mechanism and use, 188
 Amiodarone, 323
 hypothyroidism, 249
 hypothyroidism with, 341
 photosensitivity, 250
 pulmonary fibrosis, 251
 Amitriptyline, 575
 migraine headaches, 518
 Amlodipine, 318
 Ammonia
 Ornithine transcarbamylase
 deficiency, 83
 Ammonia transport, 82
 Ammonium chloride
 overdose treatment, 233
 Ammonium magnesium phosphate
 (kidney stones), 598
 Amnesia
 brain lesions, 511
 classification of, **558**
 Amnionitis
 Listeria monocytogenes, 139
 Amniotic fluid abnormalities, **641**
 Amniotic fluid emboli, 672
 Amniotic fluid tests
 AChE in, 491
 with neural tube defects, 491
 Amoxapine, 575
 Amoxicillin
 clinical use, 188
 Haemophilus influenzae, 142
 Helicobacter pylori, 146
 Lyme disease, 146
 prophylaxis, 198
 Amphetamines, 242
 intoxication and withdrawal, 570
 narcolepsy treatment, 568
 norepinephrine and, 239
 as weak bases, 233
 Amphotericin B
 clinical use, **199**
 Cryptococcus neoformans, 153
 Naegleria fowleri, 156
 opportunistic fungal infections,
 153
 systemic mycoses, 151
 Ampicillin
 Clostridium difficile, 138
 Listeria monocytogenes, 139
 mechanism and use, 188
 meningitis, 180
 prophylaxis, 198
 Ampulla of Vater, 368
 Amygdala
 lesions of, 511
 limbic system, 499
 Amylase in pancreatitis, 397
 Amylin analogs, 353
 Amyloid angiopathy
 intraparenchymal hemorrhage, 513
 Amyloidosis
 cardiomyopathy with, 308
 carpal tunnel syndrome, 459
 classification, **212**
 kidney deposition in, 597
 with rheumatoid arthritis, 466
 Amyloid precursor protein (APP), 520
 Amyotrophic lateral sclerosis (ALS)
 drug therapy for, 549, 551
 spinal cord lesions, 530
 Anabolic steroids
 hepatic adenomas and, 392
 Anaerobic metabolism
 glycolysis, 74
 pyruvate metabolism, 77
 Anaerobic organisms
 aspiration and, 179
 clindamycin, 192
 Clostridia (with exotoxins), 138
 culture requirements, **127**
 glycyclines, 192
 metronidazole, 195
 Nocardia vs *Actinomyces*, 139
 overgrowth in vagina, 148
 pneumonia caused by, 179
 Anal atresia, 614
 Anal cancer
 HIV and, 177
 oncogenic microbes and, 226
 Anal fissures, 366
 Anal wink reflex, 510
 Anaphase, 46
 Anaphylaxis, 112
 blood transfusion, 114
 complement and, 106
 epinephrine for, 242
 IgA-containing products, 116
 shock with, 310
Anaplasma spp
 Gram stain, 125
 transmission, 146, 149
 Anaplasmosis
 Anaplasma spp, 150
 Anaplastic thyroid carcinomas, 343
 Anastomoses, 503
 Anastrozole, 656
 Anatomic dead space, 664
 Anatomic snuff box, 449
 Anatomy
 endocrinal, 327–328
 gastrointestinal, 360–369
 of heart, 276, **283**
 hematologic/oncologic, 406–409
 musculoskeletal, 446–454
 nervous system, 493–510
 renal, 580
 reproductive, 624–627
 respiratory, 662–663
 “anchovy paste” exudate, 155
Ancylostoma, 159
 diseases associated with, 161
 infection routes, 158
 microcytic anemia, 161
 Androblastoma, 653

- Androgen-binding protein
Sertoli cell secretion, 628
- Androgenetic alopecia, 658
- Androgenic steroid abuse, 636
- Androgen insensitivity syndrome, **639**
- Androgen receptor defect, 639
- Androgens, source and functions, **636**
- Androstenedione, 335, 636
- Anemia, **417**
amphotericin B, 199
Ancylostoma, 161
babesiosis, 157
bacterial endocarditis, 311
blood oxygen content, 666
blood transfusion therapy, 429
blood viscosity in, 286
cephalosporins, 189
chloramphenicol, 192
colorectal cancer, 388
cytarabine and, 440
dapson, 194
Diphyllobothrium latum, 160
drug reaction and, 250
Escherichia coli, 145
ESR in, 214
extrinsic hemolytic, 423
G6PD deficiency, 79
hookworms, 159
in hypertensive emergency, 300
intrinsic hemolytic, 422
isoniazid, 197
kwashiorkor, 71
lab values, 419
macrocytic, 419
malaria, 157
megaloblastic, 420
microcytic, hypochromic, 418
nonhemolytic, normocytic, 421
normocytic, normochromic, 421
NRTIs, 203
oxygen deprivation and, 669
penicillin G, V, 189
pernicious anemia, 372, 379
pregnancy and, 633
pure red cell aplasia, 228
recombinant cytokines for, 121
renal failure, 603
sideroblastic, 67, 419
spherocytes in, 415
in sulfa drug allergies, 252
taxonomy, 417
thioamides causing, 354
trimethoprim, 194
tropical sprue, 381
vitamin B₁₂ deficiency, 69
vitamin B₉ deficiency, 68
Weil disease, 147
Wilson disease, 395
- Anemia of chronic disease, 421
rheumatoid arthritis, 466
- Anencephaly, 491
polyhydramnios and, 641
- Anergy, **110**
- Anesthetics
general principles, **549**
inhaled, 550
intravenous, 550
local, 550
- Aneuploidy, 638
- Aneurysms, **516**
atherosclerosis, 302
coarctation of aorta, 299
Ehlers-Danlos syndrome, 51
superior vena cava syndrome, 685
ventricular, 305, 307
- Angelman syndrome
chromosome association, 64
- Angina
aortic stenosis, 291
atherosclerosis, 302
cocaine causing, 571
contraindicated drugs, **323**
drug therapy for, 318, 324
glycoprotein IIb/IIIa inhibitors
for, 438
ischemic disease and, 304
unstable/NSTEMI treatment, 307
- Angina, "intestinal," 386
- Angina pectoris
β-blockers for, 245
- Angiodysplasia, 386
- Angioedema, 610
C1 esterase inhibitor deficiency,
107
scombroid poisoning, 247
- Angiogenesis
bevacizumab and, 442
wound healing and, 216
- Angiokeratomas, 88
- Angiomas, 117
- Angiosarcomas, 392, 478
carcinogens causing, 225
nomenclature for, 220
- Angiotensin-converting enzyme
(ACE) inhibitors, **610**
- Angiotensin II, 588, 590
ACE inhibitor effects on, 610
signaling pathways for, 337
- Angiotensin II receptor blockers, **610**
heart failure, 309
hypertension, 316
naming convention for, 253
preload/afterload effects, 284
- Anhedonia, 561
- Anhidrosis
Horner syndrome, 540
- Anidulafungin, 200
- Aniline dyes, 606
transitional cell carcinoma and,
606
- Anisocytosis, 407
- Anitschkow cells, 312
- Ankle sprains, **455**
- Ankylosing spondylitis, 469
HLA-B27 and, 100
therapeutic antibodies for, 122
TNF-α inhibitors for, 487
- Annular pancreas, 360
- Anopheles* mosquito, 157
- Anopia
visual field defects, 542
- Anorectal varices
portal circulation, 365
- Anorexia
hypothalamus and, 498
liver cancer/tumors, 392
pancreatic adenocarcinoma, 398
renal failure, 603
- Anorexia nervosa
anovulation with, 645
characteristics of, 567
- Anosmia
zinc deficiency, 71
- ANOVA tests, 264
- Anovulation
common causes, 645
- Antacids, **399**
- Antagonists
ADH, 254
endothelin receptor, 254
ethanol antidote, for, 235
- H₂, 254
nonselective, 245
of drugs, 235
- Anterior cerebral artery
cingulate herniation, 529
cortical distribution, 502
stroke, 514
- Anterior circulation strokes, 514
- Anterior communicating artery
saccular aneurysm, 516
- Anterior corticospinal tract, 508
- Anterior cruciate ligament (ACL)
injury
anterior drawer sign in, 454
"unhappy triad," 460
- Anterior drawer sign, 454
- Anterior hypothalamus, 498
- Anterior inferior cerebellar artery, 514
- Anterior inferior tibiofibular
ligament, 455
- Anterior pituitary gland, 327, 331
- Anterior spinal artery
complete occlusion, 530
stroke, 514
- Anterior talofibular ligament, 455
- Anterograde amnesia, 558
benzodiazepines, 550
brain lesions, 511
- Anthracois, 677
- Anthracyclines, 439
cardiomyopathy from, 248
- Anthrax, 132, 137
- Anthrax toxin
Bacillus anthracis and, 137
- Anti-ACh receptor antibody, 115
- Antiandrogen drugs, **658**
- Antianginal therapy, 307, 318, **319**,
324
- Antiapoptotic molecule
oncogene product, 224
- Antiarrhythmic drugs
mechanisms and clinical uses,
322–324
torsades de pointes, 248
- Antibiotics, 153
acne treatment, 477
Clostridium difficile with, 138
Jarisch-Herxheimer reaction with,
148
long QT interval, 294
selective growth media, 126
torsades de pointes, 248
- Antibodies
in adaptive immunity, 99
antibody diversity generation, 104
antibody specificity generation, 104
exo- and endotoxins, 131, 133
hepatitis viruses, 174
hypersensitivity mediation, 112
structure and function, **104**
therapeutic, 122
- Antibody-dependent cell-mediated
cytotoxicity, 101
- Anticardiolipin
antiphospholipid syndrome, 470
- Anticardiolipin antibody, 115
- Anti-CCP antibody, 115
- Anti-centromere antibodies
scleroderma, 473
- Anticentromere autoantibody, 115
- Anticholinergic drugs
delirium with, 558
toxicity treatment for, 248
- Anticholinesterase drugs, 240
- Anticholinesterase poisoning, **240**
- Anticipation (genetics), 56
- Anticoagulant drugs
acute coronary syndromes, 307
antiphospholipid syndrome, 470
atrial fibrillation, 295
coagulation cascade and, 412
- Anticoagulation
coagulation cascade and, 413
- Anticonvulsant drugs, 471
- Antidepressant drugs, **574–576**
atypical, 576
for fibromyalgia, 471
long QT interval with, 294
torsades de pointes, 248
- Anti-desmoglein (anti-desmosome)
autoantibody, 115
- Anti-digoxin Fab fragments, 248
for cardiac glycoside toxicity, 321
- Antidiuretic hormone (ADH), **329**,
588, 590
antagonist naming conventions,
254
antagonists, 338, 354
in diabetes insipidus, 338
function of, 328
hypothalamus synthesis, 498
pituitary gland and, 327
SIADH and, 338
signaling pathways of, 337
- Anti-DNA topoisomerase I
autoantibody, 115
- Anti-dsDNA antibody, 115
- Antiemetic drugs
aprepitant, 401
long QT interval with, 294
marijuana, 571
metoclopramide, 400
ondansetron, 400
torsades de pointes, 248
- Antiepileptic drugs
Cytochrome P-450 interactions,
252
for fibromyalgia, 471
rash from, 250
teratogenicity, 614
- Antifungal drugs
griseofulvin, 48
mechanism and use, **198–200**
seborrheic dermatitis, 476
tinea versicolor, 152
- Antigenic shift/drift, 169
- Antigen-presenting cells (APCs)
B cells as, 409
CD28, 110
MHC I and II and, 100
naive T-cell activation, 103
in spleen, 98
- Antigens
active immunity, 110
antibody structure and function,
104
chronic mucocutaneous
candidiasis, 116
for self, 102
HLA I and II, 100
lymphocyte recognition of, 98
type and memory, **105**
- Anti-glomerular basement membrane
autoantibody, 115
- Anti-glutamic acid decarboxylase
autoantibody, 115
- Antigout drugs
colchicine, 48
- Anti-HBc, 174
- Anti-HBe, 174
- Anti-HBs, 174
- Anti-helicase autoantibody, 115

- Antihelminthic drugs, **200**
 mebendazole, 48
 naming convention, 253
- Anti-hemidesmosome autoantibody, 115
- Antihistamines, 686
 for scombroid poisoning, 247
- Anti-histone antibody, 115
- Antihypertensive drugs
 hypertension in pregnancy, 643
- Anti-IgE monoclonal therapy, 687
- Anti-IL-5 monoclonal therapy, 687
- Anti-inflammatory drugs, 485
- Anti-intrinsic factor autoantibody, 115
- Anti-La/SSB autoantibody, 115
- Antileukotrienes
 for asthma, 687
- Antimetabolites, **440**
- Antimicrobial drugs
 highly resistant bacterial treatment, 198
 mechanisms of action summary, **187–204**
 naming conventions for, 253
 pregnancy contraindications, **204**
 prophylaxis, **198**
- Antimicrosomal autoantibody, 115
- Anti-mite/ouse therapy, **200**
- Antimitochondrial autoantibody, 115
- Antimuscarinic drugs
 Parkinson disease, 548
 reactions to, 251
 toxicity treatment for, 248
- Antimycin A
 electron transport chain, 78
- Antimycobacterial therapy, **196**
- Anti-NMDA receptor paraneoplastic syndrome encephalitis, 228
- Antinuclear (ANA) antibody, 115
 Sjögren syndrome, 468
- Antioxidants
 free radical elimination by, 210
- Antiparasitic drugs
 naming convention for, 253
- Antiparietal cell autoantibody, 115
- Anti-phospholipase A2 receptor autoantibody, 115
- Antiphospholipid syndrome, **470**
 autoantibody in, 115
- Antiplatelet antibodies
 abciximab as, 122
- Antiplatelet drugs
 for acute coronary syndromes, 307
- Anti-presynaptic voltage-gated calcium channel autoantibody, 115
- Antiprogesterin drugs, **657**
- Antiprotozoan drugs, **200**
- Antipseudomonal drugs
 cephalosporins, 189
 fluoroquinolones, 195
 penicillins, **188**
- Antipsychotic drugs
 adverse effects/events, 573
 antimuscarinic reaction, 251
 atypical, 573
 dopaminergic pathways, 499
 dystonia with, 569
 long QT interval with, 294
 Parkinson-like syndrome, 251
 tardive dyskinesia, 251
 torsades de pointes, 248
 Tourette syndrome, 572
 typical, 573
- Antiribonucleoprotein antibodies
 Sjögren syndrome, 468
- Anti-Ro/SSA autoantibody, 115
- Anti-Scl-70 autoantibody, 115
- Anti-Smith autoantibody, 115
- Anti-smooth muscle autoantibody, 115
- Antisocial personality disorder, 565
 early-onset disorder, 557
- Antispasmodics, 551
- Anti-SRP autoantibody, 115
- Anti-streptolysin O (ASO) titers, 312
- Antisynthetase autoantibody, 115
- Antithrombin
 coagulation cascade and, 413
 deficiency of, 428
- Antitoxins
 as passive immunity, 110
- Anti-TSH receptor autoantibody, 115
- Antitumor antibiotics, **439**
- Anti-U1 RNP antibodies, 115, 470
- Antiviral therapy
 hepatitis C, 203, 204
 mechanism and use, **201**
- Anti- β_2 glycoprotein antiphospholipid syndrome, 470
 autoantibody, 115
- Anxiety
 drug therapy, 546, 563, 575
 neurotransmitters, 495
- Aorta
 branches, 363
 coarctation of, 299, 300
 congenital heart disease, 298
 diaphragm, 663
 ECG and, 293
 retroperitoneal, 360
 in syphilitic heart disease, 312
 traumatic rupture of, 303
 “tree bark” appearance, 312
- Aortic aneurysm, 302
 Ehlers-Danlos syndrome, 51
 hypertension, 300
 Marfan syndrome, 300
 syphilitic heart disease, 312
- Aortic arch derivatives, **619**
- Aortic arch receptors, 296
- Aortic dissection, **303**
 hypertension, 300
 Marfan syndrome, 300
- Aortic insufficiency
 syphilitic heart disease, 312
- Aorticopulmonary septum, 281
 embryologic derivatives, 613
- Aortic regurgitation, 288
 heart murmurs with, 291
 Marfan syndrome, 300
- Aortic root dilation
 heart murmur with, 291
- Aortic stenosis, 288
 heart murmurs, 291
 macroangiopathic anemia, 423
 Williams syndrome, 300
- Aortic valve
 cardiac cycle, 287
 embryological development of, 281
- Aortitis
 syphilis, 147, 184
- Apalutamide, 658
- APC gene, 224
 adenomatous colonic polyps and, 387
 colorectal cancer and, 389
 familial adenomatous polyposis and, 387
 “Ape hand,” 447, 451
 Apgar score, **634**
- Aphasia
 MCA stroke, 514
 types of, **516**
- Apthous stomatitis
 Crohn disease, 382
- Apixaban
 factor Xa inhibitors, 437
- Aplasia, 613
- Aplasia cutis
 methimazole, 354
- Aplastic anemia, 421
 chloramphenicol, 192
 drug reaction and, 250
 neutropenia with, 424
 thionamides, 354
- Aplastic crisis
 hereditary spherocytosis, 422
 sickle cell anemia, 422
- Apolipoproteins
 functions of, 93
- Apoptosis, **208**
 corticosteroids, 424
- Appendages (bacterial), 124
- Appendicitis, **383**
 mittelschmerz vs, 631
- Appetite regulation, **336**, 371
 “Apple core” lesion (X-ray), 388
- Apraclonidine, 552
- Aprepitant, **401**
- Aqueous humor pathway, **535**
- Arabinofuranosyl cytidine, 440
- Arabinogalactan synthesis, 196
- Arabinosyltransferase, 197
- Arachidonic acid pathway, **485**
- Arachnodactyly, 52
- Arachnoid granulations, 503, 504, 522
- Arachnoid mater
 meninges, 496
 meningioma, 526
 ventricular system, 504
- Arcuate fasciculus
 aphasia and, 516
 diagram, 501
- Area postrema, 498
- Area under the curve, 231
- Arenaviruses, characteristics of, 167
- Argatroban, 435
- Arginine
 classification, 81
 cystinuria, 85
 kidney stones and, 598
 Argininosuccinate, 82
- Argyll Robertson pupils
 in syphilis, 184
 tabes dorsalis, 530
- Aripiprazole, 573
- Arm abduction, **446**
- Armadillos (disease vectors), 149
- Aromatase, 636
 in pathway, 335
- Aromatase inhibitors, **656**
- Aromatic amines
 carcinogenicity of, 225
- Arrhythmias
 amphotericin B, 199
 diabetic ketoacidosis, 347
 diphtheria, 139
 drug reactions and, 248
 hypokalemia and, 591
 local anesthetics and, 550
 macrolides, 193
 McArdle disease, 87
 muscular dystrophy, 61
 myocardial infarction and, 305, 307
 shock caused by, 310
 sleep apnea and, 679
- stimulants and, 570
 TCA toxicity, 569
 thyroid hormones and, 354
- Arsenic
 angiosarcomas, 392, 478
 carcinogenicity of, 225
 glycolysis and, 74
 toxicity symptoms, 76
 toxicity treatment, 248
- Artemether, 200
- Arterial oxygen saturation, 666
- Arterial PCO₂, 668
- Arteries, anatomy of, 283
- Arteriolosclerosis, 301, 346
- Arteriosclerosis, **301**
- Arteriovenous malformations (AVMs)
 hereditary hemorrhagic telangiectasia, 316
- Arteriovenous shunts, 463
- Arteritis
 giant cell (temporal), 314, 518
 headaches, 518
- Artesunate
 malaria, 157, 200
- Arthralgias
 alkaptonuria, 84
 coccidiomycosis, 151
 Henoch-Schönlein purpura, 315
 hepatitis viruses, 172
 in alkaptonuria, 84
 rubella, 169, 182
 serum sickness, 113
 vitamin A toxicity, 66
- Arthritis, 457, 466
 azathioprine for, 440
Campylobacter jejuni, 145
 carpal tunnel syndrome and, 459
 celecoxib for, 486
 chlamydiae, 148, 184
 Crohn disease, 382
 gonococcal, 468
 gonorrhea, **142**, 180, 184
 immunosuppressants, 120
 inflammatory bowel disease, 100
 lupus, 470
 Lyme disease, 146
 psoriatic, 469
 reactive arthritis, 469
 septic, 468
Staphylococcus aureus, 135
 Takayasu arteritis, 314
 therapeutic antibodies, 122
 ulcerative colitis, 382
- Arthropathy
 hemochromatosis, 395
- Arthus reaction, 113
- Arylsulfatase A
 metachromatic leukodystrophy, 88
- Arytenoids, 620
- Asbestos
 carcinogenicity, 225
- Asbestosis, **677**, 678
- Ascaris lumbricoides*, 159
- Ascaris* spp, 158
- Ascending cholangitis, 397
- Ascending colon, 360
- Aschoff bodies, 312
- Ascites
 Budd-Chiari syndrome, 392
 hepatocellular carcinoma, 392
 spontaneous bacterial peritonitis, 390
- Ascorbic acid, 69
- Asenapine, 573
- Aseptic meningitis
 mumps, 170
 picornaviruses, 167

- Asherman syndrome, 648
 Ashkenazi Jews
 disease incidence, 88
 ASO titer, 136
 Aspart, 352
 Aspartame, 84
 Aspartate
 urea cycle, 82
 Aspartate aminotransferase (AST), 390
 hepatitis, 172
 toxic shock syndrome, 135
 Aspartic acid, 81
 Aspergillosis
 Aspergillus fumigatus, 153
 bronchiectasis, 674, 675
 echinocandins, 200
Aspergillus fumigatus, 153
 HIV-positive adults, 177
Aspergillus spp
 chronic granulomatous disease, 109
 Aspiration
 ARDS and, 678
 in utero “breathing,” 660
 lung abscess, 685
 reflux-related, 359, 377
 Zenker diverticulum, 384
 Aspiration pneumonia
 alcoholics, 179
 clindamycin, 192
 nosocomial infections, 185
 Aspirin, **486**
 acute coronary syndromes, 307
 cyclooxygenase, 411
 hemolysis in G6PD deficiency, 250
 for ischemic stroke, 512
 Kawasaki disease, 314
 thrombogenesis and, 411
 uncoupling agent, 78
 as weak acid, 233
 zero-order elimination of, 232
 Asplenia
 Streptococcus pneumoniae, 136
 target cells, 415
 Asterixis, 519
 hepatic encephalopathy, 391
 renal failure, 603
 Asteroid bodies, 676
 Asthma, 674
 albuterol for, 242
 β -blockers and, 245
 breast milk and, 636
 cromolyn sodium for, 408
 drug therapy, **687**
 epinephrine for, 242
 gastroesophageal reflux disease, 377
 hypertension treatment with, 316
 immunosuppressants, 120
 muscarinic antagonists for, 241
 omalizumab for, 122
 pulsus paradoxus in, 310
 salmeterol for, 242
 type I hypersensitivity, 112
 Astigmatism, 535
 Astrocytes, **493**
 foot processes, 496
 origin of, 490
 Ataxia
 abetalipoproteinemia, 94
 cerebellar hemisphere lesions, 511
 Friedreich, 62, 64, 531
 hypnotics, 546
 lithium toxicity, 569
 metachromatic leukodystrophy, 88
 normal pressure hydrocephalus, 522
 opsoclonus-myoclonus syndrome, 228
 prion disease, 178
 psychoactive drug intoxication, 570
 streptomycin, 197
 syphilis, 147
 tabes dorsalis, 530
 trinucleotide repeat expansion disease, 62
 truncal, 511
 vitamin B₁₂ deficiency, 530
 vitamin E deficiency, 70
 Wernicke-Korsakoff syndrome, 511, 571
 Ataxia-telangiectasia, 40, **117**
 Atazanavir, 203
 Atelectasis, **680**
 Atenolol, 245, 323
 Atherosclerosis, **302**
 abdominal aortic aneurysms and, 302
 diabetes mellitus and, 346
 familial dyslipidemias, 94
 homocystinuria, 84
 renovascular disease, 604
 stable angina with, 304
 transplant rejection, 119
 Athetosis, 511, 519
 Atomoxetine, 557
 Atonic seizures, 517
 Atopic dermatitis (eczema), 477
 Atopic reactions, 112
 Atovaquone
 babesiosis, 157
 malaria, 157
 P. falciparum, 200
 for *Pneumocystis jirovecii*, 154
 ATPase, 395
 ATP production, **74**
 Atresia
 anal, 614
 duodenal, 359
 esophageal, 359
 intestinal, 359
 jejunal/ileal, 359
 Atria
 cardiac tumors, 316
 depolarization/repolarization of, 293
 embryologic development of, 281
 Atrial fibrillation
 β -blockers for, 323
 calcium channel blockers for, 324
 cardiac glycosides for, 321
 ECG tracing of, 295
 hypertension, 300
 jugular venous pulse in, 287
 potassium channel blockers for, 323
 Atrial flutter
 β -blockers for, 323
 ECG tracing of, 295
 potassium channel blockers for, 323
 “Atrial kick,” 287
 Atrial natriuretic peptide (ANP), **296**, 588, 590
 in SIADH, 338
 signaling pathways for, 337
 Atrial septal defect (ASD), 299
 congenital rubella, 300
 Down syndrome, 300
 fetal alcohol syndrome, 300
 Atrioventricular (AV) block
 β -blockers, 245, 323
 calcium channel blockers, 318, 324
 ECG tracings, 295
 Lyme disease, 146
 Atrioventricular canals, 281
 Atrioventricular node
 AV node, 292
 conduction pathway, 293
 ECG and, 293
 Atrioventricular valves
 embryologic development of, 281
 Atrophic gastritis
 gastrin in, 371
 Atrophy, 206
 motor neuron signs, 529, 531
 neurodegenerative disorders, 520
 optic disc/nerve, 536
 ventriculomegaly, 522
 Atropine, **241**
 antimuscarinic reaction, 251
 for β -blocker overdose, 323
 effects of, 241
 toxicity treatment, 248
 Attention-deficit hyperactivity disorder (ADHD), 557
 drug therapy for, 557, 572
 Tourette syndrome, 557
 Attributable risk (AR), 258
 Atypical antidepressants, **576**
 Atypical antipsychotic drugs, **573**
 postpartum psychosis, 562
 Atypical depression, 575
 Atypical pneumonias
 chlamydiae, 148
 macrolides, 193
 typical organisms, 683
 Auditory cortex
 diagram, 501
 thalamic relay, 498
 Auditory hallucinations, 559
 Auditory physiology, **533**
 Auerbach plexus, 376, 384
 Auer rods
 in AML, 432
 Auramine-rhodamine stain, 125
 Auscultation of heart, **290**
 Auspitz sign, 477
 Autism spectrum disorder, 557
 double Y males and, 638
 fragile X syndrome, 62
 Autoantibodies, **115**
 Autoclaves
 disinfection/sterilization, 204
 Autodigestion, 397
 Autoimmune diseases
 acute pericarditis, 313
 blistering skin, **480**
 collagen and, 50
 Dressler syndrome, 307
 myocarditis with, 313
 rheumatoid arthritis, 466
 self-antigen in, 102
 Sjögren syndrome, 468
 SLE, 470
 Autoimmune gastritis, 379
 Autoimmune hemolytic anemia, 423
 cephalosporins, 189
 Autoimmune hepatitis type 1
 autoantibody, 115
 Autoimmune hypothyroidism, 173
 Autoimmune lymphoproliferative syndrome, 208
 Autoimmune regulator (AIRE), 102
 Autoimmune thrombocytopenia, 121
 Autonomic drugs, **236–245**
 actions of, 239
 bladder dysfunction, action on, 237
 naming conventions for, 253
 Autonomic ganglia, 236
 Autonomic insufficiency, 242
 Autonomic nervous system (ANS)
 delirium tremens, 569
 dysregulation in inflammatory demyelinating polyradiculopathy, 524
 limbic system in, 499
 male sexual response, **627**
 receptors in, **236**
 in serotonin syndrome, 569
 Autonomy (ethics), 265
 Autoregulation of blood flow, **297**
 Autosomal dominant diseases, **60**
 ADPKD, 516
 Brugada syndrome, 294
 Charcot-Marie-Tooth disease, 524
 elastin syndrome, 52
 Huntington disease, 520
 hyper-IgE syndrome, 116
 hypertrophic cardiomyopathy, 308
 malignant hyperthermia susceptibility, 550
 neurofibromatosis, 525
 porphyrias, 425
 Romano-Ward syndrome, 294
 tuberous sclerosis, 525
 von Hippel-Lindau disease, 525
 Autosomal dominant polycystic kidney disease (ADPKD), 604
 chromosome association, 64
 Autosomal dominant tubulointerstitial kidney disease, 604
 Autosomal recessive diseases, **60**
 abetalipoproteinemia, 94
 adenosine deaminase deficiency, 117
 alkaptonuria, 84
 Chédiak-Higashi syndrome, 117
 cystic fibrosis, 60
 5 α -reductase deficiency, 639
 Friedreich ataxia, 531
 hemochromatosis, 395
 hereditary hyperbilirubinemias, 394
 IL-12 receptor deficiency, 116
 Jervell and Lange-Nielsen syndrome, 294
 Kartagener syndrome, 49
 leukocyte adhesion deficiency, 117
 maple syrup urine disease, 84
 severe combined immunodeficiency, 117
 spinal muscular atrophy, 530
 Wilson disease, 395
 Autosomal recessive polycystic kidney disease (ARPKD), 604
 Potter sequence caused by, 578
 Autosomal trisomies, **63**
 Down syndrome (trisomy 21), 63
 Edwards syndrome (trisomy 18), 63
 karyotyping for, 55
 Patau syndrome (trisomy 13), 63
 Avascular necrosis
 femoral head, 461
 Avascular necrosis, **463**
 scaphoid bone, 449
 sickle cell anemia, 422
 Aversive stimulus (positive punishment), 554
 Avoidant personality disorder, 566
 Axilla/lateral thorax, 455
 Axillary nerve
 arm abduction, 446
 injury presentation, 447
 neurovascular pairing, 455

- Axonal injury, 495
 Axonal trafficking, 48
 Axonemal dynein, 49
 Azathioprine
 antimetabolites, 440
 for Crohn disease, 382
 immunosuppressant, 120
 pancreatitis caused by, 249
 Azithromycin
 atypical pneumonia treatment, 148
 babesiosis, 157
 chlamydiae, 148
 in cystic fibrosis, 60
 gonorrhoea treatment, 142
 macrolides, 193
 Mycobacterium avium-intracellulare, 140, 196
 prophylaxis in HIV, 198
 Azoles, 153, **199**
 vaginal infections, 181
 Azotemia
 acute interstitial nephritis, 601
 Aztreonam, 190
- B**
- Babesia* spp, 146, 157
 Babesiosis, 157
 Babinski reflex, 635
 motor neuron signs, 529
 primitive reflexes, 510
 Bachmann bundle, 293
 Bacillary angiomatosis, 478
 animal transmission, 149
 HIV-positive adults, 177
Bacillus anthracis, **137**
 exotoxin production, 132
Bacillus cereus, **138**
 food poisoning, 178
 Bacitracin
 gram-positive antibiotic test, 134
 sensitivity to, 134, 136
 Baclofen, 551
 multiple sclerosis, 523
 Bacteremia
 brain abscesses, 180
 cutaneous anthrax, 137
 daptomycin, 195
 Staphylococcus gallolyticus, 137
 Streptococcus bovis, 137
 Bacteria
 biofilm-producing, 128
 culture requirements, 126
 encapsulated, 127
 genetics, **130**, 131
 hemolytic, 135
 highly resistant, 198
 normal flora, 178
 phage infection of, 130
 pigment-producing, 128
 spore-forming, 129
 stains for, 125
 structures and functions, **124**
 virulence factors, 127, **129**, 135, 143, 144, 145
 zoonotic, 149
 Bacterial capsules, 124
 Bacterial endocarditis, **311**
 daptomycin, 195
 Staphylococcus aureus, 135
 Bacterial exotoxin mechanisms
 increase fluid secretion, **132**
 inhibit phagocytic ability, 132
 inhibit protein synthesis, 132
 lyse cell membranes, 133
 superantigens causing shock, 133
 Bacterial infections
 with immunodeficiency, 118
 myocarditis with, 313
 skin, 479
 Bacterial peritonitis (spontaneous), 389, 390
 Bacterial toxin mechanisms
 inhibit release of neurotransmitter, 132
 lysogenic phage encoding of, 130
 Bacterial vaginosis
 characteristics of, 158, 181
 Gardnerella vaginalis, 148
Bacteroides fragilis, 178
Bacteroides spp
 alcoholism, 179
 clindamycin, 192
 culture requirements of, 127
 metronidazole, 195
 nosocomial infections, 185
 “Bag of worms,” 651
 Baker cyst, 460
 tibial nerve injury, 453
 BAK protein, 208
 Balancing (quality measurement), 273
 Bamboo spine, 469
 Band cells, 406
 Barbiturates
 intoxication and withdrawal, 570
 intravenous anesthetics, 550
 mechanism and use, **546**
 naming convention for, 253
 sleep alterations, 497
 Barlow maneuver, 461
 Baroreceptors, **296**
 Barr bodies, 34
 in x-inactivation, 61
 Barrett esophagus, **378**
 Bartholin cyst/abscess, 644
Bartonella henselae
 bacillary angiomatosis, 478
Bartonella quintana, 161
Bartonella spp
 animal transmission, 149
 Gram Stain, 125
 Bartter syndrome, 586
 markers in, 591
 Basal cell carcinomas, 484
 5-fluorouracil for, 440
 Basal ganglia, **500**
 intraparenchymal hemorrhage, 513
 lesions in, **511**
 movement disorders, 519
 thalamic connections, 498
 Basal lamina, 50
 Basal nucleus of Meynert, 495
 Basal plate, 490
 Base excision repair, 40
 Basement membrane, 98
 blood-brain barrier, 496
 collagen in, 50
 glomerular filtration barrier, 581
 Basic amino acids, 81
 Basilar artery
 herniation syndromes, 529
 stroke effects, 515
 Basilar membrane (cochlea), 533
 Basiliximab
 immunosuppressant, 120
 Basophilia, 408
 Basophilic stippling, 416
 lead poisoning, 419
 sideroblastic anemia, 419
 Basophils, **408**
 IgE antibody, 105
 BAX protein, 208
 B-cell lymphomas
 HIV-positive adults, 177
 B cells, **409**
 activation, 103, 105
 adaptive immunity, 99
 anergy, 110
 cell surface proteins, 110
 class switching, 103
 disorders of, 116, 117
 function of, 409
 functions of, 101
 glucocorticoid effects, 120
 immunodeficiency infections, 118
 in lymph node, 96
 neoplasms, 430
 non-Hodgkin lymphoma, 429
 sirolimus effect, 120
 spleen, 98
 BCG vaccine
 false positives from, 140
 IL-12 receptor deficiency and, 116
 BCL-2 gene, 224
 Bcl-2 protein, 208
 BCR-ABL gene, 224
 Bead-like costochondral junctions, 463
 Becker muscular dystrophy, 61
 Beck triad (cardiac tamponade), 310
 Beckwith-Wiedemann syndrome, 358, 606
 Beers criteria, **247**
 Behavioral therapy, 572
 Behavior modulation
 frontal lobe lesions and, 511
 limbic system and, 499
 Behçet syndrome, 314
 Bell palsy, 532, 676
 Bell-shaped distribution, 262
 Bendazoles, 159
 Bends, 463
 Beneficence (ethics), 265
 Benign breast disease, **649**
 Benign paroxysmal positional vertigo (BPPV), 534
 Benign prostatic hyperplasia (BPH), 237, **654**
 α_1 -blockers for, 244
 epididymitis and orchitis with, 654
 tamsulosin for, 658
 treatment of, 237
 Benign tumors, 220
 bones, 464
 breast, 649
 Benralizumab, 687
 Benzathine penicillin G, 198
 Benzene
 aplastic anemia, 250
 Benzidine as carcinogen, 225
 Benzimidazole, 158
 Benzocaine, 550
 Benzodiazepines
 addictive risk, 546
 alcohol withdrawal, 572
 Beers criteria, 247
 clinical use and adverse effects, **546**
 cocaine overdose, 571
 epilepsy treatment, 544
 intoxication and withdrawal, 570
 naming convention for, 253
 phobias, 563
 sleep effects, 497
 toxicity treatment for, 248
 Benzoyl peroxide for acne, 477
 Benzotropine, 241, 548
 Berger disease, 596
 Beriberi
 vitamin B₁ deficiency, 66
 Berkson bias, 260
 Bernard-Soulier syndrome, 411, 427
 Berry aneurysm, 516
 Berylliosis, 677
 β_1 -blockade, 284
 β_2 -agonists
 naming convention for, 253
 β_2 -agonists
 asthma, 687
 insulin and, 334
 β -adrenergic agonist
 potassium shifts, 590
 β -blockers, 245, **323**
 acute coronary syndromes, 307
 angina, 319
 aortic dissections, 303
 Cardiomyopathy (hypertrophic), 245
 cocaine overdose, 571
 diabetes and, 245
 dilated cardiomyopathy, 308
 essential tremor, 519
 for cocaine overdose, 242
 for pheochromocytomas, 350
 for thyroid storm, 342
 glaucoma treatment, 552
 heart failure, 309
 hyperkalemia, 590
 hypertension, 316
 hypertrophic cardiomyopathy, 308
 juxtaglomerular apparatus effects, 589
 migraine headaches, 518
 naming convention for, 253
 overdose treatment, 323
 phobias, 563
 selectivity, 245
 toxicity treatment for, 248
 β cells, 328
 diabetes mellitus type 1 and 2, 347
 insulinomas of, 351
 insulin production by tumors, 351
 insulin secretion by, 334
 β -dystroglycan, 61
 β -galactosidase, 144
 β -glucan, 200
 β -glucuronidase, 406
 β -hCG, 653
 as tumor marker, 226
 β -hemolysis, 133
 β -hemolytic bacteria, 135
 common colonization sites, 135
 Staphylococcus aureus, 135
 Staphylococcus epidermidis, 135
 Staphylococcus saprophyticus, 136
 Streptococcus agalactiae (Group B strep), 137
 Streptococcus pyogenes (Group A strep), 136
 β -hydroxybutyrate, 90
 β -interferon
 multiple sclerosis, 523
 β -lactam antibiotics, 187
 β -lactamase inhibitors, 189
 β_2 -microglobulin
 MHC I and II and, 100
 β -oxidation of very-long-chain fatty acids (VLCFA), 47
 β -prophage
 Corynebacterium exotoxin encoding, 139
 Beta rhythm (EEG), 497
 β -thalassemia, 418
 allelic heterogeneity, 57
 Betaxolol, 245, 552
 Bethanechol, 240

- Bevacizumab, 122, **442**
 Bezafibrate, 320
 Bias and study errors, **260–262**
 Bicalutamide, 658
 Bicarbonate
 carbon dioxide transport, 670
 GI secretion, 372
 overdose treatment, 233
 pancreatic insufficiency, 381
 salicylate toxicity, 248
 TCA toxicity, 233, 248
 Biceps brachii muscle
 Erb palsy, 448
 Biceps femoris, 452, 453
 Biceps reflex, 510
 Biceps tendon, 446
 Bicornuate uterus, 623
 Bicuspid aortic valve
 aortic dissection and, 303
 coarctation of aorta and, 299
 heart murmur with, 291
 thoracic aortic aneurysms and, 302
 Turner syndrome, 300
 Bifid ureter, 579
 Bifurcation external landmarks, 663
 Biguanide drugs, 353
 Bilaminar disc, 612
 Bilateral adenopathy, 676
 Bilateral renal agenesis
 oligohydramnios and, 641
 Potter sequence, 578
 pulmonary hypoplasia and, 660
 Bile, **374**
 hereditary hyperbilirubinemias, 394
 secretin effect on, 371
 Bile acid resins, 320
 lipid transport and, 92
 reabsorption of, 320
 synthesis of, 47
 Bile canaliculus, 367
 Bile ducts, 367, 368
 Bile salts, 374
 in cholelithiasis, 396
 Biliary atresia, **393**
 Biliary cholangitis, primary
 autoantibody, 115
 Biliary cirrhosis, 389, 393
 cystic fibrosis, 60
 Biliary colic, 396
 Biliary structures, **368**
 Biliary tract disease, **395**
 Clonorchis sinensis, 161
 gallstones, 368
 hyperbilirubinemia with, 393
 Biliary tract infections
 Enterococci, 137
 Bilirubin, **375**
 bile, 374
 cholelithiasis, 396
 hereditary hyperbilirubinemias, 394
 liver marker, 390
 toxic shock syndrome, 135
 Bimatoprost, 552
 Bimodal distribution, 262
 Binge-eating disorder, 567, 575
 Binge-eating/purging, anorexia
 nervosa, 567
 Bioavailability, 231
 area under the curve from, 231
 Biochemistry
 cellular, 46–52
 genetics, 56
 laboratory techniques, 52–94
 metabolism, 72–94
 molecular, 34
 nutrition, 65
 Biochemistry laboratory techniques
 blotting procedures, 53
 Cre-lox system, 56
 CRISPR/Cas9, 53
 enzyme-linked immunosorbent
 assay, 54
 flow cytometry, 54
 fluorescence in situ hybridization,
 55
 free light chain (FLC) assay, 431
 gene expression modifications, 56
 karyotyping, 55
 microarrays, 54
 molecular cloning, 55
 polymerase chain reaction, 52
 reverse transcriptase polymerase
 chain reaction, 52
 RNA interference, 56
 serum protein electrophoresis, 431
 Biofilm-producing bacteria, 128
 Staphylococcus epidermidis, 135
 Biologic agents
 naming conventions for, **254**
 Biomarkers
 α -fetoprotein, 491
 astrocytes, 493
 neurons, 493
 Biostatistics/epidemiology, 256–262
 Biotin, 68
 Bipolar disorder, **561**
 drug therapy for, 572
 lithium for, 574
 Birbeck granules
 Langerhans cell histiocytosis, 434
 “Bird’s beak” sign (X-ray), 376
 Birds (disease vectors), 148, 149
 Bismuth, **399**
 Bisoprolol, 245
 Bisphosphonates, 462, **486**
 esophagitis with, 249
 naming convention for, 254
 osteogenesis imperfecta treatment,
 51
 Bitemporal hemianopia, 542
 Craniopharyngioma, 528
 Hypopituitarism, 339
 Optic chiasm compression, 516
 Bitot spots, 66
 Bivalirudin, 435
 BK virus, 164
 Black eschar, 137
 Black lung disease, 677
 Bladder, 160
 bethanechol effect on, 240
 BPH and, 654
 development of, 618
 exstrophy, 624
 outlet obstruction, 579
 placenta percreta invasion, 640
 spasm treatment, 241
 urachus, 618
 urgency in cystitis, 241
 Bladder cancer
 cisplatin/carboplatin for, 442
 hematuria with, 594
 hypercalcemia and, 228
 Schistosoma haematobium, 161
 “Blast crisis,” 433
 Blastocyst implantation, 612
Blastomyces spp
 amphotericin B, 199
 itraconazole, 199
 Blastomycosis, 151
 Bleeding, 642
 adenomatous polyps, 387
 direct factor Xa inhibitors, 437
 direct thrombin inhibitors, 435
 diverticulosis, 383
 essential thrombocythemia, 433
 glycoprotein IIb/IIIa inhibitors, 438
 heparin, 436
 inflammatory bowel disease, 382
 peptic ulcer disease, 380
 thrombolytics, 437
 variceal, 371
 warfarin, 436
 Bleeding time, 427
 Bleomycin
 antitumor antibiotics, 439
 pulmonary fibrosis, 251
 targets of, 438
 toxicity, 444
 Blepharospasm, 519
 Blindness
 Chlamydia trachomatis, 149
 giant cell arteritis, 314
 neonatal, 142
 Onchocerca volvulus, 159
 Toxocara canis, 159
 Blistering skin disorders, 480
 Blood
 chocolate-colored, 666
 coagulation and kinin pathways,
 412
 hCG detection in, 633
 oxygen content, 666
 in placenta, 617
 umbilical cord, 618
 viscosity of, 668
 Blood-brain barrier
 anesthetics, 549
 astrocytes, 493
 function and mechanism, **496**
 at hypothalamus, 498
 L-DOPA, 549
 Blood flow
 autoregulation, 297
 exercise response, 670
 Blood groups, **405**
 Blood-nerve permeability barrier, 495
 Blood pH
 diuretic effects on, 609
 Blood pressure
 angiotensin II effects, 588
 antihypertensive therapy, 319
 cortisol effect on, 336
 fenoldopam and, 318
 renal disorders and, 591
 sympathomimetic effect on, 243
 Blood-testis barrier, 628
 Blood transfusions, **429**
 reactions, **114**
 therapy, 429
 Blood vessels
 collagen in, 50
 Ehlers-Danlos syndrome, 50
 hereditary hemorrhagic
 telangiectasia, 316
 Blood volume
 atrial natriuretic peptide release,
 296
 regulation, 588
 Bloody diarrhea, 179
 Campylobacter jejuni, 145, 149
 Shigella, 144
 ulcerative colitis, 382
 Bloody stool, 366
 Blotting procedures, **53**
 Blown pupil, 541
 CN III damage, 541
 saccular aneurysms, 516
 “Blue babies,” 298
 Blueberry muffin rash
 cytomegalovirus, 182
 rubella, 169, 182
 Toxoplasma gondii, 182
 “Blue bloater,” 674
 “Blue kids,” 299
 Blue sclerae, 51
 Blumer shelf, 379
 BMPR2 gene, 679
 Body compartments, 231
 Body dysmorphic disorder, 563
 Boerhaave syndrome, 377
 Bombesin, 350
 Bone cancer, 464
 primary bone tumors, 464
 Bone cell biology, 458, 459
 Bone crises, 88
 Bone disorders
 adult T-cell lymphoma and, 430
 lab values in, 464
 Langerhans cell histiocytosis, 434
 lytic (“punched out”), 431
 osteogenesis imperfecta, 51
 Bone formation, **458**
 Bone-in-bone (x-ray), 463
 Bone marrow
 cytokine stimulation of, 121
 lymphoid functions of, 96
 suppression, 199
 Bone mineral density scan, 462
 Bone morphogenetic protein (BMP),
 490
 Bones
 collagen in, 50
 cortisol effect on, 336
 lytic/blastic metastases, 223
 primary bone tumors, 464
 PTH effect on, 332
 renal osteodystrophy, 603
 Borderline personality disorder,
 565
Bordetella pertussis, **143**
 culture requirements, 126
 exotoxin production, 132
 macrolides, 193
 vaccines, 143
 Bordet-Gengou agar, 126
Borrelia burgdorferi
 animal transmission, 149
 coinfection with, 157
 facial nerve palsy, 186
 Lyme disease, 146
 tetracyclines, 192
Borrelia recurrentis
 animal transmission, 149
 vectors, 161
Borrelia spp, 146
 Bortezomib, **443**
 Bosentan, 686
 Botulinum toxin
 lysogenic transduction, 130
 passive antibodies for, 110
 symptoms of, 138
 toxin effects, 132
 Bovine spongiform encephalopathy
 (BSE), 178
 Bowel stenosis, 383
 Bowen disease, 651
 Bowenoid papulosis, 651
 Bow legs (genu varum), 463

- Bowman capsule, 583
 Boxer's fracture, 459
 BPH (benign prostatic hyperplasia)
 hydronephrosis in, 599
 Brachial artery, 455
 Brachial plexus
 Pancoast tumor, 685
 Brachial plexus lesions, **448**
 Brachiocephalic syndrome, 685
 Brachiocephalic vein, 685
 Brachioradialis reflex, 510
 Bradycardia
 amiodarone and, 323
 atropine for, 241
 β -blockers and, 245, 323
 dopamine for, 242
 on ECG, 293
 hypermagnesemia, 591
 reflex, 588
 sympatholytic drugs and, 243
 Bradykinesia
 with antipsychotic drugs, 573
 Bradykinin
 ACE inhibitors and, 610
 C1 esterase inhibitor deficiency,
 107
 Bradykinin, 610
 BRAF gene, **224**, 387
 melanomas and, 484
 papillary thyroid carcinoma and,
 343
 vemurafenib and, 444
 Brain
 blood flow autoregulation, 297
 embryologic derivatives, 613
 embryology of, 490
 infarcts, 209
 ischemia in, 210
 metastasis to, 223
 Brain abscesses
 bacteremia, 180
 HIV-positive adults, 180
 otitis media, 180
 Staphylococcus aureus, 180
 Toxoplasma gondii, 177
 Viridans streptococci, 180
 Brain cysts, 161
 Brain death, 269, 501, 502
 Brain injury
 gastritis with, 379
 hypopituitarism from, 339
 Brain lesions (common), **511**
 Brain natriuretic peptide (BNP), 296,
 588
 in SIADH, 338
 signaling pathways for, 337
 Brain stem
 dorsal view, **504**
 ventral view, **504**
 Brain stem/cerebellar syndromes
 multiple sclerosis, 523
 Brain tumors
 adult primary, **526–527**
 childhood primary, 528
 hallucinations with, 559
 incidence and mortality, 222
 metastatic source, 223
 nitrosoureas for, 441
 Branched-chain ketoacid
 dehydrogenase, 66
 Branching enzyme (glycogen
 metabolism), 86
 BRCA1/BRCA2 genes, 224
 DNA repair in, 40
 tumor suppressor genes, 224
 Breastfeeding, 636
 Breast milk
 IgA antibodies in, 105
 oxytocin's role in, 328
 prolactin and, 330
 Breast/ovarian cancer
 BRCA1 mutation, 64
 BRCA2 mutation, 64
 incomplete penetrance, 56
 Breast pathology, **649**
 benign disorders, 649
 invasive carcinomas, **650**
 noninvasive carcinomas, **650**
 Breast cancer
 aromatase inhibitors for, 656
 breastfeeding and, 636
 hormonal contraception
 contraindication, 657
 hypercalcemia and, 228
 incidence/mortality of, 222
 oncogenes and, 224
 paclitaxel for, 441
 paraneoplastic cerebellar
 degeneration and, 228
 tamoxifen for, 443
 trastuzumab for, 443
 tumor suppressor genes and, 224
 Breathing
 mechanics of, 675
 with pneumothorax, 682
 Breath sounds
 bronchial, 680, 682
 diminished, 682
 physical findings, 680
 Brenner tumor, 646
 Brief psychotic disorder, 560
 Brimonidine, 552
 Brittle bone disease, gene defects
 in, 51
 Broad-base budding (blastomycosis),
 151
 Broad ligament, 625
 Broca area, 501
 aphasia, 516
 MCA stroke, 514
 Bromocriptine, 548
 Bronchi, 662
 Bronchial carcinoid tumor, 684
 Bronchiectasis
 Aspergillus fumigatus, 153
 cystic fibrosis, 60
 Kartagener syndrome, 49
 Bronchioles, 662
 histamine receptors and, 238
 Bronchiolitis obliterans, 119, 683
 organizing pneumonia (BOOP),
 683
 Bronchitis
 cystic fibrosis, 60
 Haemophilus influenzae, 142
 Bronchoconstriction, 687
 Bronchodilation, 687
 sympathetic receptors and, 238
 Bronchogenic carcinomas
 asbestosis and, 677
 carcinogens causing, 225
 Bronchogenic cysts, 660
 Bronchopneumonia, 683
 Bronchopulmonary dysplasia, 210
 free radical injury, 210
 neonatal respiratory distress
 syndrome, 661
 Brown-Séquard syndrome, **531**
 Horner syndrome, 531
 "Brown tumors," 464
Brucella spp, 127
 transmission and treatment of, **143**
 zoonotic infections, 149
 Brucellosis, 149
 Brugada syndrome, **294**, 304
 Bruising
 scurvy, 69
 Brunner glands
 bicarbonate product, 372
 duodenum, 362
 Bruton agammaglobulinemia, 61, **116**
 Bruxism, 497
 BTK gene, 116
 B-type natriuretic peptide, **296**
 Buckle (torus fracture), 462
 Budd-Chiari syndrome, **392**
 Budesonide, 687
 Buerger disease, 314
 Bugs
 affecting unvaccinated children, **186**
 causing diarrhea, **179**
 causing food-borne illness, **178**
 hints, **186**
 Bulbar (spongy) urethra injury, 627
 Bulbus cordis, 281
 Bulimia nervosa, 567
 anovulation and, 645
 drug therapy for, 572
 laxative abuse by, 401
 Mallory-Weiss syndrome and, 377
 SSRIs for, 575
 Bulk-forming laxatives, 401
 Bullae, 475
 impetigo, 479
 necrotizing fasciitis, 479
 skin lesions, 475
 Bull neck lymphadenopathy, 132
 Bullous impetigo, 479
 Bullous pemphigoid, 475, 480
 autoantibody, 115
 Bulls-eye erythema, 146
 Bumetanide, 608
 BUN (blood urea nitrogen)
 ornithine transcarbamylase
 deficiency, 83
 Bundled payment, 271
 Bundle of His, 293
 Bundle of Kent, 294
 Bunyaviruses, 167–168
 Bupivacaine, 550
 Buprenorphine, 551
 heroin detoxification, 576
 morphine and, 234
 Bupropion, 576
 seizures with, 251
Burkholderia cepacia
 cystic fibrosis, 179
 Burkitt lymphoma, 430
 chromosomal translocations and,
 434
 EBV, 165
 oncogenes and, 224
 oncogenic microbes and, 226
 Burnout (medical errors), **274**
 Burns
 classification, **483**
 shock with, 310
 sunburn, 482
 testosterone/methyltestosterone
 for, 658
 "Burr cells," 414
 Bursitis
 prepatellar, 460
 Burton line
 lead poisoning, 419
 Buspirone
 mechanism and clinical use, **574**
 Busulfan, 441
 pulmonary fibrosis and, 251
 toxicity, 444
 Butorphanol, 551, 552
 C
 C1 esterase inhibitor deficiency, 107
 C3 deficiency, 107
 C5a receptor, 406
 C5-C9 deficiencies, 107
 CA 15-3/CA27-29 (tumor markers),
 226
 CA 19-9 (tumor marker), 226, 398
 CA 125 (tumor marker), 226
 CAAT box, 41
 Cachexia, **227**
 TNF- α and, 108
 Café-au-lait spots
 McCune-Albright syndrome, 57
 Caffeine intoxication and withdrawal,
 570
 Cahill cycle, 82
 Calcarine sulcus
 thalamic relay to, 498
 Calciferol (vitamin D), 589
 Calcification, 211
 dystrophic, 227
 Calcineurin, 120
 Calcinosis cutis, 473
 Calcitonin, **333**
 medullary thyroid carcinoma
 production, 343
 tumor marker, 226
 Calcitriol, 589
 Calcium
 in bone disorders, 464
 calcitonin and, 333
 in cardiac muscle, 292
 in osteomalacia/rickets, 463
 in Paget disease of bone, 463
 Vitamin D and, 337
 Calcium carbonate, 399
 Calcium channels
 ethosuximide effect on, 544
 glucose and, 334
 Lambert-Eaton myasthenic
 syndrome, 228
 myocardial action potential, 292
 opioid effect on, 551
 pacemaker action potential, 292
 Calcium channel blockers
 angina, 318
 antiarrhythmic drugs, **324**
 contractility in, 284
 cutaneous flushing, 248
 gingival hyperplasia, 250
 hypertension, 316
 hypertrophic cardiomyopathy, 308
 mechanism and clinical use, 318
 migraine headaches, 518
 Raynaud phenomenon, 472
 Calcium homeostasis, 333
 Calcium (kidney stones), 598
 calcium oxalate nephrolithiasis, 69
 Calcium pyrophosphate deposition
 disease, **467**
 Calcium-sensing receptor (CaSR), 355
 Calculation
 bioavailability of, 231
 reabsorption and secretion rate, **584**
 Calculous cholecystitis, 396
 Caliciviruses, 163
 characteristics of, 167

- California encephalitis, 167
 Calluses (dermatology), 475
 cAMP (cyclic adenosine monophosphate)
 endocrine hormone messenger, 337
 fructose biphosphatase-2 and, 76
 heat-labile/heat-stable toxin effects, 132
 hyperparathyroidism, 345
 PTH effect on, 332
 Vibrio cholerae, 146
 cAMP factor, 137
Campylobacter spp
 animal transmission, 149
 bloody diarrhea, 179
 jejuni, **145**
 reactive arthritis and, 469
 Canagliflozin, 353
 Canalicular stage (development), 660
 Cancer
 common metastases, 223
 deaths from, 272
 ESR in, 214
 hallmarks of, 221
 immune evasion in, 221
 mortality of, 222
 pneumoconioses, 676, 677
 Cancer drugs
 cell cycle, **438**
 targets, **438**
 Cancer epidemiology, **222**
 Candesartan, 610
Candida albicans, 153
 HIV-positive adults, 177
 T cell dysfunction, 116
Candida spp
 amphotericin B, 199
 azoles, 199
 echinocandins, 200
 immunodeficiency infections, 118
 osteomyelitis, 180
 tricuspid valve endocarditis and, 311
 vulvovaginitis, 181
 Candidiasis
 Candida albicans, 153
 chronic mucocutaneous, 116
 cortisol and, 336
 nystatin, 199
 Cannibalism, 178
 “Cannonball” metastases, 642
 Capecitabine
 5-F-dUMP, 36
 “Cape-like” sensory loss, 492
 Capillary fluid exchange, **297**
 Capitate bone, 449
 Capitation, 271
 Caplan syndrome, 466, 677
 Capsules (bacterial), 124
 Captain’s wheel
 Paracoccidioidomycosis, 151
 Captopril, 610
 Caput medusae, 365
 Carbachol, 240, 552
 Carbamazepine
 agranulocytosis, 250
 aplastic anemia, 250
 cytochrome P-450 and, 252
 epilepsy, 544
 SIADH and, 249
 Carbamoyl phosphate, 82
 Carbamoyl phosphate synthetase, 73
 Carbapenems
 mechanism and use, **190**
 Pseudomonas aeruginosa, 143
 Carbidoopa, **549**
 Carbohydrate absorption, **373**
 Carbohydrate metabolism
 inborn errors of, 80
 Carbol fuchsin, 125
 Carbon dioxide (CO₂)
 production in tissues, 127
 retention, 679
 transport, **670**
 Carbonic anhydrase, 670
 Carbon monoxide (CO)
 vs cyanide poisoning, 667
 electron transport inhibition, 78
 poisoning, 666
 teratogenicity, 614
 toxicity treatment, 248
 Carbon tetrachloride
 free radical injury and, 210
 Carboplatin
 mechanism and clinical use, **442**
 toxicities of, 444
 Carboplatin toxicity, 444
 Carboxylases, 73
 Carboxypeptidase, 373
 Carcinoembryonic antigen (CEA)
 (tumor marker), 226
 Carcinogens, **225**
 griseofulvin, 200
 Carcinoid syndrome, **352**
 bronchial carcinoid tumors, 684
 somatostatin in treatment, 371
 Carcinoid tumors
 biomarkers for, 226
 immunohistochemical stains for, 227
 octreotide for, 400
 stomach, 379
 Carcinoma in situ, 219
 cervical dysplasia, 645
 ductal, 650
 neoplastic progression, 219
 penis, 651
 vulvar, 644
 Carcinomas
 bone, 464
 colorectal, 388
 invasive, 219
 metastases of, 219, 223
 nomenclature of, 220
 thyroid, 343
 Cardiac arrest
 antacid adverse effects, 399
 hypermagnesemia, 591
 Cardiac cycle, 287
 Cardiac depression, 318
 Cardiac function curves, **286**
 Cardiac glycosides
 mechanism and clinical use, **321**
 sodium-potassium pump inhibition, 49
 Cardiac looping, 280
 Cardiac output
 exercise and, 670
 in pregnancy, 633
 variables in, **284**
 V/Q mismatch and, 669
 Cardiac output equations, 285
 Cardiac pressures (normal), 297
 Cardiac tamponade, **310**
 aortic dissection and, 303
 jugular venous pulse in, 287
 MI, 305, 307
 shock, 310
 Cardiac therapy, **317**
 Cardiac tumors, **316**
 Cardinal veins, 281
 Cardiogenic shock, 310
 etiology, 310
 Cardiomegaly
 Pompe disease, 87
 Cardiomyopathy, **308**
 β-blockers, 245
 Chagas disease, 158
 drug reaction and, 248
 familial amyloid, 212
 glycogen storage diseases, 67
 heart failure with, 309
 hemochromatosis and, 395
 hypertrophic, 245
 Kussmaul sign in, 316
 Starling curves, 285
 sudden cardiac death, 304
 Cardiotoxicity
 doxorubicin, 439
 drugs causing, 444
 methylxanthines, 687
 TCA adverse effects, 575
 trastuzumab, 443
 Cardiovascular drugs
 naming conventions for, 253
 reactions to, 248
 Cardiovascular system, 281–323
 anatomy, 283
 embryology, 281–283
 pathology, 298–313
 pharmacology, 316–322
 physiology, 284–299
 sclerosis of, 473
 Carditis
 Lyme disease, 146
 rheumatic fever, 312
 Carfilzomib, **443**
 Carina (trachea), 663
 Carmustine, 441
 pulmonary fibrosis, 251
 Carnitine acyltransferase I, 73, 89
 Carotid artery
 atherosclerosis in, 302
 cavernous sinus, 542
 emboli from, 538
 embryonic development, 619
 giant cell arteritis and, 314
 Carotid massage, 296
 Carotid sinus, 296
 Carpal bones, 449
 Carpal tunnel syndrome, 459
 lunate dislocation, 449
 nerve injury, 447
 rheumatoid arthritis, 466
 Carteolol, 552
 Cartilage
 collagen in, 50
 fluoroquinolone damage to, 250
 Cartilage damage, 204
 Carvedilol, 245, 323
 Casal necklace, 67
 Caseating granulomas
 in tuberculosis, 140
 Case-control studies, 266
 Caseous necrosis, 209
 Caspases, 208
 Caspofungin
 echinocandins, 200
 Casts in urine, **594**
 Catabolism of amino acids, 82
 Catalase, 210
 Catalase-positive organisms, **128**
 Cataplexy, 568
 Cataracts, **535**
 corticosteroid toxicity, 120
 diabetes mellitus and, 346
 galactosemia, 80
 muscular dystrophy, 61
 rubella, 182
 sorbitol, 81
 Catecholamines
 adrenal medulla secretion, 327
 amphetamines and, 242
 contractility effects of, 284
 ephedrine and, 242
 heart contractility, 284
 pacemaker action potential, 292
 pheochromocytoma and, 350
 Catecholamine synthesis
 tyrosine catabolism, **83**
 Cats (disease vectors)
 Campylobacter jejuni, 145
 Cat scratch disease, 149
 Pasteurella multocida, 149, 186
 Tinea corporis, 152
 Toxoplasma gondii, 156, 182
 Cauda equina, 507
 Cauda equina syndrome, 530
 Caudal fold closure defects, 358
 Caudal regression syndrome, 614
 Caudate
 basal ganglia, 500
 Huntington disease, 520
 Cavernous hemangiomas
 liver, 392
 Cavernous hemangiomas (liver), 392
 Cavernous sinus, **542**
 dural venous sinuses, 503
 syndrome, 542
 thrombosis with mucormycosis, 153
 CCR5 protein
 HIV and, 175
 maraviroc, 203
 viral receptor, 166
 CD4+ cell count
 disease associations by levels, 177
 CD4 protein, 101
 viral receptor, 166
 CD4+ T cells (HIV), 176
 CD5 protein, 432
 CD8 protein, 101
 CD16 protein, 101
 CD20 protein, 110
 in CLL, 432
 CD21 protein, 110
 viral receptor, 166
 CD25 protein
 cell surface protein, 110
 regulatory T cells and, 102
 CD28 protein, 110
 CD34 protein, 110
 leukocyte extravasation and, 215
 CD40 protein, 110
 CDKN2A gene, 224
 CEA tumor marker, 388
 Cefaclor, 189
 Cefazolin
 mechanism and use, 189
 prophylaxis, 198
 Cefepime
 mechanism and use, 189
 Pseudomonas aeruginosa, 143
 Cefotaxime, 189
 Cefotetan
 mechanism and use, 189
 Cefoxitin
 mechanism and use, 189
 Cefpodoxime
 mechanism and use, 189
 Ceftaroline
 mechanism and use, 189
 MRSA, 198

- Ceftazidime
mechanism and use, 189
Pseudomonas aeruginosa, 143
- Ceftriaxone
Chlamydia spp, 148
for gonococci, 142
for *Haemophilus influenzae*, 142
mechanism and use, 189
meningitis, 180
meningococci, 142
typhoid fever, 144
- Cefuroxime
mechanism and use, 189
- Celecoxib, 252, **486**
- Celiac artery
mesenteric ischemia, 386
structures supplied, 364
- Celiac disease, 381
autoantibody, 115
HLA genes and, 100
IgA deficiency, 116
- Celiac sprue, 381
- Celiac trunk, **364**
- Cell biology of bone, **459**
- Cell cycle phases, **46**
- Cell envelope (bacterial), **124**
- Cell injury, **207**
- Cell lysis, 590
- Cell membrane in apoptosis, 208
- Cell surface proteins
association and functions, **110**
leukocyte adhesion deficiency, 117
T cells and, 101
- Cell trafficking, **47**
- Cell types
labile, 46
permanent, 46
stable (quiescent), 46
- Cellular biochemistry, 46–52
- Cellular injury
cellular adaptations, **206**
irreversible, 207
reversible, 207
- Cellulitis, 136, 479
Pasteurella multocida, 149
- Cell walls (bacterial), 124
- Central clearing
nuclei, 343
rash, 152
- Central diabetes insipidus, 338
- Central/downward transtentorial
herniation, 529
- Central nervous system (CNS)
anesthetic principles for, 549
antiarrhythmic adverse effects,
322, 323
cancer epidemiology, 222
cell types in, 493–494
depression, 546
drug name conventions, 253
nitrosoureas effect on, 441
origins of, **490**
posterior fossa malformations, 491,
492
regional specification of, 506
shock from injury, 310
- Central nervous system stimulants,
572
- Central pontine myelinolysis, 524
- Central post-stroke pain syndrome,
515
- Central precocious puberty, 637
- Central retinal artery occlusion, **538**
- Central sleep apnea, 679
- Central sulcus, 501
- Central tendency measures, 262
- Central tendon (diaphragm), 663
- Central vertigo, 534
- Centriacinar emphysema, 674
- Cephalexin
mechanism and use, 189
- Cephalosporins
disulfiram-like reaction, 251
mechanism and use, **189**
pseudomembranous colitis, 249
Pseudomonas aeruginosa, 143
- Ceramide trihexoside
in sphingolipidoses, 88
- Cerebellar degeneration
paraneoplastic, 228
- Cerebellar lesions
hemisphere, 511
lateral, 499
medial, 499
tonsillar herniation, 492, 529
vermis lesions, 511
- Cerebellum
development of, 490
input/output of, **499**
tonsils, 492
- Cerebral aqueduct of Sylvius, 504
- Cerebral artery distributions, **502**
- Cerebral cortex
aphasia, 514
arterial distribution, **502**
dominant parietal lesions, 511
functional areas of, **501**
hemineglect, 514
nondominant parietal lesions, 511
visual field defects, 514
- Cerebral edema
diabetic ketoacidosis and, 347
therapeutic hyperventilation, 501
- Cerebral hemispheres, 490
- Cerebral palsy, 551
- Cerebral perfusion pressure (CPP),
501
- “Cerebriform” nuclei, 430
- Cerebrospinal fluid (CSF)
albuminocytologic dissociation,
524
blood-brain barrier and, 496
circulation of, 496, 503
findings in meningitis, **180**
Guillain-Barré syndrome, 524
hydrocephalus, 522
multiple sclerosis, 523
neurodegenerative disorders, 521
origins, 490
poliomyelitis, 531
production, 493
ventricular system, 504
- Cerebrovascular disease
diabetes mellitus, 346
- Certolizumab, 487
- Ceruloplasmin
free radical elimination by, 210
- Cervical cancer, **645**
carcinogens causing, 225
epidemiology of, 643
HIV-positive adults, 177
hydronephrosis with, 599
oncogenic microbes and, 226
papillomaviruses, 164
- Cervical rib, 448
- Cervicitis
sexually transmitted infections, 184
- Cervix
anatomy of, 625
epithelial histology, 626
lymphatic drainage of, 624
pathology of, 645
- Cestodes, **160**
- Cetirizine, 686
- Cetuximab, **442**
- CFTR gene
chronic pancreatitis and, 397
- cGMP (cyclic guanosine
monophosphate)
atrial natriuretic peptide and, 296
endocrine hormone messenger,
337
male sexual response, 627
- Chagas disease, 158
achalasia in, 376
- Chalk-stick fractures, 463
- Chancroids, 184
- Changes in the elderly, **270**
- Chaperone protein, **45**
- Charcoal yeast extract culture
Legionella pneumophila, 126, 143
- Charcot-Bouchard microaneurysm,
516
- Charcot joints
syphilis, 147
tabes dorsalis and, 530
- Charcot-Leyden crystals, 674
- Charcot-Marie-Tooth disease, 524
- Charcot triad, 397
- Charging, tRNA, 44
- Chédiak-Higashi syndrome, 117
- Cheilosis, 67
- Chelation
hemochromatosis, 395
lead poisoning, 419
- Chemokines, 108
delayed hypersensitivity, 112
- Chemoreceptors, **296**
- Chemoreceptor trigger zone (CTZ),
496
- Chemotherapy
AML and, 432
MDR1 and responsiveness to,
227
neutropenia with, 424
ondansetron, 400
paclitaxel, 48
pancreatic cancer, 398
readmissions with, 272
treatments for vomiting, 401
vincristine/vinblastine, 48
- Chemotoxicities, 444
- Cherry hemangiomas, 478
- “Cherry red” epiglottitis, 142
- Cherry-red spot (macula), 538
lysosomal storage disease, 88
- Chest pain
panic disorder, 563
pneumothorax, 682
- Chest wall
elastic properties, 665
- Chest X-rays
aortic dissections on, 303
balloon heart on, 308
eggshell calcification, 677
notched ribs on, 299
Wegener granulomatosis on, 315
widened mediastinum on, 137
- Cheyne-Stokes respirations, 679
- Chiari malformations, 492
- Chickenpox
rash, 183
VZV, 165
- Chief cells (parathyroid), 332
- Chief cells (stomach), 372
- Child abuse, **556**
osteogenesis imperfecta and, 51
reporting requirements, 269
- Childbirth
brachial plexus injury in, 448
Budd-Chiari syndrome and, 392
Graves disease and, 342
low birth weight, 635
misoprostol induction, 399
neonatal flora, 178
oxytocin and uterine contractions,
328, 636
postpartum mood disturbances,
562
preterm, as common cause of
death, 272
progesterone levels after, 630
Sheehan syndrome after, 339
stress incontinence and, 599
- Childhood disorders, **557**
- Childhood musculoskeletal
conditions, **461**
- Childhood primary brain tumors, **528**
- Child neglect, **556**
- Children
causes of death, 272
- Chi-square tests, 264
- Chlamydia* spp, **148**, 184
atypical infections, 179
diagnostic tests for, 148
Giemsa stain, 125
Gram stain, 125
intracellular organism, 127
macrolides, 193
pneumonia, 683
reactive arthritis, 469
sulfonamides for, 194
tetracyclines, 192
- Chlamydia trachomatis*
eosinophilia, 149
pelvic inflammatory disease, 149
pneumonia, 179
serotypes, **149**
UTIs, 600
- Chlamydia pneumoniae*, 148
pneumonia, 179
- Chlamydia psittaci*
atypical pneumonia, 148
transmission, 149
- Chloasma (melasma), 476
- Chloramphenicol
aplastic anemia and, 250
gray baby syndrome, 250
mechanism and use, **192**
pregnancy contraindications, 204
protein synthesis inhibition, 191
- Chlordiazepoxide, 546
alcohol withdrawal, 572
- Chlorhexidine, 204
- Chloride channels
cystic fibrosis, 60
- Chlorine, 204
- Chlorprocaine, 550
- Chloroquine, **200**
malaria, 157
- Chlorpheniramine, 686
- Chlorpromazine, 573
- Chlorthalidone, 609
- Chocolate agar
Haemophilus influenzae, 126, 142
- Cholangiocarcinomas
Clonorchis sinensis, 160, 161
hyperbilirubinemia, 393
oncogenic microbes and, 226
sclerosing cholangitis, 395
- Cholangitis, 368, 382, 393, 397
- Cholecalciferol, 70
- Cholecystectomy, 396, 397
- Cholecystitis, 396

- Cholecystokinin (CCK)
secretory cell location, 373
- Cholelithiasis, 396
- Cholelithiasis, 395, **396**, 397
acute pancreatitis, 397
bile ducts and, 368
biliary cirrhosis and, 396
Crohn disease, 382
hyperbilirubinemia and, 393
octreotide and, 400
- Cholera toxin
lysogenic phage infection, 130
mechanism, 132
- Cholestasis serum markers, 390
- Cholesteatomas, **533**
- Cholesterol
atherosclerosis, 302
in bile, 374
cholelithiasis and, 396
lipid-lowering agents, 320
synthesis of, 47, 73, 79
vitamin B₃ effects, 67
- Cholesteryl ester transfer protein, 93
- Cholestyramine, 320
- Cholinergic agonists, 253
- Cholinergic effects, 321
cardiac glycosides, 321
- Cholinesterase inhibitors
diarrhea with, 249
neuromuscular blockade reversal,
551
- Cholinomimetic agents, **240**
glaucoma treatment, 552
- Chondrocalcinosis, 467
- Chondrocytes, 458, 462
osteoarthritis, 466
- Chondroma, 464
- Chondrosarcoma, 465
- Chordae rupture, 291
- Chorea
brain lesions, 511
Huntington disease, 520
movement disorders, 519
- Choriocarcinoma, **642**
hCG in, 633
methotrexate for, 440
testicular, 653
theca-lutein cysts and, 646
- Chorionic plate, 617
- Chorionic somatomammotropin, 634
- Chorionic villi
hydatidiform moles, 642
placenta, 617
- Chorioretinitis
congenital toxoplasmosis, 182
- Choristomas, 220
- Choroid layer (ophthalmology)
inflammation, 536
neovascularization, 536
- Choroid plexus (CNS), 504
- Christmas tree distribution, 482
- Chromaffin cells
diagram, 327
pheochromocytomas, 350
- Chromatin structure, **34**
- Chromatolysis, **495**
- Chromogranin, 226, 684
- Chromosomal translocations, **434**
- Chromosome abnormalities
congenital microdeletion, 64
gene associations with, 64
hemochromatosis, 395
karyotyping for, 55
nephroblastoma, 606
nondisjunction (meiosis), 63
omphaloceles, 358
polyposis syndrome, 387
renal cell carcinoma, 605
Robertsonian translocation, 64
von Hippel-Lindau disease, 525
Wilson disease, 395
- Chronic bronchitis, 674
- Chronic disease, anemia of, 421
- Chronic gastritis, 379
- Chronic granulomatous disease
(CGD)
catalase-positive microbes, 186
immunodeficiencies and, 117
recombinant cytokines for, 121
respiratory burst in, 109
- Chronic inflammation, **216**
- Chronic ischemic heart disease, 304
- Chronic kidney disease
erythropoietin in, 589
hypertension and, 300
- Chronic lymphocytic leukemia
(CLL), 432
immunosuppressants, 120
rituximab for, 443
- Chronic mesenteric ischemia, 386
- Chronic mucocutaneous candidiasis,
116
- Chronic myelogenous leukemia
(CML), 433
basophilia caused by, 408
busulfan for, 441
chromosomal translocations and,
434
imatinib for, 443
oncogenes and, 224
- Chronic myeloproliferative disorders,
433
- Chronic obstructive pulmonary
disease (COPD)
albuterol for, 242
 β -blockers and, 245
muscarinic antagonists for, 241
salmeterol for, 242
- Chronic pancreatitis, **397**
pancreatic insufficiency from, 381
- Chronic placental insufficiency,
578
- Chronic pyelonephritis, 600
- Chronic renal failure, 603
hyperphosphatemia with, 345
- Chronic respiratory diseases
bronchitis, 674
with chronic inflammatory
diseases, 683
death in children, 272
obstructive diseases, 674
pneumoconioses, 675
sinusitis, 674
thromboembolism, 679
- Chronic thromboembolic pulmonary
hypertension, 679
- Chronic transplant rejection, 119
- Churg-Strauss syndrome, 315
autoantibody, 115
- Chvostek sign
hypocalcemia, 591
hypoparathyroidism, 344
- Chylomicrons, 92, 94
- Chymotrypsin, 373
- Cidofovir, **202**
- Cigarette smoke (carcinogen), 225
- Ciguatera, 247
- Cilastatin
imipenem and, 190
seizures with, 251
- Ciliary ganglia, 539
- Cilia structure, **49**
- Ciliated cells, 662
- Cimetidine, 399
cytochrome P-450 and, 252
gynecomastia and, 649
- Cinacalcet, **355**
- Cinchonism
antiarrhythmic causing, 322
neurologic drug reaction, 251
- Cingulate gyrus
limbic system, 499
- Cingulate (subfalcine) herniation,
529
- Ciprofloxacin
for Crohn disease, 382
cytochrome P-450 and, 252
fluoroquinolones, 195
meningococci, 142
*Mycobacterium avium-
intracellulare*, 196
prophylaxis, 198
Pseudomonas aeruginosa, 143
- Circadian rhythm
hypothalamic control, 498
sleep physiology, 497
- Circle of Willis, **503**
saccular aneurysms, 516
- Circulatory system
fetal, 282
kidneys and, 580
- Circumflex femoral artery, 463
- Circumflex pallor
group A streptococcal pharyngitis,
136
- Cirrhosis
 α_1 -antitrypsin deficiency, 392
alcoholic, 71, 391
bacterial peritonitis (spontaneous),
390
cholelithiasis and, 396
cystic fibrosis, 60
encephalopathy with, 391
esophageal varices and, 377
fructose intolerance, 80
gynecomastia, 649
hemochromatosis, 395
with hepatitis, 389
hepatocellular carcinomas, 392
hyperbilirubinemia in, 393
non-alcoholic fatty liver disease,
391
portal hypertension, **389**
serum markers for, 390
- Cisplatin
mechanism and clinical use, **442**
targets of, 438
toxicities of, 251, 444
- Citalopram, 575
- Citrate synthase, 74
- Citrulline, 82
- c-KIT gene, 224
- CK-MB, 304, 306
- Cladribine, **440**
for hairy cell leukemia, 432
- Clara cells, 661, 662
- Clarithromycin
Helicobacter pylori, 146
HIV prophylaxis, 198
macrolides, 193
*Mycobacterium avium-
intracellulare*, 196
pregnancy use, 204
- Clasp knife spasticity, 529
- Class IA antiarrhythmics, 322
- Class IB antiarrhythmics, 322
- Class IC antiarrhythmics, 322
- Class II antiarrhythmics, 323
- Class III antiarrhythmics, 323
- Class IV antiarrhythmics, 324
- Classical conditioning, **554**
- Class switching
CD40, 103
thymus-dependent antigens, 105
- Clathrin, 47
- Claudication
atherosclerosis, 302
Buerger disease, 314
giant cell arteritis, 314
- Clavicle fractures, **460**
- Clavulanate
Haemophilus influenzae, 142
- Clavulanic acid, 189
- Clawing (hand), 451
Klumpke palsy, **448**
- Clearance (CL) of drugs, 231
- Clear cell adenocarcinoma, 644
DES and, 656
- Cleavage in collagen synthesis, 50
- Cleft lip and palate
development, **621**
Patau syndrome, 63
Pierre Robin sequence, 620
- Clevidipine, 318
for hypertensive emergency, 318
- Clindamycin
bacterial vaginosis, 148
Clostridium difficile and, 138
mechanism and use, **192**
metronidazole vs, 192
protein synthesis inhibition, 191
pseudomembranous colitis with,
249
- Clinical reflexes, **510**
- Clinical trials, **256**
- Clinical vignette strategies, 23
- “Clock-face” chromatin, 409, 431
- Clofazimine
Hansen disease, 141
Mycobacterium leprae, 196
- Clomiphene
estrogen receptor modulators,
656
hot flashes with, 249
- Clomipramine, 575
- Clonidine, 240, 243
- Cloning methods (laboratory
technique), 55
- Clonorchis sinensis*
cholangiocarcinoma, 226
diseases association, **161**
trematodes, 160
- Clopidogrel, 437
acute coronary syndromes, 307
for ischemic stroke, 512
thrombogenesis and, 411
- Closed-angle glaucoma, 536
pilocarpine for, 240
- Clostridium botulinum*
exotoxin production, 132
food poisoning, 178
therapeutic uses, 138
- Clostridium difficile*
antibiotic use, 185
metronidazole, 195
nosocomial infection, 185
PPI association, 138
proton pump inhibitor use, 399
toxins and effects of, 138
vancomycin, 190
watery diarrhea, 179
- Clostridium perfringens*
clindamycin, 192
exotoxin production, 133

- food poisoning, 178
 toxins produced, 138
 traumatic open wound, 186
 watery diarrhea, 179
- Clostridium* spp.**, **138**
 anaerobic organism, 127
 exotoxins, 138
- Clostridium tetani***, 138
 exotoxin production, 132
- Clotrimazole, 199
- Clotting factors, 71
- Clozapine, 573
 agranulocytosis with, 250
- Clubbing (digital), 675
 cystic fibrosis, 60
 Eisenmenger syndrome, 299
- Club cells, **661**
- Clue cells
 bacterial vaginosis, 148, 181
- Cluster A personality disorders
 characteristics of, **565**
 schizoid, 565
 schizotypal, 565
- Cluster B personality disorders
 antisocial, **565**
 borderline, 565
 histrionic, 565
 narcissistic, 565
- Cluster C personality disorders
 avoidant, **566**
 dependent, 566
 obsessive-compulsive, 566
- Cluster headaches, 518, 547
- c-MYC gene, 224
- CN III, IV, VI palsies, **541**
- CNS (central nervous system)
 cancer epidemiology, 222
- CNS lymphomas
 HIV-positive adults, 177
 oncogenic microbes and, 226
- CO₂
 production in tissues, 127
- Coagulation, 71
- Coagulation disorders, **426**
 hemophilia, 426
 hereditary thrombosis syndromes,
 428
 mixed platelet/coagulation, 428
 vitamin K and, 426
- Coagulation pathways, **412**
- Coagulative necrosis, 209
 MI, 305
- Coagulopathy
 postpartum hemorrhage, 641
 preeclampsia, 643
 uterine bleeding with, 633
- Coal workers' pneumoconiosis, 677
- CoA production, 67, 72
- Coarctation of aorta, 299, 300
- Cobalamin, 69
- Cobblestone mucosa, 382
- Cocaine
 β -blockers and, 245
 coronary vasospasm, 248
 intoxication and withdrawal, 571
 liver processing of, 367
 sympathomimetic action, 242
 teratogenicity, 614
- Coccidioides* spp.**
 silver stain, 125
 treatment, 199
- Coccidioidomycosis, 151
 erythema nodosum and, 482
- Coccobacilli, 141
- Cocci bacteria
 antibiotic tests, 134
- Cochlea
 CN VIII, 506
 inner ear, 533
 presbycusis, 533
- Codeine, 551
- Codman triangle (x-ray), 465
- Codominance, 56
- Codons
 amino acid specification, 37
 genetic code features, 37
 start and stop, 44
- Cofactors
 apolipoproteins, 93
 biotin, 68, 73
 cobalamin, 69
 copper, 51
 Menkes disease, 51
 pantothenic acid, 67
 phenylketonuria, 84
 pyridoxine, 67
 pyruvate dehydrogenase complex,
 76
 riboflavin, 67
 TCA cycle, 77
 thiamine, 74
 vitamin K, 71
- "Coffee bean sign" (X-ray), 386
- Cognitive behavioral therapy (CBT),
 557, 572
- ADHD, 557
- anxiety disorders, 562
- atypical depression, 561
- bipolar disorder, 561
- body dysmorphic disorder, 563
- obsessive-compulsive disorder, 563
- panic disorder, 563
- phobias, 563
- postpartum depression, 562
- PTSD, 563
- Cohort studies, 256
- Coin lesion (X-ray), 684
 x-ray signs, 684
- Cola-colored urine, 596
- Colchicine, 55
 agranulocytosis, 250
 calcium pyrophosphate deposition
 disease, 467
 diarrhea with, 249
 gout, 467, 487
 microtubules and, 48
 myopathy with, 250
- "Cold enrichment," 139
- Colectomy
 adenomatous polyposis, 387
 inflammatory bowel disease, 382
- Colesevelam, 320
- Colestipol, 320
- Colistin
 polymyxin E, 193
Pseudomonas aeruginosa, 143
- Colitis
Clostridium difficile, 138
 oral vancomycin, 190
 pseudomembranous, **179**, 188, 192
- Collagen, **50**
 decreased/faulty production, 51
 epithelial cell junctions and, 474
 osteoblast secretion of, 459
 polyostotic fibrous dysplasia and, 57
 scar formation, 218
 synthesis and structure, **50**
 in systemic sclerosis, 473
 types of, 50
 vitamin C in synthesis, 69
 wound healing, 216
- Collagenase in neutrophils, 406
- Collapsing pressure (alveoli), 661
- Collecting tubules
 potassium-sparing diuretics and,
 609
- Colles fracture, 462
- Colliculi, 504
- Colon
 histology of, 362
 ischemia in, 210
- Colon cancer, **388**
 adenomatous polyposis and, 387
 bevacizumab for, 442
 cetuximab for, 442
 5-fluorouracil for, 440
 incidence/mortality in, 222
 irinotecan/topotecan for, 442
 Lynch syndrome, 40
 molecular pathogenesis of, 389
 oncogenes and, 224
 serrated polyps and, 387
Staphylococcus gallolyticus and,
 137
 tumor suppressor genes and, 224
- Colonic ischemia, 363, 386
- Colonic polyps, **387**
- Colony stimulating factor, 121
- Colorado tick fever, 167
- Color blindness, 197
- Colovesical fistulas, 383
- Coltivirus, 167
- Coma
 hepatic encephalopathy, 391
 herniation syndromes, 529
 hyponatremia, 591
 rabies, 171
 thyroid storm, 342
Toxocara canis, 159
Trypanosoma brucei, 156
- Comedones, 477
- Commaless genetic code, 37
- Comma-shaped rods, 141
- Common bile duct, 361, 368
- Common cold, 168
- Common peroneal nerve, 452, 453
- Common variable immunodeficiency
 (CVID), 116
- Communicating hydrocephalus, 522
- Communication with patient, 268
- Compartment syndrome, 461
- Competence (bacterial genetics), 130
- Competitive agonists, 234
- Competitive antagonist, 234
- Competitive inhibitors, 230
- Complement
 activation pathways and functions,
106
 binding of, 104
 disorders of, **107**–122
 eculizumab, 122
 endotoxin activation, 133
 immunodeficiency infections, 118
 immunoglobulin isotypes, 105
 innate immunity, 99
 splenic dysfunction, 98
 transplant rejection, 119
- Complement activation inhibition
 β -hemolytic bacteria, 135
- Complementation (viral), 162
- Complete (third-degree) AV block,
 295
- Complex partial seizures, 517
- Complex renal cysts, 604
- Compliance (lung), 665
- Comprehensive Basic Science
 Examination (CBSE), 11
- Comprehensive Basic Science Self-
 Assessment (CBSSA), 11
- Computer-Based Test (CBT)
 environment of, 3–4
 exam schedule for, 7–8
 structure of, 3
- COMT inhibitors, 548
- Conditioning (psychological), 554
- Conduct disorder, 557
- Conducting zone (respiratory tree),
 662
- Conduction aphasia, 516
- Conductive hearing loss, 533
- Condylomata acuminata, 477
 sexual transmission, 184
- Condylomata lata
 syphilis, 147, 184
- Confidence intervals, **263**
- Confidentiality, **267**
 in abuse, 269
 behavioral science ethics, 265
 exceptions to, 267
- Confluence of the sinuses, 503
- Confounding bias, 261
- Congenital adrenal hyperplasias, **335**
- Congenital heart disease, **298**–300
 defect associations, **300**
 maternal phenylketonuria, 84
 pulmonary arterial hypertension,
 679
 rubella, 182
- Congenital hydrocele (scrotal), 652
- Congenital hypothyroidism
 (cretinism), 341
- Congenital long QT syndrome, 294
- Congenital lung malformations, **660**
- Congenital malformation mortality,
 272
- Congenital nevus, 475
- Congenital rubella
 cardiac defect associations, 300
 heart murmur, 291
- Congenital solitary functioning
 kidney, **579**
- Congenital syphilis, 147
- Congestion (respiratory)
 with lobar pneumonia, 683
- Congo red stain, 212
- Conivaptan, 354
 SIADH, 354
- Conjoined tendon, 369
- Conjugated (direct)
 hyperbilirubinemia, 393
- Conjugate vaccines, 127
- Conjugation (bacterial genetics), 130
- Conjunctival suffusion/injection
 eye disorders, 147
 Kawasaki disease, 314
- Conjunctivitis, 148, **534**
 adenovirus, 164
 chlamydia, 184
 gonococcal proctylaxia, 198
Haemophilus influenzae, 142
 reactive arthritis, 469
 rubeola, 170, 183, 186
 Zika virus, 171
- Connective tissue diseases
 aortic dissection and, 303
 pulmonary arterial hypertension, 679
 thoracic aortic aneurysms and, 302
- Connective tissue drug reactions, 250
- Conn syndrome, 349, 591
- Consensual light reflex, 539
- Consent
 healthcare proxy, 269
 minors, **265**, 268

- Consolidation (lung finding), 680
lobar pneumonia, 683
- Constipation
anal fissures, 366
calcium channel blockers, 318
Hirschsprung disease, 384
irritable bowel syndrome, 383
laxative treatments, 401
loperamide, 400
ondansetron, 400
ranolazine, 319
vincristine, 441
- Constrictive pericarditis
jugular venous pulse in, 287
Kussmaul sign, 316
- Contact dermatitis, 113
- Contemplation stage, substance addiction, 568
- Continuous heart murmurs, 291
- Contraception
isotretinoin teratogenicity, 614
methods for, **657**
parental consent for minors and, 265
progestins for, 657
- Contractility in cardiac output, 284
- Contraction alkalosis, 60, 587, 588, 609
- Coombs-positive hemolysis
 α -methyldopa, 243
anemia with, 423
- Coombs test, 423
- Cooperative kinetics, 230
- COPI/COPII proteins, 47
- Copper deficiency, 419
- Copper intrauterine device, **657**
- Copper metabolism
Wilson disease, 395
- Copper toxicity, 248
- Coprolalia, 557
- Copy number variations (CNV), 54
- Cord factor, 140
- Cori cycle, 82
- Cori disease, 87
- Corkscrew fibers, 528
- “Corkscrew” hair, 69
- Cornea
astigmatism, 535
collagen in, 50
- Corneal arcus
familial hypercholesterolemia, 94
hyperlipidemia, 301
- Corneal reflex, 507
- Corneal vascularization, 67
- Corniculate cartilage, 620
- Coronary arteries
anatomy of, **283**
atherosclerosis in, 302
- Coronary artery disease
atrial fibrillation and, 295
diabetes mellitus and, 346
hormonal contraception with, 657
hypertension and, 300
sudden cardiac death, 304
- Coronary sinus
anomalous pulmonary return, 298
development, 281
- Coronary steal syndrome, 304
- Coronary vasospasm, 248
triptans and, 547
- Coronaviruses
characteristics of, 167
genomes of, 163
- Cor pulmonale, 309, 668
from obstructive lung disease, 674
pneumoconioses, **677**
pulmonary hypertension, 679
right ventricular failure, 668
- Corpus cavernosum
lymphatic drainage of, 624
- Corpus luteum
hCG and, 633
progesterone production, 630
- Correlation coefficient, 264
- Cortical signs, 514
- Corticopapillary osmotic gradient, 588
- Corticosteroid-binding globulin, 336
- Corticosteroids
asthma, 687
cataracts, 535
Crohn disease, 382
Cushing syndrome, 348
giant cell arteritis, 314
hyperglycemia with, 249
hypopituitarism, 339
lymphopenia with, 424
microscopic polyangiitis, 315
neutrophilia from, 424
osteonecrosis, 463
osteoporosis with, 250
pancreatitis with, 249
for polymyalgia rheumatica, 470
Takayasu arteritis, 314
thyroid storm, 342
Wegener granulomatosis, 315
- Corticotropin-releasing hormone (CRH)
adrenal cortex regulation of, 327
cortisol regulation, 336
function of, 328
signaling pathways of, 337
- Cortisol, **336**
congenital adrenal hyperplasias, 335
in Cushing syndrome, 348
signaling pathways for, 337
- Corynebacterium diphtheriae*
culture requirements for, 126
exotoxin effects, **139**
exotoxin production, 132
- Costovertebral angle tenderness, 601
- Co-transporter 2 (SGLT2) inhibitors, 353
- Cough
ACE inhibitors, 251, 610
chronic bronchitis, 674
gastroesophageal reflux disease, 377
guaifenesin, 686
hypersensitivity pneumonitis, 675
lung cancer, 684
nonproductive, 140
staccato, 149
Wegener granulomatosis, 315
whooping, 132, 143
- Councilman bodies
yellow fever, 168
- Countertransference, 554
- Courvoisier sign
pancreatic cancer, 398
- Cowpox, 164
- Coxiella burnetii*
animal transmission, 149
Q fever, 150
- Coxiella* spp
intracellular organisms, 127
- Coxsackievirus
acute pericarditis, 313
picornavirus, 168
presentation, 167
rashes of childhood, **183**
- C-peptide
insulin and, 334
with insulinomas, 351
- CpG island methylator phenotype (CIMP), 387
- Crackles (physical findings), 680
- Cranial nerve palsies
osteopetrosis, 463
- Cranial nerves, **506**
common lesions, **532**
functions of, 506
locations of, 504
nerve and vessel pathways, **505**
nuclei of, **505**
pharyngeal arch derivation, 620
reflexes of, **507**
- Craniopharyngioma, 528, 613
hypopituitarism with, 339
- Craniofacial, 463
- C-reactive protein (CRP), 213
innate immunity and, 99
- Creatine kinase, 203
- Creatinine
ACE inhibitor effects, 610
glomerular filtration rate and, 582
proximal convoluted tubules, 587
- Creatinine clearance, 582
- Cre-lox system, 56
- Creomastic muscle and fascia
inguinal canal and, 369
- Creomastic reflex, 452, 510
- Crepitus
esophageal perforation, 377
in necrotizing fasciitis, 479
soft tissue, 138
- Crepitus in necrotizing fasciitis, 479
- Crescentic glomerulonephritis, 596
- CREST syndrome, 473
autoantibody, 115
sclerodermal esophageal dysmotility, 377
- Creutzfeldt-Jakob disease, 178, 521
- “Crew cut” (skull X-ray), 422
- Cricoid cartilage, 620
- Cricothyroid muscle, 620
- Cri-du-chat syndrome, **64**
- Crigler-Najjar syndrome, 393, 394
- Crimean-Congo hemorrhagic fever, 167
- CRISPR/Cas9, **53**
- Crohn disease, 382
azathioprine, 120
B₁₂ deficiency, 420
cholelithiasis and, 396
natalizumab, 122
spondyloarthritis and, 469
sulfasalazine for, 400
vitamin B₁₂ deficiency, 69
- Cromolyn, 687
- Cross-linking in collagen synthesis, 50
- Cross-over study, **256**, 261
- Cross-sectional studies, **256**
- Croup, **170**
paramyxoviruses, 167, 169
pulsus paradoxus in, 310
- CRP and ESR, 214
- Crust (skin), 475
basal cell carcinoma, 484
impetigo, 479
varicella zoster virus, 479
- Cryoprecipitate, 429
- Crypt hyperplasia, 381
- Cryptococcal meningitis, 199
- Cryptococcosis, 153
- Cryptococcus neoformans*, 153
HIV-positive adults, 177
stains for, 125
- Cryptococcus* spp
meningitis, 180
treatment, 199
urease-positive, 127
- Cryptogenic organizing pneumonia, 683
- Cryptorchidism, **651**
Sertoli cells and, 628
- Cryptosporidium* spp, 155
fluorescent antibody stain, 125
watery diarrhea, 179
- Crypts of Lieberkühn, 362
- C-section deliveries
neonatal flora, 178
neonatal flora with, 178
risk factors after, 640
- Culture requirements
bacteria, 126
- Cuneiform cartilage, 620
- Curling ulcers
gastritis, 379
- “Currant jelly” stools, 385, 386
- “Currant jelly” sputum
Klebsiella spp, 186
- Curschmann spirals, 674
- Cushing disease, 348
- Cushing-like symptoms, 203
- Cushing reflex, 296
- Cushing syndrome, **348**
anovulation with, 645
corticosteroids, 120
eosinopenia, 424
paraneoplastic syndrome, 228
small cell lung cancer, 684
- Cushing ulcers
gastritis, 379
- Cutaneous anthrax, 137
edema toxin, 132
- Cutaneous flushing
drugs causing, 248
- Cutaneous larva migrans, 159
- Cutaneous leishmaniasis, 158
- Cutaneous mycoses, **152**
- Cutaneous paraneoplastic syndromes, 228
- Cutaneous small-vessel vasculitis, 314
- Cutis aplasia
Patau syndrome, 63
- CXCR4
viral receptor, 166
- CXCR4/CCR5 protein
presence on cells, 110
- Cyanide
electron transport chain, 78
- Cyanide poisoning
vs carbon monoxide poisoning, **667**
induced methemoglobinemia, 666
nitroprusside, 318
treatment for, 248
- Cyanosis
“blue babies,” 298
“blue kids,” 299
Eisenmenger syndrome, 299
esophageal atresia, 359
methemoglobinemia, 666
patent ductus arteriosus, 299
pulmonary hypertension, 679
tetralogy of Fallot, 298
- Cyclins, 46
- Cyclobenzaprine, 551
- Cyclooxygenase
aspirin effect on, 411

- Cyclooxygenase inhibition
irreversible, 486
reversible, 485, 486
selective, 486
- Cyclophosphamide, 441
hemorrhagic cystitis with, 249
microscopic polyangiitis, 315
SIADH with, 249
toxicities of, 444
transitional cell carcinoma and, 606
Wegener granulomatosis, 315
- Cycloplegia
atropine, 241
muscarinic antagonists for, 241
- Cyclosporine
gingival hyperplasia, 250
gout, 250
immunosuppression, 120
- Cyclothymic disorder, 561
- Cystathionine, 67
- Cystathionine synthase deficiency, 84
- Cyst disorders (renal), 604
- Cysteine, 85
- Cystic duct, 368
- Cystic fibrosis, **60**
Aspergillus fumigatus, 153
bronchiectasis, 675
chromosome association, 64
common organisms, 179
meconium ileus and, 386
N-acetylcysteine, 686
pancreatic insufficiency, 381
vitamin deficiencies and, 65
- Cystine, 598
- Cystinuria, **85**
- Cystitis
acute bacterial, 594, 600
squamous cell carcinoma risk, 606
- Cytarabine, 440
- Cytochrome C, 208
- Cytochrome P-450
- azoles, 199
barbiturates and, 546
cimetidine and, 399
griseofulvin, 200
interactions with, **252**
macrolides, 193
phenobarbital effect on, 544
rifamycins, 196
ritonavir, 203
- Cytokeratin, 227
- Cytokines, 101, 108
corticosteroids and, 120
Graves disease and, 342
rejection reactions, 119
type IV hypersensitivity, 113
- Cytokinesis, 46
- Cytomegalovirus (CMV)
AIDS retinitis, 165
cholecystitis and, 396
clinical significance, 165
esophagitis and, 377
HIV-positive adults, 177
hyper-IgM syndrome and, 117
immunodeficient patients, 118
pneumonia, 683
TORCH infection, 182
treatment, 202
viral receptor, 166
- Cytoplasm
cell cycle phase, 46
cytoskeletal elements, 48
glycolysis, 76
HMP shunt, 79
metabolism in, 72
- Cytoplasmic membrane (bacterial), 124
- Cytoplasmic processing bodies (P-bodies), 41
- Cytoskeletal elements, **48**
- Cytotoxic T cells, **102**
cell surface proteins, 110
MHC I and II, 100
- Cytotrophoblast, 617
- D**
- Dabigatran, 435
- Dabrafenib, **444**
- Daclizumab
targets of, 121
- Dacrocytes, 414
- Dactinomycin, 439
RNA polymerase inhibition, 42
targets of, 438
- Dactylitis
seronegative spondyloarthritis, 469
sickle cell anemia, 422
- Dalfopristin
VRE, 198
- Danazol, **658**
pseudotumor cerebri, 521
“Dancing eyes, dancing feet,” 228
- Dandy-Walker syndrome, 492
- Dantrolene, 550, 551
- Dapagliflozin, 353
- Dapsone
Hansen disease, 141
hemolysis in G6PD deficiency, 250
mechanism and use, **194**
Mycobacterium leprae, 196
Pneumocystis jirovecii, 154
- Daptomycin
mechanism and use, **195**
MRSA, 198
- Darkfield microscopy
for *Treponema*, 146
- Darunavir
HIV therapy, 203
- Dasabuvir, 204
- Dasatinib, 443
- Datura*, 241
- Daunorubicin, 439
dilated cardiomyopathy, 248
- DCC gene, 224
- Deacetylation
histones, 34
- Dead space (lung), 664
- Deafness
Alport syndrome, 596
congenital long QT syndrome, 294
congenital syphilis, 147
rubella, 182
syphilis, 182
- Deamination
base excision repair, 40
- Death
aortic dissection in, 303
children, explaining to, 269
common causes, **272**
sudden cardiac death, 304
thyroid storm, 342
- Death receptor pathway, 207, 208
- Debranching enzyme
Cori disease, 87
glycogen metabolism, 86
- Decay-accelerating factor (DAF), 106
- Deceleration injury, 303
- Decerebrate (extensor) posturing, 511
- Decidua basalis, 617
- Decision-making capacity, **266**
- Decorticate (flexor) posturing, 511
- Decussation
in spinal tracts, 509
- Deep brachial artery, 455
- Deep inguinal lymph nodes, 624
- Deep venous thrombosis (DVT), **671**
direct factor Xa inhibitors for, 437
glucagonomas and, 351
heparin for, 436
tamoxifen/taloxifene and, 443
- Deer flies (disease vectors), 159
- Defense mechanisms
immature, 554–555
mature, 555
- Defensins, 99
- Deferasirox, 248
- Deferiprone, 248
- Deferoxamine, 248
- Deformation, 613
- Degarelix, **656**
- Degenerate/redundant genetic code, 37
- Degmacytes, 414
- G6PD deficiency, 79
- Dehydration
diabetic ketoacidosis, 347
filtration changes and, 583
gout exacerbation, 467
loop diuretics and, 608
mannitol and, 607
osmotic laxatives, 401
salivary stones with, 376
shock, 310
- Dehydrogenases, 73
- Delavirdine
HIV therapy, 203
- Delirium, **558**
barbiturate withdrawal, 570
diabetic ketoacidosis, 347
PCP, 571
thyroid storm, 342
- Delirium tremens (DTs), **569**, 570
- δ cells
endocrine pancreas
somatostatin production, 371
somatostatinomas of, 351
- Delta rhythm (EEG), 497
- Delta virus, 167
- Deltoid muscle
axillary nerve injury, 447
Erb palsy, 448
- Delusional disorder, 560
- Delusions, 559
mesolimbic pathway, 499
- Demeclocycline
diabetes insipidus and, 249, 338
- Dementia
HIV-positive adults, 177
metachromatic leukodystrophy, 88
neurodegenerative disorders, 520–521
prion disease, 178
vitamin B₃ deficiency, 67
- Demyelinating/dysmyelinating disorders, **523**
lead poisoning (adult), 425
metachromatic leukodystrophy, 88
progressive multifocal leukoencephalopathy, 524
vitamin B₁₂ deficiency, 530
- Dendritic cells, **408**
IL-10, 108
innate immunity, 99
T- and B-cell activation, 102, 103
- Dengue, 167
- Denial, 554
- Denosumab, 122
for osteoporosis, 462
- De novo synthesis
pyrimidine and purine, **36**, 73
- Dense deposit disease, 596
- Dental plaque
normal flora, 178
viridans streptococci, 128
- Dentate line, 366
- Dentate nucleus, 499
- Dentin
collagen in, 50
osteogenesis imperfecta, 51
- Dentinogenesis imperfecta, 51
- Denys-Drash syndrome, 606
- Dependent personality disorder, 566
- Depersonalization/derealization disorder, 558
panic disorder, 563
- Depolarizing neuromuscular blocking drugs, 551
- Depressants, intoxication and withdrawal, 570
- Depression
atypical features in, **561**
benzodiazepine withdrawal, 570
dissociative identity disorder, 558
drug therapy, 572
electroconvulsive therapy, 562
hyperparathyroidism, 345
metoclopramide, 400
mirtazapine for, 244
neurotransmitters for, 495
postpartum, 562
seasonal pattern with, 561
serotonin-norepinephrine reuptake inhibitors (SNRIs) for, 575
SSRIs for, 575
- Deprivation effects (infants), 556
- De Quervain tenosynovitis, 461
- De Quervain thyroiditis, 341
- Dermacentor* tick (disease vector), 149
- Dermatitis
B-complex deficiency, 65
drug reactions and, 250
glucagonomas, 351
IPEX syndrome, 102
type IV hypersensitivity reaction, 113
vitamin B₅ deficiency, 67
vitamin B₇ deficiency, 68
- Dermatitis herpetiformis, 481
celiac disease and, 381
- Dermatologic lesion terms, **475**
- Dermatomes
landmarks, 510
- Dermatomyositis, 228
autoantibody, 115
- Dermatomyositis/polymyositis, 471
- Dermatophytes, **152**
- Dermis, 473
- Descending colon, 360
- Desert bumps, 151
- Desert rheumatism
Coccidioidomycosis, 151
- Desflurane, 550
- Desipramine, 575
- Desloratadine, 686
- Desmin, 48, 227
- Desmopressin
central DI, 329
DI treatment, 338
for hemophilia, 426
- Desmosome, 474
- Desquamation
staphylococcal toxic shock syndrome, 135
- Desvenlafaxine, 575
- Detached retina, 537

- Developmental delay
fetal alcohol syndrome, 615
renal failure and, 603
- Dexamethasone
Cushing syndrome diagnosis, 348
- Dexlansoprazole, 399
- Dexrazoxane, 439
- Dextroamphetamine, 572
- Dextrocardia, 280
x-ray, 49
- Dextromethorphan, 551, **686**
- DHT (dihydrotestosterone), 622, 636, 639
- Diabetes insipidus, **338**
antidiuretic hormone in, 329
drug reaction and, 249
lithium, 574
lithium toxicity, 569
potassium-sparing diuretics for, 609
thiazides for, 609
- Diabetes mellitus, **346–347**
atherosclerosis and, 302
 β -blockers and, 245
carpal tunnel syndrome, 459
cataracts and, 535
chronic renal failure and, 603
CN III damage, 541
diabetic ketoacidosis, 347
diabetic retinopathy, 537
Friedreich ataxia, 531
fungal infections, 186
glucagonomas, 351
glucosuria in, 584
hemochromatosis, 395
hepatitis C, 173
hypertension and, 300, 316
management of, 352–353
naming conventions for, 253
nephropathy with, 597
opportunistic infections, 153
pancreatic cancer, 398
polyhydramnios and, 641
in pregnancy, 300
pyelonephritis and, 600
readmissions with, 272
tacrolimus and, 120
teratogenic potential of maternal, 614
therapy management, **352**
type 1 vs type 2, 347
urinary tract infections, 181
UTIs and, 600
- Diabetes mellitus type 1, 347
autoantibody, 115
HLA subtypes with, 100
- Diabetes mellitus type 2, 347
amyloidosis, 212
- Diabetic ketoacidosis (DKA), **347**
DM type 1 and, 346
ketone bodies, 90
- Diabetic nephropathy
ACE inhibitors for, 610
- Diabetic neuropathy, 575
- Diabetic retinopathy, **537**
- Diagnosis errors, 274
- Diagnostic criteria, psychiatric
grief, 562
panic disorder, 563
symptom duration and, **564**
- Diagnostic maneuvers/signs
Gower sign, 61
- Diagnostic test evaluation, 257
- Diagnostic tests/maneuvers
laboratory tests in bone disorders, 464
lower extremity, 491
wrist and hand injury, 459
- Dialectical behavioral therapy, 565, 572
- Dialysis-related amyloidosis, 212
- Diamond-Blackfan anemia, 420
- Diapedesis, 215
- Diaper rash
Candida albicans, 153
nystatin, 199
- Diaphoresis, 305
acromegaly, 339
- Diaphragmatic hernia, 370
- Diaphragm structures, **663**
- Diaphysis, 465
- Diarrhea, 117
B-complex deficiency, 65
bismuth/sucralfate for, 399
Campylobacter jejuni, 145
cholera toxin, 132
clindamycin, 192
Clostridium difficile, 138
Cryptosporidium, 155
drug reaction and, 249
ezetimibe, 320
giardiasis, 155
graft-versus-host disease, 119
HIV-positive adults, 177
inflammatory bowel diseases, 382
irritable bowel syndrome, 383
lactase deficiency, 81
lactose intolerance, 381
as laxative adverse effect, 401
leflunomide, 486
loperamide for, 400
magnesium deficiency from, 332
magnesium hydroxide, 399
malabsorption syndromes, 381
metoclopramide, 400
misoprostol, 399
opioids for, 551
organisms causing, 179
pellagra, 67
rice-water, 132
rotavirus, 168
Salmonella, 144
Shigella, 144
thyroid storm and, 342
Vibrio cholerae, 146
VIPomas, 371
vitamin C toxicity, 69
watery, 132
- Diastole
cardiac cycle, 287
heart failure and, 309
heart murmurs of, 291
heart sounds of, 287
- Diazepam, 546
alcohol withdrawal, 572
- Diclofenac, 486
- Dicloxacillin
mechanism and use, 188
- Dicrotic notch, 287
- Dicyclomine, 241
- Didanosine, 203
HIV therapy, 203
pancreatitis, 249
- Diencephalon, 490
- Diethylcarbamazine
antihelminthic, 200
nematode infections, 159
- Diethylstilbestrol (DES), 656
teratogenicity, 614
vaginal tumors, 644
- Differential media, 126
- Diffuse axonal injury, **515**
- Diffuse cortical necrosis, **602**
- Diffuse esophageal spasm, 377
- Diffuse gastric cancer, 379
- Diffuse glomerular disorders, 594
- Diffuse large B-cell lymphoma (DLBCL), 430, 432
- Diffuse partial seizures, 517
- Diffuse proliferative glomerulonephritis (DPCN), 596
- Diffuse scleroderma, 473
- Diffuse stomach cancer, 379
- Diffusion-limited gas exchange, 668
- DiGeorge syndrome, 344
lymph node paracortex in, 96
thymic shadow in, 98
- Digestion
malabsorption syndromes, 381
secretory products for, 372–374
ulcerative colitis and, 382
- Digestive tract
anatomy and histology, **362**
- Digitalis
arrhythmias induced by, 322
contractility effects, 284
hyperkalemia and, 590
toxicity treatment for, 248
- Digoxin
contractility effects of, 285, 286
for dilated cardiomyopathy, 308
mechanism and clinical use, 321
sodium-potassium pump inhibition, 49
therapeutic index of, 234
toxicity treatment, 324
- Dihydroergotamine, 518
- Dihydrofolic acid, 194
- Dihydroorotate dehydrogenase
leflunomide effects, 36, 486
- Dihydropyridine calcium channel blockers, 253
- Dihydropyridine receptor, 456
- Dihydrorhodamine test, 117
- Dihydrotestosterone (DHT)
 5α -reductase deficiency, 639
function, 636
sexual determination, 622
- Dihydroxyacetone-P, 80
- Dilated cardiomyopathy, 308, 309
doxorubicin, 439
drug reaction and, 248
hemochromatosis, 395
muscular dystrophy, 61
wet beriberi, 66
with myocarditis, 313
- Diltiazem, 318
- Dimenhydrinate, 686
- Dimercaprol
for arsenic toxicity, 248
for lead poisoning, 248, 419
for mercury poisoning, 248
- Dinitrophenol, 78
- Diphenhydramine, 686
- Diphenoxylate, 551
- Diphtheria
Corynebacterium diphtheriae, 139
exotoxins, 130, 131, 132
unvaccinated children, 186
vaccine for, 139
- Diphyllobothrium latum*
disease association, 161
presentation, 160
vitamin B₁₂ deficiency, 69, 420
- Diplococci, 142
- Diplopia
brain stem/cerebellar syndromes, 523
central vertigo, 534
drug toxicity, 544
in botulism, 138
intracranial hypertension, 521
myasthenia gravis, 472
osmotic demyelination syndrome, 524
- Dipyridamole
for coronary steal syndrome, 304
- Direct bilirubin, 375
- Direct cholinomimetic agonists, 240
- Direct (conjugated)
hyperbilirubinemia, 393
- Direct Coombs test
Type II hypersensitivity, 112
- Direct factor Xa inhibitors, **437**
- Direct fluorescent antibody (DFA)
microscopy
for *Treponema*, 146
- Direct inguinal hernia, 370
- Direct light reflex, 539
- Direct sympathomimetics, 242
- Direct thrombin inhibitors, **435**
- Discharge planning, 272
- Discolored teeth, 204
- Discounted fee-for-service, 271
- Disease associations, **161**
- Disease prevention, **270**
- Disease vectors
Aedes mosquitoes, 168
Anopheles mosquito, 157
armadillos, 149
birds, 148, 149
cats, 149
dogs, 145, 149
fleas, 149, 150
flies, 144, 149
horse flies, 159
Ixodes ticks, 146
rodents, 167
ticks, 146, 150
zoonotic bacteria, 149
- Disinfection/sterilization methods, **204**
- Disinhibited behavior
Klüver-Bucy syndrome, 511
- Disinhibited social engagement, 556
- Disopyramide, 322
- Disorganized thought, 559
- Dispersion measures, 262
- Displacement, 554
- Disruption (morphogenesis), 613
- Disruptive mood dysregulation disorder, 557
- Disseminated candidiasis, 153
- Disseminated intravascular coagulation (DIC), 428
acute myelogenous leukemia, 432
Ebola, 171
endotoxins, 131, 133
meningococci, 142
microangiopathic anemia, 423
Waterhouse-Friderichsen syndrome, 349
- Dissociation, 554
- Dissociative disorders, **558**
amnesia, 558
identity disorder, 558
- Distal humerus, 455
- Distal interphalangeal (DIP) joints, 451
- Distal renal tubular acidosis (type 1), 593
- Distributive shock, 310
- Disulfiram
alcoholism treatment, 571
ethanol metabolism and, 72
disulfiram-like reaction, 251

- Diuresis
atrial natriuretic peptide, 296
for shock, 310
- Diuretics
dilated cardiomyopathy, 308
electrolyte changes, **609**
glaucoma treatment, 552
hypertension treatment, 316
magnesium levels and, 332
mechanism and clinical use, **607**
pancreatitis, 249
for SIADH, 338
- Diverticula, **383**
- Diverticulitis, 383
- Diverticulosis, 383
- Diverticulum, 383
- Dizygotic (“fraternal”) twins, 616
- Dizziness
AChE inhibitors, 549
calcium channel blockers, 318
dihydropyridine, 318
nitrates, 318
ramelteon, 547
ranolazine, 319
sacubitril, 319
vertigo and, 534
- DMPK gene, 61
- DNA
cloning methods, 55
free radical effect on, 210
introns vs exons, 43
laddering in apoptosis, 208
methylation in, 34
mutations in, 39
plasmid transfer, 130
repair of, **40**
- DNA ligase
action of, 38
- DNA mutations
types of, 39
- DNA polymerase inhibitors, 253
- DNA polymerases
action of, 38
- DNA repair, **40**
- DNA replication, **38**
- DNA topoisomerases, 38
- DNA transcription
deacetylation, 34
- DNA viruses, **164**
characteristics, **163**
genomes, 163
- Dobutamine, 242
- Dofetilide, 323
- Dogs (disease vectors), 145, 149
- Dolutegravir, 203
- Dominant inheritance, 59
- Dominant negative mutations, 57
- Donepezil, 240, 549
- Do not resuscitate (DNR) order, 266
- Dopamine, 242
basal ganglia, 500
bupropion effect, 576
changes with disease, 495
function of, 328
Huntington disease, 520
kidney functions and, 589
L-DOPA, 548, 549
MAO inhibition, 549, 575
Parkinson disease, 548
PCT secretion of, 589
pheochromocytoma secretion, 350
vitamin B₆ and, 67
- Dopamine agonists, 330, 339, 548
- Dopamine receptors, 238
- Dopaminergic pathways, **499**
- Dornase alfa (DNase), 60
- Dorsal columns (spinal cord), 508, 509
thalamic relay for, 498
- Dorsal interossei muscle, 450
- Dorsal motor nucleus, 506
- Dorsal optic radiation, 542
- Dorsal pancreatic bud, 360
- Dorsiflexion
common peroneal nerve injury, 453
- Dosage calculations, **231**
- Dosage interval, 231
- Double-blinded studies, 256
- Double stranded viruses, 163
- Double Y males, 638
- Down syndrome, 63
ALL and AML in, 432
cardiac defect association, 300
cataracts and, 535
chromosome association, 64
hCG in, 633
Hirschsprung disease and, 384
intestinal atresia and, 359
- Doxazosin, 244
- Doxepin, 575
- Doxorubicin
dilated cardiomyopathy, 248
targets, 438
toxicities, 444
- Doxycycline
chlamydiae, 148
in gonorrhea treatment, 142
lymphogranuloma venereum, 149
MRSA, 198
Mycoplasma pneumoniae, 150
ricketsial/vector-borne disease, 150
tetracyclines, 192
- Doxylamine, 686
- DPP-4 inhibitors, 353
- Dressler syndrome, 305, 307, 313
- DRESS syndrome, 250
- Droling treatment, 241
- Drop metastases, 528
- Drop seizures, 517
- Drug dosages, 234
calculations, 231
lethal median, 234
liver disease, 231
loading, 231
maintenance dose, 231
median effective, 234
renal disease, 231
toxic dose, 234
- Drug elimination, 231
- Drug-induced lupus, 250
- Drug interactions
additive, type, 235
antagonistic, type, 235
permissive, type, 235
potentiation, type, 235
synergistic, type, 235
tachyphylactic, type, 235
- Drug metabolism, **232**
cytochrome P-450 dependent, 232
geriatric patients in, 232
- Drug name conventions, **253–255**
- Drug overdoses
of weak acids, 233
of weak bases, 233
- Drug reactions
cardiovascular, **248**
endocrine/reproductive, **249**
gastrointestinal, **249**
hematologic, **250**
multiorgan, **251**
musculoskeletal, **250**
neurologic, **251**
renal/genitourinary, **251**
respiratory, **251**
- Drug reaction with eosinophilia
and systemic symptoms
(DRESS), 250
- Drug-related disorders
myocarditis, 313
- Drug resistance
plasmids in, 131
- Drugs
cholinomimetic agents, 240
efficacy vs potency, 233
elimination of, 232, 233
errors in, 274
patient difficulty with, 268
phase I metabolism, 232
reactions to, 248–251
therapeutic index, 234
toxicities, 248
- Drug safety
therapeutic index, measurement, 234
- Drug trials, 256
- Drunken sailor gait, 511
- Drusen, 536
- Dry beriberi, 66
- Dry cough with ACE inhibitors, 251
- Dry mouth
Lambert-Eaton myasthenic
syndrome, 472
- Dry skin, 66
- Dubin-Johnson syndrome, 393, 394
- Duchenne muscular dystrophy, 61
inheritance, 61
- Ductal adenocarcinomas, 368
- Ductal carcinoma in situ (DCIS), 650
- Ductal carcinomas (invasive), 650
- Ductus arteriosus, 282, 619
- Ductus deferens
embryology, 622
- Ductus venosus, 282
- Duloxetine, 575
- Duodenal atresia, 359
- Duodenal ulcer, 380
- Duodenum
basal electric rhythm, 362
biliary structures and, 368
histology of, 362
location, 360
secretory cells, 373
- Duplex collecting system, **579**
- Dural venous sinuses, **503**
- Dura mater, 496
- Dwarfism
achondroplasia, 462
- D-xylose test, 381
- Dynein
movement of, 48
- Dynein motors, 171
- Dysarthria, 516
Friedreich ataxia as, 531
in botulism, 138
osmotic demyelination syndrome, 524
- Dysbetalipoproteinemia, 94
- Dysentery
Entamoeba histolytica, 179
Escherichia coli, 145
Shigella spp, 132, 144, 179
- Dysfunctional uterine bleeding, 633
- Dysgerminoma, 647
- Dysgeusia, 71
- Dyskinesia
tardive, 251
- Dyslipidemia
 β -blocker adverse effects, 323
 β -blockers, **245**
familial, 94
renal failure and, 603
vitamin B₃ for, 67
- Dysmenorrhea
copper IUD, 657
primary, 645
- Dysmetria
central vertigo, 534
with strokes, 514
- Dyspareunia, 567
- Dysphagia
achalasia, 376
esophageal pathologies and, 377–378
in botulism, 138
osmotic demyelination syndrome, 524
Plummer-Vinson syndrome, 418
stroke effects, 514
Zenker diverticulum, 384
- Dysplasia
bronchopulmonary, 210
cellular adaptive response, 206
cervical, 645
neoplastic progression, 219
- Dysplastic kidney
multicystic, 578, 579
- Dyspnea
 α_1 -antitrypsin deficiency, 392
aortic stenosis, 291
heart failure, 309
hypersensitivity pneumonitis, 675
hypertrophic cardiomyopathy, 308
in botulism, 138
Wegener granulomatosis, 315
- Dystonia
acute, 241
antipsychotics/antiepileptics, 569
benztropine for, 241
Lesch-Nyhan syndrome, 37
movement disorders, 519
nigrostriatal pathway and, 499
treatment of, 241
- Dystrophic calcification, 211, 227
- Dystrophin gene, 61
- Dysuria, 654
cystitis, 181
urinary catheterization, 185
UTIs causing, 600
- E**
- Ear
pharyngeal pouch derivation, 621
- Eardrum, 533
- Early complement deficiencies (C1–C4), 107
- Eating disorders
anovulation and, 645
binge-eating disorder, 567
bulimia nervosa, 567
characteristics of, **567**
pica, 567
- Eaton agar
culture requirements, 126
- Ebola virus, 167, **171**
- Ebstein anomaly, 281, 298, 300
lithium side effect, 574
- E-cadherin, 219, 474
- ECF
body compartments of, 231
- Echinocandins, **200**

- Echinococcus granulosus*
cestodes, 160
disease association, 161
- Echinocytes, 414
- Echothiophate, 552
- Echovirus
picornavirus, 167, 168
- Eclampsia, 300, 643
- Ecthyma gangrenosum, 143
Pseudomonas spp, 143
- Ectocervix, 626
- Ectoderm
branchial clefts, 619
derivatives of, 613
- Ectoparasites, **161**
- Ectopic pregnancy, 641
appendicitis differential diagnosis, 383
Chlamydia trachomatis, 149
hCG in, 633
Kartagener syndrome, 49
methotrexate for, 440
salpingitis and, 185
- Eculizumab, 122
- Eczema
atopic dermatitis, 477
eczematous dermatitis, 475
phenylketonuria, 84
skin scales in, 475
type I hypersensitivity, 112
Wiskott-Aldrich syndrome, 117
- Edema
acute poststreptococcal
glomerulonephritis, 596
Arthus reaction, 113
calcium channel blockers, 318
capillary fluid exchange and, 297
danazol, 658
diabetic ketoacidosis, 347
endotoxins, 133
fludrocortisone, 354
heart failure and, 309
with hyperaldosteronism, 349
immunosuppressants, 120
Kawasaki disease and, 314
kwashiorkor, 71
loop diuretics for, 608
periorbital, 159, 161
peripheral, 309, 610
pitting, 309
pseudoephedrine/phenylephrine, 686
pulmonary hypertension, 668
Trichinella spiralis, 159, 161
trichinosis, 159
vasogenic, 496
wet beriberi, 66
- Edema factor
Bacillus anthracis, 132
- Edinger-Westphal nuclei, 539
- Edrophonium, 240
- Edwards syndrome, 63
cataracts and, 535
chromosome association, 64
- Efavirenz
HIV-positive adults, 203
- Effective refractory period
Class IA antiarrhythmic effect, 322
Class IC antiarrhythmic effect, 322
myocardial action potential, 292
- Effective renal plasma flow, **582**
- Efferent/afferent nerves, 296
- Efferent arteriole, 580
ANP/BNP, 588
constriction of, 583
- Efficacy vs potency of drugs, **233**
- EGFR gene, 684
“Eggshell” calcification, 677
- Ego defenses, **554**, 555
- Ego-dystonic behavior, 563
- Egophony, 680
- Ego-syntonic behavior, 565, 566
- Ehlers-Danlos syndrome
aneurysm association with, 516
collagen in, 50
collagen synthesis in, **51**
heart murmur with, 291
- Ehrlichia* spp
animal transmission, 149
Ehrlichia chaffeensis, 149
Gram stain, 125
ricketsial/vector-borne, **150**
- Eisenmenger syndrome, 299
- Ejaculation
innervation of, 627
sperm pathway in, 626
- Ejaculatory ducts, 626
embryology of, 622
- Ejection fraction
equation for, 285
- Elastase, 373
activity in emphysema, 674
- Elastic recoil (lung and chest wall), 665
- Elastin
characteristics of, **52**
- Elbow injuries, 459, 461
- Elderly
changes in, **270**
- Electrocardiograms (ECGs), **293**
acute pericarditis on, 313
cardiac tamponade on, 310
low-voltage, 308, 310
MI diagnosis with, 306
with pulmonary embolism, 672
tracings of, **295**
- Electroconvulsive therapy (ECT)
adverse effects, **562**
postpartum psychosis, 562
- Electroencephalogram (EEG)
Creutzfeldt-Jakob disease, 521
in delirium, 558
sleep stages, 497
- Electrolytes
disturbances in, **591**
diuretic effects on, 609
- Electron acceptors (universal), 75
- Electron transport chain and
oxidative phosphorylation, **78**
inhibitors of, 78
- Electrophoresis
hemoglobin, 410
- Elek test, 139
- Elementary bodies (chlamydiae), 148
- Elephantiasis, 159
- 11 β -hydroxylase, 335
- 11-deoxycortisol
and metyrapone, 335
- 11-deoxycorticosterone, 335
- Elfin facies, 64
- Elimination constant, 231
- Elimination of drugs, **232**
urine pH and, 233
- ELISA (enzyme-linked
immunosorbent assay), **54**
- Elongation Factor 2
Corynebacterium diphtheriae, 139
- eltrombopag (TPO receptor agonist), 121
- Elvitegravir, 203
- Emancipated minors, 265
- EMB agar
Escherichia coli, 181
lactose-fermenting enterics, 144
- Emboli
atherosclerosis, 302
atrial fibrillation, 295
atrial septal defect, 299
pulmonary, 310
stroke, 512
- Emboliform nucleus, 499
- Embryogenesis
genes involved in, 612
intrinsic pathway and, 208
- Embryology
cardiovascular, 281–283
derivatives, **613**
erythropoiesis, 404
gastrointestinal, 358–359
gland derivations in, 621
hematologic/oncologic, 404
neurological, 490–492
pancreas and spleen, 359, 360
renal, 578–579
reproductive, 612–623
respiratory, 660
thyroid development, 326
USMLE Step 1 preparation, 276
- Embryonal carcinoma, 653
- Embryonic age calculation, 633
- Embryonic development, 612
- Embryonic morphogenic errors, 613
- Embryonic stage (development), 660
- Embryotoxic, 204
- Emergent care proxy, 269
- Emission
innervation of, 627
- Emollients
laxative, 401
- Emotion
neural structures and, 499
- Emotionally distraught patients, 268
- Emotional/social development
neglect and deprivation effects, 556
- Empagliflozin, 353
- Emphysema, 674
diffusion in, 668
diffusion-limited gas exchange, 668
elastin in, 52
panacinar, 392
- Empty/full can test, 446
- Empty sella syndrome, 339
- Emtricitabine, 203
HIV-positive adults, 203
- Enalapril, 610
- Encapsulated bacteria, **127**
infections with immunodeficiency, 118
- Encephalitis
anti-NMDA receptor, 228
Cryptococcus neoformans, 153
guanosine analogs, 201
herpesviruses, 165, 180
HSV identification, 166
Lassa fever, 167
neonatal, 182
West Nile virus, 180
- Encephalomyelitis
paraneoplastic, 228
paraneoplastic syndrome, 228
- Encephalopathy
hepatic, 365, 391
hypertensive emergency, 300
lead poisoning, 419
Lyme disease, 146
- prion disease, 178
renal failure, 603
Wernicke, 66
- Encephalotrigeminal angiomatosis, 525
- Endemic typhus, 149
- Endocannabinoids, 336
- Endocardial cushion, 281
- Endocardial fibroelastosis, 308
- Endocarditis
bacterial, 311
Candida albicans, 153
coarctation of aorta, 299
Coxiella burnetii, 150
daptomycin, 195
enterococci, 137
heart murmurs, 291
heroin addiction and, 576
nonbacterial thrombotic, 228
prophylaxis, 198
Staphylococcus aureus, 135
Streptococcus bovis, 137
viridans streptococci, 128
- Endocervix, 626
- Endochondral ossification, 458
- Endocrine functions
kidney, 589
- Endocrine system
anatomy, 327–328
embryology, 326
hormones acting on kidney, 589, 590
hormone signaling pathways, 337
pathology, 338–354
pharmacology, 352–354
physiology, 328–336
- Endocrine/reproductive drug
reactions, 249
- Endoderm
branchial pouch derivation, 619
derivatives of, 613
- Endodermal sinus tumor, 647, 653
- Endodermal tubules, 660
- Endometrial abnormal uterine
bleeding, 633
- Endometrial artery, 617
- Endometrial carcinoma, 648
epidemiology of, 643
estrogens and, 656
Lynch syndrome and, 388
PCOS and, 645
tamoxifen and, 443
- Endometrial hyperplasia
follicular cysts, 646
- Endometrial polyps
uterine bleeding with, 633
- Endometrial vein, 617
- Endometriosis
characteristics and treatment, 648
danazol for, 658
- Endometritis, 648
- Endometrium
hyperplasia, 648–649
maintenance of, 632
- Endoneurium, 495
- Endoplasmic reticulum, 46, 47
rough, 46
smooth, 46
- Endosomes, 47
- Endothelial cells
leukocyte extravasation and, 215
in wound healing, 216
- Endothelin receptor antagonist, 686
- naming conventions for, 254
- Endothelium-derived relaxing factor
(EDRF), 337

- Endotoxins
effects of, **133**
features of, 131
- Enflurane, 550
seizures with, 251
- Enfuvirtide, 203
HIV-positive adults, 203
- Enhancers (gene expression), 41
- Enoxacin, 195
- Entacapone, 548
- Entamoeba histolytica*
amebiasis, 155
bloody diarrhea, 179
metronidazole, 195
- Enteric gram \ominus bacteria
facultative anaerobic metabolism, 127
- Enteric nerves, 362, 401
- Enteritis
vitamin B₁₂ deficiency, 69
vitamin B₅ deficiency, 67
vitamin B₇ deficiency, 68
- Enterobacter aerogenes*, 189
- Enterobacter* spp
nosocomial infection, 185
- Enterobius* spp
diseases association, 161
infection routes, 158
- Enterobius vermicularis*, 159
- Enterochromaffin-like (ECL) cells, 350, 373
- Enterococci, **137**
penicillins for, 188
vancomycin, 190
vancomycin-resistant (VRE), 137
- Enterococcus* spp
UTIs, 181
- Enterococcus faecalis*, 137
- Enterococcus faecium*, 137
- Enterocolitis
vitamin E excess, 70
- Enterocolitis (necrotizing), 386
- Enterohemorrhagic *Escherichia coli* (EHEC), 132, 145, 178, 219
- Enteroinvasive *Escherichia coli* (EIEC), 145, 179
- Enterokinase/enteropeptidase, 373
- Enteropathogenic *Escherichia coli* (EPEC), 145
- Enterotoxigenic *Escherichia coli* (ETEC), 132, 179
- Enterovirus meningitis, 179, 180
- Entorhinal cortex, 499
- Enuresis
characteristics/treatment, **568**
sleep stages and, 497
TCA use for, 575
- Envelopes (viral), 163
- env* gene, 175
- Enzalutamide, 658
- Enzyme kinetics, **230**
antagonists, 234
partial agonists, 234
- Enzymes
glycolysis regulation, 76
lipid transport and, 91, 92
rate-determining, 73
terminology for, **73**
- Eosin-methylene blue (EMB) agar
special culture, 126
- Eosinopenia, 424
- Eosinophilia
Aspergillus fumigatus, 153
Chlamydia trachomatis, 149
drug reaction and, 250
macrolides, 193
- Eosinophilic casts (urine), 600
- Eosinophilic esophagitis, 377
- Eosinophilic granuloma, 675
- Eosinophilic granulomatosis
autoantibody, 115
- Eosinophilic granulomatosis with polyangiitis, 315
- Eosinophils, **408**
corticosteroid effects, 424
in esophagus, 377
- Ependymal cells, **493**
- Ependymoma, 528
- Ephedrine, 242
- Epicanthal folds
cri-du-chat syndrome, 64
Down syndrome, 63
- Epidemic typhus, 149
- Epidemiology/biostatistics, 256–262
- Epidermal growth factor (EGF)
signaling pathways for, 337
in wound healing, 216
- Epidermis, 473
embryologic derivatives, 613
- Epidermophyton*, 152
- Epididymis, 626
embryology of, 622
- Epididymitis, 184, **654**
- Epidural hematomas, 513
- Epidural space, 496
- Epigastric pain
chronic mesenteric ischemia, 386
Ménétrier disease, 379
pancreatitis, 397
- Epigastric veins, 365
- Epiglottitis
Haemophilus influenzae, 142
unvaccinated children, 186
- Epilepsy
confidentiality exceptions for patients with, 267
gustatory hallucinations in, 559
hallucinations in, 559
seizures, 517
- Epinephrine, 242
adrenal medulla secretion, 327
glaucoma treatment, 552
glycogen regulation by, 85
pheochromocytoma secretion, 350
unopposed secretion of, 346
vitamin B₆ and, 67
- Epineurium, 495
- Epiphysis
estrogen effects on, 459
slipped capital femoral, 461, 463
tumors in, 464
widening of, 463
- Episcleritis
inflammatory bowel disease, 382
- Epispadias, 624
- Epistaxis, **671**
hereditary hemorrhagic telangiectasia, 316
- Epithelial cell junctions, **474**
- Epithelial cells
tumor nomenclature of, 220
- Epithelial histology (female), **626**
- Epithelial hyperplasia, 649
- Eplerenone, 609
- Epley maneuver, 534
- Epoetin alfa (EPO analog), 121
- Epstein-Barr virus (EBV)
aplastic anemia, 421
Burkitt lymphoma, 430
false-positive VDRL, 148
hairy leukoplakia and, 479
- head and neck cancer, 671
- HIV-positive adults, 177
- Hodgkin lymphoma, 429
in immunodeficient patients, 118
nasopharyngeal carcinomas, 165
oncogenesis of, 226
paracortical hyperplasia in, 96
receptors for, 166
- Eptifibatide, 438
thrombogenesis and, 411
- Equations
half-life of, 231
- Erb palsy, 448
- Erectile dysfunction, 567
 β -blockers and, 245, 323
cimetidine, 399
Lambert-Eaton myasthenic syndrome, 472
Peyronie disease, 651
sildenafil, 686
- Erection
autonomic innervation, 627
ischemic priapism, 651
- Ergocalciferol, 70
- Ergosterol synthesis inhibitors, 253
- Ergot alkaloids
coronary vasospasm, 248
- Erlotinib, **442**
- Erosions (gastrointestinal), 362, 379
- Errors (medical), 274
- Erysipelas, 479
Streptococcus pyogenes, 136
- Erythema
complicated hernias, 370
Kawasaki disease, 314
- Erythema marginatum, 312
- Erythema migrans
in Lyme disease, 146
- Erythema multiforme, 481
Coccidioidomycosis, 151
- Erythema nodosum, 151, 482
inflammatory bowel disease, 382
- Erythroblastosis fetalis, 405
- Erythrocytes, **407**
blood types, 405
casts in urine, 594
Coombs test, 423
DAF deficiency and, 107
erythropoietin and, 589
glucose usage by, 334
hereditary spherocytosis, 422
myeloproliferative disorders, 433
transfusion of, 429
- Erythrocyte sedimentation rate (ESR), **214**
subacute granulomatous thyroiditis, 341
- Erythrocytosis, 407
oxygen-hemoglobin dissociation curve, 666
- Erythrocytic toxin, 136
- Erythromelalgia, 433
- Erythromycin
macrolides, 193
prophylaxis, 198
protein synthesis inhibition, 191
reactions to, 249
- erythroplasia of Queyrat, 651
- Erythropoiesis, 679
fetal, 404
- Erythropoietin (EPO), 121
anemia of chronic disease, 421
aplastic anemia, 421
high altitude, 670
with pheochromocytoma, 350
polycythemia and, 228
- release of, 589
in renal failure, 603
signaling pathways for, 337
- Eschar, 132
in cutaneous anthrax, 137
with mucormycosis, 153
- Escherichia coli*, **145**
cephalosporins, 189
culture requirements, 126
encapsulation, 127
galactosemia, 80
lac operon, 40
lactose fermentation, 144
meningitis, 180
neonatal illness, 182
nosocomial infection, 185
O157-H7, 132, 145, 178, 179
penicillins for, 188
pneumonia, 179
prostatitis, 654
urinary tract infections, 181, 600
- Escitalopram, 575
- E-selectin, 215
- Esmolol, 245, 323
- Esomeprazole, 399
- Esophageal adenocarcinoma, 378
- Esophageal atresia, 359
- Esophageal cancer, **378**
achalasia and, 376
- Esophageal dysmotility
CREST syndrome, 473
- Esophageal perforation, 377
- Esophageal strictures, 377
- Esophageal varices, 365, 377
- Esophageal webs, 418
- Esophagitis, 377
bisphosphonates, 486
drug reaction and, 249
HIV-positive adults, 177
- Esophagus
blood supply and innervation, 364
diaphragm, 663
histology of, 362
pathologies of, **377**
portosystemic anastomosis, 365
- Essential amino acids, 81
- Essential fructosuria, 80
- Essential hypertension, 316
- Essential mixed cryoglobulinemia, 173
- Essential thrombocythemia, 433
- Essential tremor, 519
- Esters (local anesthetics), 550
- Estrogen, **656**
androgen insensitivity syndrome, 639
androgen conversion to, 636
benign breast tumors, 649
bone formation, 459
epiphyseal plate closure, 636
gynecomastia, 649
menopause, 636
menstrual cycle, 632
osteoporosis, 462
ovulation, 631
premature ovarian failure, 636, 645
prolactin suppression of, 330
signaling pathways for, 337
source and function of, **630**
Turner syndrome, 638
- Estrogen receptor modulators (selective), 656
- Eszopiclone, 546
- Etanercept, 487
- Ethacrynic acid, 608
- Ethambutol, 196, **197**

- Ethanol
 as carcinogen, 225
 gluconeogenesis and, 72
 lactic acidosis and, 72
 metabolism, **72**, 232
 NADPH (nicotinamide adenine dinucleotide phosphate), 72
 zero-order elimination, 232
- Ethics, **265–268**
 confidentiality, 267
 consent, 265
 core principles of, **265–267**
 directives, 268
 religious beliefs and, 269
 situations in, 268–269, 269–270
- Ethinyl estradiol, 656, 657
- Ethosuximide
 absence seizures, 544
- Ethylenediaminetetraacetic (EDTA), 419
- Ethylene glycol
 toxicity treatment, 248
- Ethylene oxide, 204
- Etonogestrel, 657
- Etoposide/teniposide, **442**
 targets of, 438
 teniposide, 38
- Euchromatin, 34
- Eukaryotes
 DNA replication, 38
 functional gene organization, 41
 mRNA start codons, 44
 ribosomes in, 45
 RNA polymerase in, 42
 RNA processing, 41
- Eukaryotic initiation factors, 45
- Eukaryotic release factors, 45
- Eustachian tubes
 embryonic derivation, 621
- Eversion, 453
- Evolucumab, 320
- Ewing sarcoma, 465
 dactinomycin for, 439
- Exanthem subitum, 165
- “Excision” event, 130
- Excitatory pathway, 500
- Exclusive provider organization plan, 271
- Executioner caspases, 208
- Exemestane, 656
- Exenatide, 353
- Exercise
 blood flow autoregulation, 297
 peripheral resistance, 286
 respiratory response, 670
 syncope with, 308
 Tetralogy of Fallot, 298
- Exercise-induced amenorrhea, 645
- Exocrine glands, 236
- Exocytosis, 50
- Exogenous corticosteroids, 336
- Exons
 deletions in muscular dystrophies, 61
 vs introns, 43
- Exotoxins
 features of, 131
 organisms with, 132–133
Pseudomonas aeruginosa, 132
Streptococcus pyogenes, 133
- Expectorants, 686
- Expiratory reserve volume (ERV), 664
- Extension
 hip, 451, 453
- External hemorrhoids, 366
- External iliac lymph nodes, 624
- External oblique muscle
 inguinal canal and, 369
- External rotation
 arm (rotator cuff), 446
 hip, 451
- External spermatic fascia, 369
- Extinction (conditioning), 554
- Extracellular fluid (ECF)
 volume measurement, 581
 volume regulation, 588
- Extragenital germ cell tumors, **652**
- Extramammary Paget disease, 644
- Extraperitoneal tissue, 369
- Extravascular hemolysis, **421**
- Extrinsic hemolytic anemia, **423**
- Extrinsic pathway, 208
 warfarin and, 436, 437
- Exudate
 “anchovy paste,” 155
- Ex vacuo ventriculomegaly, 522
- Eye movements, 540
 bilateral, 543
 cranial nerve palsies, 541
 with stroke, 515
- Eyes
 anatomy of, 534
 aqueous humor pathway, 535
- Ezetimibe, 320
- Diarrhea, 249
- F**
- Fab region of antibodies, 104
- Fabry disease, 61
- Facial nerve (Bell) palsy, 146, 186
- Facial nerve (CN VII), 146, 186
 functions of, 506
 inflammatory demyelinating polyradiculopathy, 524
 lesions of, **532**
 pharyngeal arch derivation, 620
 tongue, 493
- Facies
 congenital syphilis, 147
 elfin, 64
 epicanthal folds, 63, 64
 “facial plethora,” 685
 in fetal alcohol syndrome, 615
 flat, 63
 leonine (lion-like), 141
 long face with a large jaw, 62
 risus sardonius, 138
 twisted face, 578
- Factitious disorder
 characteristics of, **566**
 on another, 566
 on self, 566
- Factor IX concentrate, 426
- Factor VIII concentrate, 426
- Factor V Leiden, 413, 428
 venous sinus thrombosis and, 503
- Factor Xa
 direct inhibitors of, 437
 heparin effect on, 436
 inhibitors of, 413
- Factor XI concentrate, 426
- Facultative anaerobes
 culture requirements, 127
- Facultative intracellular organisms, 127
- FADH (flavin adenine dinucleotide), 77
- Failure mode and effects analysis, 274
- Failure to thrive, 556
 galactosemia, 80
 orotic aciduria, 420
 SCID, 117
- Falciform ligament, 361
- Fallopian tubes
 anatomy, 625
 epithelial histology, 626
 fertilization, 633
- False-negative rate, 257
- False-positive rate, 257
- Famciclovir, **201**
- Familial adenomatous polyposis, 387
 APC gene and, 389
 chromosome association, 64
- Familial amyloid cardiomyopathy, 212
- Familial amyloid polyneuropathies, 212
- Familial dyslipidemias, **94**
- Familial hypercholesterolemia, 60, 94
- Familial hypocalciuric hypercalcemia, **345**
- Family discussions, 268
- Family therapy
 separation anxiety, 557
- Famotidine, 399
- Fanconi anemia, 421
 DNA repair in, 40
 nonhomologous end joining and, 40
- Fanconi syndrome, 586
 drug reaction and, 251
 renal tubular acidosis, 593
- Fascia
 collagen in, 50
- Fascia of Buck, 627
- Fasciculations, 529
- Fastigial nucleus, 499
- Fasting plasma glucose test, 346
- Fasting state, 76, 91
- Fat emboli, 672
- Fatigue
 adrenal insufficiency, 349
 heart failure and, 309
 MI signs, 305
- Fatigue, medical errors and, **274**
- Fat necrosis, 209, 649
- Fat redistribution, 250
- Fat-soluble vitamins, 65
- Fatty acids
 gluconeogenesis, 78
 metabolism of, 72, **89**, 90
 oxidation of, 72, 73
 synthesis, 73
- Fatty acid synthase, 67
- Fatty casts, 594
- Fatty liver disease
 hepatocellular carcinoma and, 392
 nonalcoholic, 391
- Fc region of antibodies, 104
- Fear
 anxiety disorder and, 562
 panic disorder and, 563
- Febrile nonhemolytic transfusion reaction, 114
- Febrile pharyngitis, 164
- Febrile seizures, 517
- Febuxostat
 gout, 487
 Lesch-Nyhan syndrome, 37
- Fecal elastase, 381
- Fecal immunochemical testing (FIT), 388
- Fecalith obstruction, 383
- Fecal microbiota transplant, 138
- Fecal occult blood testing (FOBT), 388
- Fecal retention, 558
- Feces
 explosive expulsion of, 384
- Federation of State Medical Boards (FSMB), 2
- Fed state, 76, 91
- Fee for service, 271
- Felty syndrome, 466
- Female genital embryology, 622
- Female/male genital homologs, 623
- Female reproductive anatomy, **625**
- Female reproductive epithelial histology, **626**
- Femoral artery, 368
- Femoral head
 osteonecrosis, 463
- Femoral hernia, 370
- Femoral nerve, 452
- Femoral region, **368**
- Femoral sheath, 368
- Femoral triangle, 368
- Femoral vein, 368
- Fenestrated capillaries, 496, 581
- Fenofibrate, 320
- Fenoldopam, 242, 318
- Fentanyl, 551
- Ferritin, 213
 anemia of chronic disease, 421
 iron deficiency anemia, 418
 lab values in anemia, 419
 sideroblastic anemia, 419
- Ferrochelatase, 425
- Fertility
 double Y males, 638
 GnRH and, 328
 menstrual cycle, 632
- Fertilization, 631, 633
- Fetal alcohol syndrome, 300, 614, **615**
 holoprosencephaly in, 491
- Fetal circulation, **282**
- Fetal development
 early, 612
 placental component, 617
- Fetal erythropoiesis, **404**
- Fetal hypothyroidism, 341
- Fetal lung maturity, 661
- Fetal movement, 612
- Fetal-postnatal derivatives, **282**
- Fetal respiration, 660
- Fetal tissue
 collagen in, 50
- Fever
 amphotericin B, 199
 childhood rashes, 183
 clindamycin, 192
 complicated hernias, 370
 endotoxins, 131
 epiglottitis, 186
 exotoxins, 133
 genital herpes, 184
 vs heat stroke, **517**
 high fever, 165, 168, 171, 183
 with inflammation, 213
 Jarisch-Herxheimer reaction, 148
 Legionnaires’ disease, 143
 low-grade, 143, 171
 malaria, 157
 with meningococci, 142
 mononucleosis, 165
 neuroleptic malignant syndrome, 569
 pulmonary anthrax, 137
 recurring, 156
Rickettsia rickettsii, 150
Salmonella spp, 149
Salmonella typhi, 144
 seizures with, 165
 spiking, 158

- Tetralogy of Fallot, 298
 thyroid storm causing, 342
 toxic shock syndrome, 135
Trichinella spiralis, 159
 tuberculosis, 140
 undulant, 143
 vasculitides, 314
 Waterhouse-Friderichsen syndrome, 142
 Weil disease, 147
 Fexofenadine, 686
 Fibrates, 320
 hepatitis and, 249
 myopathy and, 250
 Fibrinogen, 213
 in cryoprecipitate, 429
 ESR and, 214
 receptor for, 407
 thrombocytes, 407
 Fibrinoid necrosis, 209
 Fibrinous pericarditis, 305
 Fibroadenoma, 649
 Fibroblast growth factor (FGF)
 signaling pathways for, 337
 in wound healing, 216
 Fibroblast growth factor receptor
 (FGFR3), 462
 Fibroblasts
 cortisol and, 336
 in wound healing, 216
 Fibrocystic breast disease, 649
 Fibro fog, 471
 Fibroid (leiomyoma)
 leuprolide for, 656
 Fibromas, 647
 nomenclature for, 220
 Fibromuscular dysplasia, 300, 604
 Fibromyalgia, 470, **471**, 575
 Fibronectin
 in cryoprecipitate, 429
 thrombocytes, 407
 Fibrosarcomas, 220
 Fibrosis
 diffusion-limited gas exchange, 668
 silicosis, 677
 Fibrous plaque in atherosclerosis, 302
 Fick principle, 285
 Fidaxomicin
 Clostridium difficile, 138
 Fifth disease
 rash, 183
 50S inhibitors, 191
 Filgrastim (G-CSF), 121
 Filoviruses
 characteristics of, 167
 negative-stranded, 168
 Filtration fraction
 glomerular dynamics, **583**
 Fimbria, 124
 Financial considerations in treatment,
 269
 Finasteride, 658
 gynecomastia with, 649
 reproductive hormones and, 636
 Finger agnosia, 511
 Finger drop, 447
 Finger movements, 450
 upper extremity nerve injury, 447
 Fingernails, 478
 Finkelstein test, 461
 First-degree AV block, 295
 First-order elimination, 231, **232**
 Fish oil, 320
 Fishy smell, 148
 Fitz-Hugh-Curtis syndrome, 142, 185
 5-aminosalicylic drugs, 382, 400
 5 α -reductase
 deficiency, 622, **639**
 inhibitors for BPH, 654
 testosterone conversion, 636
 5-fluorouracil (5-FU)
 antimetabolites, 440
 photosensitivity, 250
 pyrimidine synthesis and, 36
 targets of, 438
 thymidylate synthase, 36
 5-HT_{1B/1D} agonists, 253
 5-HT agonists, 547
 5-HT
 opioid effects, 551
 MAO inhibitor effect on, 575
 trazodone effects, 576
 vilazodone effects, 576
 vortioxetine effects, 576
 Fixation, 555
 Fixed splitting, 289
 Flaccid paralysis
 botulinum toxin, 138
 LMN lesion, 531
 motor neuron signs, 529
 Flagellin, 99
 Flagellum, 124
 Flask-shaped ulcers, 155
 Flat affect, 499
 Flavin nucleotides, 75
 Flaviviruses, 163, 167
 Fleas (disease vectors), 149, 150
 Flecainide, 322
 Flexion
 foot, 453
 hand, 450
 hip, 451
 Flexor digiti minimi muscle, 450
 Flexor pollicis brevis muscle, 450
 Flies (disease vectors), 144, 159
 Floppy baby syndrome
 Clostridium botulinum, 138
 spinal cord lesions, 530
 Flow volume loops, **673**
 Fluconazole, 151
 Cryptococcus neoformans, 153
 mechanism and use, 199
 opportunistic fungal infections,
 153
 Flucytosine, **199**
 Cryptococcus neoformans, 153
 Fludrocortisone, **354**
 Fluid compartments, **581**
 Flumazenil
 benzodiazepine overdose, 248,
 546, 570
 nonbenzodiazepine hypnotics, 546
 Fluorescence in situ hybridization, **55**
 Fluorescent antibody stain
 bacteria, 125
 Fluoroquinolones, 38
 mechanism and use, **195**
 Mycoplasma pneumoniae, 150
 pregnancy contraindication, 204
 Pseudomonas aeruginosa, 143
 tendon/cartilage damage with, 250
 TOP II (DNA gyrase) and TOP IV
 inhibition in prokaryotes,
 38
 typhoid fever, 144
 Fluoxetine, 575
 Fluphenazine, 573
 Tourette syndrome, 572
 Flutamide, 658
 Fluticasone, 687
 Fluvoxamine, 575
FMR1 gene, 62
 Foam cells
 in atherosclerosis, 302
 Niemann-Pick disease, 88
 Focal glomerular disorders, 594
 Focal hepatic necrosis, 249
 Focal necrotizing vasculitis, 315
 Focal neurological deficits
 pituitary apoplexy, 339
 Focal segmental glomerulosclerosis,
 597
 Focal seizures, 517
 Folate antagonist
 teratogenicity, 614
 Folate synthesis
 inhibition/block, 194
 Folic acid
 antimicrobials and, 187
 folate, 68
 in pregnancy, 68
 neural tube defects and, 491
 Follicles (lymph), 96
 Follicle-stimulating hormone (FSH)
 clomiphene effect, 656
 hCG and, 633
 PCOS, 645
 premature ovarian failure, 636
 secretion of, 327
 signaling pathways of, 337
 Follicular conjunctivitis, 148
 Follicular cysts, 646
 Follicular lymphomas, **430**, 434
 Follicular phase (menstrual cycle), 632
 Follicular thyroid carcinomas, 343
 Fomepizole
 ethanol metabolism and, 72
 toxicity treatment with, 248
 Food-borne illness, 178
 Food poisoning
 Bacillus cereus, 138, 178
 causes of, 178
 Staphylococcus aureus, 135, 178
 toxic shock syndrome toxin, 133
 Food toxins, 247
 Foot drop, 453
 lead poisoning, 419
 Foramen cecum, 326
 Foramen of Magendie, 504
 Foramen of Monro, 504
 Foramen ovale
 atrial septal defect, 299
 embryology, 280
 fetal circulation, 282
 retained patency of, 298
 Foramen primum, 280
 Foramen secundum, 280
 Foramina of Luschka, 504
 Forced expiratory volume (FEV)
 obstructive lung disease, 674
 restrictive lung disease, 675
 Forebrain, 490
 Foregut
 blood supply/innervation of, 364
 development of, 358
 Foreign body inhalation, 663
 Formoterol, 687
 Fornix (uterus), 625
 45,XO, 638
 47,XXY, 638
 46,XX/46,XY DSD, 639
 Fosamprenavir
 HIV-positive adults, 203
 Foscarnet, **202**
 Fosphenytoin, 544
 Fossa ovalis, 282
 Fovea
 cherry-red spot, 538
 FOXP3 protein, 102
 Fractures
 bone diseases and, 51
 chalk-stick, 463
 common pediatric, 462
 compartment syndrome with, 461
 humerus, 447
 in child abuse, 556
 pathologic, 465
 scaphoid, 449
 vertebral compression, 462
 Fragile X syndrome, **62**
 chromosome association, 64
 Frameshift mutations
 deletions, 61
 muscular dystrophy and, 61
Francisella spp
 intracellular organism, **127**
Francisella tularensis
 animal transmission, 149
 Frataxin, 531
 Free fatty acids
 fast/starvation states, 91
 lipid transport and, 92
 Free light chain (FLC) assay
 plasma cell dyscrasias, 431
 Free nerve endings, 494
 Free radical injury, **210**
 Fremitus (tactile), 680, 682
 Fresh frozen plasma, 429
 "Fried egg" cells, 494, 526
 Friedrich ataxia, **531**
 chromosome association, 64
 hypertrophic cardiomyopathy, 308
 inheritance of, 60
 mechanism of, 62
 Frontal bossing, 339
 Frontal eye fields
 cortical functions, 501
 lesions in, 511
 Frontal lobe
 lesions in, 511
 stroke effects, 514
 Frontotemporal dementia, 520
 Fructokinase, 80
 Fructose-1,6-bisphosphatase, 73
 gluconeogenesis, 78
 in metabolic pathways, 74
 Fructose-2,6-bisphosphate, 76
 Fructose intolerance, 80
 Fructose metabolism
 disorders, 80
 pathways, 74
 Fructosuria, 80
 FTA-ABS, 125, 147
 Fumarate, 82
 Functional hypothalamic
 amenorrhea, **645**
 Functional neurologic symptom
 disorder, 566
 Functional residual capacity (FRC),
 664
 Fungal infections
 IL-12 receptor deficiency, 116
 infections with
 immunodeficiencies, 118
 Fungi
 culture requirements, 126
 immunocompromised patients,
 179
 necrosis and, 209
 opportunistic infections, 153
 silver stain, 125
 topical infections, 199
 Funny current, 292
 "funny" sodium channels, 324

- Furosemide, 252, 608
gout with, 250
pancreatitis, 249
- Fusion inhibitors, 203
- Fusion protein EWS-FL11, 465
- Fusobacterium* spp
alcoholism, 179
anaerobic metabolism of, 127
- G**
- G6PD
deficiency, 61, **79**
HMP shunt and, 73
in respiratory burst, 109
- G6PD deficiency, **422**
- GABA
anesthesia effects, 550
barbiturate effects, 546
basal ganglia and, 500
benzodiazepine effects, 546
changes with disease, 495
epilepsy drugs, 544
Huntington disease, 520
vitamin B₆ and, 67
- Gabapentin, 544
- GABA_B receptor agonists, 523, 551
- gag* gene, 175
- Gag reflex, 507
- Gait disorders
“steppage,” 453
Trendelenburg sign/gait, 453
- Gait disturbance
cerebellar lesions and, 499
Friedreich ataxia, 531
Parkinson disease, 520
vitamin B₁₂ deficiency, 530
waddling, 61
- Galactocerebrosidase, 88
- Galactocerebroside, 88
- Galactokinase deficiency, 80
cataracts and, 535
- Galactorrhea
antipsychotic drugs and, 328
pituitary prolactinomas, 328
tuberoinfundibular pathway, 499
- Galactose-1-phosphate
uridylyltransferase, 80
- Galactose metabolism
disorders of, 80
- Galactosemia, 80
cataracts and, 535
- Galantamine, 240, 549
- Galant reflex, 510
- Gallbladder
biliary structures, 368
blood supply and innervation of, 364
regulatory substances, 371
- Gallbladder cancer
porcelain gallbladder and, 397
sclerosing cholangitis and, 395
- Gallstone ileus, 396
- γ-glutamyltransferase (GGT)
alcohol use, 570
γ-glutamyl transpeptidase (GGT), 390
- γ-interferon, 407
- Ganciclovir, **202**
agranulocytosis, 250
- Ganglion cyst, 461
- Ganglioneuromatosis
oral/intestinal, 351
- Gangrene
Buerger disease, 314
diabetes mellitus, 346
- Gangrenous necrosis, 209
- Gap junctions, 474
- Gardener’s pupil, 241
- Gardnerella vaginalis*, **148**
- Gardner syndrome, 387
- Gargoylism, 88
- Gas gangrene
alpha toxin, 133
Clostridium perfringens, 138, 179
- Gastrectomy, 420
- Gastric acid, 372
histamine receptors and, 238
regulatory substances and, 371
- Gastric arteries
celiac trunk, 364
intraglmental, 361
- Gastric bypass surgery
ghrelin and, 371
vitamin B₁₂ deficiency, 69
- Gastric cancer, **379**
carcinogens causing, 225
Helicobacter pylori, 146
oncogenes and, 224
oncogenic microbes and, 226
sign of Leser-Trélat and, 228
trastuzumab for, 443
types of, 379
- Gastric inhibitory peptide (GIP), 351
- Gastric outlet obstruction, 359, 380
- Gastric sclerosis, 473
- Gastric ulcers, 380
NSAID toxicity, 486
- Gastric vessels, 361
- Gastrin, 371, 373
signaling pathways for, 337
somatostatinomas and, 351
- Gastrinomas, 354, 371
- Gastrin-releasing peptide (GRP), 371
- Gastritis, 146, **379**
gastrin in, 371
H₂ blockers for, 399
proton pump inhibitors for, 399
stomach cancer and, 379
- Gastrocolic ligament, 361
- Gastroenteritis
caliciviruses, 167
Listeria monocytogenes, 139
rotavirus, 168
Salmonella spp, **144**
- Gastroepiploic arteries, 361, 364
- Gastroesophageal reflux disease (GERD)
Barrett esophagus, 378
esophageal cancer and, 378
presentation, 377
- Gastrohepatic ligament, 361
- Gastrointestinal bleeding
hereditary hemorrhagic telangiectasia, 316
- Gastrointestinal drug reactions, 249
- Gastrointestinal stromal tumors (GISTs), 224
- Gastrointestinal system
anatomy, 360–369
blood supply to, 363
embryology, 358–359
innervation of, **364**
ligaments, 361
pathology, 376–397
pharmacology, 398–400
physiology, 371–375
regulatory substances, **371**
secretory cells, 373
secretory products, **372**
- Gastroschisis, 358
- Gastrosplenic ligament, 361
- Gastrulation, 612
- Gaucher disease, 88
osteonecrosis, 463
osteonecrosis in, 463
- Gaussian distribution, 262
- G cells, 371
- Gemfibrozil, 320
- Gemifloxacin, 195
- Gender dysphoria, **567**
- Gender identity, 635
- Gene expression
modifications, **56**
regulation, 41
- Generalized anxiety disorder (GAD), **563**
buspirone, 574
drug therapy for, 572
Selective serotonin reuptake inhibitors (SSRIs) for, 575
serotonin-norepinephrine reuptake inhibitors (SNRIs) for, 575
- Generalized seizures, 517
- Genes
introns vs exons, 42, 43
- Genetics, 56–65
anticipation, 62
autosomal dominant diseases, 60
autosomal recessive diseases, 60
autosomal trisomies, 63
bacterial, 130–204, 131
chromosome disorders, **64**
code features, **37**
inheritance modes, 59
muscular dystrophies, 61
terms, **56–57**
trinucleotide repeat expansion diseases, 62
viral, 162–163
X-linked recessive disorders, 61
- Geniculate nuclei (thalamus), 498
- Genital herpes, 184
- Genitalia
ambiguous, 622, 638, 639
embryology of, 612, **622**
estrogen and, 630
male/female homologs, 623
- Genital tubercles, 624
- Genital ulcers, 184
- Genital warts, 184
- Genitofemoral nerve, 452
- Genitourinary/renal drug reactions, 251
- Genome editing
reverse transcriptase polymerase chain reaction, 52
- Genotyping microarrays, 54
- Gentamicin, 191
- Genu varum (bow legs), 463
- Geriatric patients
atropine in, 241
Beers criteria in, 247
changes in, 270
colonic ischemia and, 386
colorectal cancer, 388
common cause of death, 272
drug-related delirium in, 558, 575
lipofuscin in, 211
Medicare for, 272
nosocomial infections, 185
osteoporosis, 462
PPI adverse effects, 399
respiratory system changes in, 665
vascular skin tumors, 478
Zenker diverticulum, 384
- Germ cell tumors
cryptorchidism risk for, 651
hormone levels in, 653
testicular, 652
- Germinal centers of lymph nodes, 96
- Germinal center (spleen), 98
- Gerstmann syndrome, 511
- Gestational age calculation, 633
- Gestational diabetes, 634
- Gestational hypertension, 643
- GFAP (glial fibrillary acid proteins), 48, 227
astrocyte marker, 493
cytoskeletal elements, 48
stain, 227
- Ghrelin, 336, 371
- Giant cells
with Aschoff bodies, 312
astrocytomas, 525
Warthin-Finkeldey, 170
- Giant cell (temporal) arteritis, 314, 518
polymyalgia rheumatica, 470
- Giant cell tumor, 464
- Giardia* spp
fluorescent antibody stain, 125
watery diarrhea, 179
- Giardia lamblia*, **155**
- Giardiasis, 155
- Giemsa stain, 125
Borrelia, 146
chlamydiae, 148
- Gigantism, 329, 339
- Gilbert syndrome, 393, 394
- Gingival hyperplasia
calcium channel blockers, 318
cyclosporine, 120
drug reaction and, 250
- Gingivostomatitis, 164
- Gitelman syndrome, 586
markers in, 591
- Glans penis
lymphatic drainage of, 624
- Glanzmann thrombasthenia, 427
- Glaucoma, 242
atropine, 241
β-blockers for, 245
carbachol for, 240
closed-angle, 240
diabetes mellitus and, 346
diagnosis of, 240
epinephrine for, 242
open-angle, 240, 242
pilocarpine for, 240
types of, **536**
- Glioblastoma multiforme, 526
nitrosoureas for, 441
- Glipizide, 353
- Glitazones/thiazolidinediones, 353
- Global aphasia, 516
- Global payment, 271
- Globoid cells
Krabbe disease, 88
- Globose nucleus, 499
- Globus pallidus externus, 500
- Glomerular disorders/disease
etiology and presentation, 594
nomenclature, 594
types of, **595**
- Glomerular filtration barrier, **581**
- Glomerular filtration parameters, 583
- Glomerular filtration rate (GFR), **582**
ACE inhibitor effects, 610
glomerular dynamics in, 583
juxtaglomerular apparatus, 589

- Glomerulonephritis
 azathioprine for, 120
 bacterial endocarditis, 311
 RBC casts in, 594
Streptococcus pyogenes, 133, 136
 Wegener granulomatosis, 315
- Glomerulus
 dynamics of, **583**
- Glomus tumors, 478
- Glossitis
 B-complex deficiency, 65
 megaloblastic anemia, 420
 vitamin B₃ deficiency, 67
 vitamin B₉ deficiency, 68
- Glossopharyngeal nerve (CN IX), 506
 blood flow regulation, 296
 pharyngeal arch derivative, 620
 tongue, 493
- Glossopostosis, 620
- Glove and stocking neuropathy, 346
- GLP-1 analogs, 353
- Glucagon, **333**
 for β -blocker toxicity, 323
 fructose biphosphatase-2, 76
 glucagonomas and, 351
 glycogen regulation, 85
 insulin and, 333, 334
 production of, 331
 somatostatin and, 371
 somatostatinomas and, 351
- Glucagonomas
 MEN 1 syndrome, **351**
 pancreatic cell tumor, 351
 somatostatin for, 354
- Glucocerebrosidase
 Gaucher disease, 88
- Glucocerebroside
 in sphingolipidoses, 88
- Glucocorticoids
 adrenal insufficiency, 349
 calcium pyrophosphate deposition disease, 467
 diabetes mellitus, 346
 fat redistribution with, 250
 gout, 467, 487
 immunosuppression, 120
 myopathy, 250
 rheumatoid arthritis, 466
- Glucokinase
 hexokinase vs, 75
 metabolic pathways, 74
- Glucconeogenesis
 cortisol and, 336
 ethanol metabolism and, 72
 irreversible enzymes, **78**
 metabolic site, 72
 pyruvate metabolism and, 77
 rate-determining enzyme for, 73
 thyroid hormone and, 331
- Glucose
 ATP production, 74
 blood-brain barrier and, 496
 clearance, **584**
 glycogen metabolism, 86
 insulin and, 334
 metabolism of, 40
 for porphyria, 425
 transporters, 334
- Glucose-6-phosphatase
 dehydrogenase deficiency, **79**
 gluconeogenesis, 78
 HMP shunt, 79
 Von Gierke disease, 87
- Glucose-dependent insulinotropic peptide (GIP), 351, **371**, 372
 polypeptide (GIP), 334
- Glucosuria
 glucose clearance, 584
 threshold for, 584
- Glucuronidation
 drug metabolism, 232
- Glutamine, 352
- Glutamic acid
 ammonia transport, 82
 classification of, 81
 opioid analgesics and, 551
- Glutathione
 acetaminophen and, 485
 in G6PD deficiency, 422
- Glutathione peroxidase, 109
 free radical elimination by, 210
- Glutathione reductase, 109
 NADPH and, 75
- Gluten-sensitive enteropathy, 381
- Gluteus maximus muscle, 453
- Gluteus minimus muscle, 451
- GLUT transporters, 334
- Glyburide, 353
- Glycerol
 starvation days and, 91
- Glycogen
 metabolism and storage, 73, **86**
- Glycogenesis, 73
- Glycogenolysis
 rate-determining enzyme for, 73
 thyroid hormone and, 331
- Glycogen storage diseases, **87**
- Glycogen synthase, 73, 86
- Glycolysis
 arsenic and, 74
 hexokinase/glucokinase in, 75
 metabolic site, 72
 pyruvate metabolism and, 77
 rate-determining enzyme for, 73
 regulation of, **76**
- Glycoprotein IIb/IIIa inhibitors, **438**
- Glycoproteins
 bacterial pilus/fimbria, 124
 HIV, 175
- Glycopyrrolate, 241
- Glycosylation
 collagen synthesis, 50
 protein synthesis, 45
- GNAQ gene, 525
- Goblet cells, 362, 662
- Goiter
 maternal hypothyroidism from, 341
 maternal iodine deficiency, 614
 in Riedel thyroiditis, 341
 types and causes of, 342
- Golfer's elbow, 459
- Golgi apparatus
 in plasma cells, 409
- Golgi tendon organ, 458
- Golimumab, 487
- Gonadal drainage, **624**
- Gonadal mosaicism, 57
- Gonadotropin, 646
- Gonadotropin-releasing hormone (GnRH)
 function of, 328
 neurons producing, 498
 ovulation, 631
 prolactin and, 330
 signaling pathways for, 337
 spermatogenesis, 628
- Gonads
 dysgenesis of, 606
- Gonococcal arthritis, 468
- Gonococci, vs meningococci, 142
- Gonorrhea
 ceftriaxone, 189
 gonococci, 142
 STI, 184
- Goodpasture syndrome, 50, 596
 autoantibody, 115
 HLA-DR2, 100
 restrictive lung disease, 675
 type II hypersensitivity reactions, 112
- Good syndrome
 paraneoplastic syndrome, **228**
 thymoma and, 98
- Goserelin, **656**
- Gottron papules, 228, 471
- Gout, **467**
 drug reaction and, 250
 drug therapy for, **487**
 kidney stones and, 598
 Lesch-Nyhan syndrome, 37
 loop diuretics and, 608
 Von Gierke disease, 87
- Gower maneuver/sign, 61
- gp41, 203
- G-protein-coupled receptors, 236
- G-protein-linked 2nd messengers, **238**
- Gracilis, 452
- Graft-versus-host disease, 119
 type IV hypersensitivity, 113
- Gram-negative organisms
 cell wall structure, **124**
 cephalosporins, 189
 lab algorithm, **141**
- Gram-positive organisms
 antibiotic tests, **134**
 cell wall structure, **124**
 cephalosporins, 189
 lab algorithm, **134**
 vancomycin, **190**
- Gram stain
 peptidoglycan layer and, 125
- Granular casts
 acute tubular necrosis, 602
 "muddy brown" in urine, 594
- Granulocyte-colony stimulating factor (G-CSF), 337
- Granulocytes
 morulae, 150
- Granulocytopenia
 trimethoprim, 194
- Granuloma inguinale, 184
- Granulomas, 147
 macrophages and, 407
 in systemic mycoses, 151
 TNF- α and, 110
 in tuberculosis, 140
- Granulomatosis infantiseptica
Listeria monocytogenes, 139
- Granulomatosis with polyangiitis (Wegener)
 restrictive lung disease and, 675
- Granulomatous disease
 catalase + organism infections with, 128
 excess vitamin D in, 70
 hypervitaminosis D with, 464
- Granulomatous inflammation, **217**
- Granulosa cells
 tumors of, 647
- Granzyme B
 cytotoxic T cells, 101, **102**
 extrinsic pathway and, 208
- Grapefruit juice and cytochrome P-450, 252
- Graves disease
 autoantibody, 115
 HLA subtype associations, 100
- hyperthyroidism, 342
 ophthalmopathy, 340
 thyroid cellular action in, 331
 type II hypersensitivity, 112
- Gray baby syndrome, **192**, 204, 250
- Gray hepatization, 683
- Grazoprevir, 204
- Greater omental sac, 361
- Great vein of Galen, 503
- Green twig (greenstick) fracture, 462
- Grief, **562**
- Griseofulvin, **200**
 cytochrome P-450 interaction, 252
 microtubules and, 48
 pregnancy contraindication, 204
 "Ground-glass" appearance (X-ray), 177, 661
Pneumocystis jirovecii, **154**
- Growth factors
 tumor suppressor gene mutations and, 46
- Growth hormone (GH), 354
 diabetes mellitus, 346
 ghrelin and, 336
 for hypopituitarism, 339
 insulin resistance and, **329**
 secretion of, 327
 signaling pathways for, 337
 somatostatin, 339
- Growth-hormone-releasing hormone (GHRH)
 function of, 328
- Growth media properties, 126
- Growth retardation
 with renal failure, 603
- GTPase, 224
- GTP (guanosine triphosphate), 77
- Guaifenesin, **686**
- Guanfacine, 240, 243
- Guanosine analogs
 mechanism and use, 201
- Gubernaculum, 624, 625
- Guessing during USMLE Step 1 exam, 23
- Guillain-Barré syndrome
 acute inflammatory demyelinating polyradiculopathy, 524
Campylobacter jejuni, 145
 restrictive lung disease, 675
 Schwann cell injury, 494
 Schwann cells, 524
- Gummas
 syphilis, 147, 184
- Gustatory hallucinations, 559
- Gustatory pathway
 cranial nerves in, 532
 thalamic relay for, 498
- Guyon canal syndrome, 459
- Gynecologic procedures
 ureteric damage in, 581
- Gynecologic tumor epidemiology, **643**
- Gynecomastia, 649
 antiandrogens for, 658
 azoles, 199
 cimetidine, 399
 SHBG and, 337
 spironolactone, 658
 tuberoinfundibular pathway, 499

HH₁ blockers, 251, **686**H₂

production in tissues, 127

- H₂ blockers, **399**
H₂O₂ degradation
 catalase and, 128
H₂-antagonist
 naming conventions for, 254
Haemophilus ducreyi
 sexual transmission, 184
Haemophilus influenzae, **142**
 biofilm production, 128
 cephalosporins, 189
 chloramphenicol, 192
 culture requirements, 126
 encapsulation, 127
 IgA protease, 129
 influenza, 169
 meningitis, 179, 180
 penicillins for, 188
 pneumonia, 179
 postviral infection, 179
 rifamycins, 196
 transformation, 130
 type b conjugate vaccine, 127
 unvaccinated children, 186
 vaccine, 142, 180
Hair
 “kinky,” 51
 Menkes disease, 51
 vitamin C deficiency, 69
Hairy cell leukemia, 227, **432**
 cladribine for, 440
Hairy leukoplakia, 479
 HIV-positive adults, 177
Half-life equation, 231
Halitosis
 feter hepaticus, 389
 Zenker diverticulum, 384
Hallucinations, 559
 cocaine, 571
 delirium, 558
 mesolimbic pathway, 499
 pellagra, 67
 postpartum psychosis, 562
 tricyclic antidepressants, 575
Hallucinogen intoxication and
 withdrawal, 571
Haloperidol, 573
 torsades de pointes, **294**
Halothane, 550
 hepatic necrosis, **249**
Hamartin protein, 224, 525
Hamartomas, **220**, 525
Hamartomatous colonic polyps, **387**
Hamate bone, 449
 fracture of hook, 447
Hamman sign crepitus, 672
Hammer toes, 531
Hand
 distortions of, 451
 injuries of, 459
 muscles of, **450**
 squamous cell carcinoma, 484
Hand-foot-mouth disease, 183
Hansen disease, 141
 animal transmission, 149
 dapson, 194
 erythema nodosum, 482
Hantavirus, 167
Haptens
 acute interstitial nephritis, 601
 amiodarone as, 323
Haptoglobin, 421
Hardy-Weinberg population genetics,
 57
Hartnup disease, 67
 vitamin B₃ deficiency, 67
Hashimoto thyroiditis, 341
 autoantibody, 115
 HLA subtype association, 100
Hassall corpuscles, 98
Hay fever
 association, 100
HbA_{1c} test, 346
HBcAg (hepatitis B core antigen), 174
HbC disease, 422
 target cells in, 415
HBsAg (hepatitis B surface antigen),
 174
HDL (high-density lipoprotein), 94
Headaches, **518**
 adverse effects with drugs, 195,
 199, 200
 α-blockers, 244
 bupropion toxicity, 576
 caffeine withdrawal, 570
 Chiari I malformation, 492
 cimetidine, 399
 drug adverse effects, 546
 electroconvulsive therapy, 562
 genital herpes, 184
 giant cell (temporal) arteritis, 518
 glaucoma, 536
 hydralazine, 318
 increased intracranial pressure, 521
 Jarisch-Herxheimer reaction, 148
 lead poisoning, 425
 malaria, 157
 Mucor spp and *Rhizopus* spp, 153
 nitrates, 318
 ondansetron, 400
 pituitary apoplexy, 339
 ranolazine, 319
 Rocky Mountain spotted fever, 150
 sodium-channel blockers, 322
 subarachnoid hemorrhage, **513**,
 516
 triptans for, 547
 vasculitides, 314
 venous sinus thrombosis and, 503
Head and neck cancer, **671**
 cetuximab for, 442
Head size
 Paget disease of bone, 463
Healing, wound, 216
Healthcare delivery, 270–273
Healthcare payment models, **271**
Healthcare proxy, 269
Health maintenance organization
 plan, 271
Hearing loss, 533
 conductive, 49
 cytomegalovirus, 182
 osteogenesis imperfecta, 51
 Paget disease of bone, 463
 sensorineural deafness, 596
Heart
 autoregulation of, 297
 electrocardiograms, 293
 embryology, **281**
 fetal development, 612
 ischemia in, 210
 morphogenesis of, **280–281**
 myocardial action potential, 292
 normal pressures in, 297
 sclerosis of, 473
Heartburn, 377
Heart disease
 congenital, 63, **298–299**
 death, common causes by age, 272
 Fabry disease, 88
 ischemic, **304**
Heart failure, **309**
 ACE inhibitors for, 610
 acromegaly, 339
 amiodarone, 323
 angiotensin II receptor blockers, 610
 aortic regurgitation as precursor,
 291
 atrial septal defect, 299
 β-blockers for, 245
 B-type natriuretic peptide in, 296
 calcium channel blockers, 324
 cardiac glycosides for, 321
 chronic ischemic heart disease, 304
 contractility in, 284
 diabetic ketoacidosis, 347
 disopyramide, 322
 dobutamine for, 242
 dopamine for, 242
 Ebstein anomaly, 298
 ESR in, 214
 fludrocortisone and, 354
 hypertension, **300, 316**
 in sleep apnea, 679
 jugular venous pulse in, 287
 left heart, 309
 Paget disease of bone, 463
 potassium-sparing diuretics, 609
 readmissions with, 272
 right heart, 309
 shock caused by, 310
 thiazides for, 609
 ventricular septal defect, 299
Heart murmurs, **291**
 cardiomyopathies, 308
 patent ductus arteriosus, 299
Heart rate, 243
Heart sounds, 287
 cardiac cycle, 287
 cardiac tamponade, 310
 splitting in, **289**
Heart transplant
 dilated cardiomyopathy, 308
Heart valve development, 281
Heat-labile toxin (LT)
 Clostridium botulinum, 138
 Clostridium perfringens, 138
 Cl⁻ secretion in gut, 132
Heat shock proteins, 45
Heat-stable toxin (ST)
 resorption of NaCl and H₂O in
 gut, 132
Heat stroke, 517
Heavy menstrual bleeding (AUB/
 HMB), 633
Heel pain, 461
Heinz bodies, 79, 414, **416**, 422
Helicase, 38
Helicobacter pylori, **146**
 as oncogenic microbe, 226
 disease association, 379
 metronidazole, 195
 penicillins for, 188
 silver stain, 125
 urease-positive, 127
Heliotrope rash, 228
HELLP syndrome, 643
“Helmet cells,” 423
Helminthic infections
 eosinophils and, 408
Helper T cells
 cell surface proteins, 110
 cytokine secretion, 108
Hemagglutinin
 influenza viruses, 169
 parainfluenza viruses, 170
Hemangioblastomas, 526
Hemangiomas, 220
 cavernous (liver), 392
 pyogenic granuloma, 478
 strawberry, 478
Hemarthroses
 hemophilias, 426
 Vitamin C deficiency, 69
Hematemesis, 377
Hematin, 126, 142
Hematochezia
 diverticulosis, 383
 intestinal disorders, 386
 Meckel diverticulum, 384, 618
Hematocrit
 polycythemia vera, 433
Hematologic abnormalities
 laboratory techniques for, 54
Hematologic disorders
 paraneoplastic syndromes, 228
Hematologic drug reactions, 250
Hematology/oncology
 anatomy, 406–409
 pathology, 414–434
 pharmacology, 435–443
 physiology, 410–413
Hematopoiesis, **406**, 432
Hematopoietic stem cells, 110
Hematuria, 595
 bladder cancer, 606
 hereditary hemorrhagic
 telangiectasia, 316
 interstitial nephritis, 601
 kidney stones, 598
 protease inhibitors, 203
 renal cyst disorders, 604
 renal oncocytoma and, 605
 renal papillary necrosis, 602–610
 Schistosoma haematobium, 161
 transitional cell carcinoma, 606
 UTIs, 181
 Wegener granulomatosis, 315
 Wilms tumor, 606
Heme
 bilirubin and, 375
 chloroquine, 200
 porphyria and, **425**
 synthesis of, **425**
 vitamin B₆ and, 67
Hemianopia, 515, 542
Hemiballismus, 519
 brain lesions and, 511
Hemidesmosome, 474, 480
Hemineglect, 514
Hemiparesis
 saccular aneurysms, 516
Hemochromatosis, **395**
 calcium pyrophosphate deposition
 disease, 467
 cardiomyopathy with, 308
 chromosome association, 64
 chronic, 426
 free radical injury, 210
 hepatocellular carcinoma and, 392
 HLA-A3 and, 100
Hemoglobin, **665**
 carbon dioxide transport, 670
 development of, 404
 kinetics of, 230
Hemoglobin electrophoresis, **410**
Hemoglobinuria
 acute tubular necrosis and, 602
 G6PD deficiency, 422
 intravascular hemolysis, 421
 paroxysmal nocturnal, 122

- Hemolysis
 alpha toxin, 133
Clostridium perfringens, 138
 G6PD deficiency, 250
 HELLP syndrome, 643
 sulfonamides, 194
- Hemolytic anemia, 421
 autoimmune, 189
 babesiosis, 157
 cephalosporins, 189
 direct Coombs-positive, 250
 extrinsic, 421, **423**
 folate deficiency and, 420
 G6PD deficiency, 79
 intrinsic, 421, **422**
 penicillin G, V, 187
 pyruvate kinase deficiency and, 422
 spherocytes in, 415
 sulfa drug allergies, 252
 Wilson disease, 395
- Hemolytic disease of the newborn, 112, **405**
- Hemolytic reactions and blood types newborns, 405
- Hemolytic-uremic syndrome (HUS)
Escherichia coli, **145**, 179
 exotoxins, 132
 platelet disorders, 427
- Hemophagocytic lymphohistiocytosis, **435**, 544
- Hemophilia, 426
 X-linked recessive disorder, 61
- Hemoptysis
Aspergillus fumigatus, 177
 bronchiectasis, 675
 choriocarcinomas, 642
 lung cancer, 684
 tuberculosis, 140
 Wegener granulomatosis, 315
- Hemorrhage
 acute pancreatitis, 397
 AIDS retinitis, 165
 baroreceptors and, 296
 bevacizumab, 442
 delirium caused by, 558
 Ebola virus, 171
 intracranial, 513
 intraventricular, 512
 pulmonary, 137
 shock from, 310
 subarachnoid hemorrhage, **513**, 516
 ulcers, 380
 Weil disease, 147
- Hemorrhagic cystitis
 adenovirus, 164
 drug reaction, 251
- Hemorrhagic fever
 bunyavirus, 167
 filovirus, 167
- Hemorrhagic stroke, 513
- Hemorrhoids, 366
- Hemosiderinuria, 421
- Hemostasis, 407
 platelet plug formation, **411**
- Henderson-Hasselbalch equation, 592
- Henoch-Schönlein purpura, 315
- Hepadnaviruses
 characteristics of, 164
 genome, 163
- Heparin, **436**
 acute coronary syndromes, 307
 in basophils, 408
 in coagulation cascade, 413
 mast cells and, 408
 osteoporosis, 250
 thrombocytopenia, 250
 toxicity treatment, 248
 warfarin vs, **436**, **437**
- Heparin-induced thrombocytopenia (HIT), 436
- Hepatic adenomas, 392
- Hepatic arteries, 364, 367
- Hepatic ascites, 609
- Hepatic ducts, 368
- Hepatic encephalopathy, **391**
- Hepatic fibrosis, 367
- Hepatic lipase, 93
- Hepatic necrosis, **249**, 485
- Hepatic steatosis, 391
- Hepatitis
 alcoholic, 391, 571
 drug reaction and, 249
 heroin addiction and, 576
 hyperbilirubinemia, 393
- Hepatitis A (HAV)
 characteristics of, 172
 picornavirus, 167, **168**
 serologic markers, 174
- Hepatitis antigens, 174
- Hepatitis B (HBV)
 characteristics of, 172
 extrahepatic manifestations, 173
 hepatocellular carcinomas and, 392
 medical importance, 164
 nosocomial infection, 185
 as oncogenic microbe, 226
 passive antibodies for, 110
 polyarteritis nodosa and, 314
 serologic markers, 174
 sexually transmitted infection, 184
- Hepatitis C (HCV)
 characteristics of, 172
 extrahepatic manifestations, 173
 flaviviruses, 167
 hepatocellular carcinoma and, 392
 lichen planus, 482
 as oncogenic microbe, 226
 therapy for, **204**
- Hepatitis D (HDV), 172
- Hepatitis E (HEV), 172
 hepevirus, 167
- Hepatitis viruses, **172**
 aplastic anemia, 421
 serologic markers for, **174**
- Hepatocellular carcinomas, **392**
Aspergillus fumigatus, 153
 Budd-Chiari syndrome and, 392
 carcinogens causing, 225
 cirrhosis and, 389
 hemochromatosis, 395
 non-alcoholic fatty liver disease, 391
 oncogenic microbes, 226
- Hepatocytes, 86
- Hepatoduodenal ligament, 361
- Hepatomas, **392**
- Hepatomegaly
 Budd-Chiari syndrome, 392
 galactosemia, 80
 hepatocellular carcinoma, 392
 pulmonary hypertension, 668
 right heart failure, 309
 Von Gierke disease, 87
 Zellweger syndrome, 47
- Hepatosplenomegaly
 β -thalassemia and, 418
 biliary tract disease, 395
 graft-versus-host disease, 119
 hyperchylomicronemia, 94
 leishmaniasis, 158
 lysosomal storage diseases, 88
 mononucleosis, 165
 TORCH infections, 182
- Hepatosteatorosis, 72
- Hepatotoxicity
 amiodarone, 323
 bosentan, 686
 danazol, 658
 inhaled anesthetics, 550
 isoniazid, 197
 leflunomide, 486
 methotrexate, 440
 pyrazinamide, 197
 rifamycins, 196
 ribinafine, 199
 thionamides, 354
 valproic acid, 544
 zileuton, 687
- Hepecidin, 213
 in anemia of chronic disease, 421
- Hepeviruses
 characteristics, 167
 genomes, 163
 naked viruses, 163
HER2/neu (c-erbB2), 224
 “Herald patch” (pityriasis rosea), 482
 Herceptin (trastuzumab), 443
- Hereditary amyloidosis, 212
- Hereditary angioedema, 658
 complement disorder and, 107
- Hereditary elliptocytosis, 414
- Hereditary hemorrhagic telangiectasia, **316**
 autosomal dominance of, 60
- Hereditary hyperbilirubinemias, **394**
- Hereditary motor and sensory neuropathy, 524
- Hereditary spherocytosis, 422
 spherocytes in, 415
- Hereditary thrombosis syndromes, **428**
- Hermaphrodites, 639
- Hernias, **370**
- Herniation syndromes, **529**
- Heroin, 551
 detoxification medications, 576
 intoxication and withdrawal, 570
 opioids for withdrawal, 551
- Herpes genitalis, 164
- Herpes labialis, 164
- Herpes simplex virus (HSV), **164–166**
 HSV-1/HSV-2, 164, 184
 cidofovir, 202
 clinical significance, 164
 envelope, 163
 foscarnet for, 202
 guanosine analogs, 201
 identification, 166
 meningitis caused by, 180
 skin infections, 479
 STI, 184
 TORCH infection, **182**
- Herpes zoster
 dorsal root latency, 165
 famciclovir, 201
 reactivation, 443
- Herpetic whitlow, 164
- Heterochromatin, 34
- Heterodimer, 48
- Heterodisomy, 57
- Heterogeneous nuclear RNA (hnRNA), 41
- Heteroplasmy, 57
- Heterozygosity loss, 56
- Hexokinase
 glucokinase vs, **75**
 metabolic pathways, 74
- “HF” cells (lungs), 309
- HFE gene
 hemochromatosis and, 395
- HGPRT (hypoxanthine guanine phosphoribosyltransferase), 37, 440
- Hiatal hernias, 370
- Hiccups, 519
- High altitude respiratory response, 670
- High-frequency recombination (Hfr) cells, 130
- Highly active antiretroviral therapy (HAART), 203
- High-riding prostate, 627
- Hilar lymph nodes
 calcification of, 677
- Hilar mass, 684
- Hilum (lung), 663
- lymphadenopathy, 675
- Hindbrain, 490
- Hindgut
 blood supply/innervation of, 364
 development of, 358
- Hip dislocation
 nerve injury with, 453
- Hip injuries/conditions
 common, **460**
 developmental dysplasia, 461
- Hip muscles, **451**
- Hippocampus
 lesions in, 511
 limbic system, 499
 pyramidal cells, 210
- Hippurate test, for *Streptococcus agalactiae*, 137
- Hirschsprung disease, **384**
- Hirsutism
 cyclosporine, 120
 danazol, 658
 menopause, 636
 SHBG and, 337
- Histaminase, 408
- Histamine blockers, 398, **399**
- Histamine receptors, 238
- Histamines
 in basophils, 408
 cortisol effect on, 336
 location of, 373
 mast cells and, 408
 seafood toxins, 247
 signaling pathways for, 337
 vitamin B₆ and, 67
- Histidine, 81
- Histiocytosis (Langerhans cell), 434
- Histocompatibility complex I and II, 100
- Histones
 acetylation, 34
 amino acids in, 81
 deacetylation, 34
 methylation, 34
- Histoplasma* spp
 treatment, 199
- Histoplasma capsulatum*
 HIV-positive adults, 177
 necrosis and, 209
- Histoplasmosis, 151
 erythema nodosum, 482
- Histrionic personality disorder, 565
- HIV (human immunodeficiency virus), **175**
 aplastic anemia in, 421

- HIV (*continued*)
 common disease associations, **177**
 diagnosis, **175**
 flow cytometry diagnosis, 54
 hairy leukoplakia, 479
 heroin addiction and, 576
 Kaposi sarcoma, 165, 478
 lymphopenia, 424
 meningitis, 180
 non-Hodgkin lymphoma and, 429
Pneumocystis jirovecii, 154
 primary central nervous system lymphoma and, 430
 prophylaxis for HIV patients, 198
 pulmonary arterial hypertension, 679
 retrovirus, 167
 rifamycins in, 196
 STI, 184
 T cells and, 409
 therapy for, **203**
 TORCH infection, **182**
 untreated time course, 176
 viral receptor, 166
- HLA-B8
 graves disease and, 342
- HLA-DR3
 graves disease and, 342
- HLA-DR4, 466
- HLA genes
 celiac disease and, 381
 disease associations, **100**, 341
 DM type 1 association, 347
 seronegative spondyloarthritis, 469
- HMG-CoA reductase
 cholesterol synthesis, 73
 metabolic pathways, 74
- HMG-CoA synthase, 73
- HMP shunt
 metabolic site, 72
 NADPH production, 75, **79**
 rate-determining enzyme, 73
 vitamin B₁ deficiency, 66
- Hoarseness
 gastroesophageal reflux disease, 377
 lung cancer, 684
 Ortner syndrome, 283
 Pancoast tumor, 685
 thyroid cancer, 343
- Hodgkin lymphoma, **429**
 bleomycin for, 439
 non-Hodgkin vs, **429**
 oncogenic microbes and, 226
 paraneoplastic cerebellar degeneration and, 228
 vinca alkaloids for, 441
- Holistic medical therapy, 269
- Holoprosencephaly, **491**
 Patau syndrome, 63
- Homatropine, 241
- Homer-Wright rosettes, 528
- Homicide, 272
- Homocysteine
 folate deficiency, 420
 vitamin B₉ deficiency, 68
 vitamin B₁₂ deficiency, 69, 420
- Homocysteine methyltransferase deficiency in, 84
 vitamin B₁₂ and, 69
- Homocystinuria
 causes of, **84**
- Homologous recombination repair, 40
- Homovanillic acid (HVA)
 in neuroblastomas, 350
- Homunculus, **502**
- Hookworms, 159
- Hormone effects on kidney, **590**
- Hormone replacement therapy, **657**
 estrogens for, 656
 for hypopituitarism, 339
 thrombotic complications, 250
- Hormone-sensitive lipase, 93
- Hormones (reproductive), 655
- Horn cysts, 477
- Horner syndrome, 514, 518, **540**
 cavernous sinus, 542
 lung cancer, 684
 Pancoast tumor, 685
- Horse flies (disease vector), 159
- Horseshoe kidney, **579**
- Hospice care, **272**
- Hospital readmission causes, **272**
- Hot flashes
 drug reaction and, 249
- “Hourglass stomach,” 370
- Howell-Jolly bodies
 postsplenectomy, 98
- Hu antigens, 228
- Human chorionic gonadotropin (hCG)
 as tumor marker, 226
 choriocarcinomas, 642
 ectopic pregnancy, 641
 hydatidiform moles, 642
 secretion of, 612, **633**
 signaling pathways, 337
 source and functions of, **633**
- Human factors design, **273**
- Human herpesvirus 4 (HHV-4), 165
- Human herpesvirus 6 (HHV-6), 165, 183
- Human herpesvirus 7 (HHV-7), 165
- Human herpesvirus 8 (HHV-8), 165, 177
 Kaposi sarcoma, 478
 as oncogenic microbe, 226
- Humanized monoclonal antibodies, 110
- Human papillomavirus 6 (HPV-6), 184
- Human papillomavirus 11 (HPV-11), 184
- Human papillomavirus 16 (HPV-16), 671
- Human papillomavirus (HPV)
 cervical pathology, 645
 HIV-positive adults, 177
 as oncogenic microbe, 226
 penile cancer, 651
 tumor epidemiology, 643
 verrucae, 477
 warts, 164
- Human placental lactogen, **634**
- Humerus fracture
 axillary nerve and, 447
 radial nerve with, 447
- Humoral immune response, 409
- Hunger/satiety regulation, 498
- Hunter syndrome, 60, 61, **88**
- Huntington disease
 drug therapy for, 549
 movement disorders, 519
 neurodegenerative disorder, 520
 neurotransmitters for, 495
 trinucleotide repeat expansion diseases, 62
- Hurler syndrome, 88
- Hürthle cells, 341
- Hutchinson teeth, 147
- Hyaline arteriosclerosis, 301
- Hyaline casts (urine), 594
- Hydatid cysts, 161
- Hydatidiform mole, **642**
 hCG in, 633
 theca-lutein cysts and, 646
- Hydralazine
 gestational hypertension, 316
 heart failure, 309
 mechanism and clinical use, **318**
- Hydrocele (scrotal)
 acquired, 652
 congenital, 652
- Hydrocephalus, **522**
 childhood tumors, 528
 headaches with, 518
 posterior fossa malformations, 492
 risk for developing, 513
Toxoplasma gondii, 182
- Hydrochlorothiazide (HCTZ), 609
 for diabetes insipidus, 338
 hyperglycemia, 249
 pancreatitis, 249
- Hydrogen peroxide, 204
- Hydronephrosis, **599**
 BPH, 654
 horseshoe kidney, 579
 kidney stones, 598
 posterior urethral valves, 579
- Hydrophobia, 171
- Hydrops fetalis
 parvovirus B19, 182, 183
 syphilis, 182
- Hydroxychloroquine
 myopathy, 250
- Hydroxylases, 73
- Hydroxylation
 collagen synthesis, 50
 in protein synthesis, 45
 Vitamin C and, 50
- Hydroxyurea, **442**
 polycythemia vera, 433
 purine synthesis, 36
 sickle cell anemia, 422
 targets of, 438
- Hyoid artery, 619
- Hyoscyamine, 241
- Hyperacute transplant rejection, 119
- Hyperaldosteronism, **349**
 hypertension with, 300
 potassium-sparing diuretics for, 609
- Hyperammonemia, **82**, 85
 fatty acid metabolism and, 89
- Hyperbilirubinemia
 conjugated (direct), 393
 hereditary, 394
 jaundice with, 393
 unconjugated (indirect), 393
- Hypercalcemia
 acute pancreatitis and, 397
 adult T-cell lymphoma, 430
 bisphosphonates for, 486
 calcium carbonate antacid effects, 399
 diabetes insipidus, 338
 hyperparathyroidism, 345
 loop diuretics for, 608
 lung cancer, 684
 paraneoplastic syndrome, 228
 succinylcholine, 551
 teriparatide, 487
 thiazides, 609
 Williams syndrome, 64
- Hypercalciuria
 hyperparathyroidism, 345
 thiazides for, 609
- Hypercapnia
 contractility in, 284
- Hypercholesterolemia, 94
 familial, 60
- Hyperchylomicronemia, 94
- Hypercoagulability, 671
 hereditary syndromes, 428
 in pregnancy, 633
 marantic endocarditis in, 311
 venous sinus thrombosis with, 503
 warfarin adverse effect, 436
- Hyperemesis gravidarum, 642
- Hyperemia
 pseudoephedrine/phenylephrine, 686
- Hyper eosinophilic syndrome, 308
- Hyperestrogenism, 646
- Hyperglycemia
 Cushing syndrome, 348
 diabetic ketoacidosis, 347
 diabetic retinopathy, 537
 drug reaction and, 249
 glucagon and, 333
 hyperkalemia, 590
 immunosuppressants, 120
 pancreatic cell tumors, 351
 protease inhibitors, 203
 thiazides, 609
 vitamin B₃ toxicity, 67
- Hypergonadotropic hypogonadism, 639
- Hypergranulosis, **475**, 482
- Hyper-IgM syndrome, 117
- Hyperinsulinemia, 645
- Hyperkalemia
 aldosterone in, 588
 aliskiren, 610
 angiotensin II receptor blockers, 610
 cardiac glycosides, 321
 causes of, 590
 depolarizing neuromuscular blocking drugs, 551
 diabetic ketoacidosis, 347
 potassium-sparing diuretics, 609
 primary adrenal insufficiency, 349
- Hyperkalemic tubular acidosis (type 4), 593
- Hyperkeratosis, **475**, 477
- Hyperlipidemia, **301**
 atherosclerosis and, 302
 immunosuppressants, 120
 thiazides, 609
- Hyperopia, 535
- Hyperosmolar hyperglycemic state, **347**
 DM type 2, 346
- Hyperosmolarity, 590
- Hyperparathyroidism, **345**
 calcium pyrophosphate deposition disease, 467
 cinacalcet for, 355
 lab values in, 464
 renal osteodystrophy and, 603
- Hyperphagia
 depression with, 561
 hypothalamus and, 498
- Hyperphosphatemia
 hyperparathyroidism (secondary), 345
 hypoparathyroidism, 344
 renal osteodystrophy and, 603
- Hyperpigmentation
 bleomycin, 439
 busulfan, 441
 fludrocortisone, 354
 hemochromatosis, 395
 Peutz-Jeghers syndrome, 387
 primary adrenal insufficiency, 349

- Hyperplasia, 206
 adrenal, 348, 349
 parathyroid, **345**, 351
 uterine bleeding with, 633
- Hyperplastic arteriosclerosis, 301
- Hyperplastic polyps, 387
- Hyperprolactinemia, 249, 328, **527**
 anovulation, 645
 calcium channel blockers and, 318
 risperidone and, 573
- Hyperpyrexia
 with TCAs, 575
- Hyperresonance (chest percussion), 682
 pneumothorax, 680, **682**
- Hypersensitivity reactions, **112–113**
 acute interstitial nephritis, 601
 C3 deficiency, 107
 cephalosporins, 189
 Graves disease, 342
 IgE antibodies, 105
 mast cells and, 408
 organ transplants, 119
 penicillins, 187–188
 pneumonitis, 675
 rheumatic fever, 312
 sulfonamides, 194
- Hypersensitivity reaction (type II)
 rapidly progressive
 glomerulonephritis, 596
- Hypersensitivity reaction (type III)
 acute poststreptococcal
 glomerulonephritis, 596
- Hypersensitivity reaction type IV
 contact dermatitis, 477
- Hypersomnia, 561
- Hypertension, **300**, 679
 ACE inhibitors for, 610
 acromegaly and, 339
 aliskiren for, 610
 α -blockers for, 244
 angiotensin II receptor blockers
 for, 610
 aortic dissection and, 303
 atherosclerosis and, 302
 atrial fibrillation and, 295
 autosomal recessive polycystic
 kidney disease, 604
 β -blockers for, 245
 Charcot-Bouchard
 microaneurysms, 516
 episodic, 350
 heart failure, 316
 hyperaldosteronism, 349
 immunosuppressants, 120
 intraparenchymal hemorrhage,
 513
 leflunomide, 486
 lipohyalinosis and, 514
 local anesthetics, 550
 loop diuretics for, 608
 MDMA, 571
 microangiopathic anemia, 423
 minoxidil, 658
 PCP, 571
 pheochromocytomas, 350
 polyarteritis nodosa, 314
 in pregnancy, 243, **643**
 pseudoephedrine/phenylephrine,
 686
 renal cyst disorders, 604
 renal failure, 603
 renovascular disease and, 604
 sleep apnea, 679
 thiazides for, 609
 thoracic aortic aneurysm and, 302
 treatment for, **316**
 tyramine, 244, 575
- Hypertensive crisis, 569
 MAO inhibitors, 575
 phenoxybenzamine for, 244
 pheochromocytoma, 350
- Hypertensive emergency, 300, **318**
 RBC casts in, 594
- Hypertensive nephropathy, 300
- Hypertensive retinopathy, **537**
- Hyperthermia
 atropine causing, 241
 ecstasy intoxication, 571
 MDMA, 571
- Hyperthyroidism, 340, **342**
 amiodarone and, 323
 β -blockers in, 245
 drug reactions, 249
 hCG elevation and, 633
 hydatidiform moles, 642
- Hypertriglyceridemia, 94
 acute pancreatitis and, 397
- Hypertrophic cardiomyopathy, 308,
 531
 Pompe disease, 87
- Hypertrophic osteoarthropathy, 684
 cancer association, 228
- Hypertrophic pyloric stenosis, **359**
- Hypertrophic scars, 218
- Hypertrophy, 206
- Hyperuricemia
 drug reaction and, 250
 gout and, 467
 kidney stones and, 598
 Lesch-Nyhan syndrome, 37
 pyrazinamide, 197
 thiazides, 609
 vitamin B₃ toxicity, 67
- Hyperventilation
 metabolic acidosis and alkalosis,
 592
 in pregnancy, 633
 therapeutic, 501
- Hypervitaminosis D, 464
- Hypnagogic hallucinations, 559
 narcolepsy, 568
- Hypnopompic hallucinations, 559
 narcolepsy, 568
- Hypoadosteronism, 593
- Hypocalcemia, 333
 acute pancreatitis and, 397
 cinacalcet causing, 355
 hypermagnesemia and, 591
 hyperparathyroidism, 345
 hypoparathyroidism, 344
 renal osteodystrophy, 603
 thyroidectomy, 343
 tumor lysis syndrome, 435
- Hypochlorhydria hypergastrinemia,
 379
- Hypocretin, 568
- Hypodermis, 473
- Hypofibrinogenemia, 214
- Hypogammaglobulinemia, 228
- Hypoglossal nerve (CN XII), 506
 lesion in, 532
 with stroke, 514
 tongue, 493
- Hypoglycemia
 fructose intolerance, 80
 glucagon production with, 333
 gluconeogenesis and, 78
 insulinomas, 351
 neonatal, 614
 Von Gierke disease, 87
- Hypogonadism
 diagnosis of, 639
 estrogens for, 656
 gynecomastia, 649
 hemochromatosis, 395
 Kallmann syndrome, 639
 pituitary prolactinomas, 328
 testosterone/methyltestosterone, 658
 zinc deficiency, 71
- Hypokalemia
 antacid use, 399
 causes of, 590
 cystic fibrosis, 60
 on ECG, 293
 loop diuretics, 608
 nephrogenic DI, 338
 VIPomas and, 371
- Hypomanic episodes, **561**
- Hyponatremia
 MDMA, 571
 as paraneoplastic syndrome, 228
 thiazides, 609
- Hypo-osmolarity, 590
- Hypoparathyroidism, **344**
- Hypophosphatemia
 hyperparathyroidism, 345
- Hypopituitarism, **339**
- Hypoplasia, 613
 pulmonary, 660
- Hypopyon, 536
- Hyporeflexia
 LMN lesions, 531
 magnesium hydroxide and, 399
- Hypospadias, 624
- Hypotension
 adrenal insufficiency, 349
 aliskiren, 610
 amphotericin B, 199
 angiotensin II receptor blockers,
 610
 baroreceptors in, 296
 cardiac tamponade, 310
 endotoxins, 131
 ephedrine for, 242
 hypermagnesemia, 591
 local anesthetics, 550
 magnesium hydroxide and, 399
 metronidazole, 195
 midodrine for, 242
 norepinephrine for, 242
 orthostatic, 349
 phenylephrine for, 242
 scombroid poisoning, 247
 sympatholytic drugs and, 243
- Hypothalamic/pituitary drugs
 clinical use and adverse effects
 of, **354**
- Hypothalamic-pituitary hormones,
328
- Hypothalamus
 ADH secretion, 329
 homeostasis and, **498**
 nuclei of, 498
 reproductive hormone control, 656
 sleep cycle role of, 497
 TRH sensitivity, 331
- Hypothenar muscles, 450
- Klumpke palsy, 448
- Hypotheses (statistical), 262
- Hypothyroidism, 340, **341**
 amiodarone and, 323
 anemia, 420
 carpal tunnel syndrome with, 459
 drug reaction and, 249
 hormone replacement, 354
 lithium, 574
- Hypotonia
 poliomyelitis, 531
 Zellweger syndrome, 47
- Hypoventilation, 592
- Hypovolemic shock, 310
- Hypoxanthine guanine
 phosphoribosyltransferase
 (HGPRT), 37
- Hypoxemia
 alveolar gas equation, 668
 oxygen deprivation, 669
 vasoconstriction, 679
- Hypoxia
 apoptosis caused by, 208
 contractility in, 284
 erythropoietin and, 589
 lung diseases, 679
 nocturnal, 679
 oxygen deprivation, 669
 regions susceptible to, 210
 renal, 666
 vasoconstriction/vasodilation and,
 297
- Hypoxia inducible factor 1 α , 224
- Hypoxic stroke, 512
- Hypoxic vasoconstriction
 (pulmonary), 668
 high altitude, 670
- Hysterectomy
 cardinal ligament in, 625
- Hysteresis (lung and chest wall), 665
- ## I
- Iatrogenic abnormal uterine bleeding,
 633
- Ibandronate, 486
- Ibuprofen, 486
 hemolysis in G6PD deficiency, 250
- Ibutilide, 323
- ICAM-1 protein
 in leukocyte extravasation, 215
 viral receptor, 166
- I cells, 371
 disease, 47
- Icosahedral viruses, 163
- Icterohemorrhagic leptospirosis, 147
- Idarucizumab, 435
- Idealization, 555
- Identification, 555
- Idiopathic intracranial hypertension,
521
- Idiopathic pulmonary fibrosis, 675
- Idiopathic thrombocytopenic purpura
 (ITP), 427
 rituximab for, 443
- IDL (intermediate-density
 lipoprotein), 94
- IFN- γ (Interferon- γ)
 cachexia and, 227
- Ifosfamide, 441
 hemorrhagic cystitis, 251
- IgA and IgG deamidated gliadin
 peptide autoantibody, 115
- IgA antibodies, 105
 anti-endomysial autoantibody, 115
 anti-tissue transglutaminase
 autoantibody, 115
 ataxia-telangiectasia, 117
 breast milk, 636
 in celiac disease, 381
 hyper-IgM syndrome, 117
 passive immunity, 110
 Peyer patches and, 374
 selective deficiency in, 116
- IgA nephropathy, 596
- IgA protease, 129

- IgD antibodies, 105
- IgE antibodies, 105
- ataxia-telangiectasia, 117
 - eczema, 477
 - hyper-IgM syndrome, 117
 - mast cells and, 408
 - type I hypersensitivity, 112
- IgG antibodies, 105
- ataxia-telangiectasia, 117
 - complement activation and, 106
 - hepatitis A (HAV), 174
 - multiple sclerosis, 523
 - as passive immunity, 110
 - type II hypersensitivity reaction, 480
 - type III hypersensitivity reactions, 113
- IgM antibodies, 105
- in biliary cirrhosis, 395
 - complement activation and, 106
 - hepatitis A (HAV), 174
 - hyper-IgM syndrome, 117
 - overproduction, 431
 - in sclerosing cholangitis, 395
 - splenic dysfunction, 98
- Ileum, 362
- basal electric rhythm, 362
- Ileus, 386
- bacterial peritonitis (spontaneous), 390
 - gallstone, 396
- Iliacus, 452
- Iliohypogastric nerve, 452
- Iliotibial band syndrome, 461
- Illness anxiety disorder, 566
- lloperidone, 573
- Imatinib, 433, **443**
- IMG registration timeframe, 6
- Imipenem
- seizures with, 251
- Imipramine, 575
- Immature ego defenses, 555
- Immature teratoma, 647
- Immune checkpoint interactions, 222
- Immune complex, 113
- Immune evasion
- in cancer, 221
- Immune responses
- acute-phase reactants, 102
 - antibody structure and function, 104–117
 - antigen type and memory, 105
 - Bordetella pertussis*, 143
 - cell surface proteins, 110
 - complement, 106
 - cytokines, 108
 - hypersensitivity types, 114–115
 - immunoglobulin, 105
 - passive vs active, 110
 - respiratory burst, 109
 - Salmonella/Shigella* spp, 144
 - transfusion reactions, 114
- Immune system organs
- cellular components, 99
 - lymph nodes, **96**
 - thymus, 98
- Immunocompromised patients
- acyclovir/famciclovir/valacyclovir, 201
 - Candida albicans* in, 153
 - common organisms affecting, 179
 - Cryptococcus neoformans*, 153
 - Cryptosporidium*, 155
 - esophagitis in, 377
 - fungal infections, 186
 - Listeria monocytogenes*, 139
 - Pneumocystis jirovecii*, 154
- Immunodeficiency syndromes
- flow cytometry diagnosis, 54
 - infections in, **118**
 - syndromes, **116–117**
- Immunoglobulin A vasculitis, 315
- Immunoglobulins
- adaptive immunity and, 99
 - breast milk and, 636
 - isotypes of, **105**
 - for Kawasaki disease, 314
- Immunohistochemical stains, 227
- Immunology, 96–122
- cellular components, 98
 - immune responses, 104–117
 - immunosuppressants, **120–122**
 - lymphoid structures, 96–98
 - pathogen recognition in, 99
- Immunomodulator signaling pathways, 337
- Immunophenotype assessment, 54
- Immunosuppressants
- for aplastic anemia, 421
 - targets, **121**
 - transplant rejection, **120**
- Immunosuppression
- vitamin A deficiency, 66
 - vitamin C deficiency, 69
- Immunotherapy, 121
- Impaired colleague, 269
- Imperforate hymen, **644**
- Impetigo, 136, 475
- crusts with, 479
- Imprinting disorders, 58
- Inactivated (killed) vaccine, 111
- Incidence vs prevalence, **259**
- in medical error corrections, 259
- Inclusions
- Cowdry A, 166
 - mulberry-like (morulae), 150
 - Negri bodies, 171
 - “owl eye,” 165
 - reticulate bodies, 148
- Incomplete penetrance, 56
- Incontinence (fecal/urinary), 453
- Incus, 533, 620
- Incus (ossicles)
- pharyngeal arch derivative, 620
- India ink stain, 125
- Indicator media, 126
- Indinavir
- HIV therapy, 203
- Indirect bilirubin, 375
- Indirect cholinomimetic agonists, 240
- Indirect Coombs test
- unbound antibody detection, 112
- Indirect inguinal hernia, 370
- Indirect sympathomimetics, 242
- Indirect (unconjugated)
- hyperbilirubinemia, 393
- Indomethacin, 486
- for diabetes insipidus, 338
- IFN- α (Interferon- α)
- clinical uses, 121
 - myopathy, 250
 - natural killer cells, 101
- Infant development, **635**
- Infarction
- blood-brain barrier effects, 496
 - of bone, 463
- Infarcts
- atherosclerosis, 302
 - calcification in, 211
 - pituitary, 339
 - regions susceptible to, 210
 - types of, 210
- IFN- β (Interferon- β)
- clinical uses, 121
 - natural killer cells, 101
- Infections
- ESR in, 214
 - in immunodeficiency, 118
- Inferior colliculi, 504
- Inferior gluteal nerve, 453
- Inferior mesenteric artery, **363**, 364
- horseshoe kidney, 579
- Inferior oblique muscle, 540
- Inferior rectal artery, 366
- Inferior rectal vein, 365
- Inferior rectus muscle, 540
- Inferior sagittal sinus, 503
- Inferior vena cava, 360
- diaphragm, 663
 - gonadal drainage and, 624
- Infertility
- clomiphene, 656
 - cystic fibrosis, 60
 - ectopic pregnancy, 641
 - Kallmann syndrome, 639
 - Kartagener syndrome, 49
 - leuprolide for, 656
 - mumps, 170
 - salpingitis, 185
 - septate uterus, 623
 - varicoceles, 651
- IFN- γ (Interferon- γ), **108**, 116
- clinical uses, 121
- Infiltrative cardiomyopathy, 308
- Inflammasome, 214
- Inflammation
- acute, 214
 - in atherosclerosis, 302
 - cardinal signs, **213**
 - chronic, **216**
 - ESR in, 214
 - Extrinsic (death receptor) pathway, 208
 - IL-1 in, 108
 - Intrinsic (mitochondrial) pathway, 208
 - wound healing, 216
- Inflammatory bowel disease (IBD), **382**
- azathioprine for, 440
 - colorectal cancer and, 388
 - erythema nodosum, 482
 - infliximab/adalimumab for, 487
 - methotrexate for, 440
 - sclerosing cholangitis and, 395
 - spondyloarthritis, 469
 - therapeutic antibodies, 122
- Inflammatory breast disease, **649**, 650
- Inflammatory demyelinating polyradiculopathy, 524
- Infliximab, 122, **487**
- for Crohn disease, 382
 - for ulcerative colitis, 382
- Influenza, **169**
- orthomyxovirus, 167
 - pneumonia, 683
 - treatment/prevention, 201
- Informed consent, **265**
- Infraspinatus muscle
- Erb palsy, 448
 - rotator cuff, 446
- Infundibulopelvic ligament, 625
- Ingested seafood toxins, **247**
- Inguinal canal, **369**
- Inguinal hernia, 370
- Inguinal ligament, 368, 369
- Inguinal triangle, 370
- Inhalational injury, **676**
- Inhaled anesthetics, **550**
- Inheritance modes, 59
- Inhibin
- Sertoli cell secretion of, 628
- Inhibitors of complement activation, 106
- Inhibitory pathway, 500
- Injury (unintentional), 272
- Innate immune system
- in acute inflammation, 214
- Innate immunity, **99**
- Inositol trisphosphate (IP₃), 337
- Inotropes, 310
- Inotropy, 286
- INR (international normalized ratio), 426
- Insomnia
- barbiturates for, 546
 - nonbenzodiazepine hypnotics, 546
 - ramelteon for, 547
 - stimulants causing, 570
 - suvorexant, 547
- Inspiratory capacity (IC), 664
- Inspiratory reserve volume (IRV), 664
- Insulin, **334**
- anabolic effects of, 334
 - diabetic ketoacidosis, 347
 - fructose biphosphatase-2 and, 76
 - GIP effect on, 371
 - glucagon and, 333
 - glycogen regulation, **73**, **85**
 - hypokalemia from, 590
 - potassium shifts wit, 590
 - in pregnancy, 334
 - production of, 329
 - secretion of, 334
 - somatostatin and, 371
 - somatostatinomas and, 351
- Insulin deficiency, 590
- diabetes mellitus diagnosis, 346
- Insulin-like growth factor 1 (IGF-1)
- acromegaly, 339
 - signaling pathways for, 337
- Insulinoma, 351
- insulin and C-peptide in, 334
 - MEN 1 syndrome, 351
 - pancreatic cell tumor, 351
- Insulin preparations, 352
- Insulin resistance, 633
- acanthosis nigricans and, 482
 - acromegaly, 339
 - cortisol, 336
 - DM type 2, 347
 - GH, 329
 - non-alcoholic fatty liver disease, 391
 - polycystic ovarian syndrome, 645
- Insurance
- disregarding in treatment, 269
 - Medicare/Medicaid as, 272
 - types of plans, 271
- Integrase inhibitors, 203
- Integrins
- epithelial cells, 474
 - viral receptor, 166
- Intellectual disability, 557
- autism and, 557
 - cri-du-chat syndrome, 64
 - Down syndrome, 63
 - Lesch-Nyhan syndrome, 37
 - Patau syndrome, 63

- phenylketonuria, 84
Williams syndrome, 64
- Intellectualization, 555
- Intention tremor, 519
cerebellar lesions, 511
- Interdigital tinea pedis, 152
- Interferons, **109**
- Interferon- γ release assay (IGRA), 140
for tuberculosis, 140
- Interleukin 1 (IL-1), 108
cachexia and, 227
endotoxins, 133
- Interleukin 2 (IL-2), 108
cyclosporine and, 120
natural killer cells and, 101
sirolimus and, 120
tacrolimus and, 120
- Interleukin 2 receptor (IL-2R), 120
- Interleukin 3 (IL-3), 108
- Interleukin 4 (IL-4), 108
- Interleukin 5 (IL-5), 108
- Interleukin 6 (IL-6), 108
cachexia and, 227
endotoxins, 133
- Interleukin 8 (IL-8), 108
neutrophils and, 406
- Interleukin 10 (IL-10), 108
- Interleukin 12 (IL-12), 108
natural killer cells and, 101
receptor deficiency, 116
- Interleukin receptor modulators
naming conventions for, 254
- Intermediate filaments
cytoskeletal element, 48
- Intermenstrual bleeding (IMB), 633
- Internal capsule
intraparenchymal hemorrhage, 513
stroke effects, 514
- Internal carotid artery
cavernous sinus, 542
- Internal hemorrhoids, 366
- Internal iliac artery, 282
- Internal iliac lymph nodes, 624
- Internal inguinal ring, 370
- Internal jugular vein, 503
- Internal oblique muscle, 369
- Internal rotation
arm (rotator cuff), 446
hip, 451
- Internal spermatic fascia, 369
- International Foundations of
Medicine (IFOM), 12
- Internuclear ophthalmoplegia, 511,
543
- Interossei muscles, 450
Klumpke palsy, 448
ulnar nerve, 447
- Interpersonal therapy, 572
- Interpreting study results, 261
- Intersex, 639
- Interstitial fluid, 297
- Interstitial lung disease, 466, 675
- Interstitial nephritis
acute, 601
as drug reaction, 251
NSAID toxicity, 486
penicillins, 188
- Interstitial pneumonia, 683
- Interstitium
leukocyte extravasation and, 215
- Interventricular foramen, 281
- Interventricular septal rupture, 305,
307
- “Intestinal angina,” 386
- Intestinal atresia, **359**
- Intestinal gastric cancer, 379
- Intestinal obstruction
hemias, 370
superior mesenteric artery
syndrome, 363
- Intimate partner violence, 269
- Intoxication (psychoactive drugs), 570
- Intracellular bacteria, **127**
- Intracellular fluid (ICF), 581
- Intracellular receptors
endocrine hormone and, 337
- Intracranial hemorrhage, **513**
eclampsia, 643
- Intracranial hypertension
vitamin A toxicity, 66
idiopathic, **521**
- Intracranial pressure (ICP)
cerebral ischemia, 296
hydrocephalus, 522
papilledema, 538
superior vena cava syndrome, 685
venous sinus thrombosis, 503
- Intraductal papilloma, 649
- Intraepithelial adenocarcinoma, 644
- Intraocular pressure (IOP), 536
- Intraparenchymal hemorrhage, 513
- Intrauterine device (IUD)
copper, 657
- Intrauterine growth restriction
(IUGR)
low birth weight, 635
substance abuse, 614
- Intravascular hemolysis, 421
paroxysmal nocturnal
hemoglobinuria, 107
- Intravenous anesthetics, **550**
- Intraventricular hemorrhage, 512
neonatal respiratory distress
syndrome, 661
- Intrinsic factor, **372**, 373
- Intrinsic hemolytic anemias, **422**
- Intrinsic pathway, 208
coagulation defects of, 426
heparin and, 437
- Intrinsic renal failure, 601
- Introns
splicing out, 41
vs exons, **43**
- Intussusception, **385**
Meckel diverticulum, 384
- Inulin
glomerular filtration rate and, 582
in proximal convoluted tubules,
587
- Inulin clearance, 582
- Invariant chain, 100
- Invasive carcinoma, 219
cervix, 645
- Invasive lobular carcinoma (breast),
650
- Inversion, 453
- In vivo biofilm-producing bacteria,
128
- Involuntary treatment, 267
- Iodine
deficiency in, 341, 342
infection control, 204
teratogenicity, 614
- Iodophors, 204
- IPEX syndrome, 102
- Ipratropium, 241, **687**
- Irinotecan/topotecan, **442**
targets of, 438
topoisomerase (TOP) I inhibition,
38
- Iritis, 536
- Iron
absorption of, 69, 374
anemia, 388, **418**
anemia of chronic disease, 421
colorectal cancer, 388
excess, 67
in hemochromatosis, 395
lab values in anemia, 419
sideroblastic anemia, 419
toxicity of, 69
toxicity treatment, 248
- Iron poisoning, **426**
- Iron studies
interpretation of, 419–444
- Irritable bowel syndrome (IBS)
antispasmodic drugs, 241
criteria and symptoms for, **383**
- Isavuconazole
mucormycosis treatment, 153
- Ischemia, **210**, 669
acute tubular necrosis from, 602
atherosclerosis, 302
colonic, 386
digital, 472
in gastrointestinal tract, 386
mesenteric, 386
necrosis and, 209
watershed areas, 210
- Ischemic brain disease, **512**
- Ischemic heart disease
contraindicated antiarrhythmics,
322
heart murmurs in, 291
manifestations of, **304**
- Ischemic priapism, 651
- Islet cell cytoplasmic antibodies, 115
- Islets of Langerhans, 325
- Isocarboxazid, 575
- Isocitrate dehydrogenase
metabolic pathways, 74
rate determining enzyme, 73
- Isodisomy, 57
- Isoflurane, 550
- Isolation of affect, 555
- Isoleucine
classification of, 81
maple syrup urine disease and, 84
- Isoniazid, **197**
cytochrome P-450, 252
hemolysis in G6PD deficiency, 250
hepatitis, 249
Mycobacterium tuberculosis, 196
seizures, 251
- Isoproterenol
norepinephrine vs, 243
sympathomimetic action, 242
- Isosorbide dinitrate, 318
- Isosorbide mononitrate, 318
- Isotretinoin
cystic acne, 66
teratogenicity, 614
- Isovolumetric contraction, 287
- Isovolumetric relaxation, 287
- Itraconazole
azoles, 199
Sporothrix schenckii, 154
systemic mycoses, 151
- Ivabradine, **324**
- IV drug use
common organisms, 179
- Ivermectin, 200
- “Ivory white” plaques, 677
- IV phlebitis, 199
- Ixodes* ticks, 146, 149
- J**
- JAK2 gene, 224
myeloproliferative disorders, 433
- Janeway lesions, 311
- Jarisch-Herxheimer reaction, **148**
- Jaundice, **393**
biliary tract disease, 395
cholangitis, 368, 397
drug reaction and, 249
fructose intolerance, 80
galactosemia, 80
graft-versus-host disease, 119
hepatitis B, 184
hepatocellular carcinoma, 392
hereditary hyperbilirubinemias,
394
with leptospirosis, 147
newborn hemolytic disease, 405
pancreatic cancer, 398
TORCH infections, 182
yellow fever, 168
- Jaw jerk reflex, 507
- JC virus (John Cunningham virus)
HIV-positive adults, 177
immunocompromised patients,
118
polyomaviruses, 164
- Jejunal and ileal atresia, 359
- Jejunum, 362
- Jervell and Lange-Nielsen syndrome,
294
- Jimson weed, 241
- Job syndrome, 116
- Jod-Basedow phenomenon, 342
- Joint hypermobility, 51, 62
- J point in ECG, 293
- Jugular foramen, 503
- Jugular venous distention (JVD),
309, 685
- Jugular venous pulse, 287
- Justice (ethics), 265
- Juvenile polyposis, 387
- Juxtaglomerular apparatus (JGA), **589**
renin secretion, 588
- Juxtaglomerular cells
tumors in, 349
- K**
- Kala-azar, 158
- Kallikrein
C1 esterase inhibitor deficiency,
107
neutrophil chemotaxis and, 406
- Kallmann syndrome, 498, **639**
- Kaposi sarcoma, 478
AIDS and, 184
bacillary angiomatosis vs, 478
HHV-8, 165
HIV-positive adults, 177
oncogenic microbes and, 226
- Kartagener syndrome, **49**, 280
obstructive lung disease, 675
- Karyotyping, **55**
- KatG, 197
- Kawasaki disease, 314
- Kayser-Fleischer rings, 395
- K cells, 371
- K complexes/sleep spindles, 497
- Kegel exercises, 599
- Keloid scars, 218
- Keratinocytes, 216
- Keratin pearls, 684
- Keratoacanthoma, 484
- Keratoconjunctivitis, 164

- Keratoconjunctivitis sicca, 468
Keratomalacia, 66
Keratitis
 actinic, 482
 hyperkeratosis, 475
 parakeratosis, 475
 seborrheic, 477
Kernicterus, 194, 204, **393**–394
Ketamine, 550
Ketoacidosis, 72, **90**
Ketoconazole, 658
 cytochrome P-450, 252
 gynecomastia from, 649
 mechanism and clinical use, 199
Ketogenesis
 diabetic ketoacidosis, 347
 metabolic site, 72
 rate-determining enzyme for, 73
Ketone bodies, **90**
 brain metabolism, 334
 in diabetic ketoacidosis, 347
 production of, 90
Ketorolac, 486
Kidney disease
 acute injury, 601
 hypertension, 300
 prenatal diagnosis of, 579
Kidneys
 anatomy, **580**
 blood flow regulation, 297
 calcification in, 211
 chronic graft nephropathy, 119
 embryology of, **578**
 endocrine functions, **589**
 glomerular structure, **580**
 hormones acting on, 590
 ischemia in, 210
 retroperitoneal location of, 360
 sclerosis, 473
 solitary functioning, 579
 transplant prophylaxis, 120
Kidney stones
 Crohn disease association, 382
 electrolyte disturbances, 591
 hematuria with, 594
 hydronephrosis, 599
 hyperparathyroidism, 345
 presentation and findings with, **598**
 risk factors for, 593
 UTIs, 181
Kiesselbach plexus, 671
Killian triangle, 384
Kinases, 73
Kinesin
 movement of, 48
Kinin cascade/pathways, 412
Klebsiella pneumoniae
 cephalosporins, 189
 encapsulation, 127
 UTIs caused by, 181
Klebsiella spp, 145
 alcoholism, 179
 currant jelly sputum, **145**, 186
 nosocomial infections, 185
 pneumonia, 683
 urease-positive, 127
 urinary tract infections, 600
Klinefelter syndrome, 638
 chromosome association, 64
 gynecomastia, 649
Klumpke palsy, 448
Klüver-Bucy syndrome, 511
Knee examination, **454**
Knee injuries/conditions
 Baker cyst, 460
 iliotibial band syndrome, 461
 ligament and meniscus, 460
 Osgood-Schlatter disease, 461
 prepatellar bursitis, 460
Knees
 common conditions of, **460**
Knock-out/Knock-in genes, 56
KOH preparation, 152
Koilocytes
 condylomata acuminata, 184
Koilocytosis, 477
Koplik spots, 183
Korsakoff syndrome, 66, 558, **571**
Krabbe disease, **88**, 524
KRAS gene, 224
 adenomatous colonic polyps and,
 387
 lung cancer and, 684
Krukenberg tumors, 379
Kübler-Ross grief model, 562
Kulchitsky cells, 684
Kuru, 178
Kussmaul respirations
 in diabetic ketoacidosis, 347
Kussmaul sign, **316**
Kwashiorkor, 71
Kyphoscoliosis, 531
Kyphosis
 in homocystinuria, 84
K_m, 230
L
Labetalol, 245
 hypertension in pregnancy, 316
 hypertensive emergency, 318
Labia, 625
Labile cells, 46
Lachman test, 454
Lac operons, **40**
Lacrimation reflex, 507
Lactase deficiency, **81**
Lactation, **636**
 oxytocin's role in, 328
 progesterone and, 630
 prolactin and, 330
 Sheehan syndrome and, 339
Lactational mastitis, 649
Lactic acid dehydrogenase, 77
Lactic acidosis
 ethanol metabolism and, 72
 exercise and, 670
 MELAS syndrome, 59
 pyruvate dehydrogenase complex
 deficiency, 77
Lactoferrin
 in neutrophils, 406
 in respiratory burst, 109
Lactose-fermenting enteric bacteria,
 126, **144**
Lactose intolerance, 381
Lactose metabolism
 genetic response to environmental
 change, 40
Lactulose
 for hepatic encephalopathy, 391
Lacunar infarcts, 514
Ladd bands, 385
Lambert-Eaton myasthenic syndrome,
 472
 autoantibody, 115
 as paraneoplastic syndrome,
 228
 small cell lung cancer, 684
Lamina propria
 Peyer patches in, 374
 in Whipple disease, 381
Lamins, 48
Lamivudine, 203
 HIV therapy, 203
Lamotrigine
 epilepsy, 544
 rash caused by, 250
Lancet-shaped diplococci, 136
Landmarks (anatomical)
 for dermatomes, **510**
 midclavicular line, 663
 pudendal nerve block, 453
Langerhans cell histiocytosis, **434**
 pulmonary, 675
Lansoprazole, 399
Laplace law, **284**, 661
Large cell carcinoma, 684
Large-vessel vasculitis
 presentation and pathology, 314
Larva migrans, 159
Laryngopharyngeal reflux, 377
Laryngospasm, 359
 drug-induced, 569
Larynx, 662
Larynx muscles, 620
Lassa fever encephalitis, 167
Latanoprost, 552
Latent errors, 274
Lateral cerebellar lesions, 499
Lateral collateral ligament (LCL)
 injury, 454
Lateral corticospinal tract, 508, **509**,
 514
Lateral epicondylitis, 459
Lateral femoral cutaneous nerve, 452
Lateral geniculate nucleus (LGN),
 498
Lateral medullary syndrome, 514
Lateral nucleus (hypothalamus), 498
Lateral pterygoid muscle, 507
Lateral rectus muscle, 540
Lateral spinothalamic tract, 508
Lateral thoracic artery, 455
Lateral ventricles
 optic radiation, 542
 ventricular system, 504
Laxatives, **401**
LD50 (lethal median dose), 234
LDH
 tumor burden indicator, 226
LDL (low-density lipoprotein), 94
 PCSK9 enzyme, 93
 receptor binding, 93
Lead paralysis, 561
Lead poisoning
 anemia with, 419
 foot drop, 419
 mechanism, 425
 memory loss with, 425
 sideroblastic anemia, 419
 treatment of, 248, **419**
Lead-time bias, 261
Leber hereditary optic neuropathy, 59
Lecithinase, 133, 138
Lecithin-cholesterol acetyltransferase
 (LCAT)
 activation of, 93
Lecithin-cholesterol acyltransferase,
 93
Lectin pathway (complement
 activation), 106
Ledipasvir, 204
Leflunomide, **486**
 dihydroorotate dehydrogenase
 inhibition, 36
Left anterior descending artery
 coronary circulation, 283
 myocardial infarction and, 305
Left bundle branch, 293
Left circumflex coronary artery, 283
Left heart disease, 679
Left heart failure, 309
Left horn of sinus venosus, 281
Left main coronary artery, 283
Left marginal artery, 283
Left shift, 424
Left-to-right shunts, 299
Legg-Calvé-Perthes disease, **461**, 463
Legionella pneumophila, **143**
Legionella spp
 atypical organism, 179
 culture requirements, 126
 facultative intracellular organisms,
 127
 Gram stain of, 125
 macrolides, 193
 nosocomial infection, 185
 intracellular organism, 127
 pneumonia, 683
 silver stain, 125
Legionnaires' disease, 143
Leiomyoma (fibroid), 648
 nomenclature for, 220
 uterine bleeding with, 633
Leiomyosarcoma, 220, **648**
Leishmaniasis, **158**, 200
Length-time bias, 261
Lens
 collagen in, 50
 infantile cataracts, 80
Lens subluxation
 Elastin syndrome, 52
 in homocystinuria, 52, **84**
Lenticulostriate artery, 514
Lentiform nucleus, 500
Leonine facies, 141
Leprosy (Hansen disease), **141**
Leptin, 336
 hypothalamus, 498
Leptospira spp
 Gram stain of, 125
 spirochete, 146
 zoonotic infections, 149
Leptospira interrogans, **147**
Leptospirosis, 149
 symptoms of, 147
Lesch-Nyhan syndrome
 inheritance, 61
 purine salvage deficiency, 37
Leser-Trélat sign, **228**, 477
 stomach cancer, 379
Lesser omental sac, 361
Letrozole, 656
Leucine
 classification of, 81
 maple syrup urine disease and, 84
Leucovorin, 440
Leukemias, **432**
 carcinogens, 225
 cell type, 220
 cyclophosphamide for, 441
 cytarabine for, 440
 doxorubicin for, 439
 epidemiology, 222
 etoposide/teniposide for, 442
 immunohistochemical stain for,
 227
 lymphoma comparison, **429**
 mucormycosis, 153
 nomenclature for, 220
 oncogenic microbes, 225
 suppressor genes, 224
 TRAP tumor marker, 227
 vinca alkaloids for, 441

- Leukocoria, **538**
 Leukocyte adhesion deficiency, **117**, 215
 Leukocyte alkaline phosphatase (LAP), 406
 Leukocyte esterase, 181, 600
 Leukocytes
 extravasation, 214, **215**
 leukemias, **432**
 in urine, **181**, 594, 600
 Leukocytoclastic vasculitis, 173
 Leukocytosis, 213
 diabetic ketoacidosis, 347
 nosocomial infections, 185
 Leukodystrophies, 494, 524
 Leukoerythroblastic reaction, 424
 Leukopenias, **424**
 cytarabine, 440
 ganciclovir, 202
 trimethoprim, 194
 Leukoplakia, 479
 Leukotrienes
 basophils and, 408
 cortisol effects, 336
 Levator veli palatini muscle, 620
 Levetiracetam, 544
 Levodopa, 548, **549**
 Levofloxacin
 fluoroquinolones, 195
 Pseudomonas aeruginosa, 143
 Levomilnacipran, 575
 Levonorgestrel, 657
 Levothyroxine, **354**
 Lewy bodies, 520, 521
 dementia, **521**
 Leydig cells
 cryptorchidism, 651
 endocrine function, **628**, 638
 genital embryology, 622
 tumors of, 653
 LFA-1 antigens, 215
 Libido
 in geriatric patients, 270
 testosterone and, 636
 Libman-Sacks endocarditis, 470
 Lice
 disease vectors, 149
 head/scalp, 161
 treatment, 200
 Lichen planus, 173, 475, **482**
 Lichen sclerosus, 644
 Lichen simplex chronicus, 644
 Liddle syndrome, 586
 markers in, 591
 Lidocaine, 322, 550
 Life support
 withdrawal, 269
 Li-Fraumeni syndrome
 osteosarcomas, 465
 tumor suppressor genes in, 46, 224
 Ligaments
 female reproductive anatomy, 625
 gastrointestinal anatomy, 361
 Ligamentum arteriosum, 282
 Ligamentum teres hepatis, 282, **361**
 Ligamentum venosum, 282
 Ligand receptors, 208
 Lightheadedness, 534
 Likelihood ratio (LR), **257**
 Limbic system, **499**
 Limited scleroderma (CREST syndrome), 115, 473
 Linagliptin, 353
 Lindane, 200
 Linea alba, 369
 Linear ulcers, 377
 Linear viruses, 163
 Lines of Zahn, 672
 Lineweaver-Burk plot, 230
 Linezolid, **193**
 highly resistant organisms, 198
 protein synthesis inhibition, 191
 Lingula (lung), 663
 Linkage disequilibrium, 57
 Liothyronine (T3), **354**
 Lipase
 pancreatic secretions, 373
 in pancreatitis, 397
 Lipid-lowering agents
 mechanism and adverse effects, **320–321**
 Lipid metabolism
 fatty acids, 73
 Lipids
 key enzymes in, 93
 transport of, **92–93**
 Lipodystrophy
 protease inhibitors, 203
 tesamorelin for, 328
 Lipofuscin, **211**
 Lipoic acid, 76
 Lipolysis
 cortisol and, 336
 and, 334
 sympathetic receptors and, 238
 thyroid hormone and, 331
 Lipomas, 220
 Lipophilic drug
 drug metabolism of, 232
 Lipoprotein lipase, 93
 Lipoproteins, 93
 functions of, **94**
 Liposarcomas, 220
 Lipoteichoic acid
 cytoplasmic membrane, 124
 Liquefactive necrosis, 209
 Liraglutide, 353
 Lisch nodules, 525
 Lisdexamfetamine, 572
 Lisinopril, 610
 Lispro, 352
 Lissencephaly, **491**
Listeria monocytogenes
 β -hemolysis, 135
 neonates, 182
 penicillins for, 188
 transmission of, **139**
Listeria spp
 facultative intracellular organisms, 127
 intracellular organism, 127
 Lithium
 diabetes insipidus and, 249, 338
 hypothyroidism, 341
 mechanism and use, **574**
 prenatal exposure, **298**, 300
 teratogenicity, 614
 therapeutic index of, 234
 thyroid functions with, 249
 toxicity of, 569
 Live attenuated vaccines, 111
 Liver
 blood supply and innervation of, 364
 in gastrointestinal anatomy, 361
 lipid transport and, 92
 tissue architecture, **367**
 Liver/biliary disease
 acanthocytes in, 414
 alcoholic, 391
 anemia, 420
 autoimmune, 389, 392, 395
 cirrhosis, 71, 80, **389**
 cystic fibrosis, 60
 hepatosteatorosis, 72
 hereditary, 394
 ischemia in, 210
 metastases to, 223
 serum markers, 390
 target cells in, 415
 Wilson disease and, 395
 Liver failure
 Budd-Chiari syndrome and, 392
 movement disorder in, 519
 Liver fluke
 hyperbilirubinemia with, 393
 as oncogenic microbe, 226
 Liver function tests
 cholestatic pattern of, 395
 serum markers for, 390
 Liver markers
 in alcohol use, 570
 Liver tumors, 392
 Living wills, 266
 Loa loa, **159**
 Loading dose, 231
 Lobar pneumonia
 natural history of, 683
 organisms and characteristics, 683
 physical findings with, 680
 Lobular carcinoma (breast), 650
 Local anesthetics, **550**
 Localized amyloidosis, 212
 Locked-in syndrome
 osmotic demyelination syndrome, 524
 stroke, 515
 Locus ceruleus, 495
 Locus heterogeneity, 57
 Löffler endocarditis, 308
 Löffler medium, 126
Corynebacterium diphtheriae, 139
 Lomustine, 441
 Lone Star tick (disease vector), 149
 Long-chain fatty acid (LCFA)
 metabolism of, 89
 Long QT syndrome
 congenital, 294
 ranolazine, 319
 sudden cardiac death, 304
 Long thoracic nerve
 arm abduction, 446
 neurovascular pairing, 455
 Loop diuretics, **608**
 for heart failure, 309
 toxicity of, 251
 Loop of Henle, 608
 Bartter syndrome and, 586
 ethacrynic acid effect on, 608
 “loose associations,” 559
 Looser zones (osteomalacia), 463
 Loperamide, **400**, 551
 Lopinavir
 HIV therapy, 203
 Loratadine, 686
 Lorazepam
 alcohol withdrawal, 572
 Losartan, 610
 Low birth weight, **635**
 Löwenstein-Jensen agar/medium, 126
 Lower esophageal sphincter (LES)
 achalasia and, 376
 in Barrett esophagus, 377
 nitric oxide and, 371
 Lower extremity nerves, **452–453**
 Lower left quadrant (LLQ) pain, 383
 Lower motor neuron (LMN) lesions, 530, 531
 facial nerve, 532
 LPS endotoxin, 124, 131, **133**, 145
 LTB₄ (Leukotriene B₄), 406, 485
 Lumbar puncture, 507, 521
 Lumbosacral radiculopathy, 455
 Lumbrical muscles, 450
 Klumpke palsy and, 448
 median and ulnar nerves, 447
 Lumefantrine, 200
 Lunate bone, 449
 Lung abscesses, **685**
 Lung and chest wall expansion, 665
 Lung cancer
 asbestosis and, 677
 carcinogens causing, 225
 cisplatin/carboplatin for, 442
 erlotinib for, 442
 hypercalcemia and, 228
 incidence/mortality in, 222
 metastases to, 223
 non-small cell, 684
 oncogenes and, 224
 paraneoplastic syndromes and, 228
 presentation and complications, **684**
 small cell, 684
 topotecan for, 442
 Lung diseases
 obstructive, 674–675
 restrictive, 675
 Lungs
 anatomical relationships, **663**
 blood flow regulation, 297
 development of, **660**
 physical findings, 680
 sclerosis of, 473
 transfusion-related injury, 114
 Lung volumes, **664**
 Lung zones, 669
 Lupus
 autoimmune hemolytic anemia and, 423
 azathioprine for, 440
 drug-induced, 115
 isoniazid, 197
 lymphopenia, 424
 marantic endocarditis in, 311
 microangiopathic hemolytic anemia, 423
 nephritis, 470
 neutropenia, 424
 Lupus anticoagulant, 115
 Lupus-like syndrome
 α -methyl dopa, 243
 hydralazine, 318
 procainamide, 322
 Lurasidone, 573
 Luteal phase of menstrual cycle, 632
 Luteinizing hormone (LH)
 clomiphene effect, 656
 contraception, 657
 estrogen/progesterone, 630
 hCG and, 617
 ovulation, 631
 PCOS, 645
 premature ovarian failure, 636
 secretion of, 327
 sex development disorders, 639
 signaling pathways of, 337
 spermatogenesis, 628
 testosterone, 658
 Lyme disease
 animal transmission, 149
 AV block in, 295
 Borrelia burgdorferi, **146**
 ceftriaxone, 189

- Lymphadenopathy
Corynebacterium diphtheriae, 132
 hilar, 675–676
 Lymphogranuloma venereum, 184
 mediastinal, 676
 mononucleosis, 165
 rubella, 169, 182–183
 serum sickness, 113
 syphilis, 147, 184
 tinea capitis, 152
Toxoplasma gondii, 182
Trypanosoma brucei, 156
 in viral infections, 96
- Lymphatic filariasis (elephantiasis)
Wuchereria bancrofti, 159
- Lymph drainage
 gonadal, 624
 superficial inguinal nodes, 624
- Lymph nodes
 absent or scanty, 116
 drainage sites, 96–97
 structure and function, 96
 T cell differentiation, 102
 tumor metastases, 223
- Lymphocyte-depleted lymphoma, 429
- Lymphocyte-rich lymphoma, 429
- Lymphocytes, 409
 breast milk and, 636
 CLL/small cell lymphocytic lymphoma, 432
 corticosteroid effect on, 424
 lymph nodes, 96
 non-Hodgkin lymphoma, 430
 spleen, 98
 thymus, 98
- Lymphocytic choriomeningitis virus (LCMV), 167
- Lymphocytic infiltrates
Bordetella pertussis, 143
- Lymphocytosis, 98
- Lymphogranuloma venereum, 184
Chlamydia trachomatis, 149
- Lymphoid hyperplasia, 383
- Lymphoid neoplasms, 432
- Lymphoid structures, 96–97
 Peyer patches, 362, 374
- Lymphomas
 carcinogens causing, 225
 celiac disease and, 381
 cyclophosphamide for, 441
 cytarabine for, 440
 doxorubicin for, 439
 EBV and, 165
 etoposide/teniposide for, 442
 Hodgkin, 429
 hypercalcemia and, 228
 leukemia comparison, 429
 methotrexate for, 440
 nomenclature for, 220
 non-Hodgkin, 430
 oncogene for, 208, 224
 oncogenic microbes, 226
 paraneoplastic syndromes with, 228
 of stomach, 379
 testicular, 653
- Lymphopenias, 424
 ataxia-telangiectasia, 117
 corticosteroid effect on, 424
- Lynch syndrome, 388
 mismatch repair and, 40
- Lyonization (x-inactivation)
 Barr body formation, 61
- Lysergic acid diethylamide (LSD), 571
- Lysine
 classification of, 81
 in cystinuria, 85
- kidney stones, 598
 for pyruvate dehydrogenase complex deficiency, 77
- Lysogenic phage infection, 130
- Lysosomal storage diseases, 47
 causes and effects of, 88
- Lysosomal trafficking regulator gene, 117
- Lysosomal α -1,4-glucosidase, 87
- Lysozyme
 innate immunity, 99
 in neutrophils, 406
- LYST gene, 117
- Lytic bone lesions
 adult T-cell lymphoma and, 430
 Langerhans cell histiocytosis, 434
- M**
- MacConkey agar, 126, 144
 “Machine-like” murmur, 291
- Macroangiopathic hemolytic anemia, 423
- Macrocytic anemia, 420
- Macroglobulinemia, 431
- Macrolides
 cytochrome P-450 and, 252
 hypertrophic pyloric stenosis and, 359
Legionella pneumophila, 143
 mechanism and use, 193
Mycoplasma pneumoniae, 150
 naming convention for, 253
 protein synthesis inhibition, 191
 torsades de pointes, 248
- Macroorchidism, 62
- Macro-ovalocytes, 415
- Macrophages, 407
 alveolar, 662
 apoptosis and, 208
 binding of, 104
 breast milk and, 636
 cell surface proteins, 110
 cytokine secretion, 108
 endotoxin activation, 133
 innate immunity, 99
 in lymph node, 96
 lymphocyte interaction, 102
 in MI, 305
 necrosis and, 209
 pneumoconioses, 677
 splenic, 98
 in wound healing, 216
- Macrosomia, 614
- Macula densa
 juxtaglomerular apparatus, 589
- Macular cherry-red spot, 88, 538
- Macular degeneration, 536
- Macules, 475
 melanocytic nevus, 477
- Maculopapular rash
 graft-versus-host disease, 119
 measles, 170
 syphilis, 147
- Magnesium
 antacid use, 399
 antiarrhythmic treatment, 324
 cardiac glycoside toxicity, 321
 in laxatives, 401
 PTH regulation, 332
 torsades de pointes and, 294
- Magnesium hydroxide, 399
- Magnesium sulfate
 preeclampsia/eclampsia, 643
- Maintenance dose, 231
- Maintenance stage, substance addiction, 568
- Major basic protein (MBP), 408
- Major depressive disorder (MDD)
 diagnostic symptoms for, 561
 peripartum onset, 562
 vortioxetine use, 576
- Malabsorption syndromes, 381–382
 fat-soluble vitamin deficiencies, 65
 inflammatory bowel disease, 382
- Malaria
 artesunate for, 200
Plasmodium, 157
 quinidine/quinine for, 200
- Malassezia spp, 152, 476
- Malathion, 200
- Male/female genital homologs, 623
- Male genital embryology, 622
- Male reproductive anatomy, 626
- Male sexual response, 627
- Malformation, 613
- Malignancy
 marantic endocarditis in, 311
 uterine bleeding with, 633
- Malignancy/hyperplasia
 uterine bleeding with, 633
- Malignant hypertension
 microangiopathic hemolytic anemia, 423
- Malignant hyperthermia, 550–551
- Malignant mesothelioma, 227
- Malignant tumors, 220
 bones, 465
- Malingering, 566
- Malleus (ossicles), 533, 620
- Mallory bodies
 in alcoholic hepatitis, 391
- Mallory-Weiss syndrome, 377
- Malnutrition, 71
 superior mesenteric artery syndrome and, 363
- Malrotation, 385
 “Maltese cross” appearance, 157, 594
- MALT lymphomas
Helicobacter pylori, 146
 oncogenic microbes and, 226
 Sjögren syndrome, 468
- Mammary glands, 613
- Mammillary bodies, 511
 limbic system, 499
 Wernicke-Korsakoff syndrome, 571
- Mandibular process, 620
- Mango flies (disease vector), 159
- Manic episode, 560
- Man-in-the-barrel syndrome, 502
- Mannitol, 607
- Mantle cell lymphomas, 430, 434
- Mantle zone
 spleen, 98
- Maple syrup urine disease, 84
- Marantic endocarditis, 228, 311
- Marasmus, 71
- Maraviroc, 203
- Marburg hemorrhagic fever, 167
- Marcus Gunn pupils, 539
 multiple sclerosis, 523
- Marfanoid habitus
 homocystinuria, 84
 MEN 2B syndrome and, 351
- Marfan syndrome
 aortic dissection and, 303
 cardiac defect association, 300
 cataracts, 535
 chromosome association, 64
 elastin and, 52
 heart murmur with, 291
 thoracic aortic aneurysm and, 302
- Marginal zone lymphoma, 430, 434
- Marijuana
 intoxication and withdrawal, 571
- Marine omega-3 fatty acids, 320
- Masseter muscle, 507
- Mast cells, 408
 IgE antibody and, 105
- Mast cell stabilizers, 687
- Mastectomy, 448
- Mastication muscles, 507
- Mastoid air cells, 621
- Mastoiditis
 brain abscesses, 180
 Wegener granulomatosis, 315
- Maternal diabetes
 cardiac defect association, 300
- Maternal PKU, 84
- Maternal (postpartum) blues, 562
- Maternal pregnancy complication, 272
- Mature cystic teratoma, 647
- Mature ego defenses, 555
- Maxillary artery, 619
- Maxillary process, 620
- Mayer-Rokitansky-Küster-Hauser syndrome, 622
- McArdle disease, 87
- McBurney point, 383
- McCune-Albright syndrome, 57
- McMurray test, 454
- MDMA (ecstasy), 571
- Mean (statistics), 262
- Mean arterial pressure, 501
 equation for, 285
- Measles, 183
 paramyxovirus, 167, 169
 rubella virus, 170
 unvaccinated children, 186
 vitamin A for, 66
- Measurement bias, 260
- Measures of central tendency, 262
- Measures of dispersion, 262
- Mebendazole, 200
 microtubules and, 48
- mecA gene
 penicillin resistance and, 135
- Meckel diverticulum, 384, 618
- Meconium ileus, 386
 cystic fibrosis, 60
- MECP2 gene, 62
- Medial calcific sclerosis, 301
- Medial cerebellar lesions, 499
- Medial collateral ligament (MCL)
 injury
 in “unhappy triad,” 460
- Medial epicondylitis, 459
- Medial femoral circumflex artery, 463
- Medial geniculate nucleus (MGN), 498
- Medial lemniscus, 514
- Medial longitudinal fasciculus (MLF), 511, 543
- Medial malleolus, 455
- Medial medullary syndrome, 514
- Medial meniscal tear, 460
- Medial pterygoid muscle, 507
- Medial rectus muscle, 540
- Medial tibial stress syndrome, 461
- Medial umbilical ligament, 282, 369
- Median (statistics), 262
 “Median claw,” 451
- Median nerve
 carpal tunnel syndrome, 459
 injury to, 447
 neurovascular pairing, 455
- Median umbilical ligament, 369

- Mediastinal lymphadenopathy, 676
 Mediastinal pathology, **672**
 Mediastinitis, 672
 in pulmonary anthrax, 137
 Medical abortion
 ethical situations, 268
 methotrexate for, 440
 Medical errors
 analysis of, **274**
 assessment of, 268
 types of, 274
 Medical insurance plans, 271
 Medical power of attorney, 266
 Medicare/Medicaid, **272**
 Medication errors, 274
 Medication noncompliance, 268
 Medium-chain acyl-CoA
 dehydrogenase deficiency, 89
 Medium-vessel vasculitis
 presentation and pathology, 314
 Medroxyprogesterone, 657
 Medulla (brain)
 adrenal cortex and, **327**
 cranial nerves and nuclei, 505
 development of, 490
 spinal tracts and, 509
 strokes in, 514–515
 Medulla (lymph nodes)
 lymph nodes, 96
 thymus, 102
 Medullary breast carcinomas, 650
 Medullary cystic kidney disease, 604
 Medullary thyroid carcinomas, 343, 351
 Medulloblastoma, 350, 528
 “Medusa head” appearance, 137
 Mefloquine, 157
 Megacolon
 Chagas disease, 158
 in Hirschsprung disease, 384
 Megakaryocytes in essential
 thrombocytopenia, 433
 Megaloblastic anemia, 420
 cytarabine, 440
 Diphyllobothrium latum, 160
 drug reaction and, 250
 orotic aciduria, 420
 trimethoprim, 194
 tropical sprue, 381
 vitamin B₉ deficiency, 68
 vitamin B₁₂ deficiency, 69
 Megestrol, 657
 Meglitinides, 353
 Meissner corpuscles, 494
 Meissner plexus, 384
 Melanocytes
 tumor nomenclature in, 220
 Melanocyte-stimulating hormone
 (MSH)
 function of, 328
 secretion of, 327
 signaling pathways of, 337
 Melanocytic nevus, 477
 Melanoma, 484
 common metastases, 223
 immunohistochemical stain for, 227
 nomenclature for, 220
 oncogene, 224
 origin of, 220
 recombinant cytokines for
 metastatic, 121
 tumor suppressor gene, 224
 Melarsoprol, 156, 200
 Melasma (chloasma), 476
 MELAS syndrome, 59
 Melatonin
 circadian rhythms and, 497
 Melatonin receptor agonist
 Ramelteon as, 547
 Melena
 Meckel diverticulum, 384, 618
 polyarteritis nodosa, 314
 Meloxicam, 486
 Memantine, 549
 Membrane attack complex (MAC), 104
 complement and, 106
 Membranoproliferative
 glomerulonephritis
 (MPGN), 596
 hepatitis B and C, 173
 Membranous glomerular disorders, 594
 hepatitis B and C, 173
 Membranous interventricular
 septum, 281
 Membranous nephropathy, 597
 primary autoantibody, 115
 Membranous ossification, 458
 Membranous urethra injury, 627
 Membranous ventricular septum, 281
 Memory
 neural structures and, 499
 Memory loss
 anti-NMDA receptor encephalitis, 228
 lead poisoning, 425
 Wernicke-Korsakoff syndrome, 511
MEN1 gene, 224, 352
 Ménétrier disease, **379**
 Ménière disease, 534
 Menin, 224
 Meninges, **496**
 Meningiomas, 526
 Psammoma bodies in, 227
 Meningitis
 ceftriaxone, 189
 chloramphenicol, 192
 coccidioidomycosis, 151
 common causes, **180**
 Cryptococcus neoformans, 153
 CSF findings in, 180
 fluconazole, 199
 flucytosine, 199
 Haemophilus influenzae, 142
 headaches with, 518
 HIV-positive adults, 177
 Listeria monocytogenes, 139
 lyssosomal storage diseases, 142
 mumps, 170
 in neonates, 182
 rifamycin prophylaxis, 196
 Streptococcus pneumoniae, 136
 Streptococcus agalactiae, 137
 unvaccinated children, 186
 Meningocele, 491
 Meningococcal prophylaxis, 198
 Meningococcal vaccine, 127
 Meningococcemia
 endotoxins, 131
 Meningococci, 131, 142
 Meningoencephalitis
 HSV-2, 182
 Naegleria fowleri, 156
 West Nile virus, 167
 Meningomyelocele, 491
 Menkes disease
 mechanism and symptoms, **51**
 protein (ATP7A), 51
 Menopause, **636**
 hormone replacement therapy, 657
 primary ovarian insufficiency, 645
 Turner syndrome, 638
 Menorrhagia, 633
 anemia with, 418
 Menstrual cycle
 estrogens for, 656
 heavy bleeding (AUB/HMB), 633
 phases of, **632**
 Meperidine, 551
 Mepivacaine, 550
 Mercury poisoning, 248
 Merkel discs, 494
 Merlin protein, 224
 MERS (Middle East respiratory
 syndrome), 167
 Mesalamine, 382
 Mesangial cells
 juxtaglomerular apparatus, 589
 Mesencephalon, 490
 Mesenchymal tumors
 nomenclature of, 220
 Mesenteric arteries, 363
 Mesenteric ischemia, 386
 Mesocortical pathway, 499
 Mesoderm, 490
 branchial arches derivation, 619
 derivatives of, 613
 Mesolimbic pathway, 499
 Mesometrium, 625
 Mesonephric (Wolffian) duct, 622
 Mesonephros, 578
 Mesosalpinx, 625
 Mesothelioma, **678**
 carcinogens causing, 225
 Psammoma bodies in, 227
 Mesovarium, 625
 Mestranol, 656
 Meta-analysis, 263, **264**
 Metabolic acidosis, 592
 adrenal insufficiency, 349
 renal failure, 603
 symptoms of, 593
 Metabolic alkalosis, 592
 hyperaldosteronism, 349
 in hypertrophic pyloric stenosis, 359
 loop diuretics, 608
 nephron transport, 586
 thiazides, 609
 Metabolic disorders
 fructose, 80
 galactose, 80
 glycogen storage, 87
 lysosomal storage diseases, 88–89
 Metabolic fuel use, **91**
 Metabolic syndrome
 with antipsychotic drugs, 573
 non-alcoholic fatty liver disease
 and, 391
 Metabolism, 72–94
 amino acid derivatives, 83
 amino acids, 81
 apolipoproteins, 93
 cellular sites of, **72**
 disorders of, 81, 84–85
 drugs, 232
 dyslipidemias, 94
 ethanol, 72, 232
 fatty acid, 89
 fuel use, 91
 gluconeogenesis, 78
 glycogen and, 86
 lipoprotein functions, 93, 94
 pyruvate, 77
 rate-determining enzymes, 73
 summary of pathways, 74
 TCA cycle, 77
 tyrosine catabolism, 83
 urea cycle, 82
 Metacarpal neck fracture, 459
 Metacarpophalangeal (MCP) joints, 451
 Metachromatic granules
 Corynebacterium diphtheriae
 diagnosis, 139
 Metachromatic leukodystrophy, 88, 524
 Metalloproteinases, 216
 Metal storage diseases, 210
 Metanephros, 578
 Metaphase, 46
 Metaplasia, 206
 benign breast disease, 649
 esophagus, 378
 intestinal, 379
 specialized intestinal, 378
 Metastases
 common, **223**
 Metastasis, 219, 223
 gastric cancer, 379
 heart tumors from, 316
 liver cancer, 392
 lung, cancer, 684
 Metastatic calcification, 211
 Metastatic melanoma
 vemurafenib for, 444
 Metatarsophalangeal (MTP) joints
 gout, 467
 Metencephalon, 490
 Metformin, 353
 diarrhea with, 249
 Methacholine, 240
 Methadone, 551
 heroin addiction, 576
 intoxication and withdrawal, 570
 Methamphetamine, 572
 Methanol toxicity, 248
 Methemoglobin, **666**
 toxicity treatment, 248
 Methemoglobinemia, 666
 local anesthetics and, 550
 Methicillin, 249
 Methimazole, 354
 agranulocytosis, 250
 aplastic anemia, 250
 teratogenicity, 614
 Methionine
 classification of, 81
 start codons, 44
 sulfonamides and, 194
 Methotrexate, 440
 folate deficiency, 420
 hydatidiform moles, 642
 megaloblastic anemia, 250
 pulmonary fibrosis, 251
 pyrimidine synthesis and, 36
 rheumatoid arthritis, 466
 targets of, 438
 teratogenicity, 614
 vitamin B₉ deficiency, 68
 as weak acid, 233
 Methoxyflurane, 550
 Methylation
 histones, 45
 Methyllopa
 Coombs-positive hemolytic
 anemia, 250
 hypertension in pregnancy, 316
 Methylene blue, 248

- Methylenetetrahydrofolate reductase (MTHFR) deficiency, 84
- Methylmalonic acid
vitamin B₁₂ deficiency, 69
vitamin B₉ deficiency, 68
- Methylmalonyl-CoA mutase, 69
- Methylmercury teratogenicity, 614
- Methylnaltrexone, 551
- Methylphenidate
ADHD, 557, 572
CNS stimulant, 572
- Methyltestosterone, 658
- Methylxanthines, 687
- Metoclopramide, **400**
Parkinson-like syndrome, 251
tardive dyskinesia, 251
- Metolazone, 609
- Metoprolol, 245, 323
- Metronidazole
bacterial vaginosis, 148
clindamycin vs., 192
Clostridium difficile, 138
for Crohn disease, 382
disulfiram-like reaction, 251
Giardia lamblia, 155
Helicobacter pylori, 146
mechanism and use, **195**
vaginal infections, 181
vaginitis, 158
- Mevalonate synthesis, 320
- Mexiletine, 322
- Meyer loop, 542
- MHC (major histocompatibility complex) I and II, **100**
- Micafungin, 200
- Michaelis-Menten kinetics, 230
- Miconazole, 199
- Microalbuminuria, 346
- Microangiopathic hemolytic anemia (MAHA), 423
hypertensive emergency and, 300
intravascular hemolysis in, 421
- Microarrays, **54**
- Microbiology, 123–204
antimicrobials, 187–204
bacteriology, 124–134
clinical bacteriology, 134–150
mycology, 151–154
oncogenic organisms, 226
parasitology, 155–161
systems, 178–186
virology, 162–177
- Microbiome
in innate immunity, 99
- Microcephaly, 63
cri-du-chat syndrome, 64
fetal alcohol syndrome, 615
maternal phenylketonuria, 84
maternal X-ray exposure, 614
Patau syndrome, 63
with Zika virus, 171
- Microcytic, hypochromic anemia, **418–419**
Ancylostoma, 161
- Microcytosis, 214
- Microdeletion
congenital, 64
fluorescent in situ hybridization and, 55
22q11, 116
- Microfilaments, 48
- Microglia, 490, **493**
- Micrognathia
Edwards syndrome, 63
Pierre Robin sequence, 620
- Micromelia, 614
- Microphthalmia, 63
- MicroRNA (miRNA), 42, **56**
- Microscopic polyangiitis, 315
- Microsporium*, 152
- Microtubule inhibitors, **441**
- Microtubules, **48**
drugs acting on, 48
structure and function of, 48
- Micturition center, **237**
- Midazolam, 546, 550
- Midbrain
brain, 490
cranial nerve nuclei of, 505
lesions in, 511
- Middlebrook medium, 126
- Middle cerebral artery (MCA)
cortical distribution, 502
saccular aneurysms, 516
stroke effects, 514
- Middle meningeal artery
epidural hematoma and, 513
- Middle rectal vein, 365
- Midgut
blood supply/innervation of, 364
development of, 358
Midgut volvulus, 386
- Midodrine, 242
- Mifepristone, 657
- Miglitol, 353
- Migraine headaches, 518
hormonal contraception
contraindication, 657
triptans for, 547
- Migrating motor complexes (MMC), 371
- Migratory polyarthritis, 312
- Milnacipran, 575
- Mineralocorticoids
adrenal insufficiency, 349
adrenal steroids and, 335
- Mineral oil, 65
- Minimal alveolar concentration, 549
- Minocycline, 192
- Minors, consent for, 265
- Minoxidil, **658**
- Minute ventilation, 664
- Miosis
cholinomimetic agents, 552
Horner syndrome, 531, 540
pupillary control, 539
sympatholytic drugs, 243
- Mirabegron, 242
- Mirtazapine, 244, 576
major depressive disorder, 561
- Mismatch repair, 40
- Misoprostol, **399**
- Missense mutations, 39
- Mites/louse treatment, 200
- Mitochondria
high altitude and, 670
metabolism in, 72
- Mitochondrial encephalopathy, 59
- Mitochondrial inheritance, 59
- Mitochondrial myopathies, 59
- Mitosis, 46
griseofulvin, 200
- Mitral regurgitation, 288
in MI, 305
murmurs caused by, 290, 291
- Mitral stenosis, 288
murmurs caused by, 291
- Mitral valve
in cardiac cycle, 287
regurgitation in, 312
- Mitral valve prolapse, 291
fragile X syndrome, 62
Marfan syndrome, 52
renal cyst disorders and, 604
- Mittelschmerz, 631
- Mixed cellularity lymphoma, 429
- Mixed connective tissue disease, 470
autoantibody, 115
Raynaud phenomenon, 472
- Mixed cryoglobulinemia, 315
- Mixed transcortical aphasia, 516
- MMR vaccine, 170
- Mobitz AV blocks, 295
- Moccasin distribution (tinea pedis), 152
- Modafinil, 568
- Mode (statistics), 262
- Molecular mimicry
autoimmune response in
rheumatic fever, 129
- Molecular motor proteins, 48
- Molluscum contagiosum, 164, 479
- Mönckeberg sclerosis, 301
- “Monday disease,” 318
- Monoamine oxidase (MAO) inhibitors
atypical depression, 561
mechanism and clinical use, **575**
Parkinson disease, 549
selegiline/rasagiline, 549
tyramine and, 244
- Monobactams, **190**
Pseudomonas aeruginosa, 143
- Monoclonal antibodies
drug names for, 254
- Monoclonal immunoglobulin (Ig)
overproduction, 431
- Monocytes, **407**
innate immunity, 99
morulae in, 150
- Monospot test, 165
- Monozygotic (“identical”) twins, 616
- Montelukast, 687
- Mood
oxytocin’s role in regulating, 328
- Mood disorder, **560**
readmissions with, 272
- Mood stabilizing drugs, 561
- Morbidly adherent placenta, 640
- Moro reflex, 510, 635
- Morphine, 551
for acute coronary syndromes, 307
intoxication and withdrawal, 570
- Morphogenesis
of heart, 280–281
errors in, 613
- Morulae, 150
- Mosaic bone architecture, 464
- Mosaicism, 57
- Mosquitoes (disease vectors)
lymphatic filariasis, 159
malaria, 157
Zika virus, 171
- Motilin, 371
- Motion sickness, 241
- Motor cortex, 514
descending spinal tracts, 509
topographic representation, 502
ventral lateral thalamus and, 498
- Motor innervation
derivation of, 620
lower extremity, 452
tongue, 493
- Motor neuron signs, **529**
- Movement disorders, **519**
dopaminergic pathways and, 499
- Moxifloxacin, 195
- M phase, 46
- MPO-ANCA/p-ANCA autoantibody, 115
- M protein
rheumatic fever and, 136
as virulence factor, 129
- mRNA
aminoglycosides, 191
hepatitis viruses, 172
pre-mRNA splicing, 42
processing, 41
protease inhibitors, 203
stop codons, 44
- MRSA (methicillin-resistant *Staphylococcus aureus*)
cephalosporins, 189
highly resistant, 198
nosocomial infections, 135
oxazolidinones, 193
vancomycin, 190
- mTOR
sirolimus (rapamycin) inhibition of, 120
- Mucicarmine stain
polysaccharide capsule staining, 125
- Mucinous cystadenocarcinoma, 646
- Mucinous cystadenoma, 646
- Mucociliary escalator, 662
- Mucocutaneous lymph node syndrome, 314
- Mucoepidermoid carcinomas, 376
- Mucopolysaccharides
Periodic acid-Schiff stain, 125
- Mucopolysaccharidoses, 88
- Mucormycosis
diabetic ketoacidosis, 347
- Mucor* spp
amphotericin B for, 199
opportunistic infection, 153
- Mucosa, 362
- Mucosal cells, 372
- Mucosal neuromas, 351
- Mucosal polyps, 387
- Mucositis
methotrexate, 440
- Mucus, 238
- “Muddy brown” casts (urine), 594
- Mulberry molars, 147
- Müllerian duct
agenesis, 622
anomalies of, 623
derivatives of, 622
- Müllerian inhibitory factor (MIF), 622
Sertoli cell production, 628
- Multicystic dysplastic kidney, 578, 579
- Multidrug resistance protein 1 (MDR1), 227
- Multifactorial pulmonary hypertension, 679
- Multiorgan drug reactions, 251
- Multiple endocrine neoplasias (MEN syndromes), **351**
Zollinger-Ellison syndrome, 351
- Multiple gestations, 633
- Multiple myeloma
bortezomib/carfilzomib in, 443
common metastases, 223
ESR in, 214
metastatic calcification, 211
monoclonal gammopathy transition to, 431
plasma cell dyscrasia, 409
- Multiple personality disorder, 558
- Multiple sclerosis, **523**
drug therapy for, 551
HLA-DR2 and, 100
internuclear ophthalmoplegia, 543

- natalizumab for, 122
recombinant cytokines for, 121
- Mumps, **170**
acute pancreatitis with, 397
paramyxovirus, 167, 169
- Munchausen syndrome, 566
- Munchausen syndrome by proxy, 566
- Murphy sign, 396
- Muscarinic acetylcholine (ACh) receptors, 236
- Muscarinic agonists, 237
- Muscarinic antagonists, 237, **241**, 687
multiple sclerosis, 523
neuromuscular blocking drugs, 550, 551
Parkinson disease, 548
- Muscarinic receptor detrusor muscle, in, 237
- Muscle conduction/contraction skeletal, 456
smooth muscle, 457
- Muscle proprioceptors, **458**
- Muscles
atrophy of, 42
in starvation, 91
metabolism in, 86
ragged red fibers in, 59
- Muscle spindled, 458
- Muscular dystrophies, **61**
frameshift mutation, 61
X-linked recessive disorder, 61
- Muscularis externa, 362
- Muscular ventricular septum, 281
- Musculocutaneous nerve injury presentation, 447
- Musculoskeletal drug reactions, 250
- Musculoskeletal paraneoplastic syndromes, 228
- Musculoskeletal system anatomy, 446–454
common conditions, **461**
pathology, 459–467
pharmacology, 485–487
- Mutase, 73
- Mutations
BRAF, 432
BRCA1 gene, 56
COL3A1, 51
de novo, 62
drug resistance mechanisms, 196
in HbS and HbC, 410
JAK2, 433
locus heterogeneity in, 57
muscular dystrophies, 61
myelodysplastic syndromes, 432
non-Hodgkin lymphoma, 430
in PBP, 187
same locus, 57
STAT3, 116
thalassemia and, 418
tumor suppressor genes, 46, 525
WT1 deletion, 606
- Myalgias
Ebola virus, 171
fluoroquinolones, 195
genital herpes, 184
Jarisch-Herxheimer reaction, 148
Leptospira interrogans, 147
Lyme disease, 146
meningitis, 186
Trichinella spiralis, 159, 161
trichinosis, 159
vasculitides, 314
- Myasthenia gravis, 472
autoantibody, 115
diagnosis of, 240
- HLA subtype association, 100
neostigmine for, 240
as paraneoplastic syndrome, 228
pyridostigmine for, 240
restrictive lung diseases, 675
- MYCL1 gene, 224
- MYCN gene, 224
- Mycobacterial cells, 196
- Mycobacterium* spp, **140**
Gram stain, 125
intracellular organism, 127
Ziehl-Neelsen stain, 125
- Mycobacterium avium-intracellulare*, 140
HIV-positive adults, 177
prophylaxis with HIV, 198
vertebral osteomyelitis, 180
- Mycobacterium leprae*
animal transmission, 149
diagnosis, 141
rifamycins/dapsone, 196
- Mycobacterium marinum*, 140
- Mycobacterium pneumoniae*, 126
- Mycobacterium scrofulaceum*, 140
- Mycobacterium* spp
facultative intracellular organisms, 127
- Mycobacterium tuberculosis*
culture requirements for, 126
osteomyelitis, 180
reactivation site, 140
symptoms of, 140
therapeutic agents, 196, 197
- Mycolic acid
isoniazid, 197
synthesis of, 196
- Mycology, 151–154
- Mycophenolate mofetil, 120
inosine monophosphate dehydrogenase inhibition, 36
- Mycoplasma* spp
atypical organisms, 179
Gram stain, 125
macrolides, 193
pneumonia caused by, **150**, 179, 683
tetracyclines, 192
- Mycoses
cutaneous, 152
systemic, 151
- Mycosis fungoides, 430
- Mydriasis
glaucoma treatment and, 552
G-protein-linked second receptor, 238
muscarinic antagonists for, 241
pupillary control, 539
saccular aneurysm, 516
- Myelencephalon, 490
- Myelin, **494**
- Myeloblasts (peripheral smear), 432
- Myelodysplastic syndromes, **432**
sideroblastic anemia, 419
- Myelofibrosis, 433
- Myeloid neoplasms, 432
- Myelomeningocele, 63, 492
- Myeloperoxidase, 109
H₂O₂ degradation, 128
in neutrophils, 406
- Myeloproliferative disorders
in AML, 432
chronic, 433
hydroxyurea for, 442
- Myeloschisis, 491
- Myelosuppression
alkylating agents, 441
antimetabolites, 440
- drugs causing, 444
hydroxyurea, 442
irinotecan/topotecan, 442
- Myenteric plexus, 362
- Mylohyoid muscle, 620
- Myocardial action potential, **292**
- Myocardial depression, 550
- Myocardial infarction (MI), 304
antiarrhythmics after, 322
β-blockers for, 245
complications of, **307**
diabetes mellitus, 346
diagnosis of, 306
on ECG, 293
ECG localization of STEMI, 306
evolution of, 305
heart failure caused by, 309
heparin for, 436
homocystinuria, 84
hypertensive emergency and, 300
shock caused by, 310
thrombolytics for, 437
- Myocardial O₂ consumption/demand, 284
angina treatment, 319
- Myocarditis, **313**
adenovirus, 164
coxsackievirus, 167
diphtheria, 139
picornaviruses, 167
Toxocara canis, 159
- Myoclonic seizures, 517
- Myoclonus, 519, 521
- Myofibroblasts, 216
- Myoglobin, 665
oxygen-hemoglobin dissociation curve, 666
- Myoglobinuria
acute tubular necrosis, 602
McArdle disease, 87
- Myonecrosis, 138
- Myopathy
daptomycin, 195
drug reaction and, 250
- Myophosphorylase, 87
- Myopia, 535
retinal detachment, 537
- Myotonic dystrophy, **61**
cataracts and, 535
- Myxedema
thyroid hormones for, 354
- Myxomas, 316
- Myxomatous degeneration, 291
- N**
- N-acetylglucosaminyl-1-phosphotransferase, 47
- N-acetylcysteine, **686**
for acetaminophen toxicity, 248
for cystic fibrosis, 60
- NADH (reduced nicotinamide adenine dinucleotide)
electron transport chain, 78
fructose metabolism, 80
TCA cycle, 77
- Nadolol, 245
- NADPH (reduced nicotinamide adenine dinucleotide phosphate)
ethanol metabolism, 72
HMP shunt and, 79
respiratory burst and, 109
universal electron acceptors, 75
- Naegleria fowleri*, 156
- Nafcillin
characteristics of, 188
- Nails
clubbing, 60
glomus tumors under, 478
hemorrhages in bed of, 311
with psoriatic arthritis, 469
- Naïve T cell activation, 103
- Naked viral genome infectivity, **163**
- Nalbuphine, 551
- Naloxone
dextromethorphan overdose, 686
heroin detoxification, 576
for opioid toxicity, 248, 551
opioid toxicity, 570
- Naltrexone
alcoholism, 571
heroin detoxification, 576
opioid toxicity, 551
- 2-naphthylamine, 225
- Naproxen, 486
acute gout drugs, 487
- Narcissistic personality disorder, 565
- Narcolepsy
amphetamines for, 242
characteristics/treatment, **568**
hypnagogic/hypnopompic hallucinations, 568
- Narrow-angle glaucoma, 536
- Nasal congestion, 686
- Nasal decongestion
ephedrine for, 242
- Nasal polyps
cystic fibrosis, 60
- Nasal septum perforation, 315
- Nasopharyngeal carcinoma
EBV and, 165
oncogenic microbes and, 226
- Natalizumab, 122
multiple sclerosis, 523
- Nateglinide, 353
- National Board of Medical Examiners (NBME), 2, 11
- Natriuresis, 588
- Natriuretic peptide, 296
- Natural killer (NK) cells, 101, **409**
cell surface proteins, 110
function of, 409
innate immunity, 99
- Nausea
adverse drug effects, 400
antiemetics for, 401
with appendicitis, 383
biliary colic, 396
with MI, 305
migraine headaches, 518
ranolazine, 319
renal failure, 603
vitamin A toxicity, 66
vitamin C toxicity, 69
- Nebivolol, 245
- Necator* spp
disease associations, 161
infection routes, 158
- Necator americanus*, 159
- Neck and head cancer, 671
cetuximab for, 442
- Necrosis, **209**
acute pancreatitis, 397
Arthus reaction, 113
benign tumors, 220
Budd-Chiari syndrome, 392
calcification, 211
enterocolitis, 386
femoral head, 120, 461, 463
fibrinoid, 466
glioblastoma multiforme, 526
hepatic, 485

- Necrosis (*continued*)
 hernias and, 370
 jaw, 486
 mesenteric ischemia, 386
 nonalcoholic fatty liver disease, 391
 saponification, 209
 scaphoid avascular, 449
 transplant reaction, 119
 warfarin, 436
- Necrotizing enterocolitis, 386
- Necrotizing fasciitis, 479
Streptococcus pyogenes (Group A strep), 136
- Necrotizing glomerulonephritis, 315
- Negative predictive value (NPV), 257
- Negative punishment, 554
- Negative reinforcement, 554
- Negative skew distribution, 262
- Negative-stranded viruses, **168**
- Neglect (child), 556
- Negri bodies, 171
- Neisseria gonorrhoeae*
 culture requirements, 126
 epididymitis and orchitis, 654
 osteomyelitis, 180
 septic arthritis, 468
 STI, 184
 UTIs with, 600
- Neisseria meningitidis*
 chloramphenicol, 192
 culture requirements, 126
 encapsulation, 127
 meningitis, 180
 penicillin G/V for, 187
 Waterhouse-Friderichsen syndrome, 349
- Neisseria* spp
 C5-C9 deficiencies, 107
 cephalosporins, 189
 fluoroquinolones, 195
 gram-negative algorithm, 142
 IgA protease, 129
 intracellular organism, 127
 transformation in, 130
- Nelson syndrome, **348**
- Nematodes, **159**
 infection routes, **158**
- Neomycin
 aminoglycosides, 191
- Neonatal abstinence syndrome, **615**
- Neonatal conjunctivitis
Chlamydia trachomatis, 149
- Neonatal pneumonia
 Group B streptococci, 179
- Neonatal respiratory distress syndrome (NRDS), **661**
- Neonates
 abstinence syndrome, 615
 Apgar score, 634
Candida albicans in, 153
 coagulation cascade in, 413
 conjunctivitis, 142, 149
 deprivation effects, 556
 esophageal atresia in, 359
 flora with C-section, 178
 galactosemia in, 80
 gastroenteritis, 168
 gray baby syndrome in, 192
 hemolytic anemia in, 422
 herpes in, 164
 hyperthermia in, 241
 hypertrophic pyloric stenosis in, 359
 indirect inguinal hernia in, 370
 intraventricular hemorrhage, **512**
- jaundice in, 393
 kernicterus, 194, 204
Listeria monocytogenes in, 139
 low birth weight, 635
 meningitis in, 182
 necrotizing enterocolitis and, 386
 obesity risk factors, 636
 pneumonia in, 149
 primitive reflexes in, 510
 sickle cell anemia in, 422
Streptococcus agalactiae in, 137
 TORCH manifestations in, 182
- Neoplasia
 pathology of, **219–226**, 518
 Neoplastic progression, **219**
- Neoplastic transformation, 216
 adenomatous polyps, 387
- Neostigmine, 240
- Nephritic-nephrotic syndrome
 etiology and presentation, 595
- Nephritic syndrome, **596–597**
 etiology and presentation, 595
- Nephritis, 608
- Nephroblastoma, **606**
- Nephrocalcinosis, 211
- Nephrogenic diabetes insipidus, 211, 338
 lithium toxicity, 569
 treatment, 609
- Nephrolithiasis, 606
 calcium oxalate, 69
- Nephron physiology, **585**
- Nephropathy
 diabetes mellitus, 346
 hypertension and, 300
 protease inhibitors, 203
 transplant rejection, 119
- Nephrotic syndrome, **597**
 early-onset, 606
 ESR in, 214
 etiology and presentation, 595
 fatty casts in, 594
 TBG and, 331
- Nephrotoxicity
 aminoglycosides, 191
 amphotericin B, 199
 cidofovir, 202
 cisplatin/carboplatin, 442
 cladribine, 440
 drug reaction and, 251
 drugs causing, 444
 immunosuppressants, 120
 inhaled anesthetics, 550
 streptomycin, 197
 sulfonamides, 194
- Nerve blockade (local anesthetics), 550
- Nerve fibers, 495
- Nerves
 lower extremity, 452, 453
 upper extremity, 446, 447
- Nerve trunk, 495
- Nesiritide, 296
- Neural crest
 derivatives of, 613
- Neural crest cells, 490, 494
- Neural development, **490**
- Neural plate, 490
- Neural tube, 490
 derivatives, 613
 formation, 612
- Neural tube defects, **491**
 maternal diabetes, 614
 prevention, 68
 valproic acid, 544
- Neuraminidase, 169–170
- Neuroblastomas, **350**
 incidence and mortality, 222
 oncogenes and, 224
 paraneoplastic syndromes with, 228
- Neurocutaneous disorders, **525**
- Neurodegenerative disorders, **520–522**
 drug therapy for, **549**
- Neuroectoderm, 490
 derivatives of, 612
 pituitary gland, 327
- Neuroendocrine tumors, **350**
- Neurofibromatosis, 535
 chromosome association, 64
 inheritance, 60
 tumor suppressor genes and, 224
 types I and II, 525
 variable expressivity, 56
- Neurofilaments
 cytoskeletal element, 48
 immunohistochemical stain for, 227
- Neurogenic ileus, 240
- Neurohypophysis, 327
 hypothalamus and, 498
- Neuroleptic malignant syndrome (NMS), 551, 569
- Neurological signs
 proximal upper and lower extremity, 502
- Neurologic drug reactions, 251
- Neurology and special senses, 490–544
 anatomy/physiology, 493–515
 embryology, 490–492
 ophthalmology, 534–541
 otology, **533–534**
 pathology, 511–518
 pharmacology, 544–551
- Neuromuscular blocking drugs, **551**
- Neuromuscular disorders
 paraneoplastic syndromes, 228
- Neuromuscular junction
 skeletal muscle, 236
 diseases of, **472**
- Neurons, **493**
 in ascending spinal tracts, 509
 local anesthetics, 550
 origins of, 490
 Parkinson disease, 548
 primary motor cortex, 509
- Neuron-specific enolase, 226, 350
- Neuropathic pain, 515
- Neuropathy
 diabetes mellitus, 346
- Neurosyphilis, 147
- Neurotoxicity
 cladribine, 440
 immunosuppressants, 120
 methylmercury exposure, 614
 methylxanthines, 687
 vincristine, 441
- Neurotransmitters
 bacterial toxin effects, 132
 changes with disease, **495**
 vomiting center receptors, 496
- Neurovascular pairing, **455**
- Neutralization (antibody), 104
- Neutropenia, 424
 ganciclovir, 202
 rheumatoid arthritis, 466
 ticlopidine, 437
- Neutrophils, **406**
 chemotaxis, 44, 106, 133, 406, 485, 487, 691
 corticosteroid effect on, **424**
- IL-8 and, 108
 innate immunity, 99
 left shift, **424**
 in leukocyte adhesion deficiency, 117
 LTB4, 485
 megaloblastic anemia, 420
 in MI, 305
 necrosis and, 209
 nonmegaloblastic anemia, 420
 pseudo-Pelger-Huet anomaly, 432
 stimulation of, 44
 wound healing, 216
- Nevi, 220
- Nevirapine
 cytochrome P-450 and, 252
 HIV therapy, 203
- NF1/NF2* genes, 224
- N-formylmethionine (fMet), 44
- NF- κ B, 120
- Niacin
 cutaneous flushing, 248
 gout, 250
 hyperglycemia, 249
 myopathy caused by, 250
- Nicardipine, 318
- Nicotinamides, 75
- Nicotine
 intoxication and withdrawal, **571**
 teratogenicity, 614
- Nicotinic acetylcholine receptors, 166, **236**
- Niemann-Pick disease, 88
- Nifedipine, 316, **318**, 643
- Nifurtimox, 158, 200
- Night terrors
 benzodiazepines for, 497
- Nigrostriatal pathway, 499
- Nikolsky sign
 blistering skin disorders, 480
 scalded skin syndrome, 479
- Nilotinib, 443
- Nimodipine, **318**, 513
- Nipple
 intraductal papilloma, 649
 lactational mastitis, 649
- Nissl bodies, 46
- Nissl substance
 chromatolysis, 495
 neurons, 493
- Nitazoxanide, 155
- Nitrates, 319
 mechanism and clinical use, **318**
- Nitric oxide, 371
 free radical injury and, 210
- Nitrites
 urinary tract infections, 181
- Nitroblue tetrazolium dye reduction test, 117
- Nitrofurantoin
 hemolysis in G6PD deficiency, 250
 pulmonary fibrosis, 251
- Nitroglycerin, 318
 acute coronary syndromes, 307
 angina, 304
- Nitroprusside, 318
- Nitrosamines
 as carcinogens, 225
 stomach cancer and, 379
- Nitrosoureas, 441
- Nitrous oxide, 550
- Nizatidine, 399
- NMDA receptor antagonist
 ketamine as, 550
 memantine as, 549
- N-*myc* oncogene, 350

- NNRTIs, 203
Nocardia spp
 aerobic culture requirements, 126
 comparison with *Actinomyces* spp, **139**
 effects and treatment, 139
 necrosis and, 209
 sulfonamides for, 194
 urease-positive, 127
 Ziehl-Neelsen stain, 125
 Nocturia, 654
 Nocturnal enuresis, 329
 Nodes of Ranvier, 494
 Nodular phlebitis, 314
 Nodular sclerosis lymphoma, 429
 Noise-induced hearing loss, 533
 Nonadherent patients, 268
 Nonalcoholic fatty liver disease, 389, 390, **391**, 392
 Nonbacterial endocarditis, 311
 Nonbacterial thrombotic endocarditis, 228
 Nonbenzodiazepine hypnotics, **546**
 Noncaseating granulomas
 sarcoidosis, 676
 Noncommunicating hydrocephalus, 522
 Noncompetitive agonists, 234
 Noncompetitive antagonist, 234
 Noncompetitive inhibitors, 230
 Noncompliant patients, 268
 Nondepolarizing neuromuscular blocking drugs, 551
 Nondominant parietal cortex lesions, 511
 Non-frameshift mutations
 deletions, 61
 Nonhemolytic, normocytic anemia, **421**
 Non-Hodgkin lymphoma, **430**
 corticosteroids, 120
 Hashimoto thyroiditis and, 341
 hepatitis C, 173
 HIV-positive adults, 177
 Hodgkin lymphoma vs, 429
 oncogenes and, 224
 rituximab for, 443
 vinca alkaloids for, 441
 Nonhomologous end joining, 40
 Nonmaleficence (ethics), 265
 Nonmegaloblastic macrocytic anemia, 420
 Nonnormal distributions, 262
 Nonoverlapping genetic code, 37
 Nonreceptor tyrosine kinase, 337
 Non-REM sleep stages, 497
 Nonselective antagonists, 245
 Nonsense mutations, 39
 Nonsteroidal anti-inflammatory drugs (NSAIDs), **486**
 acute pericarditis, 313
 aplastic anemia, 250
 Beers criteria, 247
 calcium pyrophosphate deposition disease, 467
 colorectal cancer
 chemopreventative, 389
 esophagitis from, 377
 gastritis with, 379
 GFR effects of, 589
 gout, **467**, 487
 headaches, 518
 interstitial nephritis, 249
 loop diuretics and, 608
 misoprostol use with, 399
 osteoarthritis, 466
 peptic ulcer disease and, 380
 rheumatoid arthritis, 466
 for sialadenitis, 376
 Non-ST-segment elevation MI (NSTEMI)
 diagnosis of, 306
 STEMI vs, 304
 treatment, 307
 Noradrenergic drugs, 574
 Norepinephrine (NE)
 adrenal medulla secretion, 327
 amphetamines and, 239
 bupropion effect on, 576
 changes with disease, 495
 circadian rhythm, 497
 direct sympathomimetic, 242
 isoproterenol vs, **243**
 male sexual response, 627
 MAO inhibitor effects, 575
 opioid effect on, 551
 pheochromocytoma secretion, 350
 release regulation, 239
 REM sleep and, 497
 vitamin B₆ and, 67
 Norethindrone, 657
 Norfloxacin, 195
 Normal distribution, 262
 Normal flora
 colonic, 137
 female genital tract, 136
 GI tract, 127
 neonates, **178**
 oropharynx, 136
 skin, 135
 Normal pressure hydrocephalus, 522
 Normal splitting, 289
 Normocytic anemia, **421**
 Norovirus, 167
 Northern blot, 53
 Nortriptyline, 575
 Nosocomial infections, 274
 Acinetobacter baumannii, 142
 Ebola, 171
 enterococci, 137
 Klebsiella, 145
 MRSA, 135
 pneumonias, 179
 Pseudomonas aeruginosa, 143
 risk factors, **185**
 UTIs as, 181
 Notochord, **490**, 612–613
 postnatal derivative of, 282
 Novobiocin
 gram-positive antibiotic test, 134
 Staphylococcus epidermidis, 135
 NRTIs, 203
 NS3/4A inhibitors, 204
 NS5A inhibitors, 204
 NS5B inhibitors, 204
 Nuchal translucency, 63
 Nucleic acids
 pathogen-associated molecular pattern (PAMP), 99
 Nucleosome, 34
 Nucleotide excision repair, 40
 Nucleotides, **35**
 deamination reactions, 35
 synthesis, 72
 Nucleus accumbens, 495
 Nucleus ambiguus, 506
 Nucleus cuneatus, 509
 Nucleus pulposus
 collagen in, 50
 fetal precursor, 282
 Nucleus tractus solitarius (NTS), 496
 Null hypothesis, 263
 Number needed to harm (NNH), 258
 Number needed to treat (NNT), 258
 Nursemaid's elbow, 461
 Nutcracker syndrome, 363
 Nutmeg liver, 309, 392
 Nutrition, 65–72
 Nyctalopia, 66
 Nystagmus
 cerebellum, 499
 common lesions with, 511
 Friedreich ataxia, 531
 internuclear ophthalmoplegia, 543
 PCP, 571
 phenytoin, 544
 stroke and, 514
 Nystatin, **199**
O
 Obesity
 amphetamine for, 242
 anovulation with, 645
 cholelithiasis and, 396
 DM type 2 and, 347
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300
 hypoventilation syndrome, 679
 lateral femoral cutaneous nerve, 452
 osteoarthritis/rheumatoid arthritis, 466
 renal cell carcinoma association, 605
 sleep apnea, 679
 stress incontinence and, 599
 Obligate intracellular organisms, 127
 Observational studies, **256**
 errors in, 260–261
 Observer-expectancy bias, 260
 Obsessive-compulsive disorder (OCD)
 diagnostic criteria/treatment, **563**
 drug therapy for, 572
 SSRIs for, 575
 Tourette syndrome and, 557
 venlafaxine for, 575
 Obsessive-compulsive personality disorder, 566
 Obstructive jaundice, 398
 Obstructive lung diseases, **674**–675
 flow volume loops in, 673
 Obstructive shock, 310
 Obstructive sleep apnea, 679
 pulsus paradoxus in, 310
 Obturator nerve, 452
 Occipital cortex, 515
 Occipital lobe, 501
 Occipital sinus, 503
 Occult bleeding, 387
 FOBT for, 388
 Octreotide, 371, **400**
 acromegaly, 339
 GH excess, 329
 hypothalamic/pituitary drugs, 354
 Ocular motility, **540**
 Oculomotor nerve (CN III), 506
 ocular motility, **540**
 palsy of, 513, **541**
 pupillary contraction, 539
 Odds ratio (OR), 256, **258**
 Ofloxacin, 195
 Okazaki fragments, 38
 “OK gesture,” 451
 Olanzapine, 573
 Olfaction
 hallucinations, 559
 limbic system in, 499
 Olfactory nerve (CN I), 506
 Oligoclonal bands, 523
 Oligodendrocytes, **494**
 Oligodendroglia, 490
 in multiple sclerosis, 494
 Oligodendroglomas, 526
 Oligohydramnios, 578, **641**
 Oligomenorrhea, 633
 Oligomycin, 78
 Oligospermia, 400
 Olive-shaped mass, 359
 Omalizumab, 122, **687**
 Ombitasvir, 204
 Omental foramen, 361
 Omeprazole, 399
 Omphalocele, 358
 Omphalomesenteric cysts, 384
 Omphalomesenteric (vitelline) duct, 618
Onchocerca volvulus, 159
 Oncocytoma (renal), 605
 Oncogenes, **224**
 esophageal cancer and, 378
 hypertension risk factors, 300

- Opsonization, **104**, 106
antibodies in, 112
complement in, 106
encapsulated organisms, 98
- Optic disc
papilledema in, 538
- Optic nerve (CN II), 506
embryologic derivation, 613
- Optic neuritis, 523
- Optic neuropathy, 197
- Optochin
gram-positive antibiotic test, 134
- Oral advance directives, 266
- Oral contraceptives (OCPs)
hepatic adenomas and, 392
prolactin effects on, 330
reproductive hormones, 656
SHBG effects on, 337
venous sinus thrombosis with, 503
- Oral glucose tolerance test, 346
- Oral hairy leukoplakia, 177
- Oral/intestinal ganglioneuromatosis, 351
- Oral rehydration therapy, 146
- Oral thrush, 177
- Orange body fluids, 196
- Orchiectomy, 651
- Orchiopexy, 651
- Orchitis, 170, **654**
- Orexigenic effect, 336
- Orexin, 568
hypocretin receptor antagonist, Suvorexant as, 547
- Organ failure, in acute pancreatitis, 397
- Organic acidemias, **85**
- Organ of Corti, 533
- Organogenesis
embryologic derivatives, 612, **613**
errors in, 613
fetal development, 612
teratogens, 614
- Organophosphates
toxicity treatment, 248
- Organ transplants
azathioprine for, 440
cytomegalovirus, 186
hairy leukoplakia and, 479
kidneys, 580
WBC casts, 594
- Organum vasculosum of the lamina terminalis (OVLT), 498
- Orientation, **557**
- Origin of replication, 38
- Orlistat, **400**
diarrhea, 249
- Omithine
cystinuria, 85
kidney stones and, 598
urea cycle, 82
- Omithine transcarbamylase, 74
- Omithine transcarbamylase deficiency, 61, **83**
- Orotic acid, 83
- Orotic aciduria, 420
- “Orphan Annie” eyes (nuclei), 343
- Orthomyxoviruses, 168
characteristics of, **167**, 168
influenza viruses, 169
- Orthopedic conditions, 460
lower extremity, 460, 461
- Orthopnea, 309
- Orthostatic hypotension
adrenal insufficiency, 349
 α -blockers, 244
phenoxybenzamine, 244
- Ortolani maneuver, 461
- Oseltamivir, **201**
- Osgood-Schlatter disease, 461
- Osler nodes, 311
- Osler-Weber-Rendu syndrome, 316
- Osmotic demyelination syndrome, 524
SIADH and, 338
- Osmotic diarrhea, 381
- Osmotic laxatives, 401
- Ossicles, 533
conductive hearing loss and, 51
- Ossification, 458
- Osteitis deformans, **463**
- Osteitis fibrosa cystica, **345**, 459, 464
- Osteoarthritis, 466
celecoxib for, 486
vs rheumatoid arthritis, **466**
- Osteoarthropathy, hypertrophic cancer association, 228
- Osteoblastoma, 464
- Osteoblasts, 459
bone formation, 458
cortisol effect on, 336
Paget disease of bone, 463
teriparatide effect on, 487
- Osteochondroma, 464
- Osteoclasts, 459
bisphosphonate effects, 486
bone formation, 458
osteopetrosis, 463
Paget disease of bone, 463
- Osteodystrophy
renal, 345, **603**
- Osteogenesis imperfecta
bisphosphonates, 486
collagen and, 50
collagen synthesis and structure in, **51**
- Osteogenic sarcoma, 463, 465
- Osteoma, 220, 464–465
nomenclature for, 220
- Osteomalacia
hypophosphatemia, 591
lab values with rickets, 464
rickets, **463**
- Osteomyelitis, **180**
Pseudomonas aeruginosa, 143
sickle cell anemia, 422
Staphylococcus aureus, 135
- Osteonecrosis, **463**, 486
- Osteopenia, 463
- Osteopetrosis, **463**, 464
- Osteophytes, 466
- Osteoporosis, **462**
bisphosphonates, 486
corticosteroids, 120
denosumab, 122
drug reaction and, 250
estrogen, 459
Gaucher disease, 88
heparin, 436
homocystinuria, 84
hormone replacement therapy, 657
lab values in, 464
pituitary prolactinomas, 328
raloxifene, 443, 656
teriparatide for, 487
thiazides for, 609
- Osteosarcoma, 220, 464–465
- Otitis media
brain abscesses with, 180
Haemophilus influenzae, 128, **142**
Langerhans cell histiocytosis, 434
Streptococcus pneumoniae, 136
Wegener granulomatosis and, 315
- Otology, 533
- Ototoxicity
aminoglycosides, **191**, 204, 614
cisplatin/carboplatin, 442
drug reaction and, 251
ethacrynic acid, 608
loop diuretics, 608
- Ouabain
sodium-potassium pump and, 49
- Outcome (quality measurement), 273
- Outer membrane, 124
- Outflow tract formation, 281
- Outpatient follow-up, 272
- “Oval fat bodies,” 594
- Ovarian artery, 625
- Ovarian cancer
breastfeeding and, 636
cisplatin/carboplatin for, 442
epidemiology of, 643
hypercalcemia and, 228
irinotecan/topotecan for, 442
Lynch syndrome and, 388
oncogenes and, 224
paclitaxel for, 441
Psammoma bodies in, 227
tumor suppressor genes and, 224
- Ovarian cycle, 632
- Ovarian cysts, **646**
- Ovarian ligament, 625
- Ovarian neoplasms
classification and characteristics, **646**
epithelial tumors, 646
germ cell tumors, 647
sex cord stromal tumors, 647
- Ovarian teratomas
paraneoplastic syndrome, 228
- Ovaries
anatomy of, 625
descent of, 624
epithelial histology, 626
estrogen production, 630
lymphatic drainage, 624
- Overflow incontinence, 599
- Overuse injury
elbow, **459**
knee, 461
radial nerve, 447
wrist, 459
- Ovotesticular disorder, 638
- Ovulation
anovulation causes, 645
progesterone and, 630
prolactin effect on, 330
regulation of, **631**
- Ovulatory uterine bleeding, 633
- “Owl eyes” inclusions, 165, 429
- Oxacillin
characteristics of, 188
- Oxaliplatin, **442**
- Oxazepam, 546
- Oxazolidinones
mechanism and use, 193
- Oxidative burst, 109
- Oxidative phosphorylation, 78
metabolic site, 72
poisons, 78
- Oxybutynin, 237, **241**
- Oxygen
deprivation, **669**
in blood, **666**
for carbon monoxide poisoning, 248
cluster headaches, 518
exercise and, 670
hemoglobin, 665
toxicity, 210
- Oxygen-hemoglobin dissociation curve, **666**
- Oxytocin
functions of, 328
hypothalamus production, 498
lactation and, 636
pituitary gland and, 327
signaling pathways for, 337
- P**
- P-450, 197
- P2Y12 receptor
inhibitors and, 437
thrombogenesis and, 411
- P53 gene mutation
dominant negative mutation, 56
- Pacemaker action potential, **292**
- Pacinian corpuscles, 494
- Paclitaxel, 441
microtubules and, 48
targets of, 438
- Paget disease
in breast, 650
- Paget disease (extramammary), 644
- Paget disease of bone, 463
bisphosphonates, 486
lab values in, 464
osteosarcomas and, 465
woven bone in, 458
- Pain
with headaches, 518
loss in syringomyelia, 492
post-stroke syndrome, 515
receptors for, 494
spinal tracts for, 509
thalamic nuclei and, 498
treatment in multiple sclerosis, 523
unilateral visual loss and, 523
- Pale (anemic) infarct, 210
- Paliperidone, 573
- Palivizumab, 122
pneumonia prophylaxis, 169
- Pallor in aplastic anemia, 421
- Palmar interossei, 450
- Palmar reflex, 510
- PALM-COEIN uterine bleeding classification, 633
- PALP, 653
- Panacinar emphysema, 392, **674**
- p-ANCA
sclerosing cholangitis and, 395
ulcerative colitis, 382
- Pancoast tumor, **685**
lung cancer, 684
superior vena cava syndrome, 685
thoracic outlet syndrome, 448
- Pancreas
annular, 360
biliary structures and, 368
blood supply and innervation of, 364
buds, 360
divisum, 360
embryology, **360**
- Pancreatic cancer
5-fluorouracil for, 440
adenocarcinomas, **398**
biliary cirrhosis and, 395
carcinogens causing, 225
hyperbilirubinemia with, 393
oncogenes and, 224
paraneoplastic syndromes with, 228
tumor suppressor genes and, 224

- Pancreatic ducts, 360, 368
 Pancreatic insufficiency, **381**, 397
 Pancreatic islet cell tumors, **351**
 Pancreatic lipase, 93
 Pancreatic secretions, **373**
 Pancreatitis
 acute, 397
 acute respiratory distress syndrome and, 678
 alcoholism, 571
 chronic, 397
 corticosteroids and, 249
 drug reactions and, 249
 hyperchylomicronemia, 94
 hyperparathyroidism, 345
 hypertriglyceridemia, 94
 mumps, 170
 necrosis and, 209
 NRTIs, 203
 valproic acid, 544
 Pancytopenia, 421
 Chédiak-Higashi syndrome, 117
 cytarabine, 440
 Gaucher disease, 88
 leishmaniasis, 158
 osteopetrosis and, 463
 with immunosuppressants, 120
 Panic disorder
 drug therapy for, 572
 SSRIs for, 563, **575**
 symptoms and treatment, **563**
 venlafaxine for, 575
 Panitumumab, **442**
 Panniculitis, 482
 Pantoprazole, 399
 Pantothenic acid, 67
 Papillary carcinomas, 220
 Papillary cystadenoma
 lymphomatosum, 376
 Papillary muscle
 blood supply to, 307
 rupture, 305, **307**
 Papillary thyroid carcinomas, 343
 carcinomas for, 225
 Psammoma bodies in, 227
 Papilledema, 521, **538**
 hypertensive emergency and, 300
 Papillomas, 220
 Papillomaviruses
 characteristics of, 164
 DNA viruses, 163
 genome, 163
 Pappenheimer bodies, 416
 Papules, 475
 capillary, 478
 molluscum contagiosum, 479
 Para-aminohippuric acid (PAH), 582
 Para-aortic lymph nodes, 624
 Paracoccidiodomycosis, 151
 Paracortex (lymph node), 96
 Paracrine, 589
 Paradoxical splitting, 289
 Paraesophageal hiatal hernia, 370
 Parainfluenza
 croup, 170
 paramyxovirus, 167, 169
 Parakeratosis, 475
 Paralysis
 conversion disorder and, 566
 of face, 514
 Guillain-Barré syndrome, 524
 poliovirus, 186
 rabies, 171
 stroke effects, 514
 unvaccinated children, 186
 Paralytic ileus, 441
 Paramedian pontine reticular formation
 extraocular movements, 497
 Paramedian pontine reticular formation lesions, 511
 Paramesonephric (Müllerian) duct, 622
 Paramyxoviruses, **169**
 characteristics of, 167, 168
 croup, 170
 mumps, 170
 Paraneoplastic syndromes, **228**, 605
 Paranoia
 LSD, 571
 Parasites
 infections with immunodeficiency, 118
 Parasitic infections
 myocarditis with, 313
 Parasitology, 155–**161**
 Parasympathetic nervous system
 cranial nerves, supply of, 236
 male erection, 627
 receptor targets, 236
 VIP and, 371
 Parathyroid adenomas
 hyperparathyroidism caused by, 345
 MEN 1/MEN 2A syndromes, 351
 Parathyroid disease diagnosis, 343
 Parathyroid glands
 pharyngeal pouch derivation, 621
 Parathyroid hormone (PTH), **332**, 590
 bone disorders, 464
 bone formation, 459
 calcitonin and, 333
 calcium homeostasis and, 333
 in hyperparathyroidism, 345
 osteomalacia/rickets, 463
 Paget disease of bone, 463
 signaling pathways of, 337
 thymic aplasia, 116
 Paraumbilical vein, 365
 Paraventricular nucleus, 498
 Parental consent, 265
 Paresthesias
 vitamin B₁₂ deficiency, 69, 530
 Parietal cells (stomach), 372
 Parietal cortex lesions, 511
 Parietal lobe, 501
 Parietal peritoneum, 369
 Parinaud syndrome, **511**, 528
 Parkinson disease, 520
 benzotropine for, 241
 dopaminergic pathways, 499
 Lewy bodies, 520
 metoclopramide contraindication, 400
 nigrostriatal pathway and, 499
 resting tremor in, 519
 seborrheic dermatitis association, 476
 therapy for, **548**
 trihexyphenidyl, 241
 ubiquitin-proteasome system in, 48
 Parkinson-like syndrome, 251
 Parotid gland
 embryologic derivation, 613
 enlargement of, 468
 stones in, 376
 tumors in, 376
 Parotitis
 mumps, 170
 Paroxetine, 575
 Paroxysmal nocturnal hemoglobinuria, 422
 CD55 deficiency, 107
 eculizumab for, 122
 flow cytometry diagnosis, 54
 intravascular hemolysis in, 421
 Partial agonists, 234
 Partial (focal) seizures, 517
 drug therapy for, 544
 Partial thromboplastin time (PTT), 426
 Parvovirus
 characteristics of, 164
 DNA viruses, 163
 naked viruses, 163
 Parvovirus B19
 hereditary spherocytosis, 422
 hydrops fetalis, 182
 rash, 183
 Passive aggression, 555
 Passive immunity, **110**
Pasteurella multocida
 osteomyelitis, 180
 transmission, **149**, 186
 Patau syndrome, 63
 cataracts, 535
 chromosome association, 64
 Patches (skin)
 pityriasis rosea, 482
 psoriatic arthritis, 469
 Patellar reflex, 510
 Patellofemoral syndrome, 461
 Patent ductus arteriosus (PDA)
 congenital rubella, 300
 fetal alcohol syndrome, 300
 heart murmur with, 291
 indomethacin for, 486
 mechanism and treatment, 299
 neonatal respiratory distress syndrome and, 661
 Patent foramen ovale
 atrial septal defect vs, 299
 septal fusion failure, 280
 Patent urachus, 618
 Pathogen-associated molecular patterns (PAMPs), 99
 Pathogen recognition features, 99
 Pathologic grief, 562
 Pathology
 cardiovascular, 298–312
 endocrine, 338–354
 gastrointestinal, 376–397
 hematologic/oncologic, 404–424, 414–434
 musculoskeletal/skin/connective tissue, 459–467
 neoplasia, 219–226
 neurological, 511–518
 psychiatric, 556–570
 renal, 594–605
 reproductive, 638–652
 respiratory, 671–681
 USMLE Step 1 preparation for, 277
 Pautrier microabscess, 430
 Pavlovian (classical) conditioning, 554
 Payment models for healthcare, 271
 P-bodies, 41
 PCP (phencyclidine)
 intoxication and withdrawal, 571
 PCSK9, 93
 PCSK9 inhibitors, 320
 PCV13 (pneumococcal conjugate vaccine), 127
 PDSA cycle, **273**
 Pearson correlation coefficient(r), **264**
 Peau d'orange, 650
 Pectinate line, **366**
 Pectineus, 451, 452
 Pectoriloquy (whispered), 680
 Pediatric patients
 arthritis in, 468
 brachial plexus injury, 448
 common causes of death, 272
 common fractures, **462**
 common orthopedic conditions, 461
 cystic fibrosis, 60
 dactinomycin for, 439
 failure to thrive, 556
 growth retardation in, 603
 hemolytic disease of newborn, 405
 hyperbilirubinemia (newborns), 393
 infant deprivation effects, 556
 intraventricular hemorrhage, 512
 juvenile polyposis syndrome in, 387
 leukocoria in, 538
 Munchausen syndrome by proxy, 566
 neglect in, 556
 neuroblastomas in, 350
 precocious puberty, 57, 335
 primary brain tumors, 528
 rashes, 183
 renal malignancy in, 606
 rhabdomyomas in, 316
 scalded skin syndrome, 479
 sleep terror disorder in, 568
 strawberry hemangiomas in, 478
 tetracycline side effects, 192
 unvaccinated, 186
 Wilms tumors in, 606
 Pegloticase, 487
 Pegvisomant, 339
 Pellagra, 67
 Pelvic inflammatory disease (PID), **185**
Actinomyces spp, 139
 chlamydia, 148, 184
Chlamydia trachomatis, 149
 clinical features, 185
 copper IUD, 657
 ectopic pregnancy, 641
 gonorrhoea, 184
Neisseria spp, 142
 Pelvic splanchnic nerves, 236
 Pelvis
 fracture and nerve injury, 452
 nerve injury with surgery, 452
 Pemphigus vulgaris, 480
 acantholysis and, 475
 autoantibody, 115
 type II hypersensitivity, 112
 Pencil-in-cup deformity (X-ray), 469
 Penicillamine
 for lead poisoning, 248
 myopathy, 250
 for Wilson disease, 395
 Penicillin
Actinomyces spp, 139
 antipseudomonal, 188
 Coombs-positive hemolytic anemia, 250
 penicillinase-resistant, 188
 penicillinase-sensitive, 188
 prophylaxis, 198
 rash, 250
 for rheumatic fever, 312
 Penicillinase-resistant penicillins, **188**

- Penicillinase-sensitive penicillins, **188**
 Penicillin-binding proteins (PBPs)
 in bacteria, 124
 Penicillin G, V, **187**
 meningococci, 142
 prophylaxis, 198
 Penile cancer, 226
 Penis
 congenital abnormalities, **624**
 female homolog, 624
 lymphatic drainage, 624
 pathology of, **651**
 Pentamidine, 154
 Pentazocine, 551, **552**
 Pentobarbital, 546
 Pentose phosphate pathway
 functions of, **79**
 Pentostatin, 432
 PEP carboxykinase, 74
 Pepsin, 372
 Pepsinogen
 location of, 373
 somatostatin and, 371
 Peptic ulcer disease, **380**
 glycopyrrolate for, 241
H. pylori and, 146
 H₂ blockers for, 399
 misoprostol for, 399
 proton pump inhibitors for, 399
 Zollinger-Ellison syndrome, 352
 Peptidoglycan
 in gram negative bacteria, 124
Peptostreptococcus spp
 alcoholism, 179
 Percussion (chest), 680
 Perforation (GI), 380
 necrotizing enterocolitis, 386
 Perforin
 cytotoxic T cells and, 102
 extrinsic pathway and, 208
 natural killer cells and, 101
 Performance anxiety, 567
 Perfusion, and ventilation, 669
 Perfusion-limited gas exchange, 668
 Perfusion pressure regulation, 297
 Periartertiolar lymphatic sheath
 (PALS), 98
 Pericardial effusion, 684
 Pericarditis
 acute, 313
 fibrinous, 305
 jugular venous pulse in, 287
 Kussmaul sign in, 316
 picornaviruses, 167
 postinfarction, 305, 307
 pulsus paradoxus in, 310
 renal failure, 603
 rheumatoid arthritis, 466
 Pericardium
 anatomy of, 283
 calcification in, 211
 Perineal straddle injury, 627
 Perinephric abscesses, 600
 Perineurium, 495
 Periodic acid-Schiff stain, 125
 glycogen storage diseases, 87
 Periorbital edema
 thyroid disease and, 340
Trichinella spiralis, 161
 Peripartum mood disturbances, **562**
 Peripheral blood smear, 416
 basophilic stippling, 419
 postsplenectomy, 98, 423
 Rouleaux formation, 431
 schistocytes, 423
 Schüffner stippling, 157
 smudge cells, 432
 spherocytes and agglutinated
 RBCs, 423
 Peripheral edema
 calcium channel blockers, 318
 heart failure, 309
 Peripheral facial palsy, 532
 Peripheral nerves, **495**
 Peripheral nervous system (PNS)
 origins of, **490**
 Peripheral neuropathy
 alcoholism, 571
 drug reactions and, 251
 Fabry disease, 88
 isoniazid, 197
 Krabbe disease, 88
 NRTIs, 203
 oxazolidinones, 193
 sorbitol, 81
 vincristine, 444
 vitamin B₆ deficiency, 67
 Peripheral precocious puberty, 637
 Peripheral resistance, 243
 Peripheral vascular disease, 302
 Peripheral vertigo, 534
 Perioplasm
 in bacteria, 124
 Peristalsis
 motilin receptor agonists and, 371
 visible, 359
 Peritoneum, 360
 hernias and, 370
 irritation with Mittelschmerz, 631
 Peritonitis
 appendicitis, 383
 diverticulitis, 383
 spontaneous bacterial, 389, 390
 Periventricular plaques, multiple
 sclerosis, 523
 Permethrin, 200
 for scabies, 161
 Permissive action
 catecholamine responsiveness, 235
 Pernicious anemia, 372
 autoantibody, 115
 B₁₂ deficiency, 69, **420**
 Peroneus brevis, 453
 Peroneus longus, 453
 Peroxisome
 metabolic processes, **47**
 Persistent cervical sinus, 619
 Persistent depressive disorder
 (dysthymia), 561
 Persistent thyroglossal duct, 326
 Persistent truncus arteriosus, 281, 298
 Personality, **565**
 Personality disorders, 565
 Cluster A, 565
 Cluster B, 565
 Cluster C, 565, 566
 Pertussis, 126
 Pertussis toxin, 132, 143
 Pes cavus
 Friedreich ataxia, 531
 Petechiae
 aplastic anemia, 421
 Peutz-Jeghers syndrome, 220, **387**
 PEX genes, 47
 Peyer patches, 362, **374**
 IgA antibody production, 105
Salmonella/Shigella invasion, 144
 Peyronie disease, 651
 PGI₂, 485
 P-glycoprotein, **227**
 Phage
 bacterial transduction, 130
 Phagocytes, 117
 Phagocytosis
 dendritic cells, 408
 eosinophils, 408
 group A streptococcal inhibition,
 136
 M protein prevention of, 129
 β-hemolytic bacteria inhibition
 of, 135
 Phalen maneuver, 459
 Pharmaceutical company
 sponsorship, 269
 Pharmacokinetics, **231**
 Pharmacology, 230–254
 autonomic drugs, 236–245
 cardiovascular, 316–322
 endocrine, 352–354
 gastrointestinal, 398–400
 hematologic/oncologic, 435–443
 musculoskeletal/skin/connective
 tissue, 485–487
 neurology, 544–551
 pharmacodynamics, 232–234
 pharmacokinetics, 230–231
 psychiatric, 572–576
 renal, 607–610
 reproductive, 655–658
 respiratory, 686–687
 toxicities and side effects, 248–251
 USMLE Step 1 preparation for,
 277
 Pharyngeal apparatus, **619**
 Pharyngeal arch derivatives, **620**
 1st pharyngeal arch, 620
 2nd pharyngeal arch, 620
 4th–6th pharyngeal arches, 620
 Pharyngeal cleft derivatives, **619**
 Pharyngeal pouch derivatives, **621**
 1st pharyngeal pouch, 621
 2nd pharyngeal pouch, 621
 4th pharyngeal pouch, 621
 Pharyngitis
 adenovirus, 164
 diphtheria, 139
 mononucleosis, 165
 prophylaxis (rheumatic fever), 198
Streptococcus pyogenes, 136
 unvaccinated children, 186
 Pharyngoesophageal false
 diverticulum, 384
 Pharynx, 662
 blood supply and innervation of,
 364
 Phenacetin, 606
 Phenelzine, 575
 Phenobarbital, 546
 epilepsy, 544
 as weak acid, 233
 Phenotypic mixing, 162
 Phenoxybenzamine, 244
 for pheochromocytomas, 350
 Phentolamine, 244
 Phenylalanine
 classification of, 81
 tyrosine catabolism, 83
 Phenylephrine, 242, **686**
 Phenylketones, 84
 Phenylketonuria
 tyrosine in, **84**
 Phenytion
 cytochrome P-450 and, 252
 epilepsy, 544
 folate deficiency caused by, 420
 gingival hyperplasia, 250
 megaloblastic anemia, 250
 peripheral neuropathy, 251
 vitamin B₉ deficiency, 68
 zero-order elimination of, 232
 Pheochromocytomas, **350**
 MEN 2A/MEN 2B and, 351
 phenoxybenzamine for, 244
 Philadelphia chromosome
 in myeloproliferative disorders, 433
 translocations of, 434
 Phlebitis
 IV amphotericin B, 199
 Phlebotomy
 for hemochromatosis, 395
 Phobias, **563**
 agoraphobia, 563
 social anxiety disorder, 563
 Phocomelia, 614
 Phonophobia, migraine headache,
 518
 Phosphatases, 73
 Phosphodiesterase (PDE) inhibitors,
246, 658, 686
 Phosphoenolpyruvate carboxykinase,
 78
 Phosphofructokinase-1 (PFK-1)
 glycolysis and, 73
 metabolic pathways, 74
 Phospholipid bilayer sac
 in bacteria, 124
 Phospholipids, 374
 Phosphorus in Paget disease of bone,
 463
 Phosphorylases, 73
 Phosphorylation, 45
 Photophobia
 leptospirosis, 147
 migraine headache, 518
 rabies, 171
 Photosensitivity
 drugs causing, 192, 194, 250
 Phototherapy for jaundice, 393
 Phrenic nerve, 663
 Phylloides tumor, 649
 Physical abuse (child), 556
 Physician-assisted suicide, 268
 Physician-patient relationship, 268
 Physiologic dead space, 664
 Physiologic neonatal jaundice, **393**
 Physiology
 cardiovascular, 284–298
 endocrine, 328–336
 gastrointestinal, 371–375
 hematologic/oncologic, 410
 neurological, 493–515
 renal, 581–592
 reproductive, 629–636
 respiratory, 664–669
 USMLE Step 1 preparation for,
 276
 Physostigmine
 anticholinergic toxicity treatment,
 248
 anticholinesterase, 240
 glaucoma, 552
 Pia mater, 496
 Pica, 567
 Pick disease, 520
 bodies, 520
 Pickwickian syndrome, 679
 Picornaviruses, 163, **168**
 characteristics, 167
 genomes, 163
 Pierre Robin sequence, 620
 Pigmented skin disorders, **476**
 Pigment-producing bacteria, **128**
 Pigment stones, 396
 Pill-rolling tremor, 519

- Pilocarpine, **240**, 552
 Pilocytic astrocytoma, 528
 Pilus, 124
 Pimozide, 572
 Pindolol, 245, 319
 Pineal gland, 504
 Pinealoma, 528
 Pinworms, 159
 Pioglitazone, 353
 Piperacillin
 characteristics of, 188
 Pseudomonas aeruginosa, 143
 Piroxicam, 486
 Pisiform bone, 449
 Pitting edema, 309
 Pituitary adenoma, 339, 527
 Pituitary apoplexy, 339
 Pituitary drugs, 354
 Pituitary gland, **327**
 Pituitary prolactinomas, 328
 Pituitary tumors
 diabetes insipidus, 338
 MEN 1 and, 351
 Pityriasis rosea, 482
Pityrosporum spp, 152
 pKa, 233
 PKD genes
 renal cyst disorders and, 604
 Placebo, 256
 Placenta
 estrogen production, 630
 fetal component, **617**
 hormone secretion by, 633
 maternal component, 617
 progesterone production, 630
 Placenta accreta/increta/percreta, 640
 Placental aromatase deficiency, **639**
 Placental insufficiency
 oligohydramnios and, 641
 Potter sequence, 578
 preeclampsia, 643
 Placenta previa, 640
 Plague, 149
 Plantar aponeurosis, 461
 Plantar fasciitis, 461
 Plantar flexion, 453
 Plantaris, 453
 Plantar reflex, 510
 Plaques (skin), 475
 actinic keratosis, 482
 basal cell carcinoma, 484
 hairy leukoplakia, 479
 lichen planus, 482
 pityriasis rosea, 482
 psoriasis, 477
 seborrheic dermatitis, 476
 squamous cell carcinoma, 484
 Plasma cell dyscrasias, **431**
 Plasma cells, **409**
 Plasmalogens, 47
 Plasma membrane
 sodium-potassium pump, 49
 structure of, 49
 Plasma osmolality
 DI treatment, 338
 Plasmapheresis, 524, 596
 Plasma protein concentration, 583
 Plasma volume measurement, 581
 Plasminogen, 437
Plasmodium spp
 chloroquine, 200
 Plasmodium falciparum, 157, 200
 Plasmodium malariae, 157
 Plasmodium ovale, 157
 Plasmodium vivax, 157
 Platelet-activating factor, 406
 Platelet-derived growth factor (PDGF)
 in wound healing, 216
 signaling pathways for, 337
 Platelet disorders, 426, **427**
 transfusion for, 429
 Platysma muscle, 620
 Pleiotropy, 56
 Pleomorphic adenomas, 376
 Pleural effusion, **681**
 asbestosis, 677
 lung cancer, 684
 mesothelioma, 678
 physical findings, 680
 Pleuritis, 466
 Pleuroperitoneal membrane, 370
 Plicae circulares, 362
 Plummer-Vinson syndrome, 377, 418
 Pneumatosis intestinalis, 386
 Pneumococcal vaccine, 127
 Pneumoconiosis, 675, **677**
Pneumocystis jirovecii, **154**
 dapsone, 194
 fluorescent antibody stain, 125
 HIV-positive adults, 177
 immunocompromised patients, 179
 prophylaxis, 198
 silver stain for, 125
 TMP-SMX, 194
 Pneumocytes, 661
 Pneumomediastinum, 672
 Pneumonia, **683**
 acute respiratory distress syndrome, 678
 adenovirus, 164
 chlamydiae, 148
 coccidioidomycosis, 151
 common causes, **179**
 Haemophilus influenzae, 142
 measles-associated death, 170
 Pneumocystis jirovecii, 154
 PPI adverse effects, 399
 Q fever, 150
 readmissions with, 272
 Staphylococcus aureus, 135
 Streptococcus pneumoniae, 136
 Streptococcus agalactiae, 137
 VZV, 165
 Pneumoperitoneum, 380
 Pneumothorax, 680, **682**
 Podocytes
 glomerular filtration barrier and, **581**, 595
 Poikilocytosis, 407
 Point of service plan, 271
pol gene, 175
 Poliomyelitis, **531**
 restrictive lung disease, 675
 Poliovirus, 531
 immunodeficient patients, 118
 medical importance, 167
 picornavirus, 168
 unvaccinated children, 186
 Polyadenylation signal, 41
 Polyangiitis, microscopic
 autoantibody, 115
 Polyarteritis nodosa, 173, **314**
 Polyarthralgias
 gonococcal arthritis, 468
 rubella, 182
 Polycystic disease (kidney), 604
 Polycystic ovarian syndrome (PCOS), **645**
 anovulation, 645
 antiandrogens, 658
 clomiphene, 656
 Polycythemia, **434**
 blood oxygen in, 666
 Eisenmenger syndrome, 299
 ESR in, 214
 paraneoplastic syndromes, 228
 Polycythemia vera, 433
 Budd-Chiari syndrome and, 392
 hepatocellular carcinoma, 392
 Polydactyly, 63
 Polydipsia, 346
 Polyhydramnios, 491, **641**
 esophageal atresia and, 359
 Polymerase chain reaction (PCR), **52**
 Polymorphic ventricular tachycardia, 294
 Polymyalgia rheumatica, **470**
 ESR in, 214
 giant cell arteritis and, 314
 Polymyositis
 autoantibody, 115
 Polymyositis/dermatomyositis, **471**
 Polymyxin B, 143, **193**, 198
 Polymyxins, 198
 mechanism and use, **193**
 Polyneuritis, 66
 Polyneuropathy, 425
 familial amyloid, 212
 Polyomaviruses
 characteristics of, 164
 DNA viruses, 163
 genome, 163
 naked viruses, 163
 Polyostotic fibrous dysplasia, 57
 Polyposis syndromes, **387**
 Polyps
 adenomatous, 387
 APC gene, 387
 colonic, 387
 hyperplastic, 387
 inflammatory pseudopolyps, 387
 KRAS gene, 387
 mucosal, 387
 nasal (cystic fibrosis), 60
 neoplastic transformation of, 387
 non-neoplastic, 387
 serated, 387
 submucosal, 387
 uterine, 650
 Polysaccharide capsule antigens
 carrier proteins with, 127
 Polyuria
 diabetes insipidus, 338
 diabetes mellitus, 346
 hyperparathyroidism, 345
 lithium, 574
 Pompe disease, 87
 Pons
 cranial nerve nuclei of, 505
 development of, 490
 Pontiac fever, 143
 Pontine syndrome, 514
 "Pope's blessing" (median nerve injury), 451
 Popliteal artery, 455
 atherosclerosis in, 302
 Popliteal fossa, 455
 Popliteus, 453
 Porcelain gallbladder, 396, 397
 Porphobilinogen deaminase, 425
 Porphyrin, 425, 546
 Porphyrin cutanea tarda, 425
 Portal hypertension
 ARPKD, 604
 cirrhosis and, 389
 pulmonary arterial hypertension, 679
Schistosoma spp, 161
 serum markers for, 390
 varices and, 365
 Portal triad, 361, **367**
 Portal vein, 361, **367**
 in fetal circulation, 282
 Portosystemic anastomoses, **365**
 Positive predictive value (PPV), 257, 259
 Positive punishment (aversive stimulus), 554
 Positive reinforcement, 554
 Positive skew distribution, 262
 Posterior cerebral artery, 502, 514
 Posterior circulation strokes, 514
 Posterior circumflex artery, 455
 Posterior communicating artery
 in saccular aneurysm, 516
 Posterior cruciate ligament (PCL)
 injury, 454
 Posterior descending artery (PDA), 283
 Posterior drawer sign, 454
 Posterior fossa
 malformations, **492**
 Posterior hypothalamus, 498
 Posterior inferior cerebellar artery
 stroke effects, 514
 Posterior malleolus, 455
 Posterior pituitary gland, 327
 Posterior tibial artery, 455
 Posterior urethral valves, **579**
 Postherpetic neuralgia, 165
 Postinfectious encephalomyelitis, 524
 Postoperative ileus, 240
 Postpartum hemorrhage, 641
 Postpartum mood disturbances, 562
 Postpartum psychosis, 562
 Postpartum thyroiditis, 341
 Postprandial pain, 363
 Postrenal azotemia, 601
 Poststreptococcal glomerulonephritis (acute), 596
 Post-traumatic stress disorder (PTSD)
 acute stress disorder, 564
 diagnostic criteria/treatment, 564
 dissociative identity disorder, 558
 drug therapy for, 572
 prazosin for, 244
 SSRIs for, 575
 venlafaxine, 575
 Postural hypotension
 midodrine for, 242
 trazodone, 576
 Postviral infections, 179
 Potassium
 amphotericin B, 199
 in cardiac muscle, 292
 diabetic ketoacidosis, 347
 shifts in, 590
 torsades de pointes and, 294
 Potassium channel blockers, **323**
 Potassium channels
 myocardial action potential, 292
 opioid effect, 551
 Potassium chloride, 249
 Potassium iodide
 Sporothrix schenckii, 154
 for thyroid storm, 342
 Potassium-sparing diuretics, **609**
 Potency of drugs
 vs efficacy, 233
 Potentiation
 of drugs, 235
 Pott disease, 180

- Potter sequence (syndrome)
 ARPKD, 604
 oligohydramnios and, **578**, 641
 pulmonary hypoplasia, 660
- Poxvirus, 164, 479
- PPD test
 for tuberculosis, 140
- PPSV23 (pneumococcal polysaccharide vaccine), 127
- PR3-ANCA/c-ANCA autoantibody, 115
- Practice tests, 21
- Prader-Willi syndrome
 chromosome association, 64
 ghrelin in, **336**, 371
- Pramlintide, 249, 353
- Prasugrel, 411, 437
- Praziquantel
 antihelminthic therapy, 200
 tapeworms, 160
 trematodes, 160
- Prazosin, 244
- Precision vs accuracy, **259**
- Precocious puberty
 adrenal steroids and, 335
 leuprolide, 656
 McCune-Albright syndrome, 57
 pinealoma, 528
 types, **637**
- Precontemplation stage, substance addiction, 568
- Predictive value, 257
- Prednisolone
 for thyroid storm, 342
- Preeclampsia, 643
 hydatidiform moles, 642
- Preferred provider organization plan, 271
- Prefrontal cortex, 501
- Pregnancy
 advanced maternal age, 63
 aliskiren contraindication, 610
 amniotic fluid abnormalities, 641
 anemia caused by, 418
 carpal tunnel syndrome in, 459
 choriocarcinomas and, 642
 contraindicated antimicrobials, 204
 ESR in, 214
 estrogen in, 630
 ethical situations, 268–269
 fetal circulation, 282
 fetal respiration, 660
 folate deficiency caused by, 420
 folic acid supplementation, 68
 heparin in, 436
 hypertension and treatment in, 243, 316, 643
 hypothyroidism in, 341
 insulin in, 334
Listeria monocytogenes in, 139
 lithium in, 298, 300
 maternal complications, 272
 maternal phenylketonuria, 84
 opiate use during, 615
 parental consent and, 265
 physiologic adaptations in, **633**
 pituitary infarcts with, 339
 posterior urethral valve diagnosis, 579
 progesterone in, 630
 prolactin and, 330
 pyelonephritis, 600
 pyogenic granulomas and, 478
 sex hormone-binding globulin, 337
 stillbirth, 182
Streptococcus agalactiae in, 137
 syphilis in, 147
 TBG in, 331
 termination of, 657
 TORCH infections, 182
 Turner syndrome and, 638
 twinning in, 616
 urinary tract infections, 181
 venous sinus thrombosis in, 503
 vitamin B₉ deficiency, 68
 Pregnancy complications, **640–641**
- Prehn sign, 654
- Preload in cardiac output, 284
- Premature ejaculation, 575
- Premature labor and delivery
 cryptorchidism and, 651
 low birth weight with, 635
 murmur in prematurity, 291
- Premature ovarian failure, 636, 645
- Premotor cortex, 501
- Preoptic nucleus, 498
- Prepatellar bursitis, 460
- Preprocollagen, 51
- Preproinsulin, 334
- Prerenal azotemia, 601
- Presbycusis, 270, 533
- Presbyopia, **535**
- Preschool age development, 635
- Presenilin, 520
- Pressure-volume loops, **287**
- Presynaptic α_2 -autoreceptor, 239
- Pretecal nuclei, 539
- Preterm birth
 common cause of death, 272
 Pretest probability, 257
- Prevalence
 diagnostic test evaluation, 257
 incidence vs, **259**
 observational studies, 256
- Prevotella* spp, 179
- Priapism, 651
 sickle cell anemia, 422
 trazodone and, 576
- Primaquine, 157
 hemolysis in G6PD deficiency, 250
- Primary adrenal insufficiency, 349
- Primary amyloidosis, 212
- Primary bacterial peritonitis, 390
- Primary biliary cholangitis, 395
- Primary central nervous system lymphoma, 430
- Primary ciliary dyskinesia, 49
- Primary disease prevention, 270
- Primary glomerular disease, 594
- Primary hemostasis, **411**
- Primary hyperaldosteronism, 349
 hypertension with, 300
 markers in, 591
- Primary hyperparathyroidism, 345, 464
- Primary hypertension, 316
- Primary hypogonadism, 639
- Primary ovarian insufficiency, **645**
- Primary polycythemia, 433
- Primary sclerosing cholangitis, 395
- ulcerative colitis, 382
- Primary spontaneous pneumothorax, 682
- Primase
 replication initiation by, 38
- Primidone, 519
- Primitive atrium, 281
- Primitive pulmonary vein, 281
- Primitive reflexes, **510**
- Primitive ventricle, 281
- Pringle maneuver, 361
- PR interval, 293, 295
 antiarrhythmic effects, 323, 324
 prolonged, 295
 shortened, 294
- Prinzmetal angina
 calcium channel blockers for, 318
 ischemic manifestations, 304
 propranolol adverse effects, 323
- Prions, **178**
- Privacy and confidentiality, 267
- Probeneid, 252
 cidofovir with, 202
 gout, 487
- Procainamide, 322
- Procaine, 550
- Procabazine, 251, **441**
- Procedure bias, 260
- Process improvement model, 273
 quality measurement, 273
- Processus vaginalis, 624
- Procoagulation, 413
- Progesterone
 lactation and, 636
 menstrual cycle, 632
 ovulation, 632
 signaling pathways for, 337
 source and function of, **630**
- Progestins, **657**
- Progressive multifocal leukoencephalopathy (PML), 494, **524**
 HIV-positive adults, 177
 polyomaviruses, 164
 rituximab, 443
- Proguanil, 200
- Projection, 555
- Prokaryotes
 DNA replication in, 38
 mRNA start codons, 44
 RNA polymerases in, 42
- Prolactin, **330**
 circadian rhythm, 497
 function of, 328
 lactation and, 636
 secretion of, 327
 signaling pathways for, 337
 tuberoinfundibular pathway, 499
- Prolactin-inhibiting factor, 328
- Prolactinomas
 dopamine agonists for, 330
- Proliferative glomerular disorders, 594
- Prometaphase, 46
- Promoters (gene expression), 41
- Promyelocytic leukemia, 66
- Pronephros, 578
- Proopiomelanocortin, 327
- Propafenone, 322
- Proper hepatic artery, 361
- Prophase, 46
- Prophylaxis
 antimicrobial, 197, 198
 antimycobacterial drugs for, 196
 HIV/AIDS patients, **198**
 in HIV/AIDS patients, 198
Pneumocystis jirovecii, 154
 rabies postexposure, 171
 for RSV, 169
Trichomonas vaginalis, 158
- Propionyl-CoA carboxylase
 metabolic pathways, 74
 vitamin B₇ and, 68
- Propofol, 550
- Propranolol, 245, 323, 342
 essential tremor, 519
- Proprioception
 Friedreich ataxia, 531
- Propylthiouracil
 agranulocytosis, 250
 aplastic anemia, 250
 thionamides, 354
 for thyroid storm, 342
- Prostacyclin analogues, 686
- Prostaglandin analogs, 254
- Prostaglandins
 aspirin effects, 486
 cortisol effect on, 336
 glaucoma treatment, 552
 kidney functions, 589
- Prostate cancer
 adenocarcinomas, **654**
 incidence/mortality of, 222
 leuprolide for, 656
 metastases of, 223
- Prostate gland
 lymphatic drainage of, 624
 with urethral injury, 627
- Prostate-specific antigen (PSA), 227, 654
- Prostatic acid phosphatase (PAP), 654
- Prostatic adenocarcinoma, 654
- Prostatitis, **654**
 gonorrhea, 184
- Prosthetic devices
Staphylococcus epidermidis, 135
- Prosthetic heart valves, 423
- Protamine sulfate, 248, 436
- Protease inhibitors
 fat redistribution, 250
 HIV therapy, 203
 hyperglycemia, 249
 naming convention for, 253
- Proteases, 373
- Proteasome, **48**
- Protein A, 129
- Proteinases, 406
- Protein C/S deficiency, 428
- Protein-energy malnutrition, **71**
- Protein kinase A
 fructose biphosphatase-2 and, 76
- Protein metabolism
 amino acids, 81
- Proteins
 free radical effect on, 210
- Protein synthesis
 elongation, 45
 inhibitors, **191**, 253
 initiation of, 44, **45**
 insulin and, 334
 metabolic site, 72
 posttranslational modification, 45
 sequence of, 45
 termination, 45
 trimming, 45
- Proteinuria
 ACE inhibitors for, 610
 angiotensin II receptor blockers, 610
 diabetes mellitus, 346
 nephritic syndrome, 595
 nephrotic syndrome, 595, 597
 preeclampsia, 643
 renal papillary necrosis and, 602
 serum sickness, 113
- Proteolysis
 cortisol and, 336
- Proteolytic processing, in collagen synthesis, 50

- Proteus* spp
 struvite stones with, 127
 urease-positive, 127
 xanthogranulomatous
 pyelonephritis, 600
- Proteus mirabilis*
 cephalosporins, 189
 penicillins for, 188
 urinary tract infections, 181, 600
- Prothrombin
 complex concentrate transfusion,
 429
 warfarin effect on, 436
- Prothrombin gene mutation, 428
- Prothrombin time, 390
- Protofilament, 48
- Proton pump inhibitors, 254, **399**
 Beers criteria, 247
 for *Helicobacter pylori*, 146
 gastrin and, 371
- Protozoa
 CNS infections, **156**
 GI infections, **155**
 hematologic infections, **157**
 miscellaneous, **158**
 watery diarrhea, 179
- Proximal convoluted tubules
 (PCT)
 in ATN, 602
 defects in, 586
 diuretics and, 609
 dopamine secretion by, 589
 glucose clearance and, 584
 ischemia susceptibility, 210
 relative concentrations in, 587
 renal cell carcinoma and, 605
- Proximal interphalangeal (PIP) joints,
 451
- Proximal renal tubular acidosis
 (type 2), 593
- PRPP (glutamine-phosphoribosyl-
 pyrophosphate)
 amidotransferase, 73
- Pruritus
 anal, 159
 atopic dermatitis, 477
 biliary tract disease, 395
 chloroquine, 200
 cutaneous mycoses, 152
 ectoparasites, 161
 histamine receptors and, 238
 hyperchylomicronemia, 94
 lichen planus, 482
 pseudofolliculitis barbae, 477
 urticaria, 477
- Prussian blue stain, 677
- Psammoma bodies, 211, **227**
 mesotheliomas, 678
 papillary thyroid carcinoma, 343
- Pseudoappendicitis
Yersinia enterocolitica, 144
- Pseudocysts, 397
- Pseudoephedrine, **686**
- Pseudofolliculitis barbae, 477
- Pseudofractures, 463
- Pseudoglandular stage
 (development), 660
- Pseudogout, 467
- Pseudohermaphroditism, 639
- Pseudohypoparathyroidism, 344
- Pseudomembranous colitis
 clindamycin, 192
Clostridium difficile, 138
 drug reaction and, 249
 penicillins, 188
- vancomycin for, 190
 watery diarrhea, 179
- Pseudomembranous pharyngitis
 diphtheria, 139
- Pseudomonas* spp
aeruginosa, **143**
 biofilm production, 128
 ceftazidime, 189
 culture requirements for, 126
 cystic fibrosis, 60, 179
 encapsulated, 127
 epididymitis and orchitis, 654
 exotoxin production, 132
 fluoroquinolones, 195
 immunodeficient patients, 118
 multidrug-resistant, 198
 nosocomial infection, 179, 185
 osteomyelitis, 180
 penicillins for, 188
 pigment production, 128
 pyocyanin of, 109
 tricuspid valve endocarditis, 311
 UTIs, 181
- Pseudo-Pelger-Huet anomaly, 432
- Pseudopseudohypoparathyroidism,
 344
- Pseudotumor cerebri, 521
 drug reactions and, 251
 vitamin A toxicity, 66
- Pseudovirion, 162
- Psittacosis, 149
- Psoriasis, 477
 arthritis and, 469
 cyclosporine, 120
 etanercept for, 487
 hyperkeratosis/parakeratosis, 475
 infliximab/adalimumab for, 487
 methotrexate for, 440
 skin lesions, 475
 therapeutic antibodies, 122
- Psoriatic arthritis, 469
 HLA-B27 and, 100
 leflunomide for, 486
 psoriasis and, 477
- Psychiatric emergencies
 acute dystonia, 569
 delirium tremens, 569
 hypertensive crisis, 569
 lithium toxicity, **569**
 neuroleptic malignant syndrome,
 569
 serotonin syndrome, 569
 tricyclic antidepressant overdose,
 569
- Psychiatry, 554–576
 pathology, 556–570
 pharmacology, 572–576
 psychology, 554–555
- Psychoactive drug intoxication/
 withdrawal, **570–571**
- Psychological child abuse, 556
- Psychology, 554–555
- Psychosis, **559**
 corticosteroids, 120
 diabetic ketoacidosis, 347
 drug therapy for, 573
 LSD and, 571
 major depressive disorder with, 561
 PCP and, 571
 postpartum, 562
- Psychotherapy, **572**
 oppositional defiant disorder, 557
- Psychotic disorders
 readmissions with, 272
- PTEN* gene, 224
- Pterygoid muscles, 507
- PTH-related peptide (PTHrP), 332
- PTHrP (parathyroid hormone-related
 protein), 228
- Ptosis
 CN III damage, 541
 Horner syndrome, 540
 myasthenia gravis, 472
 saccular aneurysm, 516
- Puberty
 GnRH and, 328
 Kallmann syndrome and, 639
 precocious, 57, 335
 Tanner stages, 637
- Pubic tubercle, 370
- Public health sciences, 256–273
- Pudendal nerve, 366, 453
- Pulmonary anthrax, 137
- Pulmonary arterial hypertension
 (PAH), 679
- Pulmonary artery, 619
 fetal circulation, 282
- Pulmonary artery stenosis, 300
- Pulmonary capillary wedge pressure
 (PCWP), 297, 668
- Pulmonary circulation, **668**
- Pulmonary edema, 309
 consolidation in, 680
 LV failure, 307
 mannitol, 607
 nitrates for, 318
 opioids for, 551
 preeclampsia and, 643
- Pulmonary embolism, **672**
 chronic thromboembolism, 679
 direct factor Xa inhibitors for,
 437
 heparin for, 436
 tamoxifen/raloxifene and, 443
 thrombolytics for, 437
 ventilation/perfusion with, 669
- Pulmonary fibrosis
 amiodarone and, 323
 bleomycin, 439
 busulfan, 441
 diffusion in, 668
 drug reaction and, 251
 methotrexate, 440
 restrictive lung disease, 675
- Pulmonary hypertension, **679**
 cor pulmonale, 668
 drug therapy, **686**
Schistosoma, 160
- Pulmonary hypoplasia, 660
- Pulmonary infections
 in immunocompromised patients,
 139
- Pulmonary Langerhans cell
 histiocytosis, 675
- Pulmonary surfactant
 club cells, 660, 661
 NRDS, 661
- Pulmonary vascular resistance (PVR),
668
- Pulmonic stenosis
 wide splitting in, 289
- Pulmonic valves, 281
 “Pulseless disease,” 314
- Pulse pressure
 equation for, 285
- Pulsus paradoxus, 310
 croup, 170
- “Punched out” bone lesions (X-ray),
 431
- Punched-out ulcers, 377
- Punishment, 554
- Pupil
 Argyll Robertson, 147
 CN III palsy, 541
 control, 507, **539**
 light reflex, 539
 pupillary light reflex, 539
 syphilis, 147, 184
- Pure motor stroke, 514
- Pure red cell aplasia, 228
 thymoma and, 98
- Purines, 194
 de novo synthesis, 36, 73
 Lesch-Nyhan syndrome, 37
 salvage deficiencies, **37**
 structure of, 35
- Purkinje cells
 cerebellum, 499
 of cerebellum, 210
 in paraneoplastic cerebellar
 degeneration, 228
- Purkinje fibers, 293
- Purpura
 aplastic anemia, 421
- Pustular psoriasis, 475
- Pustules, 475
 acne, 477
 pseudofolliculitis barbae, 477
 rosacea, 477
- Putamen, 500
 neurodegenerative disorders, 520
- Pyelonephritis, **600**
 kidney stones, 598
 urinary tract infections, 181, 600
 WBC casts in, 594
- Pygmalion effect, 260
- Pyloric sphincter, 373
- Pyloric stenosis, 359
- Pyloromyotomy, 359
- Pyoderma gangrenosum
 inflammatory bowel disease, 382
- Pyogenic granulomas, 478
- Pyramidal cells, 210
- Pyramidalis muscle, 369
- Pyramidal tract demyelination,
 multiple sclerosis, 523
- Pyrantel pamoate, 200
- Pyrazinamide, **197**
 gout, 250
 hepatitis, 249
Mycobacterium tuberculosis, 196
- Pyridostigmine, 240
 myasthenia gravis treatment, 472
- Pyridoxal phosphate, 67
- Pyridoxine, 67
- Pyrimethamine, 200
 effect on purine synthesis, 36
- Pyrimidine dimers, 40
- Pyrimidines
 de novo synthesis, 36
 structure of, 35
- Pyrimidine synthesis, 486
- Pyruvate carboxylase, 77, 78
 metabolic pathways, 74
 vitamin B₇ and, 68
- Pyruvate dehydrogenase
 complex, **76**
 deficiency, **77**
 metabolic pathways, 74
 vitamin B₁ and, 66
- Pyruvate kinase, 74
 deficiency, 422
- Pyruvate metabolism, **77**
- Pyuria, 601

Q

- Q fever
 rickettsial disease, 150
 transmission, 149
- QRS complex, 293
- QT interval
 atypical antipsychotic effect on, 573
 Class IA antiarrhythmic effects, 322
 congenital long QT syndrome, 294
 drug-induced long, 294
 ECG, 293
 ondansetron effect on, 400
 in torsades de pointes, 294
- Quadrantanopia, 542
- Quadriceps, 452
- Quality measurements, **273**
- Quantifying risk, **258**
- Quaternary amines, 204
- Quaternary disease prevention, 270
- Quetiapine, 573
- Quiescent (stable) cells, 46
- Quinidine, 157, 200, 322
 cinchonism, 251
- Quinine, 200
- Quinolone
Legionella pneumophila, 143
- Quinupristin, 198
- R**
- Rabies, **171**
 active and passive immunity, 110
 rabdovirus, 167
 viral receptors, 166
- Rachischisis, 491
- Rachitic rosary, 463
- Radial head subluxation, 461
- Radial nerve, 447
 neurovascular pairing, 455
- Radiation exposure
 acute myelogenous leukemia and, 432
 aplastic anemia, 421
 apoptosis caused by, 208
 as carcinogen, 225
 free radical injury caused by, 210
 hypopituitarism, 339
- Radiation therapy
 acute pericarditis and, 313
 angiosarcomas, 478
 lymphopenia, 424
 neutropenia, 424
 osteosarcoma and, 465
 pancreatic cancer, 398
 papillary thyroid carcinoma risk, 343
 readmissions with, 272
- Radiculopathy
 lumbosacral, 455
- Radon
 as carcinogen, 225
- Ragged red muscle fibers, 59
- Rales, 309
- Raloxifene, 443, 656
- Raltegravir, 203
- Ramelteon, **547**
- Ramipril, 610
- Ranitidine, 399
- RANKL (RANK ligand), 332, 459
- Ranolazine
 mechanism and clinical use, **319**
- Raphe nucleus, 495
- Rapid automated broth cultures, 126
- Rapid-eye movement (REM) sleep, 497
- Rapid filling (cardiac cycle), 287

- Rapidly progressive
 glomerulonephritis (RPGN), 596
- Rasagiline, **549**
- Rasburicase, **444**
- RAS gene, 343
- Rashes
 “blueberry muffin,” 169
 carbapenems, 190
 childhood, 183
 cytomegalovirus, 182
 desquamating, 314
 fluoroquinolones, 195
 heliotrope, 228
 macrolides, 193
 palms and soles, 150
 penicillinase-sensitive penicillins, 188
 rickettsial infections, 150
 rubella, 169, 182
 syphilis, 147, 184
 unvaccinated children, 186
- Rathke pouch, 327, 528
 tumor, 613
- Rationalization, 555
- Raynaud phenomenon, **472**
 Buerger disease, 314
 calcium channel blockers for, 318
 scleroderma and, 473
- Razor bumps, 477
- RBC casts (urine), 594
- RBC inclusions, **416–444**
- RBC morphology (pathologic), **414**
- Rb* gene, 224
- Reabsorption/secretion rate
 calculation, 584
- Reaction formation, 555
- Reactive arthritis, 469
Campylobacter jejuni, 145
 chlamydia, 148, 184
 HLA-B27 and, 100
- Reactive attachment disorder, 556
- Readmission recurrences, 272
- Reassortment (viral), 162
- Recall bias in studies, 260
- Receiving operating characteristic curve, **260**
- Receptor binding, **234**
- Receptor fusion proteins
 naming conventions for, 254
- Receptor-mediated endocytosis, 47
- Receptors (viral), **166**
- Receptor tyrosine kinase
 hormone messenger, 337
 as oncogene product, 224
- Recessive inheritance, 59
- Recklinghausen disease, 525
- Recombinant cytokines, **121**
- Recombination (viral), 162
- Recruiting study participants, 260
- Rectal sparing, 382
- Rectosigmoid junction
 blood supply to, 363
- Rectum
 blood supply and innervation, 364
 familial adenomatous polyposis, 387
 Hirschsprung disease, 384
 ischemia susceptibility, 210
 portosystemic anastomosis, 365
- Rectus abdominis muscle, 369
- Recurrent branch (median nerve), 447
- Recurrent laryngeal nerve
 compression of, 684
 Pancoast tumor, 685

- Red cell casts, 315
- Red-green color blindness, 197
- Red hemorrhagic infarct, 210
- Red hepatization, 683
- Red nucleus (midbrain), 511–552
- Redox reactions
 free radical injury and, 210
 vitamin B₂ and, 67
- Red pulp (spleen), 98
- Red rashes of childhood, **183**
- Reduced filling (cardiac cycle), 287
- Redundant/degenerate genetic code, 37
- Reed-Sternberg cells, 429
- Refeeding syndrome, anorexia
 nervosa, 567
- Referred pain
 cholecystitis, 396
 from diaphragm, 663
- Reflex bradycardia, 588
- Reflexes
 clinical, 510
 cranial nerve, 507
 motor neuron sign, 529
 primitive, 510
- Reflex tachycardia, 244
- Refractive errors (vision), **535**
- Refractory hypertension, 658
- Refsum disease, 47
- Refusing care
 minors, 269
- Regadenoson, 304
- Regan-Lowe medium, 126
- Regional specification (brain), **490**
- Registering for exam, 5–6
- Regression, 555
- Regulation of cell cycle
 Cyclin-dependent kinases (CDKs), 46
 p53, 46
 Tumor suppressors, 46
- Regulation of gene expression, **41**
- Regulatory T cells, **102**
 cell surface proteins, 110
- Regurgitation
 in GERD, 377
- Reheated rice syndrome, 138
- Reichert cartilage, 620
- Reid index, 674
- Reinforcement, 554
- Relapse stage, substance addiction, 568
- Relapsing fever
 animal transmission, 149
 lice, 161
- Relationship with patients, 268
- Relative risk reduction (RRR), 258
- Relative risk (RR), 256, **258**, 263
- Reliability, 259
- Remodeling (tissue), 216
- REM sleep, 497
- Renal agenesis
 bilateral, 578
 unilateral, 579
- Renal artery, 580
 stenosis, 610
- Renal blood flow (RBF), 363, 580
 acute injury and, 602
 NSAID effects on, 589
 renal plasma flow and, 582
- Renal cell carcinoma, **605**
 bevacizumab for, 442
 carcinogens for, 225
 chromosome association, 64
 hypercalcemia and, 228
 immunohistochemical stain for, 227
 metastases of, 223
- recombinant cytokines, 121
 therapeutic antibodies, 122
- Renal clearance, **582**
- Renal cortex
 atrophy of, 599
- Renal disorders/failure
 consequences of, **603**
 diabetes mellitus, 346
 diffuse cortical necrosis, 602
 ESR in, 214
 Fabry disease, 88
 features of, 591
 gout and, 467
 in utero, 578
 markers for, 591
 myoclonus in, 519
 NSAIDs, 589
 preeclampsia and, 643
 renal cyst disorders, **604**
 tetracycline use in, 192
 waxy casts in, 594
 Wilson disease, 395
- Renal/genitourinary drug reactions, 251
- Renal hypoxia, 666
- Renal ischemia, 486
- Renal medulla
 hydronephrosis, 599
- Renal oncocytoma, **605**
- Renal osteodystrophy, 345, **603**
- Renal papillary necrosis, **602**
 pyelonephritis and, 600
 sickle cell anemia, 422
- Renal plasma flow, 582
 glomerular dynamics and, 583
- Renal sympathetic discharge, 588
- Renal toxicity
 ganciclovir, 202
- Renal tubular acidosis (RTA), **593**
- Renal tubular defects, 585, **586**
- Renal vascular smooth muscle, 238
- Renin, 588
 ACE inhibitor effect on, 610
 aliskiren effect on, 610
 in hyperaldosteronism, 349
 renal disorders and, 591
 sympathetic receptors and, 238
- Renin-angiotensin-aldosterone system, 327, 588
- Renovascular disease, **604**
- Renovascular hypertension, 349
- Reoviruses
 characteristics, 167
 genome, 163
 naked viruses, 163
 segmented, 168
- Repaglinide, 353
- Reperfusion injury, 210
- Reperfusion therapy, 307
- Replication fork, 38
- Reportable diseases
 confidentiality exceptions, 267
- Repression, 555
- Repressor proteins
 lactose effects on, 40
- Reproductive/endocrine drug reactions, 249
- Reproductive hormones, **655**
- Reproductive system, 612–653
 anatomy, 624–627
 embryology, 612–623
 pathology, 638–652
 pharmacology, 655–658
 physiology, 629–636
- Reptile (disease vectors), 149
- RER staining, 493

- Rescheduling exam, 6
Reserpine
 Parkinson-like syndrome, 251
Residual volume (RV), 664
Resistance in vessels, 286
Respiration
 exercise response, 670
 high altitude response, 670
 in diabetic acidosis, 347
 Kussmaul, 347
Respiratory acidosis, 592
Respiratory alkalosis, 592
 high altitude, 670
Respiratory burst, **109**
 free radical injury and, 210
Respiratory depression
 anesthetics, 550
 barbiturates, 546, 570
 benzodiazepines, 544, 570
 epilepsy drugs, 544
 inhaled anesthetics, 550
 opioids, 551, 570
 psychoactive drug intoxication, 570
 tricyclic antidepressants, 575
Respiratory drug reactions, 251
Respiratory syncytial virus (RSV)
 paramyxovirus, 167, 169
 pneumonia, 179, 683
 prophylaxis, 122
Respiratory system, 660–683
 anatomy, 662–663
 change in elderly, **665**
 embryology, 660
 pathology, 671–681
 pharmacology, 686–687
 physiology, 664–669
Respiratory tract infections
 C3 deficiency, 107
Respiratory tree, **662**
Respiratory zone, 662
Resting tremor, 519
Restless legs syndrome, 519
Restricting type of anorexia nervosa, 567
Restrictive cardiomyopathy, 308
 hemochromatosis, 395
Restrictive lung diseases, **675**
 ankylosing spondylitis, 469
 flow volume loops, 673
 sarcoidosis, 676
Reteplase (rPA), 437
Rete testis, 652
RET gene, 224
 carcinoma risks with, 343
 Hirschsprung disease, 384
Reticular activating system, 511
Reticular fibrous framework (spleen), 98
Reticulate bodies, 148
Reticulin, 50
Reticulocyte index (RI), **417**
Reticulocytes, 407
 in aplastic anemia, 421
 intravascular hemolysis, 421
Retina
 chronic hyperglycemia, 537
 embryologic derivation of, 613
Retinal hemorrhage
 hypertensive emergency, 300
Retinal pathology
 degeneration, 536
 detachment, **537**
 hemorrhage, 537
 retinitis, 536
 vein occlusion, **537**
 visual field defects, 542
Retinal vein occlusion, 537
Retinitis
 AIDS, 165
 cidofovir, 202
 foscarnet, 202
Retinitis pigmentosa, **538**
Retinoblastoma
 chromosome association, 64
 heterozygosity loss, 56
Retinoblastomas
 osteosarcomas, 465
Retinoids, 477
Retinopathy
 Alport syndrome, 596
 chloroquine, 200
 diabetes mellitus, 346
 hypertension, 300
 of prematurity, 210, 661
 sorbitol, 81
Retrograde amnesia, 558
Retroperitoneal fibrosis, 599
Retroperitoneal structures, **360**
Retrospective studies, 260
Retroviruses
 characteristics, 167
 genomes, 163
Rett syndrome, 61, **62**
Reverse transcriptase, 175
 Telomerase, 38
Reverse transcriptase polymerase
 chain reaction, **52**
Reye syndrome, **390**
Reynolds pentad, 397
Rhabdomyolysis
 daptomycin, 195
Rhabdomyomas, 316
 nomenclature for, 220
Rhabdomyosarcomas
 dactinomycin for, 439
 nomenclature for, 220
 variant, 644
Rhabdoviruses
 characteristics, 167
 negative-stranded, 168
Rhagades, 147
Rh blood classification, 405
 newborn hemolysis, 405
Rheumatic fever, **312**
 chorea with, 519
 heart murmur with, 291
 myocarditis with, 313
 Streptococcus pyogenes, 136
 streptolysin O, 133
 type II hypersensitivity, 112
Rheumatoid arthritis, 457, 466
 autoantibody, 115
 azathioprine for, 440
 carpal tunnel syndrome and, 459
 celecoxib for, 486
 etanercept for, 487
 HLA-DR4 and, 100
 immunosuppressants, 120
 infliximab/adalimumab for, 487
 leflunomide for, 486
 methotrexate for, 440
 rituximab for, 443
Rheumatoid factor, 115
Rhinitis
 phenylephrine for, 242
Rhinophyma, 477
Rhinosinusitis, **671**
Rhinovirus
 picornavirus, 167, **168**
 receptors for, 166
Rhizopus spp, 153
Ribavirin, 204
 contraindicated in pregnancy, 204
 purine synthesis, 36
Riboflavin, 67
Ribose, 79
Ribosomes, 46
Rice-water diarrhea
 organisms causing, 179
 Vibrio cholerae, 146
Richter transformation, 432
Rickets, 463
 hypophosphatemic, 591, 593
 lab values in, 464
 vitamin D and, 70
Rickettsia
 Gram stain, 125
Rickettsial diseases, **150**
Rickettsia prowazekii, 150
 transmission of, 149, 161
Rickettsia rickettsii, 150
 animal transmission, 149
 chloramphenicol, 192
Rickettsia spp
 intracellular organism, 127
 tetracyclines, 192
Rickettsia typhi, 149, 150
Riedel thyroiditis, 341
Rifabutin, 196
Rifampin, 196
 cytochrome P-450 and, 252
 Hansen disease, 141
 hepatitis, 249
 Mycobacterium leprae, 196
 Mycobacterium tuberculosis, 196
 as prophylaxis, 198
 prophylaxis with *Haemophilus influenzae* for contacts, 142
 protease inhibitors and, 203
 RNA polymerase inhibition, 42
Rifamycins, **196**
Rifaximin, 391
Rift Valley fever, 167
Right anterior cardinal vein, 281
Right bundle branch, 293
Right bundle branch block, 289
Right common cardinal vein, 281
Right coronary artery (RCA)
 coronary circulation, 283
 occlusions of, 305
Right heart failure, 309
Right horn of sinus venosus, 281
Right lower quadrant (RLQ) pain, 384
Right marginal artery, 283
Right-to-left shunts, 298
Right upper quadrant (RUQ) pain, 397
Right ventricular hypertrophy (RVH)
 high altitude, 670
 pulmonary hypertension, 679
Riluzole, 549
Ringed sideroblasts, 416
ring-enhancing lesions (MRI)
 Toxoplasma gondii, 156
Ringworm
 griseofulvin, 200
 tinea corporis, 152
Risedronate, 486
Risk assessment, 258
Risk quantification, 258
Risperidone, 573
Ristocetin, 411
risus sardonius
 tetanospasmin, 138
Ritonavir
 HIV therapy, 203
Rituximab, 122, **443**
Rivaroxaban, 437
Rivastigmine, 240, 549
RNA
 capping, 41
 interference, 56
 processing (eukaryotes), **41**
RNA polymerases
 types and functions of, **42**
RNA viruses, **167**
 genome, **163**
Robertsonian translocation, **64**
Rocker-bottom feet, 63
“Rocket tails,” 139
Rocky Mountain spotted fever, 150
 animal transmission, 149
 chloramphenicol, 192
Romaña sign, 158
Romano-Ward syndrome, 294
Romberg sign, 147, 530
Romiplostim (TPO analog), 121
Root cause analysis, 274
Rooting reflex, 510
Ropivacaine, 550
Rosacea, 477
Rose gardener’s disease, 154
Rosenthal fibers, 528
Roseola
 HHV-6/HHV-7, 165
 rash, 183
Rosiglitazone, 353
Rotator cuff muscles, **446**
Rotavirus, **168**
 diarrhea, 167
Rotenone, 78
Roth spots, 311
Rotor syndrome, 393, 394
Rough endoplasmic reticulum, **46**
Rouleaux formation, 431
Round ligament of uterus, 625
Rovsing sign, 383
Rubella, **169**
 cardiac defect association, 300
 cataracts, 535
 heart murmur with, 291
 rash, 183
 TORCH infection, 182
 unvaccinated children, 186
Rubeola (measles) virus, 170
Ruffini corpuscles, 494
Ryanodine receptor, 456
RYR1 gene, 550
S
S-100, 227
Saber shins
 congenital syphilis, 147
 syphilis, 182
Sabin poliovirus vaccine, 167
Sabouraud agar, 126
Saccular aneurysms, 516
 Ehlers-Danlos syndrome, 51
 renal cyst disorders and, 604
Saccular (development stage), 660
Sacrococcygeal teratomas, 652
Sacubitril
 mechanism and clinical use, **319**
Saddle embolus, 672
Saddle nose
 syphilis, 182
Safety culture, **273**
Salicylates
 toxicity treatment for, 248
 as weak acids, 233
Salivary gland tumors, **376**
Salivary stimulation, 240
Salmeterol, 242, 687
Salmonella, 118

- Salmonella* spp
 animal transmission, 149
 bloody diarrhea, 179
 encapsulated bacteria, 127
 food poisoning, 178
 intracellular organism, 127
Shigella spp vs, **144**
 osteomyelitis, 180
 penicillins for, 188
 reactive arthritis, 469
 TMP-SMX for, 194
- Salmonella typhi*, 144
- Salpingitis
 ectopic pregnancy and, 641
- Sampling bias, 260
- “sand”(orange) in diaper, 37
- Sandflies (disease vectors), 158
- Sandfly fever, 167
- SA node, 292
- Saponification, 209
- Saprophyticus*
 urease-positive, 127
- Saquinavir, 203
- Sarcoidosis
 erythema nodosum, 482
 restrictive lung disease, **676**
- Sarcoma botryoides, 644
- Sarcomas
 metastases of, 223
 methotrexate for, 440
 nomenclature of, 220
- Sarcoplasmic reticulum, 456
- Sargramostim (GM-CSF), 121
- SARS (sudden acute respiratory syndrome), 167
- Satiety/hunger regulation, 498
- Saturday night palsy, 447
- “Saw-tooth” crypt pattern, 387
- Saxagliptin, 353
- SBLA cancer syndrome, 224
- Scabies, 161, 200
- Scalded skin syndrome
Staphylococcus aureus, 135
 toxic shock syndrome toxin, 133
- Scales (skin), 475
 basal cell carcinoma, 484
 pityriasis rosea, 482
 psoriasis, 477
 seborrheic dermatitis, 476
- Scaphoid bone, 449
- Scar formation, **218**
- Scarlet fever
 presentation, 136
 rash with, 183
Streptococcus pyogenes, 136
- S cells, 371
- Schatzki rings, 377
- Schaumann bodies, 676
- Schilling test, 420
- Schistocytes, 414
 in extrinsic hemolytic anemias, 423
 HELLP syndrome, 643
 in intravascular hemolysis, 421
 in microangiopathic anemia, 423
- Schistosoma haematobium*
 bladder cancer, 226
 disease association, 161
 squamous cell carcinoma of bladder, 606
- Schistosoma* spp, 160, 161
- Schistosomiasis
 portal hypertension, 389
 pulmonary arterial hypertension, 679
- Schizoaffective disorder, 560
- Schizoid personality disorder, 565
- Schizophrenia, 572
 atypical antipsychotics for, 573
 diagnostic criteria, 560–576
 neurotransmitters for, 495
 readmissions with, 272
- Schizophrenia spectrum disorders, **560**
- Schizophreniform disorder, 560
- Schizotypal personality, 560, 565
- Schüffner stippling
 in blood smear, 157
- Schwann cells, **494**
 Guillain-Barré syndrome, 524
 origin of, 490
- Schwannomas, 494, 527
- Sciatic nerve, 452
- SCID (severe combined immunodeficiency), 37, 117
 adenosine deaminase deficiency, 37
 lymphopenia caused by, 424
 thymic shadow in, 98
- Sclerae
 alkaptonuria, 84
 osteogenesis imperfecta, 51
- Scleritis, 466
- Sclerodactyly, 473
- Scleroderma, **473**
- Scleroderma (diffuse)
 autoantibody, 115
- Sclerodermal esophageal dysmotility, 377
- Sclerosing adenosis, 649
- Sclerosing cholangitis, 393, 395
 ulcerative colitis association, 382
- Scombroid poisoning, 247
- Scopolamine, 241
- Scoring of USMLE Step 1 exam, 7, 8–9
- Scorpion sting, 397
- Scotoma, 542
- Scrotal hematoma, 627
- Scrotum, 627
 lymphatic drainage of, 624
 masses in, **652**
- Scurvy
 collagen synthesis and, 50
 vitamin C deficiency, 69
- Seafood toxins, 247
- Seal-like barking cough, 170
- Seborrheic dermatitis, **476**
- Sebum, 477
- Secobarbital, 546
- Secondary adrenal insufficiency, 349
- Secondary amyloidosis, 212
- Secondary biliary cholangitis, 395
- Secondary disease prevention, 270
- Secondary glomerular disease, 594
- Secondary hyperaldosteronism, 349
- Secondary hyperparathyroidism, 345, 464
- Secondary polycythemia, 433
- Secondary spontaneous pneumothorax, 682
- Second-degree AV block, 295
- Second messengers
 G-protein linked, 238
- Second-wind phenomenon, 87
- Secretin
 regulatory substances, 371
 secretory cell location, 373
 somatostatinomas and, 351
- Secretion rate calculation, 584
- Secretory (exported) protein synthesis, 46
- Segmented viruses, **168**
- Seizures
 anti-NMDA receptor encephalitis, 228
 barbiturates for, 546
 benzodiazepine withdrawal, 546, 570
 β -blockers, 245
 brain injury with recurring, **517**
 bupropion, 576
 clozapine use and, 573
 cytomegalovirus, 182
 drug reaction and, 251
 with eclampsia, 643
 electrolyte disturbances, 591
 enflurane, 251
 foscarnet, 202
 during heat stroke, 517
 herpesviruses and, 164, 165, 183
 high fever, 165
 imipenem/cilastatin, 251
 isoniazid, 197
 medium-chain acyl-CoA dehydrogenase deficiency, 89
 meropenem, 190
 nitrosourea toxicity, 441
 parasite infestation and, 161
 PCP, 571
 phenylketonuria, 84
 psychoactive drug intoxication/withdrawal, 570–571
- Rett syndrome, 62
- Sturge-Weber syndrome, 525
- Taenia solium*, 160, 161
- tramadol and, 552
- types of, 517
- venous sinus thrombosis, 503
- visceral larva migrans, 159
- vitamin B₆ deficiency, 67
- Zellweger syndrome, 47
- Selection bias, 260
- Selective estrogen receptor modulators (SERMs), 443, 462, **656**
- Selective IgA deficiency, 116
- Selective media, 126
- Selective mutism, 557
- Selectivity
 β -blockers, 245
 Selegiline, 548, **549**, 575
- Selenium sulfide
 tinea versicolor, 152
- Self-fulfilling prophecies, 260
- Self-image of patient, 268
- Self-mutilation
 Lesch-Nyhan syndrome, 37
- Semimembranosus, 451, 452
- Seminal vesicles, 622
- Seminiferous tubules, 626, **628**, 629
- Seminoma, 653
- Semitendinosus, 451, 452
- Sensitivity (diagnostic tests), 257
- Sensorineural hearing loss, 533
- Sensory cortex, 514
 topographic representation, 502
- Sensory innervation
 derivation of, 620
 lower extremity, 452, 453
 tongue, 493
 upper extremity nerve injury, 447
- Sensory loss
 conversion disorder and, 566
 stroke effects, 514
- Sensory modalities/pathways
 receptors for, **494**
 spinal tracts in, 509
 thalamus in, 498
- Separation anxiety disorder, 557
- Separation anxiety (infants), 635
- Sepsis
 ARDS, 678
 immunodeficient patients, 118
 lymphopenia with, 424
 neutropenia with, 424
 shock with, 310
Streptococcus agalactiae, 137
- Septate uterus, 623
- Septation of heart chambers, 280
- Septic arthritis, **468**
 gonococci, 142
Staphylococcus aureus, 135
- Septicemia
Listeria monocytogenes, 139
 readmissions with, 272
 Waterhouse-Friderichsen syndrome, 349
- Septic shock
 diffuse cortical necrosis (renal), 602
 macrophages and, 407
 norepinephrine for, 242
- Septum primum, 280
- Septum secundum, 280
- Sequence (morphogenesis error), 613
- Serine, 224
- Serologic markers
 hepatitis, 174
- Seronegative spondyloarthritis, **469**
- Serosa, 362
- Serotonergic drugs, 574
- Serotonin
 changes with disease, 495
 vitamin B₆ and, 67
- Serotonin syndrome, 400, 547, 552, 569
 atypical antidepressants, 576
 dextromethorphan, 686
 MAOIs, 575
 MDMA, 571
 oxazolidinones, 193
- Serous cystadenocarcinoma, 227, 646
- Serous cystadenoma, 646
- Serpentine cord, 140
- Serrated colon polyps, 387
- Serratia marcescens*, 128
 treatment of, 189
 in immunodeficiency, 128
 UTIs, 181
- Serratia* spp
 immunodeficient patients, 118
- Serratus anterior muscle, 448
- Sertoli cells
 secretions of, 622, 628
 sexual determination, 622
 tumors of, 653
- Sertoli-Leydig cell tumor, 647
- Sertraline, 575
- Serum amyloid A, 213
- Serum markers (liver pathology), **390**
- Serum osmolarity
 antidiuretic hormone regulation of, 329
- Serum protein electrophoresis (SPEP)
 in plasma cell dyscrasias, 431
- Serum tumor markers, **226**
- Sevelamer, **355**
 17 α -hydroxylase, 335
 17-hydroxyprogesterone, 335
- Sevoflurane, 550
- Sex chromosome disorders, **638**
- Sex cord stromal tumors, 647
- Sex development disorders
 phenotypic and gonadal disagreement in, 639
 physical characteristics, 639

- Sex hormone-binding globulin (SHBG), 337
- Sex hormone disorder diagnosis, 639
- Sex pilus (bacterial genetics), 130
- Sex steroid replacement, 339
- Sexual abuse, 556, 558
- Sexual behavior
hypothalamus regulation of, 498
- Sexual differentiation, **622**, 636
- Sexual dysfunction
β-blockers and, 245, 323
cimetidine, 399
differential diagnosis of, **567**
Lambert-Eaton myasthenic syndrome, 472
Peyronie disease and, 651
tuberoinfundibular pathway, 499
- Sexually transmitted infections (STIs)
clinical features, **184**
parental consent with, 265
- Sézary syndrome, 430
- Shawl and face rash, 471
- Sheehan syndrome, 339
- Sheep (disease vectors), 160
- Shiga-like toxin (SLT), 145
cytokine release and, 132
- Shiga toxin, 130, 132, 144
- Shigella boydii*, 144
- Shigella dysenteriae*, 144
- Shigella flexneri*, 144
- Shigella sonnei*, 144
- Shigella* spp
bloody diarrhea, 179
penicillinase-sensitive penicillins for, 188
reactive arthritis, 469
TMP-SMX, 194
vs *Salmonella* spp, 144
- Shingles, 165
- Shin splints, 461
- Shock, **310**
dopamine for, 242
Ebola, 171
endotoxins, 131
in pulmonary anthrax, 137
norepinephrine for, 242
Waterhouse-Friderichsen syndrome and, 349
- short-chain fatty acids
in anaerobic organisms, 127
- SIADH, 338
drug raction and, 249
markers in, 591
paraneoplastic syndrome, 228
- Sialadenitis, 376
- Sialolithiasis, **376**
- Sialyl Lewis^x, 215
- Sibling studies, 256
- Sickle cell disease, **422**, 651
ESR in, 214
osteonecrosis and, 463
postsplenectomy state in, 98
Streptococcus pneumoniae, 136
- Sickle cells, 415
- Sideroblastic anemia
causes and treatment, 419
lead poisoning, 419
vitamin B₆ deficiency, 67
- Sigmoid colon, 383
- Sigmoid sinus, 503
- Sigmoid volvulus, 386
- Signaling pathways
endocrine hormones, **337**
steroid hormones, **337**
- Signal recognition particle (SRP), 47
- Signet ring cells, 379
- Sign of Leser-Trélat, 228
- Sildenafil, 651
- Silencer (gene expression), 41
- Silicosis, 677
- Silver stain, 125, 143
- Simeprevir, 204
- Simple partial seizures, 517
- Simple pneumothorax, 680
- Simple renal cysts, 604
- Single nucleotide (point) mutation, 39
- Single nucleotide polymorphisms (SNPs), 54
- Single nucleotide substitutions, **39**
- Single-stranded binding proteins, 38
- Sinusitis
brain abscesses, 180
C3 deficiency and, 107
Kartagener syndrome, 49
Streptococcus pneumoniae, 136
Wegener granulomatosis, 315
- Sinus venosus, 281
- Sirolimus
immunosuppressant, 120
- Sister Mary Joseph nodules, 379
- Sitagliptin, 353
- Situs inversus, 49
- 6-mercaptopurine, 440
azathioprine, 120
for ulcerative colitis, 382
purine synthesis, 36
targets of, 438
- Sjögren syndrome, **468**
autoantibody, 115
pilocarpine for, 240
rheumatoid arthritis, 466
- Skeletal muscles
ACh receptors in, 236
blood flow regulation in, 297
glycogen metabolism in, 86
somatic nerve, supply of, 236
- Skewed distributions, 262
- Skin
blood flow regulation in, 297
collagen in, 50
normal flora of, 135
pigmentation, 56
wrinkles of aging, 52
- Skin anatomy, 473–483
layers of, **473**
microscopic terms, 475
morphology, 475
- Skin cancer, **484**
albinism and, 476
Lynch syndrome and, 388
- Skin drug reactions, 250
- Skin flora, 178
- Skin lesions
autoimmune disorders, 480
bulla, 475
burns, 483
café-au-lait spots, 57
cancer, 222
common disorders, **477**
crust, 475
dermatitis herpetiformis, 381
erythema multiforme, 151
Gottron papules, 228
hyperlipidemia signs, 300, **301**
hyperpigmentation, 395
inflammatory bowel disease, 382
Kaposi sarcoma, 165
kwashiorkor, 71
macule, 475
papule, 475
patch, 475
petechiae, 407
pigmentation disorders, 475, **476**
plaque, 475
pustule, 475
scale, 475
scaling, 152
scaly, 66
seborrheic keratoses, 228
splinter hemorrhages, 311
striae, 348
T-cell lymphoma, 430
telangiectasia, 316, 473
ulcers, 158
vascular tumors, 478
vasculitides, 314
verrucous, 151
vesicle, 475
wheal, 475
- Skip lesions, 382
- Skull thickening, 463
- Slapped cheek rash, 183
- Sleep
ghrelin/leptin production, 336
- Sleep disturbance
apnea, **679**
benzodiazepines and, 570
β-blockers, 245
delirium and, 558
generalized anxiety disorder, 563
in geriatric patients, 270
hypnagogic hallucinations, 559
hypnopompic, 559
paralysis, 568
paroxysmal nocturnal dyspnea, 309
pulsus paradoxus in, 310
sleep terror disorder, 567, **568**
varenicline, 576
- Sleep physiology, **497**
stages in, 497
- Sleepwalking
sleep stages and, 497
- SLE (systemic lupus erythematosus), **470**
antiphospholipid syndrome and, 470
autoantibodies, 115
DPGN, 596
HLA subtypes, 100
kidney disease with, 596
Raynaud phenomenon, 472
- Sliding hiatal hernia, 370
- Slime (S) layer, 124
- Slipped capital femoral epiphysis, 461
osteonecrosis, 463
- Slow acetylators, 232
- Small bowel disease, 374
- Small cell carcinoma of lung, **684**
carcinogens for, 225
immunohistochemical stains for, 227
oat cell carcinoma, 684
paraneoplastic syndromes, 228
Lambert-Eaton myasthenic syndrome, 472
topotecan for, 442
- Small interfering RNA (siRNA) **56**
- Small intestine, 371
- Small lymphocytic lymphoma (SLL), 432
- Small molecule inhibitors
naming conventions for, 254
- Smallpox, 164
- Small-vessel vasculitis
presentation and pathology, 314
- Smoking
abdominal aortic aneurysms and, 302
atherosclerosis and, 302
Buerger disease and, 314
bupropion for cessation, 576
carcinogenicity of, 225
cataracts, 535
colorectal cancer and, 388
emphysema, 674
esophageal cancer and, 378
head and neck cancer, 671
hormonal contraception, 657
Legionnaires' disease, 143
lung cancer, 684
pancreatic cancer and, 398
renal cell carcinoma, 605
saccular aneurysms, 516
squamous cell carcinoma of bladder, 606
stomach cancer and, 379
teratogenic effects, 614
transitional cell carcinoma, 606
varenicline for cessation, 576
- Smooth brain, 491
- Smooth endoplasmic reticulum, **46**
- Smooth muscle
BMPR2 gene, 679
contraction of, 457
glomus tumors, 478
respiratory tree, 662
tumor nomenclature in, 220
- Smooth muscle (vascular)
in arteriosclerosis, 301
atherosclerosis and, 302
calcium channel blocker action, 318
- Smudge cells, 432
- SNARE proteins
in neurotransmission, 138
- SNC (substantia nigra pars compacta), 495
- SNRIs (serotonin-norepinephrine reuptake inhibitors)
clinical use, 572
major depressive disorder, 561
mechanism and clinical use, **575**
- Snuffles, 147
- “Soap bubble” appearance/lesions
giant cell tumor, 464
Cryptococcus neoformans, 153
- Social anxiety disorder, 563
drug therapy for, 572
SSRIs for, 575
venlafaxine for, 575
- Social engagement
infant deprivation effects, 556
- Sodium channel blockers, **322**
- Sodium channels
cystic fibrosis, 60
epilepsy drug effects, 544
local anesthetic effects, 550
pacemaker action potential and, 292
permethrin, 200
- Sodium-glucose co-transporters (SGLT), 334, 353, 373, 584
- Sodium oxybate (GHB)
narcolepsy treatment, 568
- Sodium-potassium channels, 236
- Sodium-potassium pump, **49**
- Sodium stibogluconate, 158, **200**
- Sofosbuvir, 204
- Solifenacin, 241
- Solitary functioning kidney, 579
- Solitary nucleus of medulla, 296

- Somatic hypermutation, 101
 Somatic mosaicism
 Sturge-Weber syndrome, 525
 Somatic mosaicism, 57
 Somatic nerves
 male sexual response, 627
 Somatic symptom disorder, **566**
 Somatic symptoms
 characteristics of, 566
 conversion disorder, 566
 illness anxiety disorder, 566
 somatic symptom disorder, 566
 Somatomedin, 329
 Somatosensory cortex (primary), 501
 thalamic relays to, 498
 Somatostatin
 function of, 328
 glucagon and, 333
 hypothalamic/pituitary drugs, 354
 production of, 329
 regulatory substances, 371
 secretory cell locations, 373
 Somatostatinoma, 351
 pancreatic cell tumor, 351
 Somatotropin, 329
 Sorbitol metabolism, **81**
 Sotalol, 323
 Southern blot, 53
 Southwestern blot, 53
 Space of Disse, 367
 Spaghetti and meatballs appearance, 152
 Spasmolytics, **551**
 Spastic paralysis
 tetanospasmin, 138
 Spastic paresis, 529
 Special senses
 ophthalmology, 534–543
 otology, 533–534
 Specificity equation, 257
 Specific learning disorder, 557
 Spermatic cord, 369
 Spermatocele, 652
 Spermatozoa, 628
 Spermatogenesis, 628
 cryptorchidism and, 651
 process of, **629**
 prolactin effect on, 330
 Spermatogonia, 628
 Spermiogenesis, 629
 Sphenomandibular ligament, 620
 Sphenoparietal sinus, 503
 Spherocytosis, 415
 autoimmune hemolytic anemia, 415
 extrinsic hemolytic anemia, 423
 hereditary, 421, 422
 Sphincter of Oddi, **368**, 371
 Sphingolipidoses, 88
 Sphingomyelin, 88
 Sphingomyelinase, 88
 Spina bifida
 neural tube defect, 491
 Spinal cord
 embryologic derivation, 613
 lesions of, **530**
 lower extent of, **507**
 nerve nuclei of, 505
 nerves of, **507**
 tracts of, **508**, 509
 Spinal cord syndromes
 multiple sclerosis, 523
 Spinal muscular atrophy, 530
 splicing of pre-mRNA in, 42
 Spinothalamic tract, 509
 thalamic relay for, 498
 Spirochetes, **146**
 Spirinolactone, **609**, 658
 for heart failure, 309
 Splay (glucose clearance), 584
 Spleen
 bacterial clearance by, 127
 blood supply and innervation of, 364
 embryology, 360
 in gastrointestinal anatomy, 361
 ischemia susceptibility, 210
 structure and function, **98**
 thrombocytes in, 407
 Splenectomy, 422
 peripheral blood smear after, 423
 Splenic artery, 364
 Splenic flexure
 blood supply to, 363
 Splenomegaly
 anemia, 157
 cirrhosis, 389
 hairy cell leukemia, 432
 hereditary spherocytosis, 422
 malaria, 157
 myelofibrosis, 433
 rheumatoid arthritis, 466
 visceral leishmaniasis, 158
 Splenorenal ligament, 361
 Splicing of pre-mRNA, **42**
 alternative splicing, 43
 Splinter hemorrhages, 311
 Splitting, 555
 Splitting of heart sounds, 288, 289
 Spondyloarthritis (seronegative), 469
 Spongiosis, 475
 Spontaneous abortion
 antiphospholipid syndrome, 470
 Listeria monocytogenes, 139
 syphilis, 182
 vitamin A excess, 614
 Spontaneous bacterial peritonitis, 389, **390**
 Spontaneous pneumothorax, 682
 Sporadic porphyria cutanea tarda, 173
 Spore-forming bacteria, **129**, 138
 Spores, 124
Sporothrix schenckii, **154**
 Sporotrichosis, 154
 Sprain (ankle), 455
 Sprue
 fat-soluble vitamin deficiencies and, 65
 vitamin B₁₂ deficiency, 69
 “Spur cells,” 414
 Sputum
 currant jelly, 145
 Klebsiella spp, 186
 Streptococcus pneumoniae, 136
 rusty, 136
 Squalene epoxidase, 199
 Squamous cell carcinoma, 484
 anus and cervix, 177
 bladder, 160, **606**
 carcinogens in, 225
 cervix, 645
 esophagus, 378
 head and neck, 671
 hypercalcemia and, 228
 lungs, 684
 pectinate line and, 366
 penis, 651
 of skin, 482
 Squamous epithelium, 662
 SRY gene, 622
 SSRIs (selective serotonin reuptake inhibitors), **575**
 anxiety disorders, 562
 atypical depression, 561
 clinical use, 572
 major depressive disorder, 561
 mechanism and clinical use, 575
 obsessive-compulsive disorder, 563
 panic disorder, 563
 phobias, 563
 postpartum depression, 562
 SIADH caused by, 249
 Stable angina, 304
 Stable (quiescent) cells, 46
 Stab wounds and winged scapula, 448
 Staghorn calculi, 598
 Stains (bacterial), **125**
 Standard deviation
 dispersion, 262
 variability, 259
 Standard error of the mean, 262
 Stapedial artery, 619
 Stapedius muscle, 620
 Stapes (ossicles), 533, 620
 Staphylococcal scalded skin syndrome, 479
 Staphylococcal toxic shock syndrome (TSS), 135
Staphylococcus aureus, **135**
 bacterial endocarditis, 311
 β-hemolytic nature of, 135
 brain abscesses, 180
 cephalosporins, 189
 cystic fibrosis, **60**, 179
 dapsons, 195
 exotoxin production, 133
 food poisoning, 178
 immunocompromised patients, 179
 influenza, 169
 IV drug use, 179
 nosocomial infection, 179, **185**
 osteomyelitis and, 180
 penicillins for, 188
 pigment production, 128
 pneumonia, 683
 postviral infection, 179
 prophylaxis for, 198
 septic arthritis, 468
 skin infections, 479
Staphylococcus epidermidis, **135**
 gram-positive testing, 134
 in vivo biofilm production, 128
 normal flora, 178
 nosocomial infection, 185
 osteomyelitis, 180
 urease-positive, 127
 vancomycin for, 190
Staphylococcus gallolyticus, 137
Staphylococcus pyogenes
 skin infections, 479
Staphylococcus saprophyticus, **136**
 gram-positive testing, 134
 urinary tract infections, 600
 UTIs, 181
Staphylococcus spp
 antibiotic tests for, 134
 facultative anaerobic metabolism, 127
 Starling curves, **285**, 297
 Starling forces, 297
 “Starry sky” appearance of B cells, 430
 Start and stop codons, **44**
 Startle myoclonus, 521
 Starvation phases, 91
 Statins
 for acute coronary syndromes, 307
 hepatitis, 249
 myopathy, 250
 Statistical distribution, **262**
 Statistical hypotheses, **262**
 confidence interval, 263
 correct result, 263
 incorrect results, 263
 test for, 264
 Status epilepticus, 517
 treatment, 544
 Stavudine, 203
 Steady state, 231
 Steatohepatitis, 389
 Steatorrhea
 chronic pancreatitis, 397
 cystic fibrosis, 60
 malabsorption syndromes and, 381
 octreotide effect, 400
 Steatosis (hepatic), 390, **391**
 Steeple sign (x-ray), 170
 Stellate cells, 367
 Stellate ganglion, 685
 Stem cells
 in aplastic anemia, 421
 bone marrow, 108
 CD34 protein, 110
 myelodysplastic syndromes and, 431, 432
 Steppage gait, 453
 Sterilization/disinfection methods, 204
 Steroid hormone signaling pathways, 337
 Steroids
 acute pancreatitis, 397
 adrenal insufficiency, 349
 berylliosis, 677
 CRH levels in, 328
 multiple sclerosis, 523
 synthesis of, 46, 72
 TBC and, 331
 Stevens-Johnson syndrome, 194, **481**, 544
 atypical variant of, 150
 drug reaction and, 250
 sulfa drug allergies, 252
 Stimulants
 for ADHD, 557
 intoxication and withdrawal, 570
 laxative, 401
 St. John’s wort, 252
 St. Louis encephalitis, 167
 Stomach
 basal electric rhythm, 362
 blood supply to, 364
 cholecystokinin effect on, 371
 in gastrointestinal anatomy, 361
 histology of, 362
 regulatory substances, 371
 sclerosis of, 473
 secretin effect on, 371
 Stone bone, 463
 Straight sinus, 503
 Stranger anxiety (infants), 635
 Strategies
 clinical vignette, 23
 test-taking, 22–23
 Strawberry cervix, **181**, 184
 Strawberry hemangiomas, 478
 Strawberry tongue, 136
 Kawasaki, 314
 scarlet fever, 136
 Streak gonads, 622

- Streptococcus agalactiae* (group B strep), **137**
 β-hemolytic nature of, 135
 encapsulated bacteria, 127
 gram-positive testing, 134
 in neonates, 182
 prophylaxis for, 198
- Streptococcus bovis*, **137**
- Streptococcus mutans*
 biofilm production, 128
 normal flora, 178
- Streptococcus pneumoniae*, **136**
 chloramphenicol, 192
 cystic fibrosis, 179
 encapsulated bacteria, 127
 gram-positive testing, 134
 IgA protease and, 129
 influenza, 169
 IV drug use and, 179
 meningitis, 179, 180
 penicillin G/V for, 187
 pneumonia, 179, **683**
 postviral infection, 179
 transformation in, 130
 α-hemolysis, 135
- Streptococcus pyogenes* (group A strep), 130, 133, **136**
 β-hemolysis, 135
 lab testing, 134
 in renal disease, 596
 M protein and, 129
 rash, 183
 treatment of, 187, 192
- Streptococcus pyogenes* toxic shock–like syndrome
 skin infection with, 135
- Streptococcus sanguinis*, 128
- Streptococcus* spp
 antibiotic tests for, 134
 facultative anaerobic metabolism, 127
 septic arthritis, 468
- Streptogramins, 198
- Streptokinase, 437
- Streptolysin O, 133
- Streptomycin, 191, **197**
- Stress incontinence, 599
- Stress-related disorders, 564
- Striated muscle, 220
- Striatum, **500**, 514
- String sign (x-ray), 382
- Stroke, 512
 ADP receptor inhibitors for, 437
 atrial fibrillation and, 295
 central post-stroke pain syndrome, 515
 direct factor Xa inhibitors for, 437
 eclampsia, 643
 effects of, 514–515
 homocystinuria, 84
 hypertension, 300
 hypertensive emergency and, 300
 sickle cell anemia, 422
 syphilis, 147
 thrombolytics for, 437
- Stroke volume, 284
 equation for, 285
- Strongyloides* spp, 158, 159
- Structural quality measurement, 273
- Struvite stones
 with *Proteus* spp, 127
- ST segment, 293
- ST-segment elevation MI (STEMI)
 diagnosis of, 304
- Studies
 error types, 256
 Studying for USMLE Step 1 exam
 timeline for, 16–19
- Sturge-Weber syndrome, 525
- Stylohyoid ligament, 620
- Stylohyoid muscle, 620
- Styloid process, 620
- Stylopharyngeus, 620
- Subacute combined degeneration (SCD), 69, 530
- Subacute endocarditis
 enterococci, 137
Staphylococcus gallolyticus, 137
- Subacute granulomatous thyroiditis, 341
- Subarachnoid hemorrhage, **513**, 518
 aneurysms, 516
 nimodipine for, 318
- Subarachnoid space, 507
- Subclavian arteries, 619
- Subcutaneous fat
 erythema nodosum in, 482
 skin layers, 473
- Subcutis, 473
- Subdural hematomas, 513
- Subendocardium, 210
- Sublimation, 555
- Sublingual gland
 stones in, 376
- Submandibular gland
 stones in, 376
- Submucosa, 362
- Submucosal polyps, 387
- Subscapularis muscle, 446
- Substance abuse
 adult T-cell lymphoma and, 430
Candida albicans, 153
 delirium with, 558
 dissociative identity disorder and, 558
 parental consent, 265
 torsades de pointes in, 294
 tricuspid valve endocarditis and, 311
- Substance P, 551
- Substance P antagonist, 401
- Substance use disorder, **568**
 addiction, stages of change in
 overcoming, 568
- Substantia nigra pars compacta (SNc), 500
- Subthalamic nucleus, 500
 lesions in, 511
- Subunit vaccines, 111
- Succimer
 heavy metal toxicity, 248
 lead poisoning, 419
- Succinate dehydrogenase, 67, **78**
- Succinylcholine, **551**, 590
- Succinyl-CoA
 gluconeogenesis, 78
 TCA cycle, 77
- Sucking reflex, 510
- Sucralfate, 399
- Sudden death
 cardiac death, 304, 313
 cocaine use, 571
 cor pulmonale, 679
 with myocarditis, 313
 sleep apnea, 679
- Sudden infant death syndrome (SIDS), 635
- Suicidal patients, 268
 confidentiality exceptions and, 267
 elderly, 270
- Suicide
 deaths from, 272
 physician-assisted, 268
 risk factors for, 562
- Sulbactam, 189
- Sulfadiazine, 194
Toxoplasma gondii, 156
- Sulfa drugs, **252**
 megaloblastic, 250
 rash, 250
- Sulfamethoxazole (SMX), 194
- Sulfapyridine, 400
- Sulfasalazine, 252, **400**, 466
- Sulfatides, 140
- Sulfisoxazole, 194
- Sulfonamides
 cytochrome P-450 and, 252
 hemolysis in G6PD deficiency, 250
 hypothyroidism, 249
 mechanism and use, **194**
Nocardia spp, 139
 photosensitivity, 250
 pregnancy contraindication, 204
 trimethoprim, 194
 vitamin B₉ deficiency, 68
- Sulfonyleureas, 353
 disulfiram-like reaction, 251
 insulin and, 334
- Sulfur granules, 128, **139**
- Sumatriptan, 547
 cluster headaches, 518
 coronary vasospasm with, 248
- Sunburn, 482
- Sunburst pattern (X-ray), 465
- Superficial inguinal nodes, 624
- Superior colliculi, 504
- Superior gluteal nerve, 453
- Superior mesenteric artery (SMA), 364
 syndrome, 363
- Superior oblique muscle, 540
- Superior olive (nucleus), 498
- Superior ophthalmic vein, 503
- Superior rectal vein, 365
- Superior rectus muscle, 540
- Superior sagittal sinus, 503
- Superior sulcus tumor, 685
- Superior vena cava
 embryologic development of, 281
 in fetal circulation, 282
- Superior vena cava syndrome, 98, **685**
 lung cancer, 684
 Pancoast tumor, 685
- Superoxide dismutase, 109
 free radical elimination by, 210
- Supination
 Erb palsy, 448
 forearm, 447
- Supportive therapy, 572
- Suppression, 555
- Suprachiasmatic nucleus (SCN), 497–498
 sleep physiology and, 497
- Supracondylar fracture, 447
- Supraoptic nucleus, 498
- Suprascapular nerve, 446
- Supraspinatus muscle, **446**, 448
- Supraventricular tachycardia
 adenosine for diagnosing, 324
 β-blockers for, 245, 323
 calcium channel blockers for, 324
- Suramin, 200
- Surface F protein, 169
- Surfactant (pulmonary), 661
- Surgical neck of humerus, 455
- Surgical procedures
 readmissions with, 272
- Surrogate decision-maker, **266**
- Suvorexant, **547**
- Swallowing
 tongue movement in, 493
- Swan-Ganz catheter, 297
- Swarming, 181
- Sweat glands, 236
 embryologic derivation, 613
 pilocarpine effects, 240
- Swiss cheese model, **273**
- Sydenham chorea, 312, **519**
- Sylvian fissure, 501
- Sympathetic nervous system
 denervation of face, 540
 male sexual response, 627
 receptor targets, 236
 venous return and, 286
- Sympatholytic drugs, **243**
- Sympathomimetics
 direct, **242**
 indirect, **242**
- Syncope
 during exercise, 308
 pulsus parvus et tardus, 291
- Synctiotrophoblasts, 617, 633
 β-hCG and, 226
 hCG secretion by, 633
- Syndrome of apparent
 mineralocorticoid excess, 586
 markers in, 591
- Syndrome of inappropriate
 antidiuretic hormone
 secretion (SIADH), **338**
- Synergism
 Aspirin and, 235
 of drugs, 235
- Syphilis, **147**
 fluorescent antibody stain for, 125
 STI, 184
 syphilitic heart disease, **312**
 tabes dorsalis, 530
 testing for, 148
 thoracic aortic aneurysms and, 302
 TORCH infection, 182
- Syringomyelia, **492**
 spinal cord lesions, 530
- Syrinx, 492
- Systemic amyloidosis, 212
- Systemic juvenile idiopathic arthritis, **468**
- Systemic lupus erythematosus (SLE), **470**
- Systemic mycoses, **151**
- Systemic primary carnitine
 deficiency, 89
- Systemic sclerosis, 473
- Systemic senile amyloidosis, 212
- Systolic ejection, 287
- Systolic murmur, 308
- T**
- T3 (liothyronine), 354
- T4 (levothyroxine), 354
- Tabes dorsalis, **147**, 184
 spinal cord lesions, 530
- Tachyarrhythmia
 isoproterenol for evaluating, 242
 thyroid storm, 342
- Tachycardia
 β-blockers, 245
 drug-induced, 318
 MDMA, 571

- Tachycardia (*continued*)
 metronidazole, 195
 with myocarditis, 313
 PCP, 571
 phenoxybenzamine, 244
 reflex, 244
 stimulants and, 570
 supraventricular, 245
 thyroid hormones, 354
 Wolff-Parkinson-White syndrome, 294
- Tachyphylaxis, 235
- Tacrolimus
 hyperglycemia, 249
 immunosuppression, 120
- Tactile hallucinations, 559
 cocaine, 571
- Taenia solium*, 160, 161
- Takayasu arteritis, 314
- Tamm-Horsfall mucoprotein, 594
- Tamoxifen, **443**, 656
 hot flashes with, 249
- Tamsulosin, 244, **658**
- Tanner stages (sexual development), **637**
- Tardive dyskinesia
 antipsychotic drugs and, 573
 metoclopramide adverse effect, 400
 nigrostriatal pathway, 499
- Target cells, 415
 postsplenectomy, 98
- Tarsal tunnel syndrome, 453
- Taste
 cranial nerve lesions and, 532
 loss with stroke, 514
 thalamic relay for, 498
- TATA box, 41
- Taxanes, 441
- Tay-Sachs disease
 lysosomal storage disease, 88
- Tazobactam, 189
- TCA cycle, **77**
 hyperammonemia, 82
 metabolic site, 72
 pyruvate metabolism, 77
 rate-determining enzyme for, 73
- TCA toxicity
 treatment of, 233
- T cells, 127, **409**
 activation, **103**
 adaptive immunity, 99
 anergy, 110
 cell surface proteins, 110
 corticosteroid effects, 120
 cytokine production, 101, **108**
 cytotoxic, 102
 delayed (type IV) hypersensitivity, 101
 differentiation and maturation, 98, **102**
 disorders of, 116, 117
 functions, 101
 leftunomide effects, 486
 lymph nodes, 96
 macrophage interaction, 102
 major functions of, 101
 neoplasms, 430
 polysaccharide antigens and, 127
 regulatory, 102
 sirolimus effect, 120
 spleen, 98
 thymus, 98
 untreated HIV, 176
- Tea-colored urine, 425
- “Teardrop” RBCs, **414**, 433
- Tearing stimulation, 240
- Teenagers
 common causes of death, 272
- Teeth
 congenital syphilis, 147
 dentinogenesis imperfecta, 51
 discoloration, 192, 204, 250, 614
 Gardner syndrome, 387
 osteogenesis imperfecta, 51
 Sjögren syndrome and, 468
- Telangiectasias
 basal cell carcinomas, 484
 hereditary hemorrhagic, 316
- Telencephalon, 490
- Tellurite agar, 126
- Telomerase
 action of, 38
- Telophase, 46
- Temazepam, 546
- Temperature receptors, 494
- Temperature regulation, 498
- Temperature sensation
 cape-like distribution loss, **492**, 530
 loss with strokes, 514
- Temporal arteritis, 314
- Temporalis muscle, 507
- Temporal lobe, **501**, 514
- Temporal lobe encephalitis, 164
- Tendinopathy (rotator cuff), 446
- Tendinous xanthomas, 301
 familial hypercholesterolemia, 94
- Tendonitis
 drug reaction and, 250
 fluoroquinolones, 195
- Tendons
 collagen in, 50
- Tenecteplase (TNK-tPA), 437
- Teniposide, 442
- Tennis elbow, 459
- “Tennis rackets” (Birbeck) granules, 434
- Tenofovir, 203
- Tenosynovitis, 468
- Tension headaches, 518
- Tension pneumothorax, 680, **682**
- Tensor fascia latae muscle, 453
- Tensor tympani muscle, 620
- Tensor veli palatini muscle, 620
- Teratogens
 ACE inhibitors, 610
 aminoglycosides, 191
 angiotensin II receptor blockers, 610
 in fetal development, 612
 fetal effects of, **614**
 griseofulvin, **200**, 204
 leftunomide, 486
 lithium as, 574
 methimazole as, 354
 propylthiouracil in pregnancy, 354
 ribavirin, 204
 vitamin A, 66
 warfarin as, 436
- Teratoma
 immature, 647
 mature cystic, 647
 testicular, 653
- Terazosin, 244
- Terbinafine, **199**
- Terbutaline, 242
- Teres minor, 446
- Teriparatide, 462, **487**
- Terminal bronchioles, 660
- Terminal complement deficiencies (C5–C9), 107
- Terminal deoxynucleotidyl transferase (TdT), 104
- Tertiary adrenal insufficiency, 349
- Tertiary disease prevention, 270
- Tertiary hyperparathyroidism, 345
- Tesamorelin, 328
- Testes
 descent of, 624
 lymphatic drainage of, 624
 progesterone production, 630
- Testicular atrophy
 alcoholism, 571
 muscular dystrophy, 61
- Testicular cancer, 439, 442, **653**
- Testicular torsion, **651**
- Testicular tumors
 germ cell, **652–653**
 non-germ cell tumors, 653
- Testing agencies, 24
- Testis-determining factor, 622
- Testosterone, 636, **658**
 androgen insensitivity syndrome, 639
 Leydig cell secretion, 628
 Sertoli cells, 628
 SHBG effect on, 337
 signaling pathways for, 337
 spermatogenesis, 628
- Testosterone-secreting tumors, 639
- Testosterone synthesis, 199
- Test-taking strategy, 22–23
- Tetanospasmin, 132
 blocks release of GABA, 138
- Tetanus
 exotoxins, 131
- Tetanus toxin, 110
- Tetany
 hypocalcemia, 591
 hypoparathyroidism, 344
- Tetrabenazine
 Huntington disease, 549
 Tourette syndrome, 572
- Tetracaine, 550
- Tetracyclines, **192**
 esophagitis, 249
 Fanconi syndrome, 251
 photosensitivity, 250
 protein synthesis inhibition, 191
 pseudotumor cerebri and, 521
 teratogenicity, 204, **614**
 tooth discoloration, 250
- Tetrahydrobiopterin (bh4)
 in phenylketonuria, 84
- Tetrahydrofolic acid (THF), **68**, 194
- Tetralogy of Fallot, 298
 22q11 syndromes, 300
 fetal alcohol syndrome, 300
 outflow tract formation, 281
- Tetrodotoxin, 247
- TGF- β
 in wound healing, 216
 neural development, 490
 regulatory T cells, 102
- Thalamus
 development of, 490
 limbic system and, **498**
 neuropathic pain, 515
- Thalassemia, 418
 target cells in, 415
- Thalidomide
 teratogenicity, 614
- Thayer-Martin agar, 126
- Theca-lutein cysts, 642, **646**
- Thecoma, 647
- Thenar muscles, 447, 448, 450
- Theophylline, 687
 therapeutic index of, 234
- Therapeutic antibodies, 121, **122**
- Therapeutic index (TI), **234**
- Therapeutic window, 234
- Thermogenin, 78
- Theta rhythm (EEG), 497
- Thiamine, 66
- Thiazide diuretics, **609**
 in gout, 250
 in heart failure, 309
 in hypertension, 316
- Thiazides, 609
- Thionamides, **354**
- Thiopental, 546, 550
- Thioridazine, 573
- Third-degree (complete) AV block, 295
- 3rd pharyngeal pouch, 621
- 3rd pharyngeal arch, 620
- Thirst
 hypothalamus and, 498
- 30S inhibitors, 191
- Thoracic aortic aneurysm, 300, **302**
- Thoracic outlet syndrome, 448, 684
- Threadworms, 159
- Threonine, 81
- Threonine kinase, 224
- Thrombi
 atherosclerosis, 302
 mural, 305, **307**
 post-MI, 305
- Thrombin, 436
- Thromboangiitis obliterans, 314
- Thrombocytes (platelets), **407**
 disorders, 426–427
 function tests of, 426
 heparin adverse effects, 436
 leukemias, 432
 liver markers, 390
 mixed coagulation disorders, 428
 thrombolytics and, 437
 transfusion of, 421, **429**
 in wound healing, 216
- Thrombocythemia (essential), 433
- Thrombocytopenia, 407
 Class IA antiarrhythmics, 322
 cytarabine, 440
 drug reaction and, 250
Escherichia coli, 145
 ganciclovir, 202
 glycoprotein IIb/IIIa inhibitors, 438
 heparin adverse effects, 436
 oxazolidinones, 193
 protease inhibitors, 203
 recombinant cytokines, 121
 sulfa drug allergies, 252
 TORCH infections, 182
 transfusion for, 429
 Wiskott-Aldrich syndrome, 117
- Thrombocytosis
 postsplenectomy, 98
- Thromboembolic event
 atrial fibrillation, 295
- Thrombogenesis, **411**
- Thrombolytic drugs, 413, **437**
- Thrombophlebitis
 pancreatic cancer, 398
- Thrombopoietin, 121
 signaling pathways, 337
- Thrombosis
 celecoxib, 486
 contraceptive and hormone replacement, 250
 essential thrombocythemia, 433
 homocystinuria, 84

- Thrombotic stroke, 512
 Thrush, 117
 Candida albicans, 153
 hairy leukoplakia vs, 479
 HIV-positive adults, 177
 nystatin, 199
 “Thumbprint” sign (imaging) colonic ischemia, 386
 “Thumb sign” (X-ray) flu, 142
 Thymic aplasia, 116
 chromosome association, 64
 lymphopenia with, 424
 Thymic cortex
 T cell selection in, 102
 Thymic hyperplasia
 myasthenia gravis association, 472
 Thymic shadow, 117
 Thymidine, 194
 Thymidine kinase, 201
 Thymidylate, 36
 Thymoma
 disease associations with, 98
 myasthenia gravis and, 228, 472
 paraneoplastic syndromes, 228
 Thymus
 benign neoplasm, **98**
 fetal development, 326
 pharyngeal pouch derivation, 621
 T cell differentiation, 102
 T cell origination in, 409
 Thymus-dependent antigens, 105
 Thymus-independent antigens, 105
 Thyroglossal duct cyst, 326
 Thyroid adenomas, 342
 Thyroid cancer, 343
 amyloidosis in, 212
 carcinogens in, 225
 metastases to, 223
 Psammoma bodies in, 227
 Thyroid cartilage, 620
 Thyroid development, 326
 pharyngeal pouch derivation, 621
 Thyroidectomy, 343
 Thyroid hormones, **331**
 signaling pathways for, 337
 in toxic multinodular goiter, 342
 Thyroiditis, 341
 Thyroidization of kidney, 600
 Thyroid peroxidase, 331
 Thyroid-regulating hormone (TRH)
 signaling pathways for, 337
 Thyroid-stimulating hormone (TSH)
 Graves disease and, 342
 secretion of, 327
 signaling pathways of, 337
 Thyroid-stimulating immunoglobulin (TSI), 331
 Thyroid storm, 342
 Thyrotoxicosis, 331
 Thyrotropin-releasing hormone (TRH), **328**, 330
 Thyroxine, 339
 Thyroxine-binding globulin (TBC), 331
 TIBC (total iron-binding capacity)
 anemia of chronic disease, 421
 lab values in anemia, 419
 microcytic anemia, 418
 Tibialis anterior, 453
 Tibial nerve, 452–453
 neurovascular pairing, 455
 Ticagrelor, 437
 Ticarcillin
 characteristics of, 188
 Pseudomonas aeruginosa, 143
 Ticks (disease vectors), 149–150
 Ticlopidine, 411, **437**
 Tics (Tourette syndrome), 557
 Tidal volume (TV), 664
 Tigecycline, **192**
 Tight junctions, **474**, 496
 Timolol, 245, 323, **552**
 Tinea, **152**, 200
 Tinea capitis, 152
 Tinea corporis, 152
 Tinea cruris, 152
 Tinea pedis, 152
 Tinea unguium, 152
 Tinea versicolor, 152
 Tinel sign, 459
 Tinnitus
 streptomycin, 197
 Tiotropium, 241, **687**
 Tirofiban, 411, **438**
 Tissue factor activation, 133
 Tissue plasminogen activator (tPA)
 for ischemic stroke, 512
 Tizanidine, 243, **551**
 TMP-SMX, 194
 for *Pneumocystis jirovecii*, 154
 prophylaxis, 198
 TNF- α , 108, 133, 208
 TNF- α inhibitors, 487
 TNF (tumor necrosis factor), 227
 Tobramycin, 191
 Tocolytics, 657
 Toddler development, 635
 Togaviruses
 characteristics of, 167
 genomes of, 163
 rubella as, 169
 Toll-like receptors (TLRs), 99
 Tolterodine, 241
 Tolvaptan, 354
 Tongue
 development, 493
 as ectopic thyroid tissue, 326
 glossoptosis, 620
 pharyngeal arch derivation, 620
 Tonic-clonic seizures, 517
 drug therapy for, 544
 Tonic seizures, 517
 Tonsils
 agammaglobulinemia, 116
 immune system organ, 96
 pharyngeal pouch derivation, 621
 Tooth abnormalities
 opalescent teeth, 51
 Tophus formation, 467
 Topiramate
 epilepsy, 544
 migraine headaches, 518
 pseudotumor cerebri, 521
 Topoisomerases, 195
 Topotecan, 442
 TORCH infections, 169, **182**
 cataracts, 535
 neonatal manifestations, 182
 Torsades de pointes, 294
 adenosine for, 324
 Class IA antiarrhythmics, 322
 drug reaction and, 248
 hypomagnesemia, 591
 ibutilide, 323
 sotalol, 323
 Torsemide, 608
 Torticollis, 519
 Torus (buckle) fracture, 462
 Total anomalous pulmonary venous return (TAPVR), 298
 Total lung capacity (TLC), 664
 Total parenteral nutrition (TPN), 396
 Total peripheral resistance (TPR), 286
 Tourette syndrome, 557
 drug therapy for, 572
 obsessive-compulsive disorder and, 563
 sympatholytic drugs for, 243
 Toxic dose, 234
 Toxicities and side effects of drugs, 120, 248–253
 Toxic megacolon
 Clostridium difficile, 138
 Toxic multinodular goiter, 342
 Toxic shock-like syndrome, 136
 Toxic shock syndrome, 133
 exotoxin A, 133
 presentation, 135
 Staphylococcus aureus, 135
 toxin, 133
 Toxins
 myocarditis with, 313
 seafood (ingested), 247
 Toxins (bacterial)
 anthrax, 137
 endotoxins, 132
 enterotoxins, 135
 erythrogenic, 136
 exfoliative, **133**, 135
 exotoxins, 132–133
 features of, 131
 lysogenic phage encoding, 130
 toxin-mediated disease, 135
Toxocara canis, 159
Toxocara spp, 158
 Toxoid, 111
Toxoplasma gondii, 156
 HIV-positive adults, 177
 TORCH infection, 182
Toxoplasma spp, 180
 Toxoplasmosis
 primary central nervous system lymphoma vs, 430
 prophylaxis, 194, **198**
 pyrimethamine, 200
 TP53 gene, 224
 Trabecula
 spleen, 98
 Trachea
 bifurcation of, 663
 fetal development, 326
 respiratory tree, 662
 Tracheal deviation, 680, **682**
 Tracheoesophageal fistula (TEF)/anomalies, 359
 Traction apophysitis, 461
 Tractus solitarius, 506
 Tramadol, 552
 seizures, 251
 “Tram-track” appearance, 596
 Transcortical aphasia, 516
 Transcription factor, 224
 Transduction (bacterial genetics), 130
 Transference, 554
 Transferrin, 213
 free radical elimination by, 210
 indirect measure of, 419
 lab values in anemia, 419
 Transformation (bacterial genetics), 130
 Transformation zone (cervix)
 dysplasia, 645
 histology of, 626
 Transfusion reaction, 114
 in Lyme disease, 146
 Transient arthritis
 in Lyme disease, 146
 Transient ischemic attack (TIA), 512
 Transitional cell carcinomas, 225
 Transition metals and free radical injuries, 210
 Transjugular intrahepatic portosystemic shunt (TIPS), 365
 Transketolase
 metabolic pathways, 74
 vitamin B₁ and, 66
 Translocation
 Down syndrome, 63
 fluorescence in situ hybridization, 55
 in protein synthesis, 45
 Robertsonian, 64
 Transmural inflammation fistulas, 382
 Transpeptidases, 187
 Transplants
 immunosuppressants in, 120
 rejection, 101, **119**
 t(8;14), 430, **434**
 t(9;22) (Philadelphia), 434
 t(14;18), 430, **434**
 t(15;17), 434
 Transposition (bacterial genetics), 131
 Transposition of great vessels, 298
 embryologic development, 281
 maternal diabetes and, 300
 Transposon
 in bacterial genetics, 131
 Transsexualism, 567
 Transversalis fascia, 369
 Transverse sinus, 503
 Transversion mutation, 39
 Transversus abdominis, 369, 452
 Transvestism, 567
 Tr antigens, 228
 Tranylcypromine, 575
 Trapezium bone, 449
 Trapezoid bone, 449
 TRAP (tartrate-resistant acid phosphatase), 227, 342
 Trastuzumab, 122, 444–443
 Trauma
 pneumothorax, 682
 psychiatric disorders due to, 564
 Traumatic aortic rupture, 303
 Traumatic pneumothorax, 682
 Travelers’ diarrhea, 145
 Trazodone, 576, 651
 Treacher Collins syndrome, 620
 “tree bark” appearance (aorta), 312
 Trematodes, 160
 Tremor, 519
 immunosuppressants, 120
 Trench fever, 161
 Trendelenburg sign, 453
 Treponema, 146
 Gram stain, 125
Treponema pallidum
 penicillin G/V for, 187
 STI, 184
 Triamterene, 609
 Triazolam, 546
 Triceps reflex, 510
 Triceps surae, 453
Trichinella spiralis, **159**, 161
 Trichinosis, 159
Trichomonas spp
 vaginits, 181
Trichomonas vaginalis, **158**, 181, 184
 Trichomoniasis, 184
Trichophyton spp, 152

- Trichotillomania, 563
 Trichuris, 158
 Tricuspid atresia, 281, **298**
 Tricuspid regurgitation, 287
 Ebstein anomaly and, 298
 heart murmurs with, 291
 Tricuspid valve endocarditis, 311
 Tricyclic antidepressants (TCAs)
 antimuscarinic reaction, 251
 mechanism and clinical use, 575
 naming convention for, 253
 torsades de pointes, 248
 toxicity of, 569
 toxicity treatment, 248
 as weak bases, 233
 Trientine, 395
 Trifluoperazine, 573
 Trigeminal nerve (CN V), 506
 lesion of, 532
 neuralgia, 518
 pharyngeal arch derivation, 620
 thalamic relay for, 498
 tongue, 493
 Triglycerides
 hypertriglyceridemia, 94
 insulin and, 334
 Von Gierke disease, 87
 Trihexyphenidyl, 241
 acute dystonia treatment, 241
 Trimethoprim, 187, **194**
 folate deficiency with, 420
 mechanism and use, 194
 pyrimidine synthesis and, 36
 teratogenicity, 614
 Trimming (protein synthesis), 45
 Trinucleotide repeat expansion
 diseases, **62**
 Friedreich ataxia, 62
 Huntington disease, 62
 myotonic dystrophy, 62
 Triose kinase, 80
 Triple-blinded studies, 256
 Triptans, 547
 angina and, 304
 for migraine headaches, 518
 Triquetrum bone, 449
 Trismus (lockjaw)
 tetanospasmin, 138
 Trisomies, autosomal, 63
 Trisomy 13 (Patau syndrome), **63**, 64
 disease associations with, 491
 hCG in, 633
 Trisomy 18 (Edwards syndrome),
 63, 64
 disease associations with, 535
 hCG in, 633
 Trisomy 21 (Down syndrome), 62,
 63, 64
 disease associations with, 226, 299–
 300, 384, 395, 432, 535
 tRNA
 structure, 44
 Trochlea, 540
 Trochlear nerve (CN IV), 506
 ocular motility, 540
 palsy of, 541
Tropheryma whippelli, 125, **381**
 Tropical sprue, 381
 Tropicamide, 241
 Troponins, 304, 306, 456
 Trousseau sign, **344**, 591
 Trousseau syndrome
 pancreatic cancer, 398
 paraneoplastic syndrome, 228
 True-negative rate, 257
 True-positive rate, 257
 Truncal ataxia, 499
 Truncus arteriosus
 22q11 syndromes, 300
Trypanosoma brucei, **156**, 200
Trypanosoma cruzi, 158
 achalasia and, 376
 nifurtimox for, 200
 Trypsin, 373
 Trypsinogen, 373
 Trypsinase, 408
 Tryptophan, 81
 TSC1/TSC2 genes, 224
 Tsetse flies (disease vectors), 156
 TSST-1 superantigen, 135
t-tests, 264
 T-tubule membrane, 456
 Tuberculosis, 140
 Addison disease, 349
 corticosteroids and, 336
 erythema nodosum, 482
 isoniazid, 197
 necrosis and, 209
 silicosis, 677
 Tuberin protein, 224, **525**
 Tuberoinfundibular pathway, 499
 Tuberosus sclerosis
 chromosome abnormalities, 525
 tumor suppressor genes and, 224
 Tubular necrosis, 594, **602**
 Tubulointerstitial inflammation
 WBC casts in, 594
 Tubulointerstitial nephritis, 601
 Tularemia, 149
 Tumor grade vs stage, 220
 Tumor lysis syndrome, 434–435
 Tumor markers (serum), **226**
 acute lymphoblastic leukemia,
 432
 colorectal cancer, 388
 pancreatic adenocarcinomas, 398
 Tumors
 benign vs malignant, 220
 grade vs stage, 220
 immunohistochemical stains for,
 227
 nomenclature of, 220
 Tumor suppressor genes, 46, **224**
 Tunica albuginea, 651
 Tunica vaginalis, 624
 Turcot syndrome, 387
 Turner syndrome, 638
 cardiac defect association, 300
 coarctation of aorta and, 299
 T wave (ECG), 293
 21-hydroxylase, 335
 22q11 deletion syndromes, 116, 300
 Twin concordance studies, 256
 Twinning, 616
 Thromboxane A₂ (TXA), 411, 486
 aspirin effects, 486
 thrombogenesis, 411
 Tympanic membrane, 533
 Type I errors (hypothesis testing), 263
 Type I hypersensitivity, 105, **112**, 408
 Type II errors in hypothesis testing,
 263
 Type II hypersensitivity, **112**
 organ transplants, 119
 Type II hypersensitivity reactions
 rheumatic fever, 312
 Type III hypersensitivity reactions,
 113
 organ transplants, 119
 Type IV hypersensitivity
 DRESS syndrome, 250
 graft-versus-host disease, 119
 Type IV hypersensitivity reactions
 contact dermatitis, 477
 Typhoid fever, 144
 Typhus, 150
 transmission of, **149**, 161
 Tyramine, 244
 Tyramine-induced hypertensive crisis
 procarbazine, 441
 Tyrosinase, 476
 Tyrosine
 in phenylketonuria, 84
 Tyrosine kinase
 endocrine hormone messenger, 337
 insulin and, 334
 as oncogene product, 224
 Tzanck test, 166
U
 Ubiquitination, 45
 Ubiquitin-proteasome system, 48
 UDP-glucuronosyltransferase
 physiologic neonatal jaundice, 393
 Ulcerative colitis, 382
 Ulcers (gastrointestinal)
 bismuth/sucralfate for, 399
 spondyloarthritis, 469
 sulfasalazine for, 400
 Ulcers (skin)
 Raynaud syndrome, 472
 Ulipristal, 657
 Ulnar claw, 447, 451
 Ulnar nerve, **447**, 459
 Ulnar nerve injury, 449
 Ultrasonography
 fetal cardiac activity on, 612
 kidney disease/disorder diagnoses,
 579
 renal cysts on, 604
 Umbilical artery, 282, **618**
 Umbilical cord, 618
 Umbilical hernia
 congenital, 358
 Umbilical vein, 618
 blood in, 282
 postnatal derivative of, 282
 Umbilicus
 portosystemic anastomosis, 365
 UMP synthase, 420
 Unambiguous genetic code, 37
 Uncal herniation, 529
 Uncinate process, 360
 Unconjugated bilirubin, 375
 Unconjugated (indirect)
 hyperbilirubinemia, 393
 Uncoupling agents, 78
 Undifferentiated thyroid carcinomas,
 343
 Undulant fever, 149
 “Unhappy triad” (knee injuries), 460
 Unilateral renal agenesis, 579
 Uniparental disomy, 57
 Universal electron acceptors, 75
 Universal genetic code, 37
 Unnecessary procedure requests,
 268–269
 Unstable angina, 304
 Unvaccinated children, 186
 Upper extremity nerves, 447
 Upper motor neuron lesion
 facial nerve, 532
 Urachal cysts, 618
 Urachus, 282, **618**
 Urea cycle, 82
 metabolic site, 72
 ornithine transcarbamylase
 deficiency and, 83
 rate-determining enzyme for, 73
Ureaplasma spp, 125, 127
 Urease, 181
 Urease-positive organisms, 127
 Uremia
 acute pericarditis, 313
 renal failure, 603
 Ureter, 625
 bifid, 579
 constrictions in, 583
 course of, **581**
 embryology, 578
 gynecological exam damage to, 581
 obstruction of, 579, **599**
 Ureteric bud, 579
 Ureteropelvic junction
 development of, 578
 Urethra
 BPH, 654
 injury to, 627
 posterior valves in, 579
 Urethritis
 chlamydia, 148, **184**
 Chlamydia trachomatis, 149
 gonorrhea, 184
 reactive arthritis, 469
 Urge incontinence, 599
 drug therapy for, 241
 Uric acid
 Lesch-Nyhan syndrome, 37
 Von Gierke disease, 87
 Uric acid (kidney stones), 598
 Urinalysis
 in renal disease, 601
 reducing sugar, 80
 Urinary incontinence
 drug therapy for, 241
 ephedrine for, 242
 hydrocephalus, 522
 mechanisms and associations of,
 599
 urgency incontinence, 237
 Urinary retention, 237
 atropine, 241
 bethanechol for, 240
 delirium, 558
 neostigmine for, 240
 post-void residual, 599
 treatment of, 237
 Urinary tract infections (UTIs), 181
 antimicrobial prophylaxis for, 198
 BPH, 654
 cystitis, 600
 duplex collecting system and, 579
 enterococci, 137
 epididymitis and orchitis with, 654
 Klebsiella, 145
 pyelonephritis, 600
 Staphylococcus saprophyticus, 136
 sulfa drugs for, 252
 sulfonamides for, 194
 TMP-SMX for, 194
 Urinary tract obstruction, 599
 Urine
 Bence Jones proteinuria, 431
 casts in, 594

- diuretic effects on, 609
leaks with urethral injury, 627
osmolality in acute injury, 601
pregnancy test, 633
renal tubular acidosis, 593
- Urine acidification, 233
- Urine alkalization, 233
- Urine pH and drug elimination, 233
- Urobilinogen
extravascular hemolysis, 421
intravascular hemolysis, 421
- Urogenital fold, 624
- Urosepsis, 600
- Urothelial carcinoma (bladder), 606
- Urticaria, 475, **477**
ethosuximide, 544
scombroid poisoning, 247
serum sickness, 113
sulfa drug allergies, 252
as type I hypersensitivity, 112
- USMLE Step 1 exam
check-in process, 7
clinical vignette strategies, 23
content areas covered in, 3
goal-setting for, 12
leaving exam early, 8
overview of, 2
passing rates for, 10
practice exams for, 11, **21–22**
registering for, 5–6
rescheduling, 6
score notifications for, 7
scoring of, 8–9
testing agencies, 24
testing locations, 6
test-taking strategies, 22–23
time budgeting during, 7–8
types of questions on, 8
- Ustekinumab, 122
- Uterine conditions
non-neoplastic, 648
- Uterine (Müllerian duct) anomalies, 622–623
- Uterine neoplasms, 648
- Uterovaginal agenesis, 639
- Uterus
anomalies of, 623
collagen in, 50
epithelial histology, 626
genital embryology, 622
zygote implantation, 633
- Uveitis, 536
inflammatory bowel disease, 382
in sarcoidosis, 676
seronegative spondyloarthritis, 469
- U wave in ECG, 293
- V**
- Vaccines
B-cell disorders, 116
Bordetella pertussis, 143
capsular polysaccharide and protein conjugates in, 127
diphtheria, 139
encapsulated bacteria, 127
Haemophilus influenzae, **142**, 180
Poliovirus, 167
PPSV23, 105
rabies, 171
rotavirus, 168
Salmonella typhi, 144
splenectomy and, 98
thymus-independent antigens, 105
toxoids as, 131
types of, 111
- Vagal nuclei, 506
- Vagina
anatomy of, 626
drainage of, 624
epithelial histology of, 626
genital embryology, 622
- Vaginal bleeding
cervical cancer, 645
- Vaginal candidiasis
nystatin, 199
- Vaginal infections, **181**
- Vaginal tumors, **644**
- Vaginismus, 567
- Vaginitis
Trichomonas spp, **158**, 181
trichomoniasis, 184
- Vagus nerve (CN X), 506
baroreceptors/chemoreceptors and, 296
cardiac glycoside effects, 321
Curling ulcers and, 379
diaphragm innervation, 663
gastrointestinal regulation substances and, 371
lesions of, 532
pharyngeal arch derivation, 620
structures innervated, 373
tongue, 493
- Valacyclovir, **201**
- Validity, 259
- Valine
classification of, 81
maple syrup urine disease, 84
- Valproic acid
cytochrome P-450, 252
epilepsy, 544
hepatic necrosis, 249
migraine headaches, 518
pancreatitis, 249
- Valsartan, 610
- Valvular disease
pressure-volume loops, 288
- Valvular dysfunction, 310
- Vancomycin, 190
Clostridium difficile, 138
cutaneous flushing, 248
meningitis, 180
MRSA, 198
toxicity of, 251
- Vanillylmandelic acid (VMA)
in neuroblastomas, 350
- Vanishing bile duct syndrome, 119
- Varenicline, 576
- Variable expressivity, 56
- Variance, 262
- Variant angina, 304
- Variceal bleeding, 245
- Varicella zoster virus (VZV), **165**, 475, 479
guanosine analogs, 201
immunodeficient patients, 118
meningitis, 180
rash, 183
vaccine, 110
- Varices
Budd-Chiari syndrome, 392
- Varicocele, 651
- Varicocele (scrotal), 628, **651**
- Vasa previa, 641
- Vasa vasorum
syphilis, 147
- Vascular dementia, 521
- Vascular function curves, 286
- Vascular tumors of skin, 478
- Vasculitides, **314–315**
- Vasculitis
focal necrotizing, 315
immunoglobulin A, 315
intraparenchymal hemorrhage, 513
large-vessel, 314
leukoclastic, 173
medium-vessel, 314
methotrexate for, 440
small-vessel, 314
- Vasculopathy
noninflammatory, 473
- Vas deferens, 626
- Vasoactive intestinal polypeptide (VIP), 371
- Vasoconstriction, 589
- Vasoconstrictors, 550
- Vasodilation
sympathetic receptors, 238
- Vasodilators
afterload effects, 284
aortic dissections, 303
atrial natriuretic peptide as, 296
coronary steal syndrome, 304
nitrates as, 318
- Vasogenic edema, 496
- Vasopressin, 329
receptors, 238
- Vasopressors, 286
- V(D)J recombination, 99
- VDRL false positives, 148
- Vector-borne illnesses, 150
- Veganism and B₁₂ deficiency, 420
- Vegetative state
axonal injury and, 515
- VEGF (vascular endothelial growth factor), 216
- Velocardiofacial syndrome, 116
- Vemurafenib, **444**, 484
- Venlafaxine, 575
clinical use, 572
panic disorder, 563
phobias, 563
PTSD, 564
- Venodilators, 284
- Venous gonadal drainage, 624
- Venous return, 286
- Venous sinus thrombosis, 503
- Venous thrombosis
heparin for, 436
paroxysmal nocturnal hemoglobinuria, 422
- Ventilation, 664
high altitude, 670
perfusion and, 669
- Ventilation/perfusion (V/Q) defects, 664
- Ventilation/perfusion (V/Q) ratio
exercise response, 670
mismatch, 669
- Ventral lateral (VL) nucleus, 498
- Ventral pancreatic bud, 360
- Ventral posterolateral (VPL) nucleus, 498
- Ventral posteromedial (VPM) nucleus, 498
- Ventral tegmentum, 495
- Ventricles
contractility of, 285
embryology, 281
morphogenesis of, 281
- Ventricular action potential, 292
- Ventricular aneurysm
pseudoaneurysm, 307
true, 305, **307**
- Ventricular fibrillation
ECG tracing, 295
torsades de pointes, 294
- Ventricular filling
early diastole, 287
ECG and, 293
- Ventricular free wall rupture, 307
- Ventricular myocytes, 296
- Ventricular noncompliance, 287
- Ventricular septal defect (VSD), 299
congenital rubella, 300
cri-du-chat syndrome, 64
Down syndrome, 300
fetal alcohol syndrome, 300
heart murmurs, 291
outflow tract formation, 281
- Ventricular system, 504
- Ventriculomegaly, 520, **522**
- Ventromedial nucleus (hypothalamus), 498
- Verapamil, 308, **318**, 319, 321, 518
- Verrucae, 477
- Vertebral compression fractures, 462
- Vertebral landmarks
diaphragm, 663
- Vertigo, 534
posterior circulation stroke, 514
streptomycin, 197
- Vesicles (skin), 475
varicella zoster virus, 479
- Vesicourachal diverticulum, 618
- Vesicoureteral reflux, 579
hydronephrosis, 599
pyelonephritis, 600
- Vesicular monoamine transporter (VMAT), 549
- Vesicular tinea pedis, 152
- Vesicular trafficking proteins, 47
- Vestibular schwannomas, 527
- Vestibulocochlear nerve (CN VIII), 506
- VHL gene, 224
- Vibrio cholerae*, 146
exotoxin production, 132
watery diarrhea, 179
- Vibrio parahaemolyticus*, 178
- Vibrio vulnificus*, 178
- Vigabatrin, 544
- Vilazodone, 576
- Vimentin, 48, **227**
- Vinblastine, 441
microtubules and, 48
- Vinca alkaloids, 438
- Vincristine, 441
microtubules and, 48
toxicities of, 444
- Vinyl chloride
angiosarcomas, 392, **478**
as carcinogen, 225
- VIPomas
MEN 1 syndrome, 351
octreotide for, 400
regulatory substances, 371
- Viral envelopes, 163
- Virchow nodes, 379
- Viridans streptococci, 136
 α -hemolysis, 135
bacterial endocarditis, 311
biofilm production, 128
brain abscesses, 180
gram-positive algorithm, 134
normal flora, 178
- Virilization, 335
- Virology, 162–177

- Virulence factors
bacterial, 129
Bordetella pertussis, 143
Escherichia coli, 145
Salmonella/Shigella, 144
Staphylococcus aureus, 135
Streptococcus pneumoniae, 136
β-hemolytic bacteria, 135
- Viruses
diarrhea with, 179
fluorescent antibody stain, 125
genetics, 162
immunocompromised patients, 179
in immunodeficiency, 118
as cause of myocarditis, 313
negative-stranded, 168
pneumonia, 179
receptors for, 166
segmented, 168
skin, 479
structure of, 162
- Visceral leishmaniasis, 158
- Viscosity (blood), 286
- Visual cortex, **501**, 515
- Visual disturbance
drug-related, 251
- Visual field defects, 542
saccular aneurysms and, 516
with stroke, 514, 515
- Visual hallucinations, 559
- Vital capacity (VC), 664
- Vitamin A (retinol), **66**
free radical elimination by, 210
idiopathic intracranial
hypertension, 251, **521**
measles morbidity and mortality, 170
teratogenicity, 614
- Vitamin B₁ (thiamine), **66**
brain lesions and, 511
deficiency of, 66
functions of, 74
pyruvate dehydrogenase complex,
76
solubility of, 65
- Vitamin B₂ (riboflavin), **67**
pyruvate dehydrogenase complex,
76
solubility, 65
- Vitamin B₃ (niacin), **67**
pyruvate dehydrogenase complex,
76
solubility, 65
vitamin B₆ and, 67
- Vitamin B₅ (pantothenic acid), **67**
pyruvate dehydrogenase complex
and, 76
solubility of, 65
- Vitamin B₆ (pyridoxine), **67**
deficiency, 67
isoniazid, 197
sideroblastic anemia, 419
- Vitamin B₇ (biotin), **68**
activated carriers, 75
functions of, 73
pyruvate metabolism, 77, 78
solubility of, 65
- Vitamin B₉ (folate), **68**
deficiency, 406, 420
functions, 68
solubility, 65
- Vitamin B₁₂ (cobalamin), **69**
absorption of, 374
deficiency, 160, 161
solubility, 65
spinal cord lesions, **530**
- Vitamin C (ascorbic acid)
free radical elimination by, 210
functions, 69
methemoglobin treatment, 248
solubility of, 65
- Vitamin D
excess, 70
functions, 70
hyperparathyroidism, 464
hypervitaminosis lab values,
464
osteomalacia/trickets, 463, 464
osteoporosis prophylaxis, 462
signaling pathways for, 337
solubility of, 65
- vitamin D (calciferol)
calcitriol production, 589
- Vitamin E
free radical elimination by, 210
function, 70
solubility of, 65
- Vitamin K
cephalosporins, 189
coagulation cascade, 413
deficiency, **413**, 426
solubility of, 65
vitamin E interaction, 71
for warfarin toxicity, 436
- Vitamin/mineral absorption, 374
- Vitamins
fat-soluble, **65**
water-soluble, **65**
- Vitelline duct/fistula, 618
- Vitiligo, 476
- Vitreous body
collagen in, 50
- VLDL (very low-density lipoprotein),
94
- Volume contraction
alkalemia from diuretics, 609
- Volume of distribution, 231
- Volumetric flow rate (Q), 286
- Volvulus, 385, **386**
Meckel diverticulum, 384
midgut, 386
Onchocerca, 158
sigmoid, 386
- Vomiting
annular pancreas, 360
biliary colic, 396
bilious, 359, 384
diabetic ketoacidosis, 347
Ebola virus, 171
fructose intolerance, 80
glycylcyclines, 192
Histoplasma capsulatum, 177
intestinal atresia, 359
Legionella spp, 185
lithium toxicity, 569
Mallory-Weiss syndrome, 377
maple syrup urine disease, 84
metoclopramide for, 400
MI and, 305
ondansetron for, 400
posttussive, 143
pyloric stenosis, 359
receptors for, 496
Salmonella spp, 149
in stroke, 514
toxic shock syndrome, 135
treatment of, 400, 401
trichinosis, 159
vitamin C toxicity, 69
- Von Gierke disease, 87
- von Hippel-Lindau disease, 525
chromosome association, 64
renal cell carcinoma and, 605
tumor suppressor genes and, 224
- von Willebrand disease, 411, **428**
- Voriconazole, 199
- Vortioxetine, 576
- VRE (vancomycin-resistant
enterococci)
daptomycin, 195
enterococci, 137
highly resistant, 198
oxazolidinones, 193
- Vulnerable child syndrome, **556**
- Vulvar carcinoma, **644**
- Vulvar lymphatic drainage, 624
- Vulvar pathology, 644
neoplastic, 644
non-neoplastic, 644
- Vulvovaginitis, 153, **181**
- vWF (von Willebrand factor)
receptor for, 407
in thrombocytes, 407
in thrombogenesis, 411
- V_{max}, 230
- W**
WAGR complex/syndrome, 606
“Waiter’s tip” (Erb palsy), 448
Waiving right to confidentiality, 267
- Waldenstrom macroglobulinemia,
431
- Walking milestone, 635
- Wallenberg syndrome, 514
- Wallerian degeneration, 495
- Wall tension, 284, 285
- Warburg effect, 221
- Warfarin, 436
adverse effects of, 428
coagulation cascade, 413
griseofulvin and, 200
heparin vs, 436, 437
PT measurement, 426
teratogenicity, 614
therapeutic index of, 234
toxicity treatment, **248**, 429
vitamin K antagonist, 71
- Warthin-Finkeldey giant cells, 170
- Warthin tumors, 376
- Waterhouse-Friderichsen syndrome,
349
meningococci, 142
- Watershed zones, **210**, 502
- Water-soluble vitamins, 65
- Waxy casts (urine), 594
- WBC casts (urine), **594**, 600
- Weak acid overdose
treatment, 233
- Weak bases overdose treatment, 233
- “Wear and tear” pigment, 211
- Wegener granulomatosis, 315
autoantibody, 115
restrictive lung disease, 675
- Weight gain
danazol, 658
duodenal ulcer, 380
mirtazapine, 576
- Weight loss
adrenal insufficiency, 349
cholelithiasis and, 396
chronic mesenteric ischemia, 386
diabetes mellitus, 346
esophageal cancer, 378
gastric ulcers, 380
glucagonoma, 351
Histoplasma capsulatum, 177
malabsorption syndromes, 381
*Mycobacterium avium-
intracellulare*, 177
orlistat for, 400
pancreatic cancer, 398
polyarthritis nodosa, 314
polymyalgia rheumatica, 470
pseudotumor cerebri treatment,
521
renal cell carcinoma, 605
sleep apnea, 679
stomach cancer, 379
for stress incontinence, 599
tuberculosis, 140
- Weil disease, 147
- Well-patient care, 270–271
- Wenckebach AV block, 295
- Werdnig-Hoffmann disease, 530
- Wernicke aphasia, 514, **516**
- Wernicke area, 501
stroke effects, 514
- Wernicke encephalopathy, 66, **571**
- Wernicke-Korsakoff syndrome, 511, **571**
vitamin B₁ deficiency, 66
- Western blot, 53
- West Nile virus, **167**, 180
- Wet beriberi, 66
- Wharton duct, 376
- Wharton jelly, 618
- Wheal
urticaria, 477
- Wheals, 475
- Wheezing
lung cancer, 684
- Whipple disease, 381
periodic acid-Schiff stain for, 125
- Whipple procedure
for pancreatic cancer, 398
- Whipple triad
insulinomas and, 351
- Whispered pectoriloquy, 680
- White matter
axonal injury, 515
demyelinating disorders, 524
glial cells in, 494
multiple sclerosis, 523
- White pulp (spleen), 98
- Whooping cough
Bordetella pertussis, 143
pertussis toxin, 132
- Wickham striae, 482
- Wide splitting, 289
- Williams syndrome, 64
cardiac defect association, 300
- Wilms tumor, 606
dactinomycin for, 439
neuroblastomas vs, 350
tumor suppressor genes and, 224
- Wilson disease, 395
ATP7B protein in, 51
autosomal recessive inheritance, 60
chromosome association, 64
Fanconi syndrome, 586
free radical injury and, 210
- Winged scapula, 448
- Winters formula, 592

- “Wire looping” of capillaries, 596
 “Wire lupus,” 596
 Wiskott-Aldrich syndrome, 117
 X-linked recessive disorder, 61
 Withdrawal (psychoactive drugs), 570
 Wobble, 37
 Wolff-Chaikoff effect, 341
 Wolffian duct, 622
 Wolff-Parkinson-White syndrome, 294
 Woolsorter’s disease, 137
 “word salad,” 559
 Wound healing
 keratinocytes, 216
 phases of, 216
 scar/keloid formation, 218
 Woven bone, 458
 Wright-Giemsa stain, 407
 Wright stain
 Borrelia spp, 146
 Wrinkles of aging, 52
 Wrist
 injuries of, 459
 bones, 449
 Wrist drop, 447
 lead poisoning, 419
 Written advance directives, 266
 WT1/WT2 genes
 in renal disease, 224, **606**
 oncogenicity of, 224
Wuchereria bancrofti, 158, 159
- X**
 Xanthine oxidase inhibitors, 467
 Xanthogranulomatous pyelonephritis, 600
 Xanthomas
 familial dyslipidemias, 94
 hyperlipidemia and, 301
 Xeroderma pigmentosum
 DNA repair defects in, 40
 Xerosis cutis, 66
 Xerostomia, 240, 243, **468**
 X-inactivation (lyonization)
 Barr body formation, 61
 X-linked agammaglobulinemia, 116
 X-linked dominant inheritance, 59
 X-linked recessive disorders, **61**
 agammaglobulinemia, 116
 hyper-IgM syndrome, 117
 Menkes disease, 51
 NADPH oxidase defect, 117
 Wiskott-Aldrich syndrome, 117
 X-linked recessive inheritance, 59
 X-ray/imaging findings
 Apple core lesion, 388
 bamboo spine, 469
 Bird’s beak sign, 376
 Bone-in-bone, 463
 Codman triangle, 465
 Coffee bean sign, 386
 Coin lesion, 684
 Crew cut (skull x-ray), 422
 kidney stones, 598
 pencil-in-cup, 469
 punched out bone lesions, 431
 Steeple sign (x-ray), 170
 String sign, 382
 Sunburst pattern, 465
 Thumbprint sign (imaging), 386
 Thumb sign, 142
 X-ray teratogenicity, 614
- Y**
 Yellow fever, 167, **168**
 liver anatomy and, 367
Yersinia enterocolitica, 179
 transmission and treatment, 144
Yersinia pestis
 animal transmission, 149
 facultative intracellular organisms, 127
 Yo antigens, 228
 Yolk sac tumor
 ovarian, 647
 testicular, 653
Yersinia spp
 reactive arthritis, 469
- Z**
 Zafirlukast, 687
 Zaleplon, 546
 Zanamivir, 201
 Zellweger syndrome, 47
 Zenker diverticulum, 384
 Zero-order elimination, 232
 Zidovudine, 203
 Ziehl-Neelsen stain, 125
 Zika virus, 171
 Zileuton, 687
 Zinc, 71
 Wilson disease, 395
 Ziprasidone, 573
 Zoledronic acid, 486
 Zollinger-Ellison syndrome, 352
 duodenal ulcer, 380
 gastrin in, 371
 MEN 1 syndrome, 351
 proton pump inhibitors for, 399
 Zolpidem, 546
 Zona fasciculata, **327**, 336
 Zona glomerulosa, 327
 Zona reticularis, 327
 Zoonotic bacteria, 149
 Zymogens, 373

About the Editors



Tao Le, MD, MHS

Tao developed a passion for medical education as a medical student. He currently edits more than 15 titles in the *First Aid* series. In addition, he is Founder and Chief Education Officer of USMLE-Rx for exam preparation and ScholarRx for undergraduate medical education. As a medical student, he was editor-in-chief of the University of California, San Francisco (UCSF) *Synapse*, a university newspaper with a weekly circulation of 9000. Tao earned his medical degree from UCSF in 1996 and completed his residency training in internal medicine at Yale University and fellowship training at Johns Hopkins University. Tao subsequently went on to cofound Medsn, a medical education technology venture, and served as its chief medical officer. He is currently chief of adult allergy and immunology at the University of Louisville.



Vikas Bhushan, MD

Vikas is a writer, editor, entrepreneur, and teleradiologist on extended sabbatical. In 1990 he conceived and authored the original *First Aid for the USMLE Step 1*. His entrepreneurial endeavors included a student-focused medical publisher (S2S), an e-learning company (medschool.com), and an ER teleradiology practice (24/7 Radiology). Trained on the Left Coast, Vikas completed a bachelor's degree at the University of California Berkeley; an MD with thesis at UCSF; and a diagnostic radiology residency at UCLA. His eclectic interests include technology, cryptoeconomics, information design, South Asian diasporic culture, and avoiding a day job. Always finding the long shortcut, Vikas is an adventurer, knowledge seeker, and occasional innovator. He enjoys intermediate status as a kiteboarder and father, and strives to raise his three children as global citizens.




Matthew Sochat, MD

Matthew is a third-year hematology/oncology fellow at St. Louis University in St. Louis, Missouri. He completed his internal medicine residency training at Temple University Hospital in Philadelphia. He completed medical school in 2013 at Brown University and is a 2008 graduate of the University of Massachusetts, Amherst, where he studied biochemistry and the classics. Pastimes include skiing, cooking/baking, traveling, the company of friends/loved ones (especially his wonderful wife), the Spanish language, and computer/video gaming. Be warned: Matt also loves to come up with corny jokes at (in)opportune moments.



Vaishnavi Vaidyanathan, MD

Vaishnavi is a second-year child neurology resident at Phoenix Children's Hospital in Phoenix, Arizona. She is a graduate of the University of Missouri-Kansas City School of Medicine, where she earned her bachelor's and medical degrees. Her interests include medical education and health advocacy. Outside of medicine, she loves to dance, learn new languages, and watch Bollywood movies.



Sarah Schimansky, MB BCh BAO

Sarah is a third-year ophthalmology resident in the UK. She grew up in Germany before moving to Dublin, Ireland, to study medicine at the Royal College of Surgeons in Ireland. She has a keen interest in medical education and is currently enrolled in a Masters in Surgical Education program at Imperial College London. An avid traveler, Sarah is always on the lookout for new destinations to explore and new countries to call home. When she is not on the road, she enjoys yoga, long walks, and red wine in the company of friends and family.



Jordan Abrams

Jordan is a fourth-year medical student at St. George's University School of Medicine who hopes to pursue residency training in anesthesiology. He graduated magna cum laude from the University of Delaware, earning a bachelor's degree in neuroscience with minors in medical humanities and biological sciences. Combining his creative mindset and passion for drawing, Jordan founded theHYMedicine.com, an educational website that offers free medical study guides, tutoring, and study schedules for students worldwide. Aside from medicine, Jordan enjoys traveling, reading, and playing soccer.



Kimberly Kallianos, MD

Originally from Atlanta, Kimberly graduated from the University of North Carolina at Chapel Hill in 2006 and from Harvard Medical School in 2011. She completed her radiology residency and fellowship at UCSF and is currently an Assistant Professor of Clinical Radiology at UCSF in the Cardiac and Pulmonary Imaging section.

SECTION IV

Top-Rated Review Resources

“Some books are to be tasted, others to be swallowed, and some few to be chewed and digested.”

—Sir Francis Bacon

“Always read something that will make you look good if you die in the middle of it.”

—P.J. O'Rourke

“So many books, so little time.”

—Frank Zappa

“If one cannot enjoy reading a book over and over again, there is no use in reading it at all.”

—Oscar Wilde

▶ How to Use the Database	2
▶ Question Banks and Books	4
▶ Web and Mobile Apps	6
▶ Comprehensive	10
▶ Anatomy, Embryology, and Neuroscience	12
▶ Behavioral Science	14
▶ Biochemistry	15
▶ Cell Biology and Histology	16
▶ Microbiology and Immunology	17
▶ Pathology	19
▶ Pharmacology	21
▶ Physiology	22

▶ HOW TO USE THE DATABASE

This section is a database of top-rated basic science review books, sample examination books, websites, apps, and commercial review courses that have been marketed to medical students studying for the USMLE Step 1. At the end of the section is a list of publishers and independent bookstores with addresses and phone numbers. For each recommended resource, we list (where applicable) the **Title**, the **First Author** (or editor), the **Series Name** (where applicable), the **Current Publisher**, the **Copyright Year**, the **Number of Pages**, the **ISBN**, the **Approximate List Price**, the **Format** of the resource, and the **Number of Test Questions**. We also include **Summary Comments** that describe their style and overall utility for studying. Finally, each recommended resource receives a **Rating**. Within each section, resources are arranged first by Rating and then alphabetically by the first author within each Rating group.

A letter rating scale with six different grades reflects the detailed student evaluations for **Rated Resources**. Each rated resource receives a rating as follows:

A+	Excellent for boards review.
A A-	Very good for boards review; choose among the group.
B+ B	Good, but use only after exhausting better resources.
B-	Fair, but there are many better resources in the discipline; or low-yield subject material.

The Rating is meant to reflect the overall usefulness of the resource in helping medical students prepare for the USMLE Step 1. This is based on a number of factors, including:

- The cost
- The readability of the text
- The appropriateness and accuracy of the material
- The quality and number of sample questions
- The quality of written answers to sample questions
- The quality and appropriateness of the images and illustrations
- The quality of the user interface and learning experience, for web and mobile apps
- The length of the text (longer is not necessarily better)
- The quality and number of other resources available in the same discipline
- The importance of the discipline for the USMLE Step 1

Please note that ratings do not reflect the quality of the resources for purposes other than reviewing for the USMLE Step 1. Many books with

lower ratings are well written and informative but are not ideal for boards preparation. We have not listed or commented on general textbooks available in the basic sciences.

Evaluations are based on the cumulative results of formal and informal surveys of thousands of medical students at many medical schools across the country. The summary comments and overall ratings represent a consensus opinion, but there may have been a broad range of opinion or limited student feedback on any particular resource.

Please note that the data listed are subject to change in that:

- Publisher and app store prices change frequently.
- Retail and online bookstores may set their own prices.
- New editions and app versions come out frequently, and the quality of updating varies.
- The same book may be reissued through another publisher.

We actively encourage medical students and faculty to submit their opinions and ratings of these basic science review materials so that we may update our database. In addition, we ask that publishers and authors submit for evaluation review copies of basic science review books, including new editions and books not included in our database. We also solicit reviews of new books, mobile apps, websites, flash cards, and commercial review courses.

Disclaimer/Conflict of Interest Statement

None of the ratings reflects the opinion or influence of the publisher. All errors and omissions will gladly be corrected if brought to the attention of the authors through our blog at www.firstaidteam.com. Please note that USMLE-Rx and the entire *First Aid for the USMLE* series are publications by certain authors of *First Aid for the USMLE Step 1*; the following ratings are based solely on recommendations from the student authors of *First Aid for the USMLE Step 1* as well as data from the student survey and feedback forms.

▶ QUESTION BANKS AND BOOKS

A+	<p>UWorld Qbank UWORLD www.uworld.com</p> <p>Questions demand multistep reasoning and are often more difficult than those on the actual Step 1 exam. Offers detailed explanations with figures and tables. Features a number of test customization and analysis options. Users can see cumulative results both over time and compared to other test takers. In addition to a desktop version, it can be accessed through iOS or Android mobile apps.</p>	\$249–\$749 Test/2400 q
A	<p>NBME Practice Exams NATIONAL BOARD OF MEDICAL EXAMINERS www.nbme.org/students/sas/Comprehensive.html</p> <p>The official practice exams published by the NBME are comprised of retired Step 1 questions. NBME research found that they show a “moderate correlation” with actual Step 1 performance. The exams will show you which questions you answered incorrectly, but they will not show any explanations. You will also not be able to review correctly answered questions. Students generally use these as rough gauges of their score progression over their study time. Note that you can sign up to for an in-person practice exam for an additional \$75 to be taken at Prometric, for students who want to practice the logistics of exam day.</p>	\$60 Test/200 q
A-	<p>AMBOSS AMBOSS www.amboss.com</p> <p>Integrated question bank for Step 1 and Step 2 CK exams with an additional interactive online library of medical resources. Contains numerous illustrations within the clinical vignettes. Allows for the selection of questions by difficulty level. Includes personalized study plan. Free trial available, accessible through iOS or Android mobile apps.</p>	\$9–\$365 Test/3500 q
A-	<p>USMLE-Rx Qmax USMLE-Rx www.usmle-rx.com</p> <p>Offers Step 1–style questions accompanied by thorough explanations. Omits obscure material and distills high yield information. Each explanation includes references from <i>First Aid</i>. However, the proportion of questions covering a given subject area does not always reflect the actual exam’s relative emphasis. Question stems occasionally rely on “buzzwords.” Most useful to help memorize <i>First Aid</i> facts. Provides detailed performance analyses. Free trial available, accessible through iOS or Android mobile apps.</p>	\$89–\$339 Test/2300 q



<p>B+</p>	<p><i>Kaplan Qbank</i> KAPLAN www.kaptest.com</p> <p>Covers most content found on Step 1, but sometimes emphasizes recall of low-yield details rather than integrative problem-solving skills. Test content and performance feedback can be organized by both organ system and discipline. Includes detailed explanations of all answer choices. Users can see cumulative results both over time and compared to other test takers. Accessible through iOS or Android mobile apps.</p>	<p>\$99–\$349 Test/2100 q</p>
<p>B</p>	<p><i>BoardVitals</i> www.boardvitals.com</p> <p>Comprehensive question bank modeled closely after the format of the Step 1. Covers all subject areas and includes explanations for each answer choice. Users can create custom exams and compare their performance to national averages. Contains fewer image-based questions compared to similar platforms.</p>	<p>\$59–\$179 Test/1750 q</p>
<p>B</p>	<p><i>Kaplan USMLE Step 1 Qbook</i> KAPLAN Kaplan, 2017, 468 pages, ISBN 9781506223544</p> <p>Consists of over 850 exam-like questions organized by the traditional basic science disciplines. Similar to the Kaplan Qbank, and offers USMLE-style questions with clear, detailed explanations; however, lacks classic images typically seen on the exam. Also includes access to a sample online question bank and a guide on test-taking strategies.</p>	<p>\$50 Test/850 q</p>
<p>B</p>	<p><i>Pastest</i> www.pastest.com</p> <p>Questions appear to be simpler than board-style questions, with many first- and second-order questions. Explanations are accompanied by references to <i>First Aid</i> and short video clips to reinforce information. Accessible through iOS or Android mobile apps.</p>	<p>\$79–\$249 Test/2100 q</p>
<p>B</p>	<p><i>TrueLearn Review</i> www.truelearn.com</p> <p>Includes over 2200 USMLE-style practice questions with topics mapped to the NBME blueprint. Uses national benchmarking to show students where they stand in comparison to peers.</p>	<p>\$159–\$399 Test/2200 q</p>

▶ WEB AND MOBILE APPS

A	<p>Anki www.ankisrs.net</p> <p>Flash card-making resource designed for retention of facts through spaced repetition. Free access via desktop and smartphone for Windows, Mac, and Android. The iOS app must be purchased for \$25. Available in different languages.</p>	Free	Flash cards
A	<p>Boards and Beyond www.boardsbeyond.com</p> <p>Includes over 400 videos averaging ~26 minutes each, covering the breadth of Step 1 material. Membership includes access to the companion books as PDFs. A collection of videos is offered as free samples on the website. Also includes over 1300 practice questions.</p>	\$19–\$249	Review
A	<p>Physeio www.physeio.com</p> <p>Online review containing 32 hours of review videos covering physiology. Accessible via website or mobile app. Includes a supplemental full-color PDF textbook. Videos are concise and focus on high-yield material, and board-style practice questions are included after each topic to help solidify understanding. Similar structure to Pathoma, but with physiology focus.</p>	\$30–\$150	Review
A	<p>SketchyMedical www.sketchymedical.com</p> <p>Video library of narrated lectures with thorough explanations that present microbiology, pharmacology, and pathology in a memorable style. Access to the entire gram-positive cocci section is free at signup. Additional content can be purchased on a subscription basis.</p>	\$99–\$369	Review
A-	<p>Cram Fighter www.cramfighter.com</p> <p>Helps organize a study schedule. Highly flexible with customizable settings. Supports more than 650 of the most popular books, video lectures, question banks, and flash cards. Mobile apps available for iOS and Android.</p>	\$29–\$159	Study plan
A-	<p>First Aid Step 1 Express www.usmle-rx.com</p> <p>More than 80 hours of high-yield videos explaining material from <i>First Aid for the USMLE Step 1</i>. Videos include more than 600 extra images and multimedia clips. Step-by-step analysis of USMLE-style questions with each video. Subscription includes a color workbook with over 200 pages.</p>	\$69–\$299	Review/Test
B+	<p>First Aid Step 1 Flash Facts www.usmle-rx.com</p> <p>Access to 12,000+ flash cards with intelligent spaced repetition integrated with <i>First Aid for the USMLE Step 1</i>, of which 3500+ are case based. Updated each year to reflect the newest edition of the book; students can access the past 3 editions' worth of flash cards. Searchable by organ system, discipline, and topic.</p>	\$29–\$149	Flash cards

B+	<p>Medbullets www.medbullets.com</p> <p>Free online learning and collaboration community for students preparing for their exams. Supplements medical school coursework and Step 1 studying with simplified, to-the-point online search platform that is best used as a reference. Recently added premium content for \$80-\$250 includes an online question bank and adaptive learning system.</p>	Free	Review/ Test/1000 q
B+	<p>Medical School Pathology www.medicalschoolpathology.com</p> <p>Offers lectures and slides based on the Robbins <i>Pathology</i> textbook. Lectures can be downloaded.</p>	Free	Review
B+	<p>OnlineMedEd www.onlinemeded.org</p> <p>A video lecture series covering primarily clinical science material, with recent addition of biochemistry, cell biology, and immunology topics. Video access is free with registration. A subscription of \$10-\$70/month gains access to ad-free videos, lecture notes, flash cards, question bank, and downloadable audio lectures.</p>	Free	Review
B+	<p>Osmosis www.osmosis.org</p> <p>Web platform that includes exam study scheduling tool, 27,000+ variable quality multiple choice questions, flash cards with spaced repetition, and 3000+ curated concept cards with videos, memory anchors, and reference articles. Includes a curriculum analysis and search engine, collaboration features for study groups, and a mobile app with quizzes and videos.</p>	\$179–\$279	Test
B+	<p>USMLE Step 1 Mastery builtbyhlt.com/medical/usmle-step-1-mastery</p> <p>Question bank accessible through website or via free mobile app. Covers all USMLE topics and includes vignettes, images, and mnemonics. Question formatting is generally less representative of actual USMLE questions compared with other widely used question banks. Mobile app contains supplemental flash cards for integrated learning.</p>	\$2–\$10	Test/1400 q
B+	<p>WebPath: The Internet Pathology Laboratory webpath.med.utah.edu</p> <p>Features more than 2700 gross and microscopic images, clinical vignette questions, and case studies. Includes nine general pathology exams and 11 system-based pathology exams with approximately 1300 questions. Also features 170 questions associated with images. Questions are useful for reviewing boards content but are typically untimed, easier, and shorter. No multimedia practice questions. Not regularly updated with regard to high-yield Step 1 material.</p>	Free	Review/ Test/1300 q
B	<p>Blue Histology www.lab.anhb.uwa.edu.au/mb140</p> <p>Provides access to 400+ histologic images with thorough explanations. Images searchable by topic, stain, keyword. Website also contains multiple choice practice questions.</p>	Free	Review/Test

B	<p>Digital Anatomist Project: Interactive Atlases UNIVERSITY OF WASHINGTON da.si.washington.edu/da.html</p> <p>Contains an interactive neuroanatomy course along with a three-dimensional atlas of the brain, thorax, and knee. Atlases have computer-generated images and cadaver sections. Each atlas also has a quiz in which users identify structures in the slide images. However, questions do not focus on high-yield anatomy for Step 1.</p>	Free	Review
B	<p>Dr. Najeeb Lectures www.drnajeeblectures.com</p> <p>Hundreds of hours of video lectures covering basic medical sciences and clinical medicine with thousands of hand-drawn illustrations and mnemonics. Website provides mobile video support on smartphones and tablets. Free lectures accessible at www.drnajeeblectures.com/free-medical-videos.html.</p>	\$99	Review
B	<p>Firecracker FIRECRACKER INC. firecracker.lww.com</p> <p>Learning platform divided into modules. The Step 1 module is divided into organ systems and includes review of preclinical lecture material, periodic quizzes on flagged reviewed material, and USMLE-style questions in interface simulating the actual exam. Contains page references to <i>First Aid for the USMLE Step 1</i> and high-yield diagrams from various textbooks. Users can grade how well they remember the quiz answers (1–5), which allows the program to customize future quizzes. Features detailed performance analysis and a calendar for personalized study plans. Accessible on all smartphones and tablets. Comprehensive; best if started early in preclinical years.</p>	\$39–\$660	Review/ Test/2800 q
B	<p>KISSPrep www.kissprep.com</p> <p>Online lecture videos focused on select subjects from the Step 1 exam. Focuses on harder-to-learn content and teaches it in a simple, easy-to-understand manner. Quizzes and other interactive tools are available to help with knowledge retention. Not all Step 1 content is covered on this platform.</p>	\$99–\$135	Review
B	<p>Lecturio www.lecturio.com</p> <p>Online platform for comprehensive exam preparation, including over 250 hours of lectures, a flash card deck, quizzes, and a question bank. Organized by subject matter and allows users to customize their learning experience. Some content may be beyond the scope of the exam and better suited for medical school coursework. Lectures and quizzes may be accessed for free. iOS and Android apps are available.</p>	\$50–\$300	Review/ Test/2150 q
B	<p>Memorang MEMORANG INC. www.memorangapp.com</p> <p>Platform utilizing spaced repetition, available both in website and app form. Utilizes custom and/or premade flash card “study sets” derived from 15,000 flash cards that focus on specific subject areas and are then tested via various games and quizzing methods. Free 7-day trial, or monthly/annual membership.</p>	\$19–\$239	Flash cards



B	<p><i>Picmonic</i> www.picmonic.com</p> <p>Helpful resource for visual learners. Unique images and stories with daily quizzes and spaced repetition. Contains 1400 images and includes study guides, webinars, and infographics that help cover 15,000+ facts of Step 1 material. Offered via both web and mobile platforms.</p>	<p>\$25–\$480 Review</p>
B–	<p><i>Radiopaedia.org</i> www.radiopaedia.org</p> <p>A user-friendly website with thousands of well-organized radiology cases and articles. Encyclopedia entries contain high-yield bullet points of anatomy and pathology. Images contain detailed descriptions but no arrows to demarcate findings. Quiz mode allows students to make a diagnosis based on radiographic findings. Content may be too broad for boards review but is a good complement to classes and clerkships.</p>	<p>Free Cases/Test</p>
B–	<p><i>The Pathology Guy</i> FRIEDLANDER www.pathguy.com</p> <p>Contains extensive but poorly organized information on a variety of fundamental concepts in pathology. A high-yield summary intended for USMLE review can be found at www.pathguy.com/meltdown.txt, but the information given is limited by a lack of images and frequent digressions.</p>	<p>Free Review</p>

▶ COMPREHENSIVE

A	<p><i>First Aid for the Basic Sciences: General Principles</i> \$55 Review</p> <p>LE McGraw-Hill, 2019, 816 pages, ISBN 9781260143676</p> <p>Comprehensive review of the basic sciences covered in year 1 of medical school. Similar to the first part of <i>First Aid</i>, organized by discipline, and includes hundreds of color images and tables. Best if started with first-year coursework and then used as a reference during boards preparation.</p>
A	<p><i>First Aid Cases for the USMLE Step 1</i> \$50 Cases</p> <p>LE McGraw-Hill, 2018, 496 pages, ISBN 9781260143133</p> <p>A recently updated series of hundreds of high-yield cases organized by organ system. Each case features a clinical vignette with relevant images, followed by questions and short, high-yield explanations. Offers coverage of many frequently tested concepts, and integrates subject matter in the discussion of the vignette. Helpful in reviewing material outlined in <i>First Aid for the USMLE Step 1</i>.</p>
A-	<p><i>First Aid for the Basic Sciences: Organ Systems</i> \$72 Review</p> <p>LE McGraw-Hill, 2017, 912 pages, ISBN 9781259587030</p> <p>A comprehensive review of the basic sciences covered in year 2 of medical school. Similar to the second part of <i>First Aid</i>, organized by organ system, and includes hundreds of color images and tables. Each organ system contains discussion of embryology and anatomy, physiology, pathology, and pharmacology. Best if started with second-year coursework and then used as a reference during boards preparation.</p>
A-	<p><i>Crush Step 1: The Ultimate USMLE Step 1 Review</i> \$45 Review</p> <p>O'CONNELL Elsevier, 2017, 704 pages, 9780323481632</p> <p>Detailed, text-heavy review book with practice questions included. Coverage of many high-yield topics but includes some outdated information. Best if used with coursework, but also recommended as a supplemental reference for boards review. Limited student feedback.</p>
A-	<p><i>Cracking the USMLE Step 1</i> \$45 Review</p> <p>PRINCETON REVIEW Princeton Review, 2013, 832 pages, ISBN 9780307945068</p> <p>Comprehensive review book with hundreds of illustrations, charts, and diagrams along with 2 full-length practice tests with detailed answer explanations available online. Limited student feedback.</p>
B+	<p><i>USMLE Step 1 Secrets in Color</i> \$43 Review</p> <p>BROWN Elsevier, 2016, 800 pages, ISBN 9780323396790</p> <p>Clarifies difficult concepts in a concise, readable manner. Uses a case-based format and integrates information well. High-quality clinical images. Complements other boards study resources, with a focus on understanding preclinical fundamentals rather than on rote memorization. Slightly lengthy for last-minute studying.</p>

B+	<p><i>Step-Up to USMLE Step 1 2015</i> JENKINS Lippincott Williams & Wilkins, 2014, 528 pages, ISBN 9781469894690</p> <p>An organ system–based review text with clinical vignettes that is useful for integrating the basic sciences covered on Step 1. Composed primarily of outlines, charts, tables, and diagrams. Limited scope of material covered. Includes access to a sample online question bank.</p>	\$50	Review
B+	<p><i>USMLE Step 1 Lecture Notes 2018</i> KAPLAN Kaplan Medical, 2018, ~2700 pages, ISBN 9781506221229</p> <p>Extremely comprehensive review of Step 1 topics through videos and lecture notes. Split into individual sections covering pathology, pharmacology, physiology, biochemistry and medical genetics, immunology and microbiology. Generally best used to fill gaps in understanding and to review unfamiliar topics that one has not come across, and therefore the notes are commonly used by foreign medical graduates. Some sections are quite detailed and go beyond the scope of the Step 1 exam.</p>	\$330	Review
B+	<p><i>USMLE Images for the Boards: A Comprehensive Image-Based Review</i> TULLY Elsevier, 2012, 296 pages, ISBN 9781455709038</p> <p>Contains more than 300 color images of content likely to be tested on the Step 1. Contains a wide variety of images including ECGs and radiographic studies. Some images may be low yield for boards studying, but still excellent as a supplement to preclinical courses.</p>	\$42	Review
B	<p><i>USMLE Step 1 Made Ridiculously Simple</i> CARL MedMaster, 2017, 416 pages, ISBN 9781935660224</p> <p>Concise, succinct text. Online access to more than 1000 practice questions. Uses a table and chart format organized by subject, but some charts are poorly labeled. Consider as an adjunct to more comprehensive sources.</p>	\$30	Review/Test 1000 q
B	<p><i>medEssentials for the USMLE Step 1</i> MANLEY Kaplan, 2012, 588 pages, ISBN 9781609780265</p> <p>A comprehensive review divided into general principles and organ systems, organized using high-yield tables and figures. Helpful for visual learners, but can be overly detailed and time consuming. Includes color images in the back along with a monthly subscription to online interactive exercises, although these are of limited value for Step 1 preparation. Comes with a free mobile version.</p>	\$55	Review

▶ ANATOMY, EMBRYOLOGY, AND NEUROSCIENCE

A-	<p><i>High-Yield Gross Anatomy</i> DUDEK Lippincott Williams & Wilkins, 2014, 320 pages, ISBN 9781451190236</p> <p>A good review of gross anatomy with some clinical correlations. Contains color clinical photos and well-labeled, high-yield radiographic images, but often goes into excessive detail that is beyond the scope of the boards.</p>	\$43	Review
A-	<p><i>Clinical Anatomy Made Ridiculously Simple</i> GOLDBERG MedMaster, 2016, 175 pages, ISBN 9780940780972</p> <p>An easy-to-read text offering simple diagrams along with numerous mnemonics, helpful charts, and amusing associations. The humorous style has variable appeal to students, so browse the content before purchasing. Offers good coverage of selected topics. Includes a CD-ROM atlas of normal radiographic anatomy. Best if used during coursework. Includes more detail than typically tested on Step 1.</p>	\$30	Review
B+	<p><i>High-Yield Embryology</i> DUDEK Lippincott Williams & Wilkins, 2013, 176 pages, ISBN 9781451176100</p> <p>A concise review of a relatively less-tested subject. Offers excellent organization with clinical correlations. Includes a high-yield list of embryologic tissue origins and USMLE-style case studies after each chapter. May not be suitable for dedicated Step 1 studying.</p>	\$56	Review
B+	<p><i>High-Yield Neuroanatomy</i> FIX Lippincott Williams & Wilkins, 2015, 208 pages, ISBN 9781451193435</p> <p>An easy-to-read, straightforward format with excellent diagrams and illustrations. Features a useful atlas of brain and spinal cord images, a glossary of important terms, and an appendix of neurologic lesions. Overall, a great resource and quick read, but more detailed than what is required for Step 1.</p>	\$40	Review/ Test/50 q
B+	<p><i>Anatomy—An Essential Textbook</i> GILROY Thieme, 2017, 528 pages, ISBN 9781626234390</p> <p>A thorough, visually appealing approach to learning anatomy. Contains over 650 colorful, helpful illustrations. Presents material in bullet-point format and tables. Includes over 160 clinical correlates and 400 USMLE-style questions, with the opportunity to complete practice questions online.</p>	\$48	Text/ Test/400 q
B+	<p><i>Netter's Anatomy Flash Cards</i> HANSEN Saunders, 2018, 688 flash cards, ISBN 9780323530507</p> <p>Netter's illustrations in a question/answer column format that allows for self-testing. Each card includes commentary on the structures with a clinical correlation. More effective as a supplement to coursework, and much too detailed for boards preparation. Lack of embryology correlates limits Step 1 usefulness. Includes online access with additional bonus cards and more than 400 multiple choice questions. Note: an iOS app has a similar cost and additional functionality.</p>	\$40	Flash cards

B+	<p><i>Crash Course: Anatomy</i> STENHOUSE Elsevier, 2015, 288 pages, ISBN 9780723438540</p> <p>Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Contains an up-to-date self-assessment section. Provides a solid review of anatomy for Step 1. Best if started early.</p>	\$45 Review
B	<p><i>BRS Embryology</i> DUDEK Lippincott Williams & Wilkins, 2014, 336 pages, ISBN 9781451190380</p> <p>An outline-based review of embryology that is typical of the BRS series. Offers a good review and includes much more detail than is required for Step 1. A discussion of congenital malformations is included at the end of each chapter, along with over 220 USMLE-style questions with answers and explanations. The comprehensive exam at the end of the book is high yield. Includes access to a searchable online text on the free companion website, which also features interactive quizzing.</p>	\$56 Review/ Test/220 q
B	<p><i>Anatomy Flash Cards: Anatomy on the Go</i> GILROY Thieme, 2013, 752 flash cards, ISBN 9781604069105</p> <p>Flash card deck containing high-quality illustrations and a question/answer format that allows for self-testing. Occasional radiographic image. Best if used with coursework; too long for efficient boards preparation.</p>	\$60 Flash cards
B	<p><i>Clinical Neuroanatomy Made Ridiculously Simple</i> GOLDBERG MedMaster, 2014, 90 pages + CD-ROM, ISBN 9781935660194</p> <p>An easy-to-read, memorable, and simplified format with clever diagrams. Offers a quick, high-yield review of clinical neuroanatomy, but not a comprehensive resource for boards review. Places appropriate emphasis on clinically relevant pathways, cranial nerves, and neurologic diseases. Includes a CD-ROM with CT and MR images, a tutorial on neurologic localization, and interactive quizzes covering classic neurology cases.</p>	\$26 Review/Test/ Few q
B	<p><i>Netter's Anatomy Coloring Book</i> HANSEN Elsevier, 2018, 392 pages, ISBN 9780323545037</p> <p>An easy-to-understand, detailed, interactive book that is an excellent companion to traditional textbooks during preclinical anatomy coursework. Provides multiple views and magnifications of anatomic structures as well as dissection layers. The coloring aspect of the book can be highly beneficial for visual learners. Contains few clinical correlations, which limits its usefulness during dedicated studying for Step 1.</p>	\$20 Review
B	<p><i>Case Files: Anatomy</i> TOY McGraw-Hill, 2014, 416 pages, ISBN 9780071794862</p> <p>Review text that includes 58 well-chosen cases with discussion, comprehension questions, and take-home pearls. Tables are helpful, but schematics are black and white and not representative of Step 1. A reasonable book to work through for those who benefit from problem-based learning.</p>	\$35 Cases

B- *Case Files: Neuroscience* **\$35** Cases
 TOY
 McGraw-Hill, 2014, 432 pages, ISBN 9780071790253

Similar to other *Case Files* books, it includes 49 clinical cases with lengthy discussion and 3–5 multiple choice questions at the end of each case. Cases are helpful, but the discussion is too lengthy. Questions are not the most representative of those seen on boards. Limited student feedback.

▶ BEHAVIORAL SCIENCE

A- *BRS Behavioral Science* **\$52** Review/
Test/700 q
 FADEM
 Lippincott Williams & Wilkins, 2016, 384 pages, ISBN 9781496310477

An easy-to-read outline-format review of behavioral science. Offers detailed coverage of mostly high-yield topics, but at a level of depth that often exceeds what is tested on Step 1. Better used prior to dedicated study period. Incorporates tables and charts as well as a short but complete statistics chapter. Features over 700 review questions, including a 100-question comprehensive exam. References DSM-V criteria.

B+ *High-Yield Biostatistics, Epidemiology, and Public Health* **\$43** Review
 GLASER
 Lippincott Williams & Wilkins, 2013, 168 pages, ISBN 9781451130171

A well-written, easy-to-read text that offers extensive coverage of epidemiology and biostatistics. Includes helpful review questions and tables, but somewhat lengthy given the low-yield nature of this subject on Step 1.

▶ BIOCHEMISTRY

A⁻	<p><i>Pixorize</i> www.pixorize.com</p> <p>Visual mnemonic system focusing primarily on biochemistry. Step-by-step videos and interactive images aid studying and review. Compare to Sketchy and Picmonic.</p>	\$100–\$130	Review
B⁺	<p><i>Medical Biochemistry—An Illustrated Review</i> PANINI Thieme, 2013, 441 pages, ISBN 9781604063165</p> <p>A comprehensive medical biochemistry study guide with an emphasis on images. Very detailed and may be better as a supplement to preclinical courses than as a review resource for the Step 1. Images and diagrams are helpful for solidifying knowledge. Online access available for additional content, including 400 USMLE-style practice questions.</p>	\$40	Review/ Test/400 q
B	<p><i>Lange Flash Cards Biochemistry and Genetics</i> BARON McGraw-Hill, 2017, 196 flash cards, ISBN 9781259837210</p> <p>Flash card deck featuring clinical vignettes on one side and concise discussions on the other. Each section contains 2–3 cards on biochemistry principles. High level of detail may make this less ideal for dedicated boards studying. Note that no carrying case for the cards is included.</p>	\$40	Flash cards
B	<p><i>Lippincott Illustrated Reviews: Biochemistry</i> FERRIER Lippincott Williams & Wilkins, 2017, 560 pages, ISBN 9781496344496</p> <p>An integrative and comprehensive review of biochemistry that includes good clinical correlations and effective color diagrams. Extremely detailed and requires significant time commitment, so it should be started with first-year coursework. High-yield summaries at the end of each chapter. Comes with access to the companion website, which includes over 200 USMLE-style questions.</p>	\$78	Review/ Test/200 q
B	<p><i>BRS Biochemistry, Molecular Biology, and Genetics</i> LIEBERMAN Lippincott Williams & Wilkins, 2013, 432 pages, ISBN 9781451175363</p> <p>A highly detailed review featuring many figures and clinical correlations highlighted in colored boxes. The biochemistry portion includes much more detail than required for Step 1, but may be useful for students without a strong biochemistry background or as a reference text. The molecular biology section is more focused and high yield. Also offers a chapter on laboratory techniques and a comprehensive, 120-question exam. Questions are clinically oriented.</p>	\$54	Review/Test
B	<p><i>Case Files: Biochemistry</i> TOY McGraw-Hill, 2014, 480 pages, ISBN 9780071794886</p> <p>Includes 51 clinical cases with comprehensive discussion and summary box, albeit too much depth and not enough breadth for boards preparation. Some cases will almost certainly <i>not</i> be tested. Questions at the end of each case are not representative of those on Step 1.</p>	\$35	Cases

B	<p><i>PreTest Biochemistry and Genetics</i> WILSON McGraw-Hill, 2017, 592 pages, ISBN 9780071791441</p> <p>500 questions with detailed, well-referenced explanations. Features a high-yield introduction and appendix, but may be overly detailed in some cases. A solid supplement to preclinical courses and board studying.</p>	\$38 Test/500 q
----------	---	------------------------

▶ CELL BIOLOGY AND HISTOLOGY

B+	<p><i>BRS Cell Biology and Histology</i> GARTNER Lippincott Williams & Wilkins, 2018, 448 pages, ISBN 9781496396358</p> <p>Covers concepts in cell biology and histology in an outline format. Can be used alone for cell biology study, but may have fewer histology images than some other resources. Includes more detail than is required for Step 1, and information is less high yield than that of other books in the BRS series. Interactive quizzes on the free companion website provide additional practice.</p>	\$54 Review/ Test/320 q
-----------	--	-----------------------------------

B+	<p><i>Crash Course: Cell Biology and Genetics</i> STUBBS Elsevier, 2015, 216 pages, ISBN 9780723438762</p> <p>Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Includes online access. High level of detail makes this resource best suited for coursework.</p>	\$47 Review/Print + online
-----------	---	--------------------------------------

B	<p><i>Wheater's Functional Histology</i> YOUNG Elsevier, 2013, 464 pages, ISBN 9780702047473</p> <p>A color atlas with more than 900 high-quality illustrations of normal histology with image captions and accompanying text. Far too detailed to use for boards studying given the low-yield nature of the material, but useful as a coursework text or boards reference. Provides online access to the entire atlas and USMLE-style self-assessment questions.</p>	\$83 Text
----------	--	------------------

▶ MICROBIOLOGY AND IMMUNOLOGY

A⁻	<p><i>Basic Immunology</i> ABBAS Elsevier, 2019, 336 pages, ISBN 9780323549431</p> <p>A useful text that offers clear explanations of complex topics in immunology. Best if used in conjunction with coursework and later skimmed for quick Step 1 review. Includes colorful diagrams, images, tables, and a glossary for further study. Features online access.</p>	\$70	Review
A⁻	<p><i>Clinical Microbiology Made Ridiculously Simple</i> GLADWIN MedMaster, 2019, 418 pages, ISBN 9781935660330</p> <p>An excellent, easy-to-read, detailed review of microbiology that includes clever and memorable mnemonics. The sections on bacterial disease are most high yield, less emphasis placed on pharmacology. Recommended to read during coursework and review the concise charts at the end of each chapter during boards review. All images are cartoons; no microscopy images that appear on boards. Requires a supplemental source for immunology.</p>	\$38	Review
A⁻	<p><i>Medical Microbiology and Immunology Flash Cards</i> ROSENTHAL Elsevier, 2016, 192 flash cards, ISBN 9780323462242</p> <p>Flash cards covering the microorganisms most commonly tested on Step 1. Each card features color microscopic images and clinical presentations on one side and relevant bug information in conjunction with a short case on the other side. Also includes Student Consult online access for extra features. Overemphasizes “trigger words” related to each bug. Not a comprehensive resource.</p>	\$40	Flash cards
B⁺	<p><i>Lippincott Illustrated Reviews: Immunology</i> DOAN Lippincott Williams & Wilkins, 2012, 384 pages, ISBN 9781451109375</p> <p>A clearly written, highly detailed review of basic concepts in immunology. Features many useful tables and review questions at the end of each chapter. More than 300 color annotated illustrations. Offers abbreviated coverage of immunodeficiencies and autoimmune disorders. Best if started with initial coursework and used as a reference during Step 1 study.</p>	\$75	Reference/Test/ Few q
B⁺	<p><i>Microcards: Microbiology Flash Cards</i> HARPAVAT Lippincott Williams & Wilkins, 2015, 312 flash cards, ISBN 9781451192353</p> <p>A well-organized and complete resource for students who like to use flash cards for review. Cards feature the clinical presentation, pathobiology, diagnosis, treatment, and high-yield facts for a particular organism. Some cards also include excellent flow charts organizing important classes of bacteria or viruses. Overall, a good review resource, but at times it is overly detailed, requiring a significant time commitment. Also useful as an aid with coursework. Includes access to online USMLE-style questions with answers.</p>	\$53	Flash cards

B+	<p><i>Review of Medical Microbiology and Immunology</i> LEVINSON McGraw-Hill, 2018, 832 pages, ISBN 9781259644498</p> <p>A clear, comprehensive text with outstanding diagrams and tables. Includes an excellent immunology section. Contains a chapter summarizing details on medically important organisms. Can be used as reference for reviewing immunology concepts. Can be detailed and dense at points, so best if started early with coursework. Includes practice questions of mixed quality and does not provide detailed explanation of answers. Compare with <i>Lippincott Illustrated Reviews: Microbiology</i>.</p>	\$63	Review/ Test/654 q
B+	<p><i>How the Immune System Works</i> SOMPAYRAC Wiley-Blackwell, 2019, 168 pages, ISBN 9781119542124</p> <p>A short overview of high-yield immunology designed for those with no prior immunology knowledge. Analogies and images create a “storybook” feel to spruce up a relatively dry subject. The 15 chapters offer a general overview with good supporting details.</p>	\$50	Review
B	<p><i>Case Studies in Immunology: Clinical Companion</i> GEHA W. W. Norton & Company, 2016, 384 pages, ISBN 9780815345121</p> <p>A text that was originally designed as a clinical companion to <i>Janeway’s Immunobiology</i>. Provides a great synopsis of the major disorders of immunity in a clinical vignette format. Integrates basic and clinical sciences. Features excellent images and illustrations from Janeway, as well as questions and discussions.</p>	\$62	Cases
B	<p><i>Pretest: Microbiology</i> KETTERING McGraw-Hill, 2013, 480 pages, ISBN 9780071791045</p> <p>Includes a short section on high-yield facts followed by 500 questions in a clinical vignette format. Questions are more difficult than encountered on the boards and some topics discussed are not likely to be tested. A good book to work through with coursework but too low yield for review purposes.</p>	\$38	Test/500 q
B	<p><i>Case Files: Microbiology</i> TOY McGraw-Hill, 2014, 416 pages, ISBN 9780071820233</p> <p>Provides 54 clinical microbiology cases followed by a clinical correlation, a discussion with boldfaced buzzwords, and questions. Cases are well chosen, but the text lacks the high-yield charts and tables found in other books in the Case Files series. Images are sparse, black and white, and of poor quality.</p>	\$36	Cases
B	<p><i>Lange Microbiology and Infectious Diseases Flash Cards, 3e</i> Somers, 2017</p> <p>Clinical vignettes presented on one side of the card as a mini-case study of the disease and the flip side presents the etiology and epidemiology, pathogenesis, clinical manifestations, laboratory diagnosis, and treatment and prevention of the disorder. Good for reviewing clinical aspects of many infectious diseases including those caused by bacteria, viruses, and fungi.</p>	\$46	Flash cards

B-	<p><i>Lippincott Illustrated Reviews: Microbiology</i> CORNELISSEN</p> <p>Lippincott Williams & Wilkins, 2019, 448 pages, ISBN 9781496395856</p> <p>A comprehensive, highly illustrated review of microbiology that is similar in style to other titles in the Illustrated Reviews series. Has more than 400 color illustrations and color-coded summaries to help visual learners. Contains several hundred USMLE-style review questions to help with exam preparation. Compare with Levinson's <i>Review of Medical Microbiology and Immunology</i>.</p>	<p>\$73 Review/Test/ Few q</p>
-----------	---	---

▶ PATHOLOGY

A+	<p><i>Pathoma: Fundamentals of Pathology</i> SATTAR</p> <p>Pathoma, 2019, 218 pages, ISBN 9780983224631</p> <p>Integrated approach to pathology review, combining a focused textbook with 35+ hours of online lectures. Book contains more than 350 color images. Videos combine “chalk talk” and slide formats to explain pathogenesis in an easy-to-understand manner. Online subscription needed for full access.</p>	<p>\$85–\$120 Review/Lecture</p>
A-	<p><i>Rapid Review: Pathology</i> GOLJAN</p> <p>Elsevier, 2018, 864 pages, ISBN 9780323476683</p> <p>A comprehensive source for key concepts in pathology, presented in a bulleted outline format with many high-yield tables and color figures. Features detailed explanations of disease mechanisms. Integrates concepts across disciplines with a strong clinical orientation. Lengthy, so best if started early with coursework. Includes access to online question bank with more than 500 questions. Covers material for both Step 1 and Step 2 exams.</p>	<p>\$65 Review/ Test/500 q</p>
A-	<p><i>Robbins and Cotran Review of Pathology</i> KLATT</p> <p>Elsevier, 2014, 504 pages, ISBN 9781455751556</p> <p>A question book that follows the main Robbins textbooks. Questions are more detailed, difficult, and arcane than those on the actual Step 1 exam, but the text offers a great review of pathology integrated with more than 1100 images. Thorough answer explanations reinforce key points. Requires significant time commitment, so best if started with coursework. 2014 edition table of contents closely follows the organization of <i>Robbins and Cotran Pathologic Basis of Disease</i>, 8th edition.</p>	<p>\$55 Test/1100 q</p>
A-	<p><i>Crash Course: Pathology</i> XIU</p> <p>Elsevier, 2019, 438 pages, ISBN 9780702073540</p> <p>Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1 review questions. Includes online access. Best if started during coursework.</p>	<p>\$40 Review</p>

B	<p><i>High-Yield Histopathology</i> DUDEK Lippincott Williams & Wilkins, 2017, 320 pages, ISBN 9781496353344</p> <p>Reviews the relationship of basic histology to the pathology, physiology, and pharmacology of clinical conditions that are tested on Step 1. Includes case studies, numerous light and electron micrographs, and pathology photographs. Given its considerable length, should be started with coursework or used as a reference to better identify images.</p>	\$36 Review
B	<p><i>Pathophysiology of Disease: Introduction to Clinical Medicine</i> HAMMER McGraw-Hill, 2018, 832 pages, ISBN 9781260026504</p> <p>An interdisciplinary text useful for understanding the pathophysiology of clinical symptoms. Effectively integrates the basic sciences with mechanisms of disease. Features great graphs, diagrams, and tables. In view of its length, most useful if started during coursework. Includes 120 case studies, checkpoint questions that appear in every chapter, and a few non-boards-style questions. The text's clinical emphasis nicely complements <i>BRS Pathology</i>.</p>	\$90 Text
B	<p><i>Haematology at a Glance</i> MEHTA Blackwell Science, 2014, 136 pages, ISBN 9781119969228</p> <p>A resource that covers common hematologic issues. Includes color illustrations. Presented in a logical sequence that is easy to read. Good for use with coursework.</p>	\$49 Review
B	<p><i>Pocket Companion to Robbins and Cotran Pathologic Basis of Disease</i> MITCHELL Elsevier, 2016, 896 pages, ISBN 9781455754168</p> <p>A condensed version of <i>Robbins and Cotran Pathologic Basis of Disease</i> that is good for reviewing keywords associated with most important diseases. Presented in a highly condensed format, but the text is complete and easy to understand. Contains no photographs or illustrations but does include tables. Useful as a quick reference.</p>	\$40 Review
B	<p><i>BRS Pathology</i> SCHNEIDER Lippincott Williams & Wilkins, 2013, 480 pages, ISBN 9781451115871</p> <p>An excellent, concise review with appropriate content emphasis. Chapters are organized by organ system and feature an outline format with boldfacing of key facts. Includes good questions with explanations at the end of each chapter plus a comprehensive exam at the end of the book. Offers well-organized tables and diagrams as well as photographs representative of classic pathology. Contains a chapter on lab testing and "key associations" with each disease. Contains excellent color images and access to an online test and interactive question bank. Most effective if started early in conjunction with coursework, as it does not discuss detailed mechanisms of disease pathology.</p>	\$54 Review/ Test/450 q

▶ PHARMACOLOGY

B+	<p><i>Crash Course: Pharmacology</i> \$40 Review</p> <p>BATTISTA Elsevier, 2019, 336 pages, ISBN 9780702073441</p> <p>Part of the Crash Course review series for basic sciences, integrating clinical topics. Offers two-color illustrations, handy study tools, and Step 1–style review questions with a self-assessment section. Includes online access. Gives a solid, easy-to-follow overview of pharmacology.</p>
B+	<p><i>Master the Boards USMLE Step 1 Pharmacology Flashcards</i> \$55 Flash cards</p> <p>FISCHER Kaplan, 2015, 200 flash cards, ISBN 9781618657947</p> <p>Easy-to-read flash cards with drug and questions on one side and discussion on the other. Useful for a quick pharmacology review. Some drugs/material may be beyond the scope of the Step 1, or more appropriate at the Step 2 level.</p>
B+	<p><i>BRS Pharmacology</i> \$55 Review/ Test/200 q</p> <p>ROSENFELD Lippincott Williams & Wilkins, 2019, 384 pages, ISBN 9781975105495</p> <p>Features two-color tables and figures that summarize essential information for quick recall. A list of drugs organized by drug family is included in each chapter. Too detailed for boards review; best used as a reference. Also offers end-of-chapter review tests with Step 1–style questions and a comprehensive exam with explanations of answers. An additional question bank is available online.</p>
B	<p><i>Lange Pharmacology Flash Cards</i> \$39 Flash cards</p> <p>BARON McGraw-Hill, 2017, 266 flash cards, ISBN 9781259837241</p> <p>A total of 230 pocket-sized flash cards of relevant drugs formatted with clinical vignettes on one side and relevant information on the other side (eg, mode of action, adverse effects, clinical uses). Particularly high-yield information is highlighted in bold. Mainly useful as a supplement for pharmacology knowledge, rather than as a primary resource. Printed on less durable material.</p>
B	<p><i>Pharmacology Flash Cards</i> \$45 Flash cards</p> <p>BRENNER Elsevier, 2017, 230 flash cards, ISBN 9780323355643</p> <p>Flash cards for more than 200 of the most commonly tested drugs. Cards include the name of the drug (both generic and brand) on the front and basic drug information on the back, with occasional cards covering high-yield pharmacology pathways. Divided and color coded by class, and comes with a compact carrying case. Lacks figures and clinical vignettes.</p>

B	<p><i>Katzung & Trevor's Pharmacology: Examination and Board Review</i> TREVOR McGraw-Hill, 2018, 592 pages, ISBN 9781259641022</p> <p>A well-organized text with concise explanations. Features good charts and tables; the crammable list in Appendix I is especially high yield for Step 1 review. Also good for reviewing drug interactions and toxicities. Offers two 100-question practice exams. Text includes many low-yield/obscure drugs. Compare with <i>Lippincott Illustrated Reviews: Pharmacology</i>, both of which are better suited to complementing coursework than last-minute studying for boards.</p>	<p>\$54 Review/ Test/800 q</p>
B	<p><i>Lippincott Illustrated Reviews: Pharmacology</i> WHALEN Lippincott Williams & Wilkins, 2018, 576 pages, ISBN 9781496384133</p> <p>A resource presented in outline format with practice questions, many excellent illustrations, and comparison tables. Effectively integrates pharmacology and pathophysiology. Best started alongside coursework, as it is highly detailed and requires significant time commitment. Focuses on basic principles.</p>	<p>\$75 Review/ Test/380 q</p>

▶ PHYSIOLOGY

A-	<p><i>BRS Physiology</i> COSTANZO Lippincott Williams & Wilkins, 2018, 304 pages, ISBN 9781496367617</p> <p>A clear, concise review of physiology that is both comprehensive and efficient, making for fast, easy reading. Includes excellent high-yield charts and tables, but lacks some figures from Costanzo's <i>Physiology</i>. Features high-quality practice questions with explanations in each chapter along with a clinically oriented final exam. An excellent reference during times of focused Step 1 studying, but best if started early in combination with coursework. Respiratory and acid-base sections are comparatively weak.</p>	<p>\$54 Review/ Test/350 q</p>
A-	<p><i>Pathophysiology of Heart Disease</i> LILLY Lippincott Williams & Williams, 2015, 480 pages, ISBN 9781451192759</p> <p>Great resource that outlines an in-depth explanation of both cardiac physiology and pathology. Best used as a supplement when learning the material for the first time, as it helps build a strong foundation. Because the book itself is rather dense, it is not recommend as a primary resource during focused boards studying period.</p>	<p>\$57 Review</p>
A-	<p><i>PreTest Physiology</i> METTING McGraw-Hill, 2013, 528 pages, ISBN 9780071791427</p> <p>Contains questions with detailed, well-written explanations. One of the best of the PreTest series. Best for use by the motivated student after extensive review of other sources. Includes a high-yield facts section with useful diagrams and tables.</p>	<p>\$38 Test/500 q</p>

A⁻	<p><i>Color Atlas of Physiology</i> SILBERNAGL Thieme, 2015, 472 pages, ISBN 9783135450070</p> <p>Contains more than 180 high-quality illustrations of disturbed physiologic processes that lead to dysfunction. An alternative to standard texts, but not high yield for boards review.</p>	\$50 Review
B⁺	<p><i>BRS Physiology Cases and Problems</i> COSTANZO Lippincott Williams & Wilkins, 2012, 368 pages, ISBN 9781451120615</p> <p>Presents 62 classic cases in vignette format with several questions per case. Includes exceptionally detailed explanation of answers along with supplemental diagrams. For students interested in an in-depth discussion of physiology concepts.</p>	\$58 Cases
B⁺	<p><i>Physiology</i> COSTANZO Saunders, 2017, 528 pages, ISBN 9780323478816</p> <p>A comprehensive, clearly written text that covers concepts outlined in <i>BRS Physiology</i> in greater detail. Offers excellent color diagrams and charts. Each systems-based chapter features a detailed summary of objectives and a Step 1–relevant clinical case. Includes access to online interactive extras. Requires time commitment, but helps develop a strong foundation in physiology concepts. Best if started alongside coursework. Practice questions at end of each chapter.</p>	\$60 Text
B⁺	<p><i>Vander's Renal Physiology</i> EATON McGraw-Hill, 2018, 224 pages, ISBN 9781260019377</p> <p>Well-written text on renal physiology, with helpful but sparse diagrams and practice questions at the end of each chapter. Too detailed for Step 1 review, however. Best if used with organ-based coursework to understand the principles of renal physiology.</p>	\$49 Text
B⁺	<p><i>Acid-Base, Fluids, and Electrolytes Made Ridiculously Simple</i> PRESTON MedMaster, 2017, 166 pages, ISBN 9781935660293</p> <p>A resource that covers major acid-base and renal physiology concepts. Provides information beyond the scope of Step 1, but remains a useful companion for studying kidney function, electrolyte disturbances, and fluid management. Includes scattered diagrams and questions at the end of each chapter. Consider using after exhausting more high-yield physiology review resources.</p>	\$24 Review
B⁺	<p><i>Pulmonary Pathophysiology: The Essentials</i> WEST Lippincott Williams & Wilkins, 2017, 264 pages, ISBN 9781496339447</p> <p>A volume offering comprehensive coverage of respiratory physiology. Clearly organized with useful charts and diagrams. Review questions at the end of each chapter provide answers but no explanations. Best used as a course supplement during the second year, less ideal for use immediately prior to Step 1.</p>	\$57 Review/ Test/75 q

B	<p><i>Rapid Review: Physiology</i> BROWN Elsevier, 2011, 384 pages, ISBN 9780323072601</p> <p>Offers a good review of physiology in a format typical of the Rapid Review series, albeit with more images. Includes online access to 350 questions with concise explanations, along with other extras. Compare with Robbins <i>Physiology</i>.</p>	\$39 Test/350 q
B	<p><i>Endocrine Physiology</i> MOLINA McGraw-Hill, 2018, 320 pages, ISBN 9781260019353</p> <p>Questions at the end of each chapter are helpful solidify knowledge, but some are not representative of Step 1 questions. Provides more detailed explanations of endocrine physiology than Costanzo review offers, but much too lengthy for Step 1 review. May be useful as a coursework adjunct.</p>	\$59 Review
B-	<p><i>Netter's Physiology Flash Cards</i> MULRONEY Saunders, 2015, 450 flash cards, ISBN 9780323359542</p> <p>Flash cards contain a high-quality illustration on one side with question and commentary on the other. Good for self-testing, but too fragmented for learning purposes and not comprehensive enough for boards.</p>	\$40 Flash cards

**Get more e-books from www.ketabton.com
Ketabton.com: The Digital Library**